

Problem set 2

1. Let R be a commutative ring and let M be an R -module. Show that for every exact sequence of R -modules $U \xrightarrow{f} V \xrightarrow{g} W \rightarrow 0$ the sequence

$$M \otimes U \xrightarrow{\text{id} \otimes f} M \otimes V \xrightarrow{\text{id} \otimes g} M \otimes W \rightarrow 0$$

is exact. *Hint:* To prove exactness at $M \otimes V$, construct a left-inverse for an appropriate map $M \otimes V / \text{im}(\text{id} \otimes f) \rightarrow M \otimes W$.

2. Let R and M be as in Problem 1 and assume additionally that M is a free R -module. Show that for every short exact sequence $0 \rightarrow U \xrightarrow{f} V \xrightarrow{g} W \rightarrow 0$ the sequence

$$0 \rightarrow M \otimes U \xrightarrow{\text{id} \otimes f} M \otimes V \xrightarrow{\text{id} \otimes g} M \otimes W \rightarrow 0$$

is exact.

3. Think about how to sensibly define an action of $\text{Ext}(-, G)$ and $\text{Tor}(-, G)$ on morphisms and prove that this makes them functors $\mathbf{Ab} \rightarrow \mathbf{Ab}$ (the first contravariant, the second covariant).
4. Prove that the sequences in the universal coefficients theorems are natural with respect to chain maps.
5. Let C_*, D_* be chain complexes of free Abelian groups and assume that $f : C_* \rightarrow D_*$ is a quasi-isomorphism, i.e. a chain map such that $f_* : H_*(C) \rightarrow H_*(D)$ is an isomorphism. Let G be an Abelian group. Prove the following statements using naturality of the sequences in the universal coefficient theorems.

(a) $f \otimes \text{id} : C_* \otimes G \rightarrow D_* \otimes G$ is a quasi-isomorphism.

(b) $f^* : \text{Hom}(D_*, G) \rightarrow \text{Hom}(C_*, G)$ is a quasi-isomorphism.

6. Show that the splitting $H^n(X; G) \cong \text{Ext}(H_{n-1}(X); G) \oplus \text{Hom}(H_n(X); G)$ whose existence is asserted by the universal coefficient theorem for cohomology *cannot* be natural in X .

Hint: Consider the map $\phi : \mathbb{R}P^2 \rightarrow S^2$ given by collapsing $\mathbb{R}P^1 \subset \mathbb{R}P^2$ to a point.

7. The Klein bottle K has $H_0(K; \mathbb{Z}) \cong \mathbb{Z}$, $H_1(K; \mathbb{Z}) \cong \mathbb{Z} \oplus \mathbb{Z}_2$ and all other homology groups vanish. Use this to compute the homology and cohomology of K with coefficients in \mathbb{Z} and \mathbb{Z}_p for p prime.
8. Let X be a topological space and let $A, B \subset X$ be subsets. Denote by $C_k(A + B) \subset C_k(X)$ the subspace of chains which are sums of simplices entirely contained in A or B . Show that the quotient $C_k(X) / C_k(A + B)$ is free.