

Interest Rate Theory Solutions Sheet 1

1. Note that

$$P(t, S)F(t; T, S) = \frac{P(t, T) - P(t, S)}{S - T},$$
$$F(T, S) = \frac{1}{S - T} \left(\frac{1}{P(T, S)} - 1 \right).$$

We build a self-financing trading strategy with time $t \leq T$ cost

$$\frac{P(t, T) - P(t, S)}{S - T},$$

yielding the value $F(T, S)$ at time S :

- At time $t \leq T$ buy one zero-coupon bond with maturity T at cost $P(t, T)$ and sell one zero-coupon bond with maturity S at cost $P(t, S)$.
Total cost at time $t \leq T$: $P(t, T) - P(t, S)$.
- At time T receive 1 dollar for the maturing zero-coupon bond.
Reinvest this 1 dollar to buy precisely $\frac{1}{P(T, S)}$ bonds with maturity S .
- At time S we receive $\frac{1}{P(T, S)} - 1$.

Scaling the strategy with $1/(S - T)$ yields the assertion.

2. For each $i = 1, \dots, n$ we have

$$F(T_{i-1}, T_i) = \frac{1}{\delta} \left(\frac{1}{P(T_{i-1}, T_i)} - 1 \right),$$

Bitte wenden!

which yields the cash flow at time T_i for each caplet

$$\begin{aligned}\delta(F(T_{i-1}, T_i) - \kappa)^+ &= \delta \left(\frac{1}{\delta} \left(\frac{1}{P(T_{i-1}, T_i)} - 1 \right) - \kappa \right)^+ \\ &= \left(\frac{1}{P(T_{i-1}, T_i)} - (1 + \delta\kappa) \right)^+, \end{aligned}$$

and hence, the time T_{i-1} value of $\delta(F(T_{i-1}, T_i) - \kappa)^+$ paid out at T_i is given by

$$\begin{aligned}P(T_{i-1}, T_i)\delta(F(T_{i-1}, T_i) - \kappa)^+ &= P(T_{i-1}, T_i) \left(\frac{1}{P(T_{i-1}, T_i)} - (1 + \delta\kappa) \right)^+ \\ &= (1 + \delta\kappa) \left(\frac{1}{1 + \delta\kappa} - P(T_{i-1}, T_i) \right)^+ . \end{aligned}$$

This is $(1 + \delta\kappa)$ times the cash-flow at date T_{i-1} of a put option on a T_i -bond with strike price $\frac{1}{1 + \delta\kappa}$ and maturity T_{i-1} .

3. a) Applying the definition of the $AR(1)$ process recursively N times we get

$$\begin{aligned}X_t &= c + \varphi(c + \varphi \cdot X_{t-2} + \varepsilon_{t-1}) + \varepsilon_t \\ &= \dots \\ &= c \cdot \sum_{k=0}^{N-1} \varphi^k + \varphi^N X_{t-N} + \sum_{k=0}^{N-1} \varphi^k \varepsilon_{t-k} . \end{aligned}$$

Therefore, if X is stationary and has finite second moment, then

$$C(0) := \text{Var}[X_t] = \varphi^{2N} C(0) + \sum_{k=0}^{N-1} \varphi^{2k} \cdot \sigma_\varepsilon^2 \quad (1)$$

which in turn implies that $\varphi^2 < 1$. On the other hand, if we assume that $|\varphi| < 1$ holds. Then, by letting $N \rightarrow \infty$ we obtain

$$X_t = c \cdot \sum_{k=0}^{\infty} \varphi^k + \sum_{k=0}^{\infty} \varphi^k \varepsilon_{t-k} = \frac{c}{1 - \varphi} + \sum_{k=0}^{\infty} \varphi^k \varepsilon_{t-k} . \quad (2)$$

Since the ε_t are assumed to be i.i.d, the stationarity follows.

b) The assertions are immediate consequences of (2) and (1).

c) Using Itô's formula the SDE can be solved explicitly

$$Y_t = y \cdot \exp(\mu t + \sigma W_t - \sigma^2 t/2).$$

Siehe nächstes Blatt!

Therefore, we have

$$\begin{aligned}\mathbb{E}[Y_t] &= y \cdot \exp(\mu t), \\ \text{Var}[Y_t] &= y^2 \exp(2\mu t)(\exp(\sigma^2 t) - 1).\end{aligned}$$

Since the right hand side diverges for $t \rightarrow \infty$, the process can not be stationary.

4. (i) see *driftedBM.m*, *AR1.m*, *poissonprocess.m* and *compoundpoissonprocess.m*

(ii) (a) We first show that $\mathbb{E}[e^{\sigma W_t}] = e^{\frac{1}{2}\sigma^2 t}$. Indeed,

$$\begin{aligned}\mathbb{E}[e^{\sigma W_t}] &= \int_{-\infty}^{\infty} \frac{1}{\sqrt{2\pi t}} e^{-\frac{1}{2t}x^2} \cdot e^{\sigma x} dx \\ &= \int_{\mathbb{R}} \frac{1}{\sqrt{2\pi t}} e^{-\frac{1}{2t}((x-\sigma t)^2 - \sigma^2 t^2)} dx \\ &= e^{\frac{1}{2}\sigma^2 t}.\end{aligned}$$

Hence,

$$\mathbb{E}[e^{X_1^{(a)}}] = e^{3+4/2} = 148.41$$

(b) Using Question 3 we get

$$\mathbb{E}[X_1^{(b)}] = \frac{c}{1 - \varphi} = 0.5/(1 - 0.6) = 1.25$$

(c) Apply the definition of expectation we have

$$\begin{aligned}\mathbb{E}[e^{X_1^{(c)}}] &= \sum_{k=0}^{\infty} e^{-\lambda} \frac{\lambda^k}{k!} e^k \\ &= e^{-\lambda} \sum_{k=0}^{\infty} \frac{(\lambda e)^k}{k!} \\ &= e^{\lambda(e-1)}.\end{aligned}$$

Plugging in $\lambda = 2$ we get $\mathbb{E}[e^{X_1^{(c)}}] = 31.08$.

(d) For the compound Poisson process we have to condition on N_1 , i.e.,

$$\begin{aligned}\mathbb{E}[e^{X_1^{(d)}}] &= \mathbb{E}[\mathbb{E}[e^{X_1^{(c)}} | N_1]] \\ &= \sum_{k=0}^{\infty} \mathbb{E}[e^{X_1^{(c)}} | N_1 = k] \mathbb{P}[N_1 = k] \\ &= \sum_{k=0}^{\infty} e^{-\lambda} \frac{\lambda^k}{k!} \mathbb{E}[\prod_{j=1}^k e^{Z_j}] \\ &= e^{-\lambda} \sum_{k=0}^{\infty} \frac{(\lambda\beta)^k}{k!} \\ &= e^{\lambda(\beta-1)},\end{aligned}$$

Bitte wenden!


where $\beta = \mathbb{E}[e^{Z_1}] = 0.5(e + e^{-1})$. Plugging in $\lambda = 2$ we get $\mathbb{E}[e^{X_1^{(d)}}] = 2.96$.

5. see nelsonsiegel.m

6. Matlab Files

```
1 function [simulatedvalue , theoreticalvalue]= driftedBM
2 % Matlab simulation drifted BM X= 1+2t+2W_t
3 tic
4 %% parameter input
5 % horizon
6 T=1;
7 % sample size
8 Nplot=10;
9 Nsimu=10^5;
10 % grid points
11 M=10^3;
12 % volatility
13 sigma=2;
14 driftcoeff=2;
15
16 % theoretical value
17 theoreticalvalue= exp(1+driftcoeff*T+sigma^2/2*T);
18 %% Simulation
19 % BM
20 BM = [zeros(1 , Nplot) ; sqrt( T/M) *cumsum( randn(M, Nplot) ) ];
21 % the process X^(a)
22 timegrid= 0:T/M:T;
23 drift=repmat( timegrid ' , 1 , Nplot );
24 Xa=1+driftcoeff*drift+sigma*BM;
25
26 %plot the sample paths
27 plot(timegrid ,Xa(: ,:))
28
29 %compute simulated value
30 normalrv=sqrt(T)*randn(1 , Nsimu) ;
31 simulatedvalue= mean(exp(1+ driftcoeff*T+sigma*normalrv))
    ;
32
33 %estimated variance
34 estvar= var(exp(1+ driftcoeff*T+sigma*normalrv)) ;
35 % confidence interval using CLT
```

Siehe nächstes Blatt!



```

36 cfplus=simulatedvalue+1.96*sqrt(estvar/Nsimu);
37 cfminus=simulatedvalue-1.96*sqrt(estvar/Nsimu);
38
39 disp('Exact value:')
40 disp(theoreticalvalue)
41 disp('Estimated value: ')
42 disp(simulatedvalue)
43 disp('Confidence interval: ')
44 disp([cfminus,cfplus])
45 toc

```

```

1 function [simulatedvalue,theoreticalvalue]= AR1
2 % In this exercise we simulated N paths of a AR(1)
   process
3 %  $X_t = c + \varphi X_{t-1} + \varepsilon_t$ ,  $\varepsilon_t \sim i.i.d.$ 
4 %  $N(0, \sigma^2)$ 
5 tic
6 %% parameter input
7 % horizon
8 T=1;
9 % sample size
10 Nsimu=10^5;
11 Nplot=Nsimu;
12 % grid points
13 M=10^3;

```

Bitte wenden!


```

14 % volatility stdnrv = sigma
15 stdnrv=sqrt(0.2);
16 c=0.5;
17 phi=0.6;
18 AR0= 0.1;
19 % time step
20 dt= T/M;
21
22 % theoretical value
23 theoreticalvalue= c/(1-phi);
24 %% Simulation
25 AR = [AR0*ones(M, Nplot) ];
26 % the process X^(b)
27 for i =1:M
28     AR(i+1 ,:)=c+ phi*AR(i ,:)+stdnrv.*randn(1 ,Nplot);
29 end
30
31 %plot the first sample path
32 timegrid= 0:dt:T;
33 plot(timegrid ,AR(: ,1))
34
35 %compute simulated value
36 simulatedvalue= mean(AR(end ,:));
37
38 %estimated variance
39 estvar= var(AR(end ,:));
40 % confidence interval using CLT
41 cfplus=simulatedvalue+1.96*sqrt(estvar/Nsimu);
42 cfminus=simulatedvalue -1.96*sqrt(estvar/Nsimu);
43
44 disp('Exact value:')
45 disp(theoreticalvalue)
46 disp('Estimated value: ')
47 disp(simulatedvalue)
48 disp('Confidence interval: ')
49 disp([cfminus ,cfplus])
50 toc

1 function [simulatedvalue ,theoreticalvalue]=
    poissonprocess
2 % In this exercise we simulated 10 paths of Poisson
    process with
3 % intensity lambda=2

```

Siehe nächstes Blatt!



```

4 tic
5 %% parameter input
6 % horizon
7 T=1;
8 % sample size
9 Nplot=10;
10 Nsimu=10^5;
11 % grid points
12 M=10^3;
13 % intensity
14 lambda=2;
15
16 % theoretical value
17 theoreticalvalue= exp(lambda*(exp(1)-1));
18 %% Simulation
19 % method 1
20 figure(1)
21 % number of total jumps at T
22 NT= poissrnd(lambda*T,1,Nplot);
23 temp1= zeros(max(NT),Nplot);
24 for i =1:Nplot
25     %condition on N_T, the jump times are ordered
        uniformly distributed
26     temp1(1:NT(i),i)= sort(T*rand(1,NT(i)));
27     stairs([0;temp1(1:NT(i),i)],(0:NT(i))');
28     hold on

```

Bitte wenden!


```

29 end
30 hold off
31
32 % method 2
33 PP = [zeros(1 , Nplot); cumsum( poissrnd( lambda*T/M,M, Nplot)
    )];
34 % the process X^(c)
35 timegrid= 0:T/M:T;
36 Xc=PP;
37
38 figure(2)
39 %show the first 10 sample paths
40 plot(timegrid ,Xc(:, :))
41
42
43 %compute simulated value
44 poissonrv=poissrnd( lambda*T,1 , Nsimu);
45 simulatedvalue= mean(exp( poissonrv));
46
47 %estimated variance
48 estvar= var(exp( poissonrv));
49 % confidence interval using CLT
50 cfplus=simulatedvalue+1.96*sqrt( estvar /Nsimu);
51 cfminus=simulatedvalue -1.96*sqrt( estvar /Nsimu);
52
53 disp( 'Exact value: ' )
54 disp(theoreticalvalue)
55 disp( 'Estimated value: ' )
56 disp(simulatedvalue)
57 disp( 'Confidence interval: ' )
58 disp( [ cfminus , cfplus ])
59 toc

1 function [simulatedvalue ,theoreticalvalue]=
    compoundpoissonprocess
2 % In this exercise we simulated 10 paths of compound
    Poisson process
3 % X^(c) _t= \sum_{i=1}^N_t Z_i with intensity lambda=2 and
    Z_i = 1/-1 with
4 % prob=0.5
5 tic
6 %% parameter input
7 % horizon

```

Siehe nächstes Blatt!



```

8 T=1;
9 % sample size
10 Nplot=10;
11 Nsimu=10^5;
12 % grid points
13 M=10^3;
14 % intensity
15 lambda=2;
16 % expectation of e^Z
17 beta = 0.5*(exp(1)+exp(-1));
18
19 % theoretical value
20 theoreticalvalue= exp(lambda*T*(beta-1));
21 %% Simulation
22 % compound poisson process
23 figure(1)
24 % method 1
25 % number of total jumps at T
26 NT= poissrnd(lambda*T,1,Nplot);
27 temp1= zeros(max(NT),Nplot);
28 for i =1:Nplot
29     %condition on N_T, the jump times are ordered
        uniformly distributed
30     temp1(1:NT(i),i)= sort(T*rand(1,NT(i)));
31     % simulate Z
32     stairs([0;temp1(1:NT(i),i)],cumsum([0,2*unidrnd(2,1,

```


Bitte wenden!

```

        NT(i) - 3])')
33     hold on;
34 end
35 hold off;
36
37
38 % method 2
39 figure (2)
40 CPP = [zeros(1, Nplot); cumsum(poissrnd(lambda*T/M, M, Nplot
        ).*(2*unidrnd(2, M, Nplot) - 3))];
41 % the process X^(a)
42 timegrid = 0:T/M:T;
43 Xc = CPP;
44
45 % show the first 10 sample paths
46 plot(timegrid, Xc(:, 1:10))
47
48 % compute simulated value
49 % the poisson process N
50 poissonrv = poissrnd(lambda*T, 1, Nsimu);
51 maxN = max(poissonrv);
52 % the jump distribution Z
53 bernoullirv = 2*unidrnd(2, maxN, Nsimu) - 3;
54 % place holder
55 temp = zeros(1, Nsimu);
56 for i = 1:maxN
57     % we only consider those numbers for which the
        % poisson process is > 0
58     index = (poissonrv > 0);
59     temp(index) = temp(index) + bernoullirv(i, index);
60     poissonrv(index) = poissonrv(index) - 1;
61 end
62 simulatedvalue = mean(exp(temp));
63
64
65 % estimated variance
66 estvar = var(exp(temp));
67 % confidence interval using CLT
68 cfplus = simulatedvalue + 1.96*sqrt(estvar/Nsimu);
69 cfminus = simulatedvalue - 1.96*sqrt(estvar/Nsimu);
70
71 disp('Exact value:')

```

Siehe nächstes Blatt!



```

72 disp(theoreticalvalue)
73 disp('Estimated value: ')
74 disp(simulatedvalue)
75 disp('Confidence interval: ')
76 disp([cfminus, cfplus])
77 toc

```

```

1 function [zoptns, zoptsv, resns, ressv]=nelsonsiegel
2 % Given a finite number of bond prices we plot the
3 % calibrated forward curve
4 %using Nelson Siegel family/ Svensson family.
5 %% Input
6 %p= [n*1] bond price; n = number of bonds
7 % Table 3.2 first 3 bonds
8 %C= [n*N] CF matrix; N = number of cash flow dates (e.g
9 % .., coupon payments)
9 %T= [N*1] payment tenor
10 p=[103.82;106.04;118.44];
11 C=zeros(length(p),1+3+6);
12 C(3,1)=6.125;
13 C(1,2)=105;
14 C(2,3)=4.875;
15 C(3,4)=6.125;
16 C(2,5)=4.875;
17 C(3,6)=6.125;

```


Bitte wenden!

```

18 C(2,7)=104.875;
19 C(3,8)=6.125;
20 C(3,9)=6.125;
21 C(3,10)=106.125;
22
23 % number of days between today 04/09/96 and the next CF
    date
24 T=[22;72;137;203;318;387;502;568;752;933];
25 % day count convention= actual/365
26 T=T/365;
27
28 %% Optimization
29 % bond prices d(z) using Nelson-Siegel Ansatz
30 % z is four-dimensional
31 funns = @(z,C) C*(exp(-(z(1)*T+z(2)/z(4)*(1-exp(-z(4)*T)
    )-z(3)/z(4)*T.*exp(-z(4)*T)+z(3)/(z(4)^2)*(1-exp(-z(4)
    *T))));
32
33 % bond prices d(z) using Svensson Ansatz
34 % z is six-dimensional
35 funsv = @(z,C) C*(exp(-(z(1)*T+z(2)/z(5)*(1-exp(-z(5)*T)
    )-z(3)/z(5)*T.*exp(-z(5)*T)+z(3)/(z(5)^2)*(1-exp(-z(5)
    *T))-z(4)/z(6)*T.*exp(-z(6)*T)+z(4)/(z(6)^2)*(1-exp(-z
    (6)*T))));
36
37 % numerical search for the least squared problem
38 % min || C*d(z) - p || over z
39
40 z0ns=0.05*ones(4,1);
41 z0sv=0.1*ones(6,1);
42 [zoptns, resns]= lsqcurvefit(@(z,C) funns(z,C), z0ns, C, p);
43 [zoptsv, ressv]= lsqcurvefit(@(z,C) funsv(z,C), z0sv, C, p);
44
45
46 % plot the forward curve
47 figure(1)
48 time = 0:0.01:T(end);
49 NSforward= zoptns(1)+(zoptns(2)+zoptns(3)*time).*exp(-
    zoptns(4)*time);
50 SVforward= zoptsv(1)+(zoptsv(2)+zoptsv(3)*time).*exp(-
    zoptsv(5)*time)+zoptsv(4)*time.*exp(-zoptsv(6)*time);
51 plot(time, NSforward, 'r-', time, SVforward, 'b-')

```

Siehe nächstes Blatt!



```
52 legend('Nelson-Siegel', 'Svensson')
53 xlabel('time')
54 ylabel('instantaneous forward rate')
55 title('calibrated forward curve -- today is 04/09/96')
56 end
```