

PERIODIC SOLUTIONS FOR A CLASS OF FUNCTIONAL DIFFERENTIAL SYSTEM

WEIBING WANG AND BAISHUN LAI

ABSTRACT. In this paper, we study the existence of periodic solutions to a class of functional differential system. By using Schauder's fixed point theorem, we show that the system has aperiodic solution under given conditions. Finally, four examples are given to demonstrate the validity of our main results.

1. INTRODUCTION

In this article, we study the existence of ω -periodic solutions to the following functional differential system

$$(1.1) \quad x'_i(t) = a_i(t)g_i(x_i(t)) - f_i(t, x_1(t - \tau_1(t)), \dots, x_n(t - \tau_n(t))), \quad i = 1, 2, \dots, n,$$

where $a_i, \tau_i: R \rightarrow R$ are ω -periodic continuous functions and $a_i(t) > 0$ for any $t \in [0, \omega]$, $f_i(t, u_1, \dots, u_n): R^{n+1} \rightarrow R$ is ω -periodic in t and $g_i: R \rightarrow R$.

When $n = 1$, the problem (1.1) reduces to the functional differential equation

$$(1.2) \quad x'(t) = a(t)g(x(t)) - h(t, x(t - \tau(t))).$$

The existence of periodic solutions for the special cases of (1.2) have been considered extensively by many authors, because (1.2) includes many important models in mathematical biology, such as, Hematopoiesis models; Nicholson's blowflies models; models for blood cell production, see [2, 3, 4, 8, 9, 7] and the references therein. Recently, Wang [5] investigated existence, multiplicity and nonexistence of positive periodic solutions for the periodic differential equation

$$(1.3) \quad x'(t) = a(t)p(x(t))x(t) - \lambda h(t)h(x(t - \tau(t))).$$

His approach depended on fixed point theorem in a cone. An essential condition on the function p in [5] is that p is bounded above and below by positive constants on $[0, +\infty)$. Hence, the method in [5] is not necessarily suitable for functional differential equation with general nonlinear term p . For example, to our best knowledge, results about periodic solutions for the following functional differential equation

$$(1.4) \quad x'(t) = a(t)x^\alpha(t) - \lambda h(t)f(x(t - \tau(t)))$$

2010 *Mathematics Subject Classification*: primary 34K13.

Key words and phrases: functional differential equation, periodic solution, fixed point theorem.

Received September 15, 2011, revised January 2012. Editor O. Došlý.

DOI: 10.5817/AM2012-2-139

are few, here $\alpha \neq 0$ is a constant and $\lambda > 0$ is a positive real parameter.

In the paper, we obtain sufficient conditions for the existence of periodic solutions for the system (1.1) by using Schauder's fixed point theorem. Our results improve and generalize the corresponding results of [1, 6, 10].

2. MAIN RESULTS

The following well-known Schauder's fixed point theorem is crucial in our arguments.

Lemma 2.1. *Let X be a Banach space with $D \subset X$ closed and convex. Assume that $T: D \rightarrow D$ is a completely continuous map, then T has a fixed point in D .*

Put $C_\omega = \{u \in C(R, R) : u(t + \omega) = u(t), t \in R\}$ with the norm defined by $\|u\|_{C_\omega} = \max_{0 \leq t \leq \omega} |u(t)|$ and

$$E = \{x = (x_1(t), \dots, x_n(t)) : x_i \in C_\omega\}, \quad \|x\|_E = \sum_{i=1}^n \|x_i\|_{C_\omega}.$$

Then C_ω and E are Banach spaces.

Let $p, q \in C_\omega$ and consider the following two differential equations

$$(2.1) \quad x'(t) = -p(t)x(t) + q(t),$$

$$(2.2) \quad x'(t) = p(t)x(t) - q(t).$$

Lemma 2.2. *Assume that $\int_0^\omega p(t)dt \neq 0$, then (2.1) has a unique ω -periodic solution*

$$x(t) = \int_t^{t+\omega} \frac{\exp \int_t^s p(r)dr}{\exp \int_0^\omega p(r)dr - 1} q(s) ds$$

and (2.2) has a unique ω -periodic solution

$$x(t) = \int_t^{t+\omega} \frac{\exp \int_s^{t+\omega} p(r) dr}{\exp \int_0^\omega p(r)dr - 1} q(s) ds.$$

Let $M \in R, m \in R: M > m$ and define

$$\prec_{[m, M]} = \{i : g_i(m) \leq g_i(M), 1 \leq i \leq n\},$$

$$\succ_{[m, M]} = \{i : g_i(m) > g_i(M), 1 \leq i \leq n\}.$$

By using Schauder's fixed point theorem, we obtain the following existence result on the periodic solution for (1.1).

Theorem 2.1. *Assume that there exist constants $M_i > m_i, i = 1, 2, \dots, n$ such that $g_i \in C^1([m_i, M_i], R), f_i \in C(R \times \Lambda, R)$, here $\Lambda = [m_1, M_1] \times \dots \times [m_n, M_n]$, and for any $u_i \in [m_i, M_i]$ and $t \in [0, \omega]$,*

$$(2.3) \quad g_i(M_i) \leq \frac{f_i(t, u_1, \dots, u_n)}{a_i(t)} \leq g_i(m_i) \quad \text{if } i \in \succ_{[m_i, M_i]}$$

and

$$(2.4) \quad g_i(m_i) \leq \frac{f_i(t, u_1, \dots, u_n)}{a_i(t)} \leq g_i(M_i) \quad \text{if } i \in \prec_{[m_i, M_i]} .$$

Then (1.1) has at least one periodic solution $(x_1^*(t), \dots, x_n^*(t)) \in E$ with $m_i \leq x_i^* \leq M_i (1 \leq i \leq n)$.

Proof. Without loss of the generality, we assume that there exists a $k: 0 \leq k \leq n$ such that

$$i \in \succ_{[m_i, M_i]} \quad \text{for } 1 \leq i \leq k, \quad i \in \prec_{[m_i, M_i]} \quad \text{for } k + 1 \leq i \leq n,$$

here if $i \leq 0, \succ_{[m_i, M_i]} = \phi$, if $i \geq n + 1, \prec_{[m_i, M_i]} = \phi$.

Since $g_i \in C^1([m_i, M_i], R)$, there exist $l_i > 0$ such that

$$(2.5) \quad 1 + \frac{1}{l_i} g'_i(u) > 0, \quad u \in [m_i, M_i], \quad i = 1, 2, \dots, k,$$

$$(2.6) \quad 1 - \frac{1}{l_i} g'_i(u) > 0, \quad u \in [m_i, M_i], \quad i = 1 + k, \dots, n.$$

Assume that $(x_1(t), \dots, x_n(t)) \in E$ is a solution of (1.1), then

$$x'_i(t) = -l_i a_i(t) x_i(t) + a_i(t) \left[g_i(x_i(t)) + l_i x_i(t) - \frac{f_i(t, X(t-\tau(t)))}{a_i(t)} \right], \quad i = 1, 2, \dots, k,$$

$$x'_i(t) = l_i a_i(t) x_i(t) - a_i(t) \left[l_i x_i(t) - g_i(x_i(t)) + \frac{f_i(t, X(t-\tau(t)))}{a_i(t)} \right], \quad i = k + 1, \dots, n$$

and

$$x_i(t) = \int_t^{t+\omega} \frac{a_i(s) \exp \int_t^s l_i a_i(r) dr}{\exp \int_0^\omega l_i a_i(r) dr - 1} \left[g_i(x_i(s)) + l_i x_i(s) - \frac{f_i(s, X(s-\tau(s)))}{a_i(s)} \right] ds, \quad i = 1, 2, \dots, k,$$

$$x_i(t) = \int_t^{t+\omega} \frac{a_i(s) \exp \int_s^{t+\omega} l_i a_i(r) dr}{\exp \int_0^\omega l_i a_i(r) dr - 1} \left[l_i x_i(s) - g_i(x_i(s)) + \frac{f_i(s, X(s-\tau(s)))}{a_i(s)} \right] ds, \quad i = k + 1, \dots, n,$$

where $f_i(t, X(t - \tau(t))) = f_i(t, x_1(t - \tau_1(t)), \dots, x_n(t - \tau_n(t)))$.

Define a set Ω in E and an operator $T: E \rightarrow E$ by

$$\Omega = \{x \in E : m_i \leq x_i \leq M_i, i = 1, 2, \dots, n\},$$

$$(Tx)(t) = ((Tx_1)(t), (Tx_2)(t), \dots, (Tx_n)(t)), \quad x = (x_1(t), \dots, x_n(t)) \in E,$$

where

$$(Tx_i)(t) := \int_t^{t+\omega} \frac{a_i(s) \exp \int_t^s l_i a_i(r) dr}{\exp \int_0^\omega l_i a_i(r) dr - 1} \\ \times \left[g_i(x_i(s)) + l_i x_i(s) - \frac{f_i(s, X(s - \tau(s)))}{a_i(s)} \right] ds, \quad 1 \leq i \leq k,$$

$$(Tx_i)(t) := \int_t^{t+\omega} \frac{a_i(s) \exp \int_s^{t+\omega} l_i a_i(r) dr}{\exp \int_0^\omega l_i a_i(r) dr - 1} \\ \times \left[l_i x_i(s) - g_i(x_i(s)) + \frac{f_i(s, X(s - \tau(s)))}{a_i(s)} \right] ds, \quad k+1 \leq i \leq n.$$

First, we show that $T(\Omega) \subset \Omega$. Using (2.5) and (2.6), we obtain that for $x \in \Omega$,

$$m_i + \frac{1}{l_i} g_i(m_i) \leq x_i(t) + \frac{1}{l_i} g_i(x_i(t)) \leq M_i + \frac{1}{l_i} g_i(M_i), \quad i = 1, 2, \dots, k,$$

$$m_i - \frac{1}{l_i} g_i(m_i) \leq x_i(t) - \frac{1}{l_i} g_i(x_i(t)) \leq M_i - \frac{1}{l_i} g_i(M_i), \quad i = k+1, \dots, n.$$

Using (2.3) and (2.4), we have

$$(Tx_i)(t) = \int_t^{t+\omega} \frac{l_i a_i(s) \exp \int_t^s l_i a_i(r) dr}{\exp \int_0^\omega l_i a_i(r) dr - 1} \left[\frac{1}{l_i} g_i(x_i(s)) + x_i(s) - \frac{f_i(s, X(s - \tau(s)))}{l_i a_i(s)} \right] ds \\ \in \left[m_i \int_t^{t+\omega} \frac{l_i a_i(s) \exp \int_t^s l_i a_i(r) dr}{\exp \int_0^\omega l_i a_i(r) dr - 1} ds, M_i \int_t^{t+\omega} \frac{l_i a_i(s) \exp \int_t^s l_i a_i(r) dr}{\exp \int_0^\omega l_i a_i(r) dr - 1} ds \right] \\ = [m_i, M_i], \quad i = 1, 2, \dots, k,$$

$$(Tx_i)(t) = \int_t^{t+\omega} \frac{a_i(s) \exp \int_s^{t+\omega} l_i a_i(r) dr}{\exp \int_0^\omega l_i a_i(r) dr - 1} \left[l_i x_i(s) - g_i(x_i(s)) + \frac{f_i(s, X(s - \tau(s)))}{a_i(s)} \right] ds \\ \in \left[m_i \int_t^{t+\omega} \frac{l_i a_i(s) \exp \int_s^{t+\omega} l_i a_i(r) dr}{\exp \int_0^\omega l_i a_i(r) dr - 1} ds, M_i \int_t^{t+\omega} \frac{l_i a_i(s) \exp \int_s^{t+\omega} l_i a_i(r) dr}{\exp \int_0^\omega l_i a_i(r) dr - 1} ds \right] \\ = [m_i, M_i], \quad i = k+1, k+2, \dots, n.$$

Next, we show that $T: \Omega \rightarrow \Omega$ is completely continuous. Obviously, $T(\Omega)$ is a uniformly bounded set and T is continuous on Ω , so it suffices to show $T(\Omega)$ is equi-continuous by Ascoli-Arzela theorem. For any $x \in \Omega$, we have

$$(Tx_i)'(t) = -l_i a_i(t)(Tx_i)(t) + a_i(t) \left[g_i(x_i(t)) + l_i x_i(t) - \frac{f_i(t, X(t - \tau(t)))}{a_i(t)} \right], \\ i = 1, 2, \dots, k,$$

$$(Tx_i)'(t) = l_i a_i(t)(Tx_i)(t) - a_i(t) \left[l_i x_i(t) - g_i(x_i(t)) + \frac{f_i(t, X(t - \tau(t)))}{a_i(t)} \right], \\ i = k+1, \dots, n.$$

Since $T(\Omega)$ is bounded and f_i, g_i, a_i are continuous, there exists $\rho > 0$ such that

$$|(Tx_i)'(t)| \leq \rho, \quad x \in \Omega, \quad i = 1, 2, \dots, n,$$

which implies that $T(\Omega)$ is equi-continuous. So T is a completely continuous operator on Ω . Clearly, Ω is a close and convex set in E . Therefore, T has a fixed point $x^* \in \Omega$ by Lemma 2.1. Furthermore, $m_i \leq x_i^*(t) \leq M_i$, which means $(x_1^*(t), \dots, x_n^*(t)) \in E$ is a ω -periodic solution of (1.1). The proof is complete. \square

Remark 2.1. Assume that all conditions of Theorem 2.1 are satisfies. Further suppose that there exist $1 \leq i_0 \leq n$ and $t_0 \in [0, \omega]$ such that any $u_i \in [m_i, M_i]$,

$$(2.7) \quad \frac{f_{i_0}(t_0, u_1, \dots, u_n)}{a_{i_0}(t_0)} < g_{i_0}(m_{i_0}) \quad \text{if } i_0 \in \succ_{[m_{i_0}, M_{i_0}]}$$

and

$$(2.8) \quad \frac{f_{i_0}(t_0, u_1, \dots, u_n)}{a_{i_0}(t_0)} > g_{i_0}(m_{i_0}) \quad \text{if } i_0 \in \prec_{[m_{i_0}, M_{i_0}]} .$$

Then $x_{i_0}^* > m_{i_0}$ for any $t \in [0, \omega]$.

Proof. Assume that there is a $t^* \in [0, \omega]$ such that $x_{i_0}^*(t^*) = m_{i_0}$. Then

$$\begin{aligned} m_{i_0} &= \int_{t^*}^{t^*+\omega} \frac{a_{i_0}(s) \exp \int_{t^*}^s l_{i_0} a_{i_0}(r) dr}{\exp \int_0^\omega l_{i_0} a_{i_0}(r) dr - 1} \\ &\quad \times \left[g_{i_0}(x_{i_0}^*(s)) + l_{i_0} x_{i_0}^*(s) - \frac{f_{i_0}(s, X^*(s - \tau(s)))}{a_{i_0}(s)} \right] ds, \quad i_0 \leq k, \end{aligned}$$

or

$$\begin{aligned} m_{i_0} &= \int_{t^*}^{t^*+\omega} \frac{a_{i_0}(s) \exp \int_s^{t^*+\omega} l_{i_0} a_{i_0}(r) dr}{\exp \int_0^\omega l_{i_0} a_{i_0}(r) dr - 1} \\ &\quad \times \left[l_{i_0} x_{i_0}^*(s) - g_{i_0}(x_{i_0}^*(s)) + \frac{f_{i_0}(s, X^*(s - \tau(s)))}{a_{i_0}(s)} \right] ds, \quad i_0 > k, \end{aligned}$$

where $f_i(t, X^*(t - \tau(t))) = f_i(t, x_1^*(t - \tau_1(t)), \dots, x_n^*(t - \tau_n(t)))$.

On the other hand, since for $s \in [0, \omega]$,

$$\frac{g_{i_0}(x_{i_0}^*(s))}{l_{i_0}} + x_{i_0}^*(s) - \frac{f_{i_0}(s, X^*(s - \tau(s)))}{l_{i_0} a_{i_0}(s)} - m_{i_0} \geq 0 \quad \text{for } i_0 \leq k,$$

$$x_{i_0}^*(s) - \frac{g_{i_0}(x_{i_0}^*(s))}{l_{i_0}} + \frac{f_{i_0}(s, X^*(s - \tau(s)))}{l_{i_0} a_{i_0}(s)} - m_{i_0} \geq 0 \quad \text{for } i_0 > k,$$

one can obtain that for any $s \in [0, \omega]$,

$$\frac{g_{i_0}(x_{i_0}^*(s))}{l_{i_0}} + x_{i_0}^*(s) - \frac{f_{i_0}(s, X^*(s - \tau(s)))}{l_{i_0} a_{i_0}(s)} - m_{i_0} \equiv 0 \quad \text{for } i_0 \leq k$$

$$x_{i_0}^*(s) - \frac{g_{i_0}(x_{i_0}^*(s))}{l_{i_0}} + \frac{f_{i_0}(s, X^*(s - \tau(s)))}{l_{i_0} a_{i_0}(s)} - m_{i_0} \equiv 0 \quad \text{for } i_0 > k,$$

which is a contradiction since

$$\begin{aligned} 0 &\geq \frac{g_{i_0}(m_{i_0})}{l_{i_0}} - \frac{f_{i_0}(t_0, X^*(t_0 - \tau(t_0)))}{l_{i_0}a_{i_0}(t_0)} > 0 \quad \text{for } i_0 \leq k, \\ 0 &\geq -\frac{g_{i_0}(m_{i_0})}{l_{i_0}} + \frac{f_{i_0}(t_0, X^*(t_0 - \tau(t_0)))}{l_{i_0}a_{i_0}(t_0)} > 0 \quad \text{for } i_0 > k. \end{aligned}$$

□

Remark 2.2. Assume that all conditions of Theorem 2.1 are satisfied. Further suppose that there exist $1 \leq r_0 \leq n$ and $t_1 \in [0, \omega]$ such that any $u_i \in [m_i, M_i]$,

$$(2.9) \quad \frac{f_{r_0}(t_1, u_1, \dots, u_n)}{a_{r_0}(t_1)} > g_{r_0}(M_{r_0}) \quad \text{if } r_0 \in \succ_{[m_{r_0}, M_{r_0}]}$$

and

$$(2.10) \quad \frac{f_{r_0}(t_1, u_1, \dots, u_n)}{a_{r_0}(t_1)} < g_{r_0}(M_{r_0}) \quad \text{if } r_0 \in \prec_{[m_{r_0}, M_{r_0}]}$$

Then $x_{r_0}^* < M_{r_0}$ for any $t \in [0, \omega]$.

Consider the equations

$$(2.11) \quad x'(t) = -a(t)x(t) + f(t, x(t - \tau(t)));$$

$$(2.12) \quad x'(t) = a(t)x(t) - f(t, x(t - \tau(t)))$$

where f is ω -periodic in t , a, τ are ω -periodic continuous functions and $a(t) > 0$ for all $t \in R$.

Corollary 2.1. Assume that there exist constants $M > m$ such that $f \in C(R \times [m, M], R)$ and for any $u \in [m, M]$ and $t \in [0, \omega]$

$$ma(t) \leq f(t, u) \leq Ma(t).$$

Then (2.11) (or (2.12)) has at least one periodic solution $m \leq x \leq M$.

Next, we consider the existence of a positive ω -periodic solution for problem (1.4). We give explicit intervals of λ such that (1.4) has at least one positive ω -periodic solution.

In the following, we assume that $a, h, \tau: R \rightarrow R$ are ω -periodic continuous functions and $a(t) > 0, h(t) > 0$ for any $t \in [0, \omega]$. $f: (0, +\infty) \rightarrow (0, +\infty)$ is continuous.

Put

$$\begin{aligned} \bar{f}_0 &= \limsup_{t \rightarrow 0^+} \frac{f(t)}{t^\alpha}, \quad \underline{f}_0 = \liminf_{t \rightarrow 0^+} \frac{f(t)}{t^\alpha}, \quad \bar{f}_\infty = \limsup_{t \rightarrow +\infty} \frac{f(t)}{t^\alpha}, \quad \underline{f}_\infty = \liminf_{t \rightarrow +\infty} \frac{f(t)}{t^\alpha}, \\ \delta^* &= \max_{t \in [0, \omega]} \frac{h(t)}{a(t)}, \quad \delta = \min_{t \in [0, \omega]} \frac{h(t)}{a(t)}. \end{aligned}$$

Theorem 2.2. The problem (1.4) has at least one positive periodic solution if one of the following conditions holds:

$$(H_1) \quad \alpha < 0, \liminf_{t \rightarrow 0^+} f(t) > 0, \limsup_{t \rightarrow +\infty} f(t) < +\infty \text{ and} \\ (\underline{f}_\infty \delta)^{-1} < \lambda < (\bar{f}_0 \delta^*)^{-1};$$

$$(H_2) \quad \alpha > 0, \limsup_{t \rightarrow 0^+} f(t) < +\infty, \liminf_{t \rightarrow +\infty} f(t) > 0 \text{ and} \\ (\underline{f}_0 \delta)^{-1} < \lambda < (\bar{f}_\infty \delta^*)^{-1}.$$

Proof. Assume that (H_1) holds. From the definition of \bar{f}_0 , \underline{f}_∞ and (H_1) , there exist $r_1 > 0$ and $\bar{r}_1 > r_1$ such that

$$\frac{\lambda h(t)f(u)}{a(t)} \leq u^\alpha, \quad 0 < u \leq r_1, \quad \inf_{u \in (0, r_1]} f(u) > 0, \\ \frac{\lambda h(t)f(u)}{a(t)} \geq u^\alpha, \quad u \geq \bar{r}_1, \quad \sup_{u \in [\bar{r}_1, +\infty)} f(u) < +\infty.$$

Let $\lambda \in (\frac{1}{\underline{f}_\infty \delta}, \frac{1}{\bar{f}_0 \delta^*})$. It is easy to check that

$$\inf \left\{ \frac{\lambda h(t)f(u)}{a(t)} : t \in [0, \omega], u \in (0, \bar{r}_1] \right\} := \mu_1 > 0, \\ \sup \left\{ \frac{\lambda h(t)f(u)}{a(t)} : t \in [0, \omega], u \in [r_1, +\infty) \right\} := \bar{\mu}_1 < +\infty.$$

Put

$$m = \min \left\{ \frac{r_1}{2}, \bar{\mu}_1^{\frac{1}{\alpha}} \right\}, \quad M = \max \left\{ 2\bar{r}_1, \mu_1^{\frac{1}{\alpha}} \right\},$$

then

$$M^\alpha \leq \mu_1 \leq \frac{\lambda h(t)f(u)}{a(t)} \leq x^\alpha \leq m^\alpha, \quad m \leq u \leq r_1, \\ M^\alpha \leq x^\alpha \leq \frac{\lambda h(t)f(u)}{a(t)} \leq \bar{\mu}_1 \leq m^\alpha, \quad \bar{r}_1 \leq u \leq M.$$

On the other hand,

$$M^\alpha \leq \mu_1 \leq \frac{\lambda h(t)f(u)}{a(t)} \leq \bar{\mu}_1 \leq m^\alpha, \quad r_1 \leq u \leq \bar{r}_1.$$

Hence,

$$M^\alpha \leq \frac{\lambda h(t)f(u)}{a(t)} \leq m^\alpha, \quad m \leq u \leq M.$$

By Theorem 2.1, (1.4) has at least one periodic solution $x \in C_\omega : 0 < m \leq x \leq M$.

Assume that (H_2) holds. There exist $0 < r_3 < 1$ and $\bar{r}_3 > 1$ such that

$$\frac{\lambda h(t)f(u)}{a(t)} \geq u^\alpha, \quad 0 < u \leq r_3, \quad \sup_{u \in (0, r_3]} f(u) < +\infty, \\ \frac{\lambda h(t)f(u)}{a(t)} \leq u^\alpha, \quad u \geq \bar{r}_3, \quad \inf_{u \in [\bar{r}_3, +\infty)} f(u) > 0.$$

Let $\lambda \in \left(\frac{1}{\underline{f}_0 \delta}, \frac{1}{\bar{f}_\infty \delta^*}\right)$, then

$$\inf \left\{ \frac{\lambda h(t) f(u)}{a(t)} : t \in [0, \omega], u \in [r_3, +\infty) \right\} := \mu_3 > 0,$$

$$\sup \left\{ \frac{\lambda h(t) f(u)}{a(t)} : t \in [0, \omega], u \in (0, \bar{r}_3] \right\} := \bar{\mu}_3 < +\infty.$$

Put

$$m = \min \left\{ \frac{r_3}{2}, \mu_3^{\frac{1}{\alpha}} \right\}, \quad M = \max \left\{ 2\bar{r}_3, \bar{\mu}_3^{\frac{1}{\alpha}} \right\},$$

then

$$m^\alpha \leq \frac{\lambda h(t) f(u)}{a(t)} \leq M^\alpha, \quad m \leq u \leq M.$$

By Theorem 2.1, (1.4) has at least one periodic solution $x \in C_\omega$: $0 < m \leq x \leq M$. The proof is complete. \square

Corollary 2.2.

- (1) Assume that $\alpha < 0$ and $0 < \liminf_{t \rightarrow 0^+} f(t) \leq \limsup_{t \rightarrow 0^+} f(t) < +\infty$, then (1.4) has at least one positive periodic solution for sufficiently large $\lambda > 0$.
- (2) Assume that $\alpha < 0$ and $0 < \liminf_{t \rightarrow +\infty} f(t) \leq \limsup_{t \rightarrow +\infty} f(t) < +\infty$, then (1.4) has at least one positive periodic solution for sufficiently small $\lambda > 0$.
- (3) Assume that $\alpha > 0$ and $0 < \liminf_{t \rightarrow 0^+} f(t) \leq \limsup_{t \rightarrow 0^+} f(t) < +\infty$, then (1.4) has at least one positive periodic solution for sufficiently small $\lambda > 0$.
- (4) Assume that $\alpha > 0$ and $0 < \liminf_{t \rightarrow +\infty} f(t) \leq \limsup_{t \rightarrow +\infty} f(t) < +\infty$, then (1.4) has at least one positive periodic solution for sufficiently large $\lambda > 0$.

Proof. Here we only prove case (1). Since $0 < \liminf_{t \rightarrow 0^+} f(t) \leq \limsup_{t \rightarrow 0^+} f(t) < +\infty$, there exists $0 < r < 1$ such that

$$\mu := \inf_{t \in (0, r]} f(t) \leq \sup_{t \in (0, r]} f(t) := \nu < +\infty.$$

Let $\lambda > 0$ such that $(\lambda \delta \mu)^{\frac{1}{\alpha}} < r$ and set

$$m = (\lambda \delta \nu)^{\frac{1}{\alpha}}, \quad M = (\lambda \delta \mu)^{\frac{1}{\alpha}},$$

then $r > M > m > 0$ and

$$M^\alpha \leq \frac{\lambda h(t) f(u)}{a(t)} \leq m^\alpha, \quad m \leq u \leq M.$$

By Theorem 2.1, (1.4) has at least one periodic solution $x \in C_\omega$: $0 < m \leq x \leq M$. The proof is complete. \square

3. SOME EXAMPLES

In this section, we apply the main results obtained in previous section to several examples.

Example 3.1. Consider the differential equation

$$(3.1) \quad x'(t) = \frac{1}{\sqrt[3]{\sin x(t)}} + b(t),$$

where $b(t)$ is a ω -periodic continuous function.

It is easy to verify from Theorem 2.1 that (3.1) has least two periodic solutions $0 < |x_1| < 0.5\pi < |x_2| < \pi$ if $|b(t)| > 1$ for all $t \in R$. Since $x_i + 2k\pi (i = 1, 2, k \in Z)$ is also the periodic solutions of (3.1), (3.1) has infinitely many periodic solutions when $|b(t)| > 1$.

Example 3.2. Consider the differential equation

$$(3.2) \quad x'(t) = \left(1 + \frac{\sin t}{100}\right)x^3(t) - f(x(t - \cos t)),$$

where

$$f(u) = \begin{cases} 0.1, & u < \frac{2}{3}, \\ u^2 - u + \frac{5}{4}, & u > 1. \end{cases}$$

In (3.2), $a(t) = 1 + 0.01 \sin t$ and $g(x) = x^3$. Put $m_1 = 0.1, M_1 = 0.6, m_2 = 1.1, M_2 = 2$, then

$$g(m_i) \leq \frac{f(u)}{a(t)} \leq g(M_i), \quad \forall u \in [m_i, M_i], t \in R, i = 1, 2.$$

By Theorem 2.1, (3.2) has two positive 2π -periodic solutions x_1, x_2 such that $m_1 \leq x_1 \leq M_1, m_2 \leq x_2 \leq M_2$.

Example 3.3. Consider the differential equation

$$(3.3) \quad x'(t) = x^3(t) + \frac{1}{x(t)} - \lambda \left(1 + \frac{\sin t}{2}\right) (2 - \sin x(t - \cos t))$$

where $\lambda > 0$ is a positive real parameter.

In (3.3), $a(t) = 1, g(x) = x^3 + x^{-1}$ and $f(t, u) = \lambda \left(1 + \frac{\sin t}{2}\right) (2 - \sin u)$. Put

$$m_1 = \frac{2}{9\lambda}, \quad M_1 = 1, \quad m_2 = 1, \quad M_2 = \sqrt[3]{\frac{9\lambda}{2}},$$

then for sufficiently large $\lambda > 0$,

$$g(M_1) \leq f(t, u) \leq g(m_1), \quad u \in [m_1, M_1], t \in R,$$

$$g(m_2) \leq f(t, u) \leq g(M_2), \quad u \in [m_2, M_2], t \in R.$$

By Theorem 2.1, (3.3) has two positive 2π -periodic solutions $x_1 \in [m_1, M_1], x_2 \in [m_2, M_2]$ for sufficiently large $\lambda > 0$.

Example 3.4. Consider the differential system

$$(3.4) \quad \begin{cases} x'(t) = (2 - \cos t)x(t) - y^2(t), \\ y'(t) = -2 \sin y(t) + \exp(0.5x(t) - y(t)). \end{cases}$$

Put $D = [0.01, 0.29] \times [0.2, 0.53]$, then for $(u_1, u_2) \in D$ and $t \in [0, 2\pi]$,

$$0.01 \leq \frac{u_2^2}{2 - \cos t} \leq 0.29, \quad 2 \sin 0.2 \leq e^{0.5u_1 - u_2} \leq 2 \sin 0.53.$$

By Theorem 2.1, (3.4) has a 2π -periodic solution $(x(t), y(t))$ such that $0.01 \leq x(t) \leq 0.29$ and $0.2 \leq y(t) \leq 0.53$.

Acknowledgement. The authors would like to thank the referees for the comments which help to improve the paper. The work is supported by Hunan Provincial Natural Science Foundation of China.

REFERENCES

- [1] A. Wan, D. Jiang, Xu, X., *A new existence theory for positive periodic solutions to functional differential equations*, Comput. Math. Appl. **47** (2004), 1257–1262.
- [2] Cheng, S., Zhang, G., *Existence of positive periodic solutions for non-autonomous functional differential equations*, Electron. J. Differential Equations **59** (2001), 1–8.
- [3] Chow, S., *Existence of periodic solutions of autonomous functional differential equations*, Differential Equations **15** (1974), 350–378.
- [4] Freedman, H. I., Wu, J., *Periodic solutions of single-species models with periodic delay*, SIAM J. Math. Anal. **23** (1992), 689–701.
- [5] H.Wang, *Positive periodic solutions of functional differential equations*, J. Differential Equations **202** (2004), 354–366.
- [6] Jiang, D., Wei, J., Zhang, B., *Positive periodic solutions of functional differential equations and population models*, Electron. J. Differential Equations **71** (2002), 1–13.
- [7] Joseph, W., So, H., Yu, J., *Global attractivity and uniform persistence in Nicholson's blowflies*, Differential Equations Dynam. Systems **1** (1994), 11–18.
- [8] Kuang, Y., Smith, H. L., *Periodic solutions of differential delay equations with threshold-type delays, oscillations and dynamics in delay equations*, Contemp. Math. **129** (1992), 153–176.
- [9] Li, Z., Wang, X., *Existence of positive periodic solutions for neutral functional differential equations*, Electron. J. Differential Equations **34** (2006), 1–8.
- [10] Weng, P., Liang, M., *The existence and behavior of periodic solution of hematopoiesis model*, Math. Appl. **4** (1995), 434–439.

DEPARTMENT OF MATHEMATICS, HUNAN UNIVERSITY OF SCIENCE AND TECHNOLOGY
 XIANGTAN, HUNAN 411201, P.R. CHINA
E-mail: wzbhunan@yahoo.cn

SCHOOL OF MATHEMATICS, HENAN UNIVERSITY,
 KAIFENG, HENAN 475004, P.R. CHINA