

## ON RIEMANN-POISSON LIE GROUPS

BRAHIM ALIOUNE<sup>a</sup>, MOHAMED BOUCETTA<sup>b</sup>, AND AHMED SID'AHMED LESSIAD<sup>c</sup>

ABSTRACT. A Riemann-Poisson Lie group is a Lie group endowed with a left invariant Riemannian metric and a left invariant Poisson tensor which are compatible in the sense introduced in [4]. We study these Lie groups and we give a characterization of their Lie algebras. We give also a way of building these Lie algebras and we give the list of such Lie algebras up to dimension 5.

## 1. INTRODUCTION

In this paper, we study Lie groups endowed with a left invariant Riemannian metric and a left invariant Poisson tensor satisfying a compatibility condition to be defined below. They constitute a subclass of the class of *Riemann-Poisson manifolds* introduced and studied by the second author (see [2, 3, 4, 5]).

Let  $(M, \pi, \langle \cdot, \cdot \rangle)$  be smooth manifold endowed with a Poisson tensor  $\pi$  and a Riemannian metric  $\langle \cdot, \cdot \rangle$ . We denote by  $\langle \cdot, \cdot \rangle^*$  the Euclidean product on  $T^*M$  naturally associated to  $\langle \cdot, \cdot \rangle$ . The Poisson tensor defines a Lie algebroid structure on  $T^*M$  where the anchor map is the contraction  $\#_\pi: T^*M \rightarrow TM$  given by  $\langle \beta, \#_\pi(\alpha) \rangle = \pi(\alpha, \beta)$  and the Lie bracket on  $\Omega^1(M)$  is the Koszul bracket given by

$$(1) \quad [\alpha, \beta]_\pi = \mathcal{L}_{\#_\pi(\alpha)}\beta - \mathcal{L}_{\#_\pi(\beta)}\alpha - d\pi(\alpha, \beta), \quad \alpha, \beta \in \Omega^1(M).$$

This Lie algebroid structure and the metric  $\langle \cdot, \cdot \rangle^*$  define a contravariant connection  $\mathcal{D}: \Omega^1(M) \times \Omega^1(M) \rightarrow \Omega^1(M)$  by Koszul formula

$$(2) \quad \begin{aligned} 2\langle \mathcal{D}_\alpha\beta, \gamma \rangle^* &= \#_\pi(\alpha) \cdot \langle \beta, \gamma \rangle^* + \#_\pi(\beta) \cdot \langle \alpha, \gamma \rangle^* - \#_\pi(\gamma) \cdot \langle \alpha, \beta \rangle^* \\ &+ \langle [\alpha, \beta]_\pi, \gamma \rangle^* + \langle [\gamma, \alpha]_\pi, \beta \rangle^* + \langle [\gamma, \beta]_\pi, \alpha \rangle^*, \quad \alpha, \beta, \gamma \in \Omega^1(M). \end{aligned}$$

This is the unique torsionless contravariant connection which is metric, i.e., for any  $\alpha, \beta, \gamma \in \Omega^1(M)$ ,

$$\mathcal{D}_\alpha\beta - \mathcal{D}_\beta\alpha = [\alpha, \beta]_\pi \quad \text{and} \quad \#_\pi(\alpha) \cdot \langle \beta, \gamma \rangle^* = \langle \mathcal{D}_\alpha\beta, \gamma \rangle^* + \langle \beta, \mathcal{D}_\alpha\gamma \rangle^*.$$

The notion of contravariant connection was introduced by Vaisman in [13] and studied in more details by Fernandes in the context of Lie algebroids [8]. The

---

2020 *Mathematics Subject Classification*: primary 53A15; secondary 53D17, 22E05.

*Key words and phrases*: Lie group, Poisson manifolds, Riemannian metric.

Received August 22, 2019. Editor J. Slovák.

DOI: 10.5817/AM2020-4-225

connection  $\mathcal{D}$  defined above is called *contravariant Levi-Civita connection* associated to the couple  $(\pi, \langle \cdot, \cdot \rangle)$  and it appeared first in [2].

The triple  $(M, \pi, \langle \cdot, \cdot \rangle)$  is called a *Riemannian-Poisson manifold* if  $\mathcal{D}\pi = 0$ , i.e., for any  $\alpha, \beta, \gamma \in \Omega^1(M)$ ,

$$(3) \quad \mathcal{D}\pi(\alpha, \beta, \gamma) := \#_{\pi}(\alpha) \cdot \pi(\beta, \gamma) - \pi(\mathcal{D}_{\alpha}\beta, \gamma) + \pi(\beta, \mathcal{D}_{\alpha}\gamma) = 0.$$

This notion was introduced by the second author in [2]. Riemann-Poisson manifolds turned out to have interesting geometric properties (see[2, 3, 4, 5]). Let's mention some of them.

- (1) The condition of compatibility (3) is weaker than the condition  $\nabla\pi = 0$  where  $\nabla$  is the Levi-Civita connection of  $\langle \cdot, \cdot \rangle$ . Indeed, the condition (3) allows the Poisson tensor to have a variable rank. For instance, linear Poisson structures which are Riemann-Poisson exist and were characterized in [5]. Furthermore, let  $(M, \langle \cdot, \cdot \rangle)$  be a Riemannian manifold and  $(X_1, \dots, X_r)$  a family of commuting Killing vector fields. Put

$$\pi = \sum_{i,j} X_i \wedge X_j.$$

Then  $(M, \pi, \langle \cdot, \cdot \rangle)$  is a Riemann-Poisson manifold. This example illustrates also the weakness of the condition (3) and, more importantly, it is the local model of the geometry of noncommutative deformations studied by Hawkins (see [10, Theorem 6.6]).

- (2) Riemann-Poisson manifolds can be thought of as a generalization of Kähler manifolds. Indeed, let  $(M, \pi, \langle \cdot, \cdot \rangle)$  be a Poisson manifold endowed with a Riemannian metric such that  $\pi$  is invertible. Denote by  $\omega$  the symplectic form inverse of  $\pi$ . Then  $(M, \pi, \langle \cdot, \cdot \rangle)$  is Riemann-Poisson manifold if and only if  $\nabla\omega = 0$  where  $\nabla$  is the Levi-Civita connection of  $\langle \cdot, \cdot \rangle$ . In this case, if we define  $A: TM \rightarrow TM$  by  $\omega(u, v) = \langle Au, v \rangle$  then  $-A^2$  is symmetric definite positive and hence there exists a unique  $Q: TM \rightarrow TM$  symmetric definite positive such that  $Q^2 = -A^2$ . It follows that  $J = AQ^{-1}$  satisfies  $J^2 = -\text{Id}_{TM}$ , skew-symmetric with respect  $\langle \cdot, \cdot \rangle$  and  $\nabla J = 0$ . Hence  $(M, J, \langle \cdot, \cdot \rangle)$  is a Kähler manifold and its Kähler form  $\omega_J(u, v) = \langle Ju, v \rangle$  is related to  $\omega$  by the following formula:

$$(4) \quad \omega(u, v) = -\omega_J\left(\sqrt{-A^2}u, v\right), \quad u, v \in TM.$$

Having this construction in mind, we will call in this paper a Kähler manifold a triple  $(M, \langle \cdot, \cdot \rangle, \omega)$  where  $\langle \cdot, \cdot \rangle$  is a Riemannian metric and  $\omega$  is a nondegenerate 2-form  $\omega$  such that  $\nabla\omega = 0$  where  $\nabla$  is the Levi-Civita connection of  $\langle \cdot, \cdot \rangle$ .

- (3) The symplectic foliation of a Riemann-Poisson manifold when  $\pi$  has a constant rank has an important property namely it is both a Riemannian foliation and a Kähler foliation.

Recall that a Riemannian foliation is a foliated manifold  $(M, \mathcal{F})$  with a Riemannian metric  $\langle \cdot, \cdot \rangle$  such that the orthogonal distribution  $T^{\perp}\mathcal{F}$  is totally geodesic.

Kähler foliations are a generalization of Kähler manifolds (see [6]) and, as for the notion of Kähler manifold, we call in this paper a Kähler foliation a foliated manifold  $(M, \mathcal{F})$  endowed with a leafwise metric  $\langle \cdot, \cdot \rangle_{\mathcal{F}} \in \Gamma(\otimes^2 T^* \mathcal{F})$  and a nondegenerate leafwise differential 2-form  $\omega_{\mathcal{F}} \in \Gamma(\otimes^2 T^* \mathcal{F})$  such any leaf with the restrictions of  $\langle \cdot, \cdot \rangle_{\mathcal{F}}$  and  $\omega_{\mathcal{F}}$  is a Kähler manifold.

**Theorem 1.1** ([4]). *Let  $(M, \langle \cdot, \cdot \rangle, \pi)$  be a Riemann-Poisson manifold with  $\pi$  of constant rank. Then its symplectic foliation is both a Riemannian and a Kähler foliation.*

Having in mind these properties particularly Theorem 1.1, it will be interesting to find large classes of examples of Riemann-Poisson manifolds. This paper will describe the rich collection of examples which are obtained by providing an arbitrary Lie group  $G$  with a Riemannian metric  $\langle \cdot, \cdot \rangle$  and a Poisson tensor  $\pi$  invariant under left translations and such that  $(G, \langle \cdot, \cdot \rangle, \pi)$  is Riemann-Poisson. We call  $(G, \langle \cdot, \cdot \rangle, \pi)$  a *Riemann-Poisson Lie group*. This class of examples can be enlarged substantially, with no extra work, as follows. If  $(G, \langle \cdot, \cdot \rangle, \pi)$  is a Riemann-Poisson Lie group and  $\Gamma$  is any discrete subgroup of  $G$  then  $\Gamma \backslash G$  carries naturally a structure of Riemann-Poisson manifold.

The paper is organized as follows. In Section 2, we give the material needed in the paper and we describe the infinitesimal counterpart of Riemann-Poisson Lie groups, namely, Riemann-Poisson Lie algebras. In Section 3, we prove our main result which gives an useful description of Riemann-Poisson Lie algebras (see Theorem 3.1). We use this theorem in Section 4 to derive a method for building Riemann-Poisson Lie algebras. We explicit this method by giving the list of Riemann-Poisson Lie algebras up to dimension 5.

## 2. RIEMANN-POISSON LIE GROUPS AND THEIR INFINITESIMAL CHARACTERIZATION

Let  $G$  be a Lie group and  $(\mathfrak{g} = T_e G, [\cdot, \cdot])$  its Lie algebra.

- (1) A left invariant Poisson tensor  $\pi$  on  $G$  is entirely determined by

$$\pi(\alpha, \beta)(a) = r(L_a^* \alpha, L_a^* \beta),$$

where  $a \in G$ ,  $\alpha, \beta \in T_a^* G$ ,  $L_a$  is the left multiplication by  $a$  and  $r \in \wedge^2 \mathfrak{g}$  satisfies the classical Yang-Baxter equation

$$(5) \quad [r, r] = 0,$$

where  $[r, r] \in \wedge^3 \mathfrak{g}$  is given by

$$(6) \quad [r, r](\alpha, \beta, \gamma) := \langle \alpha, [r_{\#}(\beta), r_{\#}(\gamma)] \rangle + \langle \beta, [r_{\#}(\gamma), r_{\#}(\alpha)] \rangle + \langle \gamma, [r_{\#}(\alpha), r_{\#}(\beta)] \rangle, \quad \alpha, \beta, \gamma \in \mathfrak{g}^*,$$

and  $r_{\#} : \mathfrak{g}^* \rightarrow \mathfrak{g}$  is the contraction associated to  $r$ . In this case, the Koszul bracket (1) when restricted to left invariant differential 1-forms induces a Lie bracket on  $\mathfrak{g}^*$  given by

$$(7) \quad [\alpha, \beta]_r = \text{ad}_{r_{\#}(\alpha)}^* \beta - \text{ad}_{r_{\#}(\beta)}^* \alpha, \quad \alpha, \beta \in \mathfrak{g}^*,$$

where  $\prec \text{ad}_u^* \alpha, v \succ = - \prec \alpha, [u, v] \succ$ . Moreover,  $r_{\#}$  becomes a morphism of Lie algebras, i.e.,

$$(8) \quad r_{\#}([\alpha, \beta]_r) = [r_{\#}(\alpha), r_{\#}(\beta)], \quad \alpha, \beta \in \mathfrak{g}^*.$$

(2) A left invariant Riemannian metric  $\langle \cdot, \cdot \rangle$  on  $G$  is entirely determined by

$$\langle u, v \rangle(a) = \rho(T_a L_{a^{-1}} u, T_a L_{a^{-1}} v),$$

where  $a \in G$ ,  $u, v \in T_a G$  and  $\rho$  is a scalar product on  $\mathfrak{g}$ . The Levi-Civita connection of  $\langle \cdot, \cdot \rangle$  is left invariant and induces a product  $A: \mathfrak{g} \times \mathfrak{g} \rightarrow \mathfrak{g}$  given by

$$(9) \quad 2\varrho(A_u v, w) = \varrho([u, v], w) + \varrho([w, u], v) + \varrho([w, v], u), \quad u, v, w \in \mathfrak{g}.$$

It is the unique product on  $\mathfrak{g}$  satisfying

$$A_u v - A_v u = [u, v] \quad \text{and} \quad \varrho(A_u v, w) + \varrho(v, A_u w) = 0,$$

for any  $u, v, w \in \mathfrak{g}$ . We call  $A$  the *Levi-Civita product* associated to  $(\mathfrak{g}, [\cdot, \cdot], \rho)$ .

(3) Let  $(G, \langle \cdot, \cdot \rangle, \Omega)$  be a Lie group endowed with a left invariant Riemannian metric and a nondegenerate left invariant 2-form. Then  $(G, \langle \cdot, \cdot \rangle, \Omega)$  is a Kähler manifold if and only if, for any  $u, v, w \in \mathfrak{g}$ ,

$$(10) \quad \omega(A_u v, w) + \omega(u, A_u v) = 0,$$

where  $\omega = \Omega(e)$ ,  $\rho = \langle \cdot, \cdot \rangle(e)$  and  $A$  is the Levi-Civita product of  $(\mathfrak{g}, [\cdot, \cdot], \rho)$ . In this case we call  $(\mathfrak{g}, [\cdot, \cdot], \rho, \omega)$  a Kähler Lie algebra.

As all the left invariant structures on Lie groups, Riemann-Poisson Lie groups can be characterized at the level of their Lie algebras.

**Proposition 2.1.** *Let  $(G, \pi, \langle \cdot, \cdot \rangle)$  be a Lie group endowed with a left invariant bivector field and a left invariant metric and  $(\mathfrak{g}, [\cdot, \cdot])$  its Lie algebra. Put  $r = \pi(e) \in \wedge^2 \mathfrak{g}$ ,  $\varrho = \langle \cdot, \cdot \rangle_e$  and  $\varrho^*$  the associated Euclidean product on  $\mathfrak{g}^*$ . Then  $(G, \pi, \langle \cdot, \cdot \rangle)$  is a Riemann-Poisson Lie group if and only if*

$$(i) \quad [r, r] = 0,$$

$$(ii) \quad \text{for any } \alpha, \beta, \gamma \in \mathfrak{g}^*, \quad r(A_\alpha \beta, \gamma) + r(\beta, A_\alpha \gamma) = 0,$$

where  $A$  is the Levi-Civita product associated to  $(\mathfrak{g}^*, [\cdot, \cdot]_r, \varrho^*)$ .

**Proof.** For any  $u \in \mathfrak{g}$  and  $\alpha \in \mathfrak{g}^*$ , we denote by  $u^\ell$  and  $\alpha^\ell$ , respectively, the left invariant vector field and the left invariant differential 1-form on  $G$  given by

$$u^\ell(a) = T_e L_a(u) \quad \text{and} \quad \alpha^\ell(a) = T_a^* L_{a^{-1}}(\alpha), \quad a \in G, \quad L_a(b) = ab.$$

Since  $\pi$  and  $\langle \cdot, \cdot \rangle$  are left invariant, one can see easily from (1) and (2) that we have, for any  $\alpha, \beta, \gamma \in \mathfrak{g}^*$ ,

$$\begin{cases} [\pi, \pi]_S(\alpha^\ell, \beta^\ell, \gamma^\ell) = [r, r](\alpha, \beta, \gamma), \quad \#_\pi(\alpha^\ell) = (r_{\#}(\alpha))^\ell, \quad \mathcal{L}_{\#_\pi(\alpha^\ell)} \beta^\ell = (\text{ad}_{r_{\#}(\alpha)}^* \beta)^\ell, \\ [\alpha^\ell, \beta^\ell]_\pi = ([\alpha, \beta]_r)^\ell, \quad \mathcal{D}_{\alpha^\ell} \beta^\ell = (A_\alpha \beta)^\ell. \end{cases}$$

The proposition follows from these formulas, (3) and the fact that  $(G, \pi, \langle \cdot, \cdot \rangle)$  is a Riemann-Poisson Lie group if and only if, for any  $\alpha, \beta, \gamma \in \mathfrak{g}^*$ ,

$$[\pi, \pi]_S(\alpha^\ell, \beta^\ell, \gamma^\ell) = 0 \quad \text{and} \quad \mathcal{D}\pi(\alpha^\ell, \beta^\ell, \gamma^\ell) = 0. \quad \square$$

Conversely, given a triple  $(\mathfrak{g}, r, \varrho)$  where  $\mathfrak{g}$  is a real Lie algebra,  $r \in \wedge^2 \mathfrak{g}$  and  $\varrho$  a Euclidean product on  $\mathfrak{g}$  satisfying the conditions (i) and (ii) in Proposition 2.1 then, for any Lie group  $G$  whose Lie algebra is  $\mathfrak{g}$ , if  $\pi$  and  $\langle \cdot, \cdot \rangle$  are the left invariant bivector field and the left invariant metric associated to  $(r, \varrho)$  then  $(G, \pi, \langle \cdot, \cdot \rangle)$  is a Riemann-Poisson Lie group.

**Definition 2.1.** A *Riemann-Poisson Lie algebra* is a triple  $(\mathfrak{g}, r, \varrho)$  where  $\mathfrak{g}$  is a real Lie algebra,  $r \in \wedge^2 \mathfrak{g}$  and  $\varrho$  a Euclidean product on  $\mathfrak{g}$  satisfying the conditions (i) and (ii) in Proposition 2.1.

To end this section, we give another characterization of the solutions of the classical Yang-Baxter equation (5) which will be useful later.

We observe that  $r \in \wedge^2 \mathfrak{g}$  is equivalent to the data of a vector subspace  $S \subset \mathfrak{g}$  and a nondegenerate 2-form  $\omega_r \in \wedge^2 S^*$ .

Indeed, for  $r \in \wedge^2 \mathfrak{g}$ , we put  $S = \text{Im} r_\#$  and  $\omega_r(u, v) = r(r_\#^{-1}(u), r_\#^{-1}(v))$  where  $u, v \in S$  and  $r_\#^{-1}(u)$  is any antecedent of  $u$  by  $r_\#$ .

Conversely, let  $(S, \omega)$  be a vector subspace of  $\mathfrak{g}$  with a non-degenerate 2-form. The 2-form  $\omega$  defines an isomorphism  $\omega^b: S \rightarrow S^*$  by  $\omega^b(u) = \omega(u, \cdot)$ , we denote by  $\#: S^* \rightarrow S$  its inverse and we put  $r_\# = \# \circ i^*$  where  $i^*: \mathfrak{g}^* \rightarrow S^*$  is the dual of the inclusion  $i: S \hookrightarrow \mathfrak{g}$ .

With this observation in mind, the following proposition gives another description of the solutions of the Yang-Baxter equation.

**Proposition 2.2.** *Let  $r \in \wedge^2 \mathfrak{g}$  and  $(S, \omega_r)$  its associated vector subspace. The following assertions are equivalent:*

- (1)  $[r, r] = 0$ .
- (2)  $S$  is a subalgebra of  $\mathfrak{g}$  and

$$\delta\omega_r(u, v, w) := \omega_r(u, [v, w]) + \omega_r(v, [w, u]) + \omega_r(w, [u, v]) = 0$$

for any  $u, v, w \in S$ .

**Proof.** The proposition follows from the following formulas:

$$\langle \gamma, r_\#([\alpha, \beta]_r) - [r_\#(\alpha), r_\#(\beta)] \rangle = -[r, r](\alpha, \beta, \gamma), \quad \alpha, \beta, \gamma \in \mathfrak{g}^*$$

and, if  $S$  is a subalgebra,

$$[r, r](\alpha, \beta, \gamma) = -\delta\omega_r(r_\#(\alpha), r_\#(\beta), r_\#(\gamma)). \quad \square$$

This proposition shows that there is a correspondence between the set of solutions of the Yang-Baxter equation the set of symplectic subalgebras of  $\mathfrak{g}$ . We recall that a symplectic algebra is a Lie algebra  $S$  endowed with a non-degenerate 2-form  $\omega$  such that  $\delta\omega = 0$ .

3. A CHARACTERIZATION OF RIEMANN-POISSON LIE ALGEBRAS

In this section, we combine Propositions 2.1 and 2.2 to establish a characterization of Riemann-Poisson Lie algebras which will be used later to build such Lie algebras. We establish first an intermediary result.

**Proposition 3.1.** *Let  $(\mathfrak{g}, r, \varrho)$  be a Lie algebra endowed with  $r \in \wedge^2 \mathfrak{g}$  and a Euclidean product  $\varrho$ . Denote by  $\mathcal{I} = \ker r_{\#}$ ,  $\mathcal{I}^{\perp}$  its orthogonal with respect to  $\varrho^*$  and  $A$  the Levi-Civita product associated to  $(\mathfrak{g}^*, [\cdot, \cdot]_r, \varrho^*)$ . Then  $(\mathfrak{g}, r, \varrho)$  is a Riemann-Poisson Lie algebra if and only if:*

- (c<sub>1</sub>)  $[r, r] = 0$ .
- (c<sub>2</sub>) For all  $\alpha \in \mathcal{I}, A_{\alpha} = 0$ .
- (c<sub>3</sub>) For all  $\alpha, \beta, \gamma \in \mathcal{I}^{\perp}, A_{\alpha}\beta \in \mathcal{I}^{\perp}$  and
 
$$r(A_{\alpha}\beta, \gamma) + r(\beta, A_{\alpha}\gamma) = 0.$$

**Proof.** By using the splitting  $\mathfrak{g}^* = \mathcal{I} \oplus \mathcal{I}^{\perp}$ , on can see that the conditions (i) and (ii) in Proposition 2.1 are equivalent to

$$(11) \quad \begin{cases} [r, r] = 0, \\ r(A_{\alpha}\beta, \gamma) = 0, \alpha \in \mathcal{I}, \beta \in \mathcal{I}, \gamma \in \mathcal{I}^{\perp}, \\ r(A_{\alpha}\beta, \gamma) + r(\beta, A_{\alpha}\gamma) = 0, \alpha \in \mathcal{I}, \beta \in \mathcal{I}^{\perp}, \gamma \in \mathcal{I}^{\perp}, \\ r(A_{\alpha}\beta, \gamma) = 0, \alpha \in \mathcal{I}^{\perp}, \beta \in \mathcal{I}, \gamma \in \mathcal{I}^{\perp}, \\ r(A_{\alpha}\beta, \gamma) + r(\beta, A_{\alpha}\gamma) = 0, \alpha \in \mathcal{I}^{\perp}, \beta \in \mathcal{I}^{\perp}, \gamma \in \mathcal{I}^{\perp}. \end{cases}$$

Suppose that the conditions (c<sub>1</sub>)-(c<sub>3</sub>) hold. Then for any  $\alpha \in \mathcal{I}$  and  $\beta \in \mathcal{I}^{\perp}$ ,  $A_{\beta}\alpha = [\beta, \alpha]_r$  and hence  $r_{\#}(A_{\beta}\alpha) = [r_{\#}(\beta), r_{\#}(\alpha)] = 0$  and hence the equations in (11) holds.

Conversely, suppose that (11) holds. Then (c<sub>1</sub>) holds obviously.

For any  $\alpha, \beta \in \mathcal{I}$ , the second equation in (11) is equivalent to  $A_{\alpha}\beta \in \mathcal{I}$  and we have from (7) and (9)  $[\alpha, \beta]_r = 0$  and  $A_{\alpha}\beta \in \mathcal{I}^{\perp}$ . Thus  $A_{\alpha}\beta = 0$ .

Take now  $\alpha \in \mathcal{I}$  and  $\beta \in \mathcal{I}^{\perp}$ . For any  $\gamma \in \mathcal{I}$ ,  $\varrho^*(A_{\alpha}\beta, \gamma) = -\varrho^*(\beta, A_{\alpha}\gamma) = 0$  and hence  $A_{\alpha}\beta \in \mathcal{I}^{\perp}$ . On the other hand,

$$r_{\#}([\alpha, \beta]_r) = r_{\#}(A_{\alpha}\beta) - r_{\#}(A_{\beta}\alpha) \stackrel{(8)}{=} [r_{\#}(\alpha), r_{\#}(\beta)] = 0.$$

So, for any  $\gamma \in \mathcal{I}^{\perp}$ ,

$$\prec \gamma, r_{\#}(A_{\alpha}\beta) \succ = \prec \gamma, r_{\#}(A_{\beta}\alpha) \succ = r(A_{\beta}\alpha, \gamma) \stackrel{(11)}{=} 0.$$

This shows that  $A_{\alpha}\beta \in \mathcal{I}$  and hence  $A_{\alpha}\beta = 0$ . Finally, (c<sub>2</sub>) is true. Now, for any  $\alpha \in \mathcal{I}^{\perp}$ , the fourth equation in (11) implies that  $A_{\alpha}$  leaves invariant  $\mathcal{I}$  and since it is skew-symmetric it leaves invariant  $\mathcal{I}^{\perp}$  and (c<sub>3</sub>) follows. This completes the proof. □

**Proposition 3.2.** *Let  $(\mathfrak{g}, \varrho, r)$  be a Lie algebra endowed with a solution of classical Yang-Baxter equation and a bi-invariant Euclidean product, i.e.,*

$$\varrho(\text{ad}_u v, w) + \varrho(v, \text{ad}_u w) = 0, \quad u, v, w \in \mathfrak{g}.$$

Then  $(\mathfrak{g}, \varrho, r)$  is Riemann-Poisson Lie algebra if and only if  $\text{Im} r_{\#}$  is an abelian subalgebra.

**Proof.** Since  $\varrho$  is bi-invariant, one can see easily that for any  $u \in \mathfrak{g}$ ,  $\text{ad}_u^*$  is skew-symmetric with respect to  $\varrho^*$  and hence the Levi-Civita product  $A$  associated to  $(\mathfrak{g}^*, [\cdot, \cdot]_r, \varrho^*)$  is given by  $A_{\alpha}\beta = \text{ad}_{r_{\#}(\alpha)}^*\beta$ . So,  $(\mathfrak{g}, \varrho, r)$  is Riemann-Poisson Lie algebra if and only if, for any  $\alpha, \beta, \gamma \in \mathfrak{g}^*$ ,

$$\begin{aligned} 0 &= r(\text{ad}_{r_{\#}(\alpha)}^*\beta, \gamma) + r(\beta, \text{ad}_{r_{\#}(\alpha)}^*\gamma) \\ &= \langle \beta, [r_{\#}(\alpha), r_{\#}(\gamma)] \rangle - \langle \gamma, [r_{\#}(\alpha), r_{\#}(\beta)] \rangle \\ &\stackrel{(5)}{=} \langle \alpha, [r_{\#}(\beta), r_{\#}(\gamma)] \rangle \end{aligned}$$

and the result follows. □

Let  $(\mathfrak{g}, [\cdot, \cdot])$  be a Lie algebra,  $r \in \wedge^2 \mathfrak{g}$  and  $\varrho$  a Euclidean product on  $\mathfrak{g}$ . Denote by  $(S, \omega_r)$  the symplectic vector subspace associated to  $r$  and by  $\#: \mathfrak{g}^* \rightarrow \mathfrak{g}$  the isomorphism given by  $\varrho$ . Note that the Euclidean product on  $\mathfrak{g}^*$  is given by  $\varrho^*(\alpha, \beta) = \varrho(\#(\alpha), \#(\beta))$ . We have

$$\mathfrak{g}^* = \mathcal{I} \oplus \mathcal{I}^\perp \quad \text{and} \quad \mathfrak{g} = S \oplus S^\perp,$$

where  $\mathcal{I} = \ker r_{\#}$ . Moreover,  $r_{\#}: \mathcal{I}^\perp \rightarrow S$  is an isomorphism, we denote by  $\tau: S \rightarrow \mathcal{I}^\perp$  its inverse. From the relation

$$\varrho(\#(\alpha), r_{\#}(\beta)) = \langle \alpha, r_{\#}(\beta) \rangle = r(\beta, \alpha),$$

we deduce that  $\#: \mathcal{I} \rightarrow S^\perp$  is an isomorphism and hence  $\#: \mathcal{I}^\perp \rightarrow S$  is also an isomorphism.

Consider the isomorphism  $J: S \rightarrow S$  linking  $\omega_r$  to  $\varrho|_S$ , i.e.,

$$\omega_r(u, v) = \rho(Ju, v), \quad u, v \in S.$$

One can see easily that  $J = -\# \circ \tau$ .

**Theorem 3.1.** *With the notations above,  $(\mathfrak{g}, r, \varrho)$  is a Riemann-Poisson Lie algebra if and only if the following conditions hold:*

- (1)  $(S, \varrho|_S, \omega_r)$  is a Kähler Lie subalgebra, i.e., for all  $s_1, s_2, s_3 \in S$ ,
- (12) 
$$\omega_r(\nabla_{s_1}s_2, s_3) + \omega_r(s_2, \nabla_{s_1}s_3) = 0,$$

where  $\nabla$  is the Levi-Civita product associated to  $(S, [\cdot, \cdot], \varrho|_S)$ .
- (2) for all  $s \in S$  and all  $u, v \in S^\perp$ ,
- (13) 
$$\varrho(\phi_S(s)(u), v) + \varrho(u, \phi_S(s)(v)) = 0,$$

where  $\phi_S: S \rightarrow \text{End}(S^\perp)$ ,  $u \mapsto \text{pr}_{S^\perp} \circ \text{ad}_u$  and  $\text{pr}_{S^\perp}: \mathfrak{g} \rightarrow S^\perp$  is the orthogonal projection.
- (3) For all  $s_1, s_2 \in S$  and all  $u \in S^\perp$ ,
- (14) 
$$\omega_r(\phi_{S^\perp}(u)(s_1), s_2) + \omega_r(s_1, \phi_{S^\perp}(u)(s_2)) = 0,$$

where  $\phi_{S^\perp}: S^\perp \rightarrow \text{End}(S)$ ,  $u \mapsto \text{pr}_S \circ \text{ad}_u$  and  $\text{pr}_S: \mathfrak{g} \rightarrow S$  is the orthogonal projection.

**Proof.** Suppose first that  $(\mathfrak{g}, r, \varrho)$  is a Riemann-Poisson Lie algebra. According to Propositions 3.1 and 2.2, this is equivalent to

$$(15) \quad \begin{cases} (S, \omega_r) \text{ is a symplectic subalgebra,} \\ \forall \alpha \in \mathcal{I}, A_\alpha = 0, \\ \forall \alpha, \beta, \gamma \in \mathcal{I}^\perp, A_\alpha \beta \in \mathcal{I}^\perp \quad \text{and} \quad r(A_\alpha \beta, \gamma) + r(\beta, A_\alpha \gamma) = 0, \end{cases}$$

where  $A$  is the Levi-Civita product of  $(\mathfrak{g}^*, [\cdot, \cdot]_r, \varrho^*)$ .

For  $\alpha, \beta \in \mathcal{I}$  and  $\gamma \in \mathcal{I}^\perp$ ,

$$(16) \quad \begin{aligned} 2\varrho^*(A_\alpha \beta, \gamma) &= \varrho^*([\alpha, \beta]_r, \gamma) + \varrho^*([\gamma, \beta]_r, \alpha) + \varrho^*([\gamma, \alpha]_r, \beta) \\ &= \varrho^*(\text{ad}_{r_\#(\gamma)}^* \beta, \alpha) + \varrho^*(\text{ad}_{r_\#(\gamma)}^* \alpha, \beta) \\ &= -\prec \beta, [r_\#(\gamma), \#(\alpha)] \succ -\prec \alpha, [r_\#(\gamma), \#(\beta)] \succ \\ &= -\varrho(\#(\beta), [r_\#(\gamma), \#(\alpha)]) - \varrho(\#(\alpha), [r_\#(\gamma), \#(\beta)]). \end{aligned}$$

Since  $\#: \mathcal{I} \rightarrow S^\perp$  and  $r_\#: \mathcal{I}^\perp \rightarrow S$  are isomorphisms, we deduce from (16) that  $A_\alpha \beta = 0$  for any  $\alpha, \beta \in \mathcal{I}$  is equivalent to (13).

For  $\alpha \in \mathcal{I}$  and  $\beta, \gamma \in \mathcal{I}^\perp$ ,

$$(17) \quad \begin{aligned} 2\varrho^*(A_\alpha \beta, \gamma) &= \varrho^*([\alpha, \beta]_r, \gamma) + \varrho^*([\gamma, \beta]_r, \alpha) + \varrho^*([\gamma, \alpha]_r, \beta) \\ &= -\varrho^*(\text{ad}_{r_\#(\beta)}^* \alpha, \gamma) - \varrho^*(\text{ad}_{r_\#(\beta)}^* \gamma, \alpha) + \varrho^*(\text{ad}_{r_\#(\gamma)}^* \beta, \alpha) + \varrho^*(\text{ad}_{r_\#(\gamma)}^* \alpha, \beta) \\ &= -\prec \alpha, [r_\#(\beta), \#(\gamma)] \succ + \prec \gamma, [r_\#(\beta), \#(\alpha)] \succ - \prec \beta, [r_\#(\gamma), \#(\alpha)] \succ \\ &\quad - \prec \alpha, [r_\#(\gamma), \#(\beta)] \succ = \varrho(\#(\gamma), [r_\#(\beta), \#(\alpha)]) \\ &\quad - \varrho(\#(\beta), [r_\#(\gamma), \#(\alpha)]) + \prec \alpha, [r_\#(\beta), \#(\gamma)] \succ - \prec \alpha, [r_\#(\gamma), \#(\beta)] \succ \\ &= -\varrho(J \circ r_\#(\gamma), [r_\#(\beta), \#(\alpha)]) + \varrho(J \circ r_\#(\beta), [r_\#(\gamma), \#(\alpha)]) \\ &\quad + \prec \alpha, [r_\#(\beta), \#(\gamma)] \succ - \prec \alpha, [r_\#(\gamma), \#(\beta)] \succ \\ &= -\omega_r(r_\#(\gamma), \text{pr}_S([r_\#(\beta), \#(\alpha)])) - \omega_r(\text{pr}_S([r_\#(\gamma), \#(\alpha)]), r_\#(\beta)) \\ &\quad + \prec \alpha, [r_\#(\beta), \#(\gamma)] \succ - \prec \alpha, [r_\#(\gamma), \#(\beta)] \succ. \end{aligned}$$

Now,  $\#(\beta), \#(\gamma) \in S$  and  $r_\#(\beta), r_\#(\gamma) \in S$  and since  $S$  is a subalgebra we deduce that  $[r_\#(\beta), \#(\gamma)], [r_\#(\gamma), \#(\beta)] \in S$  and hence

$$-\prec \alpha, [r_\#(\beta), \#(\gamma)] \succ = -\prec \alpha, [r_\#(\gamma), \#(\beta)] \succ = 0.$$

We have also  $\#: \mathcal{I} \rightarrow S^\perp$  and  $r_\#: \mathcal{I}^\perp \rightarrow S$  are isomorphisms so that, by virtue of (17),  $A_\alpha \beta = 0$  for any  $\alpha \in \mathcal{I}$  and  $\beta \in \mathcal{I}^\perp$  is equivalent to (14).

On the other hand, for any  $\alpha, \beta, \gamma \in \mathcal{I}^\perp$ , since  $\# = -J \circ r_\#$ , the relation

$$2\varrho^*(A_\alpha \beta, \gamma) = \varrho^*([\alpha, \beta]_r, \gamma) + \varrho^*([\gamma, \beta]_r, \alpha) + \varrho^*([\gamma, \alpha]_r, \beta)$$

can be written

$$\begin{aligned} 2\varrho(J \circ r_\#(A_\alpha \beta), J \circ r_\#(\gamma)) &= \varrho(J \circ r_\#([\alpha, \beta]_r), J \circ r_\#(\gamma)) \\ &\quad + \varrho(J \circ r_\#([\gamma, \beta]_r), J \circ r_\#(\alpha)) \\ &\quad + \varrho(J \circ r_\#([\gamma, \alpha]_r), J \circ r_\#(\beta)). \end{aligned}$$



But  $r_{\#}([\alpha, \beta]_r) = [r_{\#}(\alpha), r_{\#}(\beta)]$  and hence

$$2\langle r_{\#}(A_{\alpha}, \beta), r_{\#}(\gamma) \rangle_J = \langle [r_{\#}(\alpha), r_{\#}(\beta)], r_{\#}(\gamma) \rangle_J + \langle [r_{\#}(\mathfrak{g}), r_{\#}(\beta)], r_{\#}(\alpha) \rangle_J + \langle [r_{\#}(\gamma), r_{\#}(\alpha)], r_{\#}(\beta) \rangle_J,$$

where  $\langle u, v \rangle_J = \varrho(Ju, Jv)$ . This shows that  $r_{\#}(A_{\alpha}\beta) = \nabla_{r_{\#}(\alpha)}r_{\#}(\beta)$  where  $\nabla$  is the Levi-Civita product of  $(S, [ \ , \ ], \langle \ , \ \rangle_J)$  and the third relation in (15) is equivalent to

$$\omega_r(\nabla_u v, w) + \omega_r(v, \nabla_u w) = 0, \quad u, v, w \in S.$$

This is equivalent to  $\nabla_u Jv = J\nabla_u v$ . Let us show that  $\nabla$  is actually the Levi-Civita product of  $(S, [ \ , \ ], \varrho)$ . Indeed, for any  $u, v, w \in S$ ,  $\nabla_u v - \nabla_v u = [u, v]$  and

$$\begin{aligned} \varrho(\nabla_u v, w) + \varrho(\nabla_u w, v) &= \langle J^{-1}\nabla_u v, J^{-1}w \rangle_J + \langle J^{-1}\nabla_u w, J^{-1}v \rangle_J \\ &= \langle \nabla_u J^{-1}v, J^{-1}w \rangle_J + \langle \nabla_u J^{-1}w, J^{-1}v \rangle_J \\ &= 0. \end{aligned}$$

So we have shown the direct part of the theorem. The converse can be deduced easily from the relations we established in the proof of the direct part.  $\square$

**Example 1.** Let  $G$  be a compact connected Lie group,  $\mathfrak{g}$  its Lie algebra and  $T$  an even dimensional torus of  $G$ . Choose a bi-invariant Riemannian metric  $\langle \ , \ \rangle$  on  $G$ , a nondegenerate  $\omega \in \wedge^2 S^*$  where  $S$  is the Lie algebra of  $T$  and put  $\varrho = \langle \ , \ \rangle(e)$ . Let  $r \in \wedge^2 \mathfrak{g}$  be the solution of the classical Yang-Baxter associated to  $(S, \omega)$ . By using either Proposition 3.2 or Theorem 3.1, one can see easily that  $(\mathfrak{g}, \varrho, r)$  is a Riemann-Poisson Lie algebra and hence  $(G, \langle \ , \ \rangle, \pi)$  is a Riemann-Poisson Lie group where  $\pi$  is the left invariant Poisson tensor associated to  $r$ . According to Theorem 1.1, the orbits of the right action of  $T$  on  $G$  defines a Riemannian and Kähler foliation. For instance,  $G = \text{SO}(2n)$ ,  $T = \text{Diagonal}(D_1, \dots, D_n)$  where  $D_i = \begin{pmatrix} \cos(\theta_i) & \sin(\theta_i) \\ -\sin(\theta_i) & \cos(\theta_i) \end{pmatrix}$  and  $\langle \ , \ \rangle = -K$  where  $K$  is the Killing form.

#### 4. CONSTRUCTION OF RIEMANN-POISSON LIE ALGEBRAS

In this section, we give a general method for building Riemann-Poisson Lie algebras and we use it to give all Riemann-Poisson Lie algebras up to dimension 5.

According to Theorem 3.1, to build Riemann-Poisson Lie algebras one needs to solve the following problem.

**Problem 1.** We look for:

- (1) A Kähler Lie algebra  $(\mathfrak{h}, [ \ , \ ], \varrho_{\mathfrak{h}}, \omega)$ ,
- (2) a Euclidean vector space  $(\mathfrak{p}, \varrho_{\mathfrak{p}})$ ,
- (3) a bilinear skew-symmetric map  $[ \ , \ ]_{\mathfrak{p}} : \mathfrak{p} \times \mathfrak{p} \longrightarrow \mathfrak{p}$ ,
- (4) a bilinear skew-symmetric map  $\mu : \mathfrak{p} \times \mathfrak{p} \longrightarrow \mathfrak{h}$ ,
- (5) two linear maps  $\phi_{\mathfrak{p}} : \mathfrak{p} \longrightarrow \text{sp}(\mathfrak{h}, \omega)$  and  $\phi_{\mathfrak{h}} : \mathfrak{h} \longrightarrow \text{so}(\mathfrak{p})$  where  $\text{sp}(\mathfrak{h}, \omega) = \{J : \mathfrak{h} \longrightarrow \mathfrak{h}, J^{\omega} + J = 0\}$  and  $\text{so}(\mathfrak{p}) = \{A : \mathfrak{p} \longrightarrow \mathfrak{p}, A^* + A = 0\}$ ,  $J^{\omega}$  is the adjoint with respect to  $\omega$  and  $A^*$  is the adjoint with respect to  $\varrho_{\mathfrak{p}}$ ,

such that the bracket  $[ , ]$  on  $\mathfrak{g} = \mathfrak{h} \oplus \mathfrak{p}$  given, for any  $a, b \in \mathfrak{p}$  and  $u, v \in \mathfrak{h}$ , by

$$(18) \quad [u, v] = [u, v]_{\mathfrak{h}}, [a, b] = \mu(a, b) + [a, b]_{\mathfrak{p}}, [a, u] = -[u, a] = \phi_{\mathfrak{p}}(a)(u) - \phi_{\mathfrak{h}}(u)(a)$$

is a Lie bracket.

In this case,  $(\mathfrak{g}, [ , ])$  endowed with  $r \in \wedge^2 \mathfrak{g}$  associated to  $(\mathfrak{h}, \omega)$  and the Euclidean product  $\varrho = \varrho_{\mathfrak{h}} \oplus \varrho_{\mathfrak{p}}$  becomes, by virtue of Theorem 3.1, a Riemann-Poisson Lie algebra.

**Proposition 4.1.** *With the data and notations of Problem 1, the bracket given by (18) is a Lie bracket if and only if, for any  $u, v \in \mathfrak{h}$  and  $a, b, c \in \mathfrak{p}$ ,*

$$(19) \quad \begin{cases} \phi_{\mathfrak{p}}(a)([u, v]_{\mathfrak{h}}) = [u, \phi_{\mathfrak{p}}(a)(v)]_{\mathfrak{h}} + [\phi_{\mathfrak{p}}(a)(u), v]_{\mathfrak{h}} + \phi_{\mathfrak{p}}(\phi_{\mathfrak{h}}(v)(a))(u) - \phi_{\mathfrak{p}}(\phi_{\mathfrak{h}}(u)(a))(v), \\ \phi_{\mathfrak{h}}(u)([a, b]_{\mathfrak{p}}) = [a, \phi_{\mathfrak{h}}(u)(b)]_{\mathfrak{p}} + [\phi_{\mathfrak{h}}(u)(a), b]_{\mathfrak{p}} + \phi_{\mathfrak{h}}(\phi_{\mathfrak{p}}(b)(u))(a) - \phi_{\mathfrak{h}}(\phi_{\mathfrak{p}}(a)(u))(b), \\ \phi_{\mathfrak{h}}([u, v]_{\mathfrak{h}}) = [\phi_{\mathfrak{h}}(u), \phi_{\mathfrak{h}}(v)], \\ \phi_{\mathfrak{p}}([a, b]_{\mathfrak{p}})(u) = [\phi_{\mathfrak{p}}(a), \phi_{\mathfrak{p}}(b)](u) + [u, \mu(a, b)]_{\mathfrak{h}} - \mu(a, \phi_{\mathfrak{h}}(u)(b)) - \mu(\phi_{\mathfrak{h}}(u)(a), b), \\ \mathfrak{f}[a, [b, c]_{\mathfrak{p}}]_{\mathfrak{p}} = \mathfrak{f} \phi_{\mathfrak{h}}(\mu(b, c))(a), \\ \mathfrak{f} \phi_{\mathfrak{p}}(a)(\mu(b, c)) = \mathfrak{f} \mu([b, c]_{\mathfrak{p}}, a), \end{cases}$$

where  $\mathfrak{f}$  stands for the circular permutation.

**Proof.** The equations follow from the Jacobi identity applied to  $(a, u, v)$ ,  $(a, b, u)$  and  $(a, b, c)$ . □

We tackle now the task of determining the list of all Riemann-Poisson Lie algebras up to dimension 5. For this purpose, we need to solve Problem 1 in the following four cases: (a)  $\dim \mathfrak{p} = 1$ , (b)  $\dim \mathfrak{h} = 2$  and  $\mathfrak{h}$  non abelian, (c)  $\dim \mathfrak{h} = \dim \mathfrak{p} = 2$  and  $\mathfrak{h}$  abelian, (d)  $\dim \mathfrak{h} = 2$ ,  $\dim \mathfrak{p} = 3$  and  $\mathfrak{h}$  abelian.

It is easy to find the solutions of Problem 1 when  $\dim \mathfrak{p} = 1$  since in this case  $\text{so}(\mathfrak{p}) = 0$  and the three last equations in (19) hold obviously.

**Proposition 4.2.** *If  $\dim \mathfrak{p} = 1$  then the solutions of Problem 1 are a Kähler Lie algebra  $(\mathfrak{h}, \varrho, \omega)$ ,  $\phi_{\mathfrak{h}} = 0$ ,  $[ , ]_{\mathfrak{p}} = 0$ ,  $\mu = 0$  and  $\phi_{\mathfrak{p}}(a) \in \text{sp}(\mathfrak{h}, \omega) \cap \text{Der}(\mathfrak{h})$  where  $a$  is a generator of  $\mathfrak{p}$  and  $\text{Der}(\mathfrak{h})$  the Lie algebra of derivations of  $\mathfrak{h}$ .*

Let us solve Problem 1 when  $\mathfrak{h}$  is 2-dimensional non abelian.

**Proposition 4.3.** *Let  $((\mathfrak{h}, \omega, \varrho_{\mathfrak{h}}), (\mathfrak{p}, [ , ]_{\mathfrak{p}}, \varrho_{\mathfrak{p}}), \mu, \phi_{\mathfrak{h}}, \phi_{\mathfrak{p}})$  be a solution of Problem 1 with  $\mathfrak{h}$  is 2-dimensional non abelian. Then there exists an orthonormal basis  $\mathbb{B} = (e_1, e_2)$  of  $\mathfrak{h}$ ,  $b_0 \in \mathfrak{p}$  and two constants  $\alpha \neq 0$  and  $\beta \neq 0$  such that:*

- (i)  $[e_1, e_2]_{\mathfrak{h}} = \alpha e_1, \omega = \beta e_1^* \wedge e_2^*$ ,
- (ii)  $(\mathfrak{p}, [ , ]_{\mathfrak{p}}, \varrho_{\mathfrak{p}})$  is a Euclidean Lie algebra,
- (iii)  $\phi_{\mathfrak{h}}(e_1) = 0, \phi_{\mathfrak{h}}(e_2) \in \text{Der}(\mathfrak{p}) \cap \text{so}(\mathfrak{p})$  and, for any  $a \in \mathfrak{p}$ ,  $M(\phi_{\mathfrak{p}}(a), \mathbb{B}) = \begin{pmatrix} 0 & \varrho_{\mathfrak{p}}(a, b_0) \\ 0 & 0 \end{pmatrix}$ ,
- (iv) for any  $a, b \in \mathfrak{p}$ ,  $\mu(a, b) = \mu_0(a, b)e_1$  with  $\mu_0$  is a 2-cocycle of  $(\mathfrak{p}, [ , ]_{\mathfrak{p}})$  satisfying

$$(20) \quad \mu_0(a, \phi_{\mathfrak{h}}(e_2)b) + \mu_0(\phi_{\mathfrak{h}}(e_2)a, b) = -\varrho_{\mathfrak{p}}([a, b]_{\mathfrak{p}}, b_0) - \alpha \mu_0(a, b).$$

**Proof.** Note first that from the third relation in (19) we get that  $\phi_{\mathfrak{h}}(\mathfrak{h})$  is a solvable subalgebra of  $\mathfrak{so}(\mathfrak{p})$  and hence must be abelian. Since  $\mathfrak{h}$  is 2-dimensional non abelian then  $\dim \phi_{\mathfrak{h}}(\mathfrak{h}) = 1$  and  $[\mathfrak{h}, \mathfrak{h}] \subset \ker \phi_{\mathfrak{h}}$ . So there exists an orthonormal basis  $(e_1, e_2)$  of  $\mathfrak{h}$  such that  $[e_1, e_2]_{\mathfrak{h}} = \alpha e_1$ ,  $\phi_{\mathfrak{h}}(e_1) = 0$  and  $\omega = \beta e_1^* \wedge e_2^*$ . If we identify the endomorphisms of  $\mathfrak{h}$  with their matrices in the basis  $(e_1, e_2)$ , we get that  $\mathfrak{sp}(\mathfrak{h}, \omega) = \mathfrak{sl}(2, \mathbb{R})$  and there exists  $a_0, b_0, c_0 \in \mathfrak{p}$  such that, for any  $a \in \mathfrak{p}$ ,

$$\phi_{\mathfrak{p}}(a) = \begin{pmatrix} \varrho_{\mathfrak{p}}(a_0, a) & \varrho_{\mathfrak{p}}(b_0, a) \\ \varrho_{\mathfrak{p}}(c_0, a) & -\varrho_{\mathfrak{p}}(a_0, a) \end{pmatrix}.$$

The first equation in (19) is equivalent to

$$\begin{aligned} \alpha(\varrho_{\mathfrak{p}}(a_0, a)e_1 + \varrho_{\mathfrak{p}}(c_0, a)e_2) &= -\alpha\varrho_{\mathfrak{p}}(a_0, a)e_1 + \alpha\varrho_{\mathfrak{p}}(a_0, a)e_1 \\ &\quad + \varrho_{\mathfrak{p}}(a_0, \phi_{\mathfrak{h}}(e_2)(a))e_1 + \varrho_{\mathfrak{p}}(c_0, \phi_{\mathfrak{h}}(e_2)(a))e_2, \end{aligned}$$

for any  $a \in \mathfrak{p}$ . Since  $\phi_{\mathfrak{h}}(e_2)$  is sekw-symmetric, this is equivalent to

$$\phi_{\mathfrak{h}}(e_2)(a_0) = -\alpha a_0 \quad \text{and} \quad \phi_{\mathfrak{h}}(e_2)(c_0) = -\alpha c_0.$$

This implies that  $a_0 = c_0 = 0$ . The second equation in (19) implies that  $\phi_{\mathfrak{h}}(e_2)$  is a derivation of  $[\cdot, \cdot]_{\mathfrak{p}}$ . If we take  $u = e_1$  in the forth equation in (19), we get that  $[e_1, \mu(a, b)] = 0$ , for any  $a, b \in \mathfrak{p}$  and hence  $\mu(a, b) = \mu_0(a, b)e_1$ . If we take  $u = e_2$  in the forth equation in (19) we get (20). The two last equations are equivalent to  $[\cdot, \cdot]_{\mathfrak{p}}$  is a Lie bracket and  $\mu_0$  is 2-cocycle of  $(\mathfrak{p}, [\cdot, \cdot]_{\mathfrak{p}})$ .  $\square$

The following proposition gives the solutions of Problem 1 when  $\mathfrak{h}$  is 2-dimensional abelian and  $\dim \mathfrak{p} = 2$ .

**Proposition 4.4.** *Let  $((\mathfrak{h}, \omega, \varrho_{\mathfrak{h}}), (\mathfrak{p}, [\cdot, \cdot]_{\mathfrak{p}}, \varrho_{\mathfrak{p}}), \mu, \phi_{\mathfrak{h}}, \phi_{\mathfrak{p}})$  be a solution of Problem 1 with  $\mathfrak{h}$  is 2-dimensional abelian and  $\dim \mathfrak{p} = 2$ . Then one of the following situations occurs:*

- (1)  $\phi_{\mathfrak{h}} = 0$ ,  $(\mathfrak{p}, [\cdot, \cdot]_{\mathfrak{p}}, \varrho_{\mathfrak{p}})$  is a 2-dimensional Euclidean Lie algebra, there exists  $a_0 \in \mathfrak{p}$  and  $D \in \mathfrak{sp}(\mathfrak{h}, \omega)$  such that, for any  $a \in \mathfrak{p}$ ,  $\phi_{\mathfrak{p}}(a) = \varrho_{\mathfrak{p}}(a_0, a)D$  and there is no restriction on  $\mu$ . Moreover,  $a_0 \in [\mathfrak{p}, \mathfrak{p}]_{\mathfrak{p}}^{\perp}$  if  $D \neq 0$ .
- (2)  $\phi_{\mathfrak{h}} = 0$ ,  $(\mathfrak{p}, [\cdot, \cdot]_{\mathfrak{p}}, \varrho_{\mathfrak{p}})$  is a 2-dimensional non abelian Euclidean Lie algebra,  $\phi_{\mathfrak{p}}$  identifies  $\mathfrak{p}$  to a two dimensional subalgebra of  $\mathfrak{sp}(\mathfrak{h}, \omega)$  and there is no restriction on  $\mu$ .
- (3)  $(\mathfrak{p}, [\cdot, \cdot]_{\mathfrak{p}}, \varrho_{\mathfrak{p}})$  is a Euclidean abelian Lie algebra and there exists an orthonormal basis  $\mathbb{B} = (e_1, e_2)$  of  $\mathfrak{h}$  and  $b_0 \in \mathfrak{p}$  such that  $\omega = \alpha e_1^* \wedge e_2^*$ ,  $\phi_{\mathfrak{h}}(e_1) = 0$ ,  $\phi_{\mathfrak{h}}(e_2) \neq 0$  and, for any  $a \in \mathfrak{p}$ ,  $M(\phi_{\mathfrak{p}}(a), \mathbb{B}) = \begin{pmatrix} 0 & \varrho_{\mathfrak{p}}(b_0, a) \\ 0 & 0 \end{pmatrix}$  and there is no restriction on  $\mu$ .

**Proof.** Note first that since  $\dim \mathfrak{p} = 2$  the last two equations in (19) hold obviously and  $(\mathfrak{p}, [\cdot, \cdot]_{\mathfrak{p}})$  is a Lie algebra. We distinguish two cases:

- (i)  $\phi_{\mathfrak{h}} = 0$ . Then (19) is equivalent to  $\phi_{\mathfrak{p}}$  is a representation of  $\mathfrak{p}$  in  $\mathfrak{sp}(\mathfrak{h}, \omega) \simeq \mathfrak{sl}(2, \mathbb{R})$ . Since  $\mathfrak{sl}(2, \mathbb{R})$  doesn't contain any abelian two dimensional subalgebra, if  $\mathfrak{p}$  is an abelian Lie algebra then  $\dim \phi_{\mathfrak{p}}(\mathfrak{p}) \leq 1$  and the first situation occurs. If  $\mathfrak{p}$  is not abelian then the first or the second situation occurs depending on  $\dim \phi_{\mathfrak{p}}(\mathfrak{p})$ .

(ii)  $\phi_{\mathfrak{h}} \neq 0$ . Since  $\dim \text{so}(\mathfrak{p}) = 1$  there exists an orthonormal basis  $\mathbb{B} = (e_1, e_2)$  of  $\mathfrak{h}$  such that  $\phi_{\mathfrak{h}}(e_1) = 0$  and  $\phi_{\mathfrak{h}}(e_2) \neq 0$ . We have  $\text{sp}(\mathfrak{h}, \omega) = \text{sl}(2, \mathbb{R})$  and hence, for any  $a \in \mathfrak{p}$ ,  $M(\phi_{\mathfrak{p}}(a), \mathbb{B}) = \begin{pmatrix} \varrho_{\mathfrak{p}}(a_0, a) & \varrho_{\mathfrak{p}}(b_0, a) \\ \varrho_{\mathfrak{p}}(c_0, a) & -\varrho_{\mathfrak{p}}(a_0, a) \end{pmatrix}$ . Choose an orthonormal basis  $(a_1, a_2)$  of  $\mathfrak{p}$ . Then there exists  $\lambda \neq 0$  such that  $\phi_{\mathfrak{h}}(e_2)(a_1) = \lambda a_2$  and  $\phi_{\mathfrak{h}}(e_2)(a_2) = -\lambda a_1$ .

The first equation in (19) is equivalent to

$$\phi_{\mathfrak{p}}(\phi_{\mathfrak{h}}(e_2)(a))(e_1) = 0, \quad a \in \mathfrak{p}.$$

This is equivalent to

$$\phi_{\mathfrak{p}}(a_1)(e_1) = \phi_{\mathfrak{p}}(a_2)(e_1) = 0.$$

Then  $a_0 = c_0 = 0$  and hence  $\phi_{\mathfrak{p}}(a) = \begin{pmatrix} 0 & \varrho_{\mathfrak{p}}(b_0, a) \\ 0 & 0 \end{pmatrix}$ . The second equation in (19) gives

$$\begin{aligned} \phi_{\mathfrak{h}}(e_2)([a_1, a_2]_{\mathfrak{p}}) &= [a_1, \phi_{\mathfrak{h}}(e_2)(a_2)]_{\mathfrak{p}} + \phi_{\mathfrak{h}}(e_2)(a_1), a_2]_{\mathfrak{p}} \\ &\quad + \phi_{\mathfrak{h}}(\phi_{\mathfrak{p}}(a_2)(e_2))(a_2) - \phi_{\mathfrak{h}}(\phi_{\mathfrak{p}}(a_1)(e_2))(a_2), \end{aligned}$$

and hence  $\phi_{\mathfrak{h}}(e_2)([a_1, a_2]_{\mathfrak{p}}) = 0$ . Thus  $[a_1, a_2]_{\mathfrak{p}} = 0$ . All the other equations in (19) hold obviously. □

To tackle the last case, we need the determination of 2-dimensional subalgebras of  $\text{sl}(2, \mathbb{R})$ .

**Proposition 4.5.** *The 2-dimensional subalgebras of  $\text{sl}(2, \mathbb{R})$  are*

$$\begin{aligned} \mathfrak{g}_1 &= \left\{ \begin{pmatrix} \alpha & \beta \\ 0 & -\alpha \end{pmatrix}, \alpha, \beta \in \mathbb{R} \right\}, \quad \mathfrak{g}_2 = \left\{ \begin{pmatrix} \alpha & 0 \\ \beta & -\alpha \end{pmatrix}, \alpha, \beta \in \mathbb{R} \right\}, \\ \mathfrak{g}_x &= \left\{ \begin{pmatrix} \alpha & \frac{2\beta - \alpha}{x} \\ (\alpha + 2\beta)x & -\alpha \end{pmatrix}, \alpha, \beta \in \mathbb{R} \right\} \end{aligned}$$

where  $x \in \mathbb{R} \setminus \{0\}$ . Moreover,  $\mathfrak{g}_x = \mathfrak{g}_y$  if and only if  $x = y$ .

**Proof.** Let  $\mathfrak{g}$  be a 2-dimensional subalgebra of  $\text{sl}(2, \mathbb{R})$ . We consider the basis  $\mathbb{B} = (h, e, f)$  of  $\text{sl}(2, \mathbb{R})$  given by

$$e = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \quad f = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix} \quad \text{and} \quad h = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}.$$

Then

$$[h, e] = 2e, \quad [h, f] = -2f \quad \text{and} \quad [e, f] = h.$$

If  $h \in \mathfrak{g}$  then  $\text{ad}_h$  leaves  $\mathfrak{g}$  invariant. But  $\text{ad}_h$  has three eigenvalues  $(0, 2, -2)$  with the associated eigenvectors  $(h, e, f)$  and hence its restriction to  $\mathfrak{g}$  has  $(0, 2)$  or  $(0, -2)$  as eigenvalues. Thus  $\mathfrak{g} = \mathfrak{g}_1$  or  $\mathfrak{g} = \mathfrak{g}_2$ .

Suppose now that  $h \notin \mathfrak{g}$ . By using the fact that  $\text{sl}(2, \mathbb{R})$  is unimodular, i.e., for any  $w \in \text{sl}(2, \mathbb{R})$   $\text{tr}(\text{ad}_w) = 0$ , we can choose a basis  $(u, v)$  of  $\mathfrak{g}$  such that  $(u, v, h)$  is a basis of  $\text{sl}(2, \mathbb{R})$  and

$$[u, v] = u, \quad [h, u] = au + v \quad \text{and} \quad [h, v] = du - av - h.$$

If  $(x_1, x_2, x_3)$  and  $(y_1, y_2, y_3)$  are the coordinates of  $u$  and  $v$  in  $\mathbb{B}$ , the brackets above gives

$$\begin{cases} -2(x_1y_3 - x_3y_1) - x_1 = 0, \\ 2(x_2y_3 - x_3y_2) - x_2 = 0, \\ x_1y_2 - x_2y_1 - x_3 = 0, \end{cases} \quad \begin{cases} y_1 = (2 - a)x_1, \\ y_2 = -(a + 2)x_2, \\ y_3 = -ax_3, \end{cases} \quad \text{and} \quad \begin{cases} dx_1 = (a + 2)y_1, \\ dx_2 = (a - 2)y_2, \\ dx_3 = ay_3 + 1. \end{cases}$$

Note first that if  $x_1 = 0$  then  $(x_2, x_3) = (0, 0)$  which impossible so we must have  $x_1 \neq 0$  and hence  $d = 4 - a^2$ . If we replace in the third equation in the second system and the last equation, we get  $x_3 = \frac{1}{4}$  and  $y_3 = -\frac{a}{4}$ . The third equation in the first system gives  $x_2 = -\frac{1}{16x_1}$  and hence  $y_1 = (2 - a)x_1$  and  $y_2 = \frac{(a+2)}{16x_1}$ . Thus

$$\begin{aligned} \mathfrak{g} &= \text{span} \left\{ \begin{pmatrix} \frac{1}{4} & -\frac{1}{16x_1} \\ x_1 & -\frac{1}{4} \end{pmatrix}, \begin{pmatrix} -\frac{a}{4} & \frac{(a+2)}{16x_1} \\ (2-a)x_1 & \frac{a}{4} \end{pmatrix} \right\} \\ &= \text{span} \left\{ \begin{pmatrix} 1 & -\frac{1}{x} \\ x & -1 \end{pmatrix}, \begin{pmatrix} -a & \frac{(a+2)}{x} \\ (2-a)x & a \end{pmatrix} \right\}; \quad x = 4x_1. \end{aligned}$$

But

$$\begin{pmatrix} 0 & \frac{2}{x} \\ 2x & 0 \end{pmatrix} = a \begin{pmatrix} 1 & -\frac{1}{x} \\ x & -1 \end{pmatrix} + \begin{pmatrix} -a & \frac{(a+2)}{x} \\ (2-a)x & a \end{pmatrix}$$

and hence

$$\mathfrak{g} = \text{span} \left\{ \begin{pmatrix} 1 & -\frac{1}{x} \\ x & -1 \end{pmatrix}, \begin{pmatrix} 0 & \frac{2}{x} \\ 2x & 0 \end{pmatrix} \right\} = \mathfrak{g}_x.$$

One can check easily that  $\mathfrak{g}_x = \mathfrak{g}_y$  if and only if  $x = y$ . This completes the proof.  $\square$

The following two propositions give the solutions of Problem 1 when  $\mathfrak{h}$  is 2-dimensional abelian and  $\dim \mathfrak{p} = 3$ .

**Proposition 4.6.** *Let  $((\mathfrak{h}, \omega, \varrho_{\mathfrak{h}}), (\mathfrak{p}, [\cdot, \cdot]_{\mathfrak{p}}, \varrho_{\mathfrak{p}}), \mu, \phi_{\mathfrak{h}}, \phi_{\mathfrak{p}})$  be a solution of Problem 1 with  $\mathfrak{h}$  is 2-dimensional abelian and  $\dim \mathfrak{p} = 3$  and  $\phi_{\mathfrak{h}} = 0$ . Then one of the following situations occurs:*

- (i)  $(\mathfrak{p}, [\cdot, \cdot]_{\mathfrak{p}}, \varrho_{\mathfrak{p}})$  is 3-dimensional Euclidean Lie algebra,  $\phi_{\mathfrak{p}} = 0$  and  $\mu$  is 2-cocycle for the trivial representation.
- (ii)  $\phi_{\mathfrak{p}}$  is an isomorphism of Lie algebras between  $(\mathfrak{p}, [\cdot, \cdot]_{\mathfrak{p}})$  and  $\mathfrak{sl}(2, \mathbb{R})$  and there exists an endomorphism  $L: \mathfrak{p} \rightarrow \mathfrak{h}$  such that for any  $a, b \in \mathfrak{p}$ ,

$$\mu(a, b) = \phi_{\mathfrak{p}}(a)(L(b)) - \phi_{\mathfrak{p}}(b)(L(a)) - L([a, b]_{\mathfrak{p}}).$$

- (iii) There exists a basis  $\mathbb{B}_{\mathfrak{p}} = (a_1, a_2, a_3)$  of  $\mathfrak{p}$ ,  $\alpha \neq 0, \beta \neq 0, \gamma, \tau \in \mathbb{R}$  such that  $[\cdot, \cdot]_{\mathfrak{p}}$  has one of the two following forms

$$\begin{cases} [a_1, a_2]_{\mathfrak{p}} = 0, [a_1, a_3]_{\mathfrak{p}} = \beta a_1, \\ [a_2, a_3]_{\mathfrak{p}} = \gamma a_1 + \alpha a_2, \alpha \neq 0, \beta \neq 0 \\ M(\varrho_{\mathfrak{p}}, \mathbb{B}_{\mathfrak{p}}) = I_3 \end{cases} \quad \text{or} \quad \begin{cases} [a_1, a_2]_{\mathfrak{p}} = [a_1, a_3]_{\mathfrak{p}} = 0, \\ [a_2, a_3]_{\mathfrak{p}} = \alpha a_2, \alpha \neq 0, \\ M(\varrho_{\mathfrak{p}}, \mathbb{B}_{\mathfrak{p}}) = \begin{pmatrix} 1 & \tau & 0 \\ \tau & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}. \end{cases}$$

In both cases, there exists an orthonormal basis  $\mathbb{B}_{\mathfrak{h}} = (e_1, e_2)$  of  $\mathfrak{h}$ ,  $x \neq 0$ ,  $u \neq 0$  and  $v \in \mathbb{R}$  such that  $\phi_{\mathfrak{p}}$  has one of the following forms

$$\begin{cases} M(\phi_{\mathfrak{p}}(a_2), \mathbb{B}_{\mathfrak{h}}) = \begin{pmatrix} 0 & u \\ 0 & 0 \end{pmatrix}, \\ M(\phi_{\mathfrak{p}}(a_3), \mathbb{B}_{\mathfrak{h}}) = \begin{pmatrix} -\frac{\alpha}{2} & v \\ 0 & \frac{\alpha}{2} \end{pmatrix}, \\ \phi_{\mathfrak{p}}(a_1) = 0, \end{cases} \quad \text{or} \quad \begin{cases} M(\phi_{\mathfrak{p}}(a_2), \mathbb{B}_{\mathfrak{h}}) = \begin{pmatrix} 0 & 0 \\ u & 0 \end{pmatrix}, \\ M(\phi_{\mathfrak{p}}(a_3), \mathbb{B}_{\mathfrak{h}}) = \begin{pmatrix} \frac{\alpha}{2} & 0 \\ v & -\frac{\alpha}{2} \end{pmatrix}, \\ \phi_{\mathfrak{p}}(a_1) = 0, \end{cases}$$

$$\begin{cases} M(\phi_{\mathfrak{p}}(a_2), \mathbb{B}_{\mathfrak{h}}) = \begin{pmatrix} u & -\frac{u}{x} \\ ux & -u \end{pmatrix}, \\ M(\phi_{\mathfrak{p}}(a_3), \mathbb{B}_{\mathfrak{h}}) = \begin{pmatrix} v & -\frac{2v+\alpha}{2x} \\ \frac{2v-\alpha}{2}x & -v \end{pmatrix}, \\ \phi_{\mathfrak{p}}(a_1) = 0. \end{cases}$$

Moreover,  $\mu$  is a 2-cocycle for  $(\mathfrak{p}, [\cdot, \cdot]_{\mathfrak{p}}, \phi_{\mathfrak{p}})$ .

(iv) There exists an orthonormal basis  $\mathbb{B} = (a_1, a_2, a_3)$  of  $\mathfrak{p}$  such that  $\phi_{\mathfrak{p}}(a_1) = \phi_{\mathfrak{p}}(a_2) = 0$ ,  $\phi_{\mathfrak{p}}(a_3)$  is a non zero element of  $\text{sp}(\mathfrak{h}, \omega)$  and

$$\begin{cases} [a_1, a_2]_{\mathfrak{p}} = 0, [a_1, a_3]_{\mathfrak{p}} = \beta a_1 + \rho a_2, \\ [a_2, a_3]_{\mathfrak{p}} = \gamma a_1 + \alpha a_2, \end{cases} \quad \text{or} \quad \begin{cases} [a_1, a_2]_{\mathfrak{p}} = \alpha a_2, [a_1, a_3]_{\mathfrak{p}} = \rho a_2, \\ [a_2, a_3]_{\mathfrak{p}} = \gamma a_2, \alpha \neq 0. \end{cases}$$

Moreover,  $\mu$  is a 2-cocycle for  $(\mathfrak{p}, [\cdot, \cdot]_{\mathfrak{p}}, \phi_{\mathfrak{p}})$ .

**Proof.** In this case, (19) is equivalent to  $(\mathfrak{p}, [\cdot, \cdot]_{\mathfrak{p}})$  is a Lie algebra and  $\phi_{\mathfrak{p}}$  is a representation and  $\mu$  is a 2-cocycle of  $(\mathfrak{p}, [\cdot, \cdot]_{\mathfrak{p}}, \phi_{\mathfrak{p}})$ .

We distinguish four cases:

- (1)  $\phi_{\mathfrak{p}} = 0$  and the case (i) occurs.
- (2)  $\dim \phi_{\mathfrak{p}}(\mathfrak{p}) = 3$  and hence  $\mathfrak{p}$  is isomorphic to  $\text{sp}(\mathfrak{h}, \omega) \simeq \text{sl}(2, \mathbb{R})$  and hence  $\mu$  is a coboundary. Thus (ii) occurs.
- (3)  $\dim \phi_{\mathfrak{p}}(\mathfrak{p}) = 2$  then  $\ker \phi_{\mathfrak{p}}$  is a one dimensional ideal of  $\mathfrak{p}$ . But  $\phi_{\mathfrak{p}}(\mathfrak{p})$  is a 2-dimensional subalgebra of  $\text{sp}(\mathfrak{h}, \omega) \simeq \text{sl}(2, \mathbb{R})$ , therefore it is non abelian so  $\mathfrak{p}/\ker \mathfrak{p}$  is non abelian.

If  $\ker \mathfrak{p} \subset [\mathfrak{p}, \mathfrak{p}]_{\mathfrak{p}}$  then  $\dim[\mathfrak{p}, \mathfrak{p}]_{\mathfrak{p}} = 2$  so there exists an orthonormal basis  $(a_1, a_2, a_3)$  of  $\mathfrak{p}$  such that  $a_1 \in \ker \mathfrak{p}$  and

$$[a_1, a_2]_{\mathfrak{p}} = \xi a_1, [a_1, a_3]_{\mathfrak{p}} = \beta a_1 \quad \text{and} \quad [a_2, a_3]_{\mathfrak{p}} = \gamma a_1 + \alpha a_2, \alpha \neq 0, \beta \neq 0$$

and we must have  $\xi = 0$  in order to have the Jacobi identity.

If  $\ker \mathfrak{p} \not\subset [\mathfrak{p}, \mathfrak{p}]_{\mathfrak{p}}$  then  $\ker \mathfrak{p} \subset Z(\mathfrak{p})$  and  $\dim[\mathfrak{p}, \mathfrak{p}] = 1$ . Then there exists a basis  $(a_1, a_2, a_3)$  of  $\mathfrak{p}$  such that  $a_1 \in \ker \mathfrak{p}$ ,  $a_2 \in [\mathfrak{p}, \mathfrak{p}]$ ,  $a_3 \in \{a_1, a_2\}^{\perp}$  and

$$[a_2, a_3]_{\mathfrak{p}} = \alpha a_2, [a_3, a_1]_{\mathfrak{p}} = [a_1, a_2]_{\mathfrak{p}} = 0, \alpha \neq 0.$$

The matrix of  $\varrho_{\mathfrak{p}}$  in  $(a_1, a_2, a_3)$  is given by

$$\begin{pmatrix} 1 & \tau & 0 \\ \tau & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}.$$

We choose an orthonormal basis  $(e_1, e_2)$  of  $\mathfrak{h}$  and identify  $\text{sp}(\mathfrak{h}, \omega)$  to  $\text{sl}(2, \mathbb{R})$ . Now  $\phi_{\mathfrak{p}}(\mathfrak{p}) = \{\phi_p(a_2), \phi_p(a_3)\}$  is a subalgebra of  $\text{sl}(2, \mathbb{R})$  and, according to Proposition 4.5,  $\phi_{\mathfrak{p}}(\mathfrak{p}) = \mathfrak{g}_1, \mathfrak{g}_2$  or  $\mathfrak{g}_x$ . But

$$[\mathfrak{g}_1, \mathfrak{g}_1] = \mathbb{R}e, [\mathfrak{g}_2, \mathfrak{g}_2] = \mathbb{R}f \quad \text{and} \quad [\mathfrak{g}_x, \mathfrak{g}_x] = \left\{ \begin{pmatrix} u & -\frac{u}{x} \\ ux & -u \end{pmatrix} \right\}.$$

So in order for  $\phi_{\mathfrak{p}}$  to be a representation we must have

$$\phi_{\mathfrak{p}}(a_2) = \begin{pmatrix} 0 & u \\ 0 & 0 \end{pmatrix}, \quad \text{and} \quad \phi_{\mathfrak{p}}(a_3) = \begin{pmatrix} -\frac{\alpha}{2} & v \\ 0 & \frac{\alpha}{2} \end{pmatrix} \quad \text{and} \quad \phi_{\mathfrak{p}}(a_1) = 0,$$

$$\phi_{\mathfrak{p}}(a_2) = \begin{pmatrix} 0 & 0 \\ u & 0 \end{pmatrix}, \quad \phi_{\mathfrak{p}}(a_3) = \begin{pmatrix} \frac{\alpha}{2} & 0 \\ v & -\frac{\alpha}{2} \end{pmatrix} \quad \text{and} \quad \phi_{\mathfrak{p}}(a_1) = 0,$$

or

$$\phi_{\mathfrak{p}}(a_2) = \begin{pmatrix} u & -\frac{u}{x} \\ ux & -u \end{pmatrix}, \quad \phi_{\mathfrak{p}}(a_3) = \begin{pmatrix} p & -\frac{2p+\alpha}{2x} \\ \frac{2p-\alpha}{2}x & -p \end{pmatrix} \quad \text{and} \quad \phi_{\mathfrak{p}}(a_1) = 0.$$

(4)  $\dim \phi_{\mathfrak{p}}(\mathfrak{p}) = 1$  then  $\ker \phi_{\mathfrak{p}}$  is a two dimensional ideal of  $\mathfrak{p}$ . Then there exists an orthonormal basis  $(a_1, a_2, a_3)$  of  $\mathfrak{p}$  such that

$$[a_1, a_2]_{\mathfrak{p}} = \alpha a_2, \quad [a_3, a_1]_{\mathfrak{p}} = pa_1 + qa_2 \quad \text{and} \quad [a_3, a_2]_{\mathfrak{p}} = ra_1 + sa_2.$$

The Jacobi identity gives  $\alpha = 0$  or  $(p, r) = (0, 0)$ . We take  $\phi_{\mathfrak{p}}(a_1) = \phi_{\mathfrak{p}}(a_2) = 0$  and  $\phi_{\mathfrak{p}}(a_3) \in \text{sl}(2, \mathbb{R})$ . □

**Proposition 4.7.** *Let  $((\mathfrak{h}, \omega, \varrho_{\mathfrak{h}}), (\mathfrak{p}, [\cdot, \cdot]_{\mathfrak{p}}, \varrho_{\mathfrak{p}}), \mu, \phi_{\mathfrak{h}}, \phi_{\mathfrak{p}})$  be a solution of Problem 1 with  $\mathfrak{h}$  is 2-dimensional abelian,  $\dim \mathfrak{p} = 3$  and  $\phi_{\mathfrak{h}} \neq 0$ . Then there exists an orthonormal basis  $(e_1, e_2)$  of  $\mathfrak{h}$ , an orthonormal basis  $(a_1, a_2, a_3)$  of  $\mathfrak{p}$ ,  $\lambda > 0$ ,  $\alpha, p, q, \mu_1, \mu_2, \mu_3 \in \mathbb{R}$  such that*

$$\phi_{\mathfrak{h}}(e_1) = 0, \quad \phi_{\mathfrak{h}}(e_2)(a_1) = \lambda a_2, \quad \phi_{\mathfrak{h}}(e_2)(a_2) = -\lambda a_1 \quad \text{and} \quad \phi_{\mathfrak{h}}(e_2)(a_3) = 0,$$

$$[a_1, a_2]_{\mathfrak{p}} = \alpha a_3, \quad [a_1, a_3]_{\mathfrak{p}} = pa_1 + qa_2, \quad [a_2, a_3]_{\mathfrak{p}} = -qa_1 + pa_2 \quad \text{and}$$

$$\phi_{\mathfrak{p}}(a_i) = \begin{pmatrix} 0 & \mu_i \\ 0 & 0 \end{pmatrix}, \quad i = 1, 2, 3$$

and one of the following situations occurs:

(i)  $p \neq 0, \alpha = 0$  and

$$\mu(a_1, a_2) = 0, \quad \mu(a_2, a_3) = -\lambda^{-1}(p\mu_1 + q\mu_2)e_1 \quad \text{and} \quad \mu(a_1, a_3) = \lambda^{-1}(-q\mu_1 + p\mu_2)e_1.$$

(ii)  $p = 0, \mu_3 \neq 0, \alpha = 0$  and

$$\mu(a_1, a_2) = ce_1, \quad \mu(a_2, a_3) = -\lambda^{-1}(p\mu_1 + q\mu_2)e_1 \quad \text{and} \quad \mu(a_1, a_3) = \lambda^{-1}(-q\mu_1 + p\mu_2)e_1.$$

(iii)  $p = 0, \mu_3 = 0$  and

$$\begin{aligned} \mu(a_1, a_2) &= c_1 e_1 + c_2 e_2, \mu(a_2, a_3) = -\lambda^{-1}(p\mu_1 + q\mu_2)e_1 \quad \text{and} \\ \mu(a_1, a_3) &= \lambda^{-1}(-q\mu_1 + p\mu_2)e_1. \end{aligned}$$

**Proof.** Since  $\phi_{\mathfrak{h}} \neq 0$  then  $\phi_{\mathfrak{h}}(\mathfrak{h})$  is a non trivial abelian subalgebra of  $\mathfrak{so}(\mathfrak{p})$  and hence it must be one dimensional. Then there exists an orthonormal basis  $(e_1, e_2)$  of  $\mathfrak{h}$  and an orthonormal basis  $(a_1, a_2, a_3)$  of  $\mathfrak{p}$  and  $\lambda > 0$  such that  $\phi_{\mathfrak{h}}(e_1) = 0$  and

$$\phi_{\mathfrak{h}}(e_2)(a_1) = \lambda a_2, \phi_{\mathfrak{h}}(e_2)(a_2) = -\lambda a_1 \quad \text{and} \quad \phi_{\mathfrak{h}}(e_2)(a_3) = 0.$$

The first equation in (19) is equivalent to

$$\phi_{\mathfrak{p}}(\phi_{\mathfrak{h}}(e_2)(a))(e_1) = 0, \quad a \in \mathfrak{p}.$$

This is equivalent to

$$\phi_{\mathfrak{p}}(a_1)(e_1) = \phi_{\mathfrak{p}}(a_2)(e_1) = 0.$$

Thus  $\phi_{\mathfrak{p}}(a_i) = \begin{pmatrix} 0 & \mu_i \\ 0 & 0 \end{pmatrix}$  for  $i = 1, 2$  and  $\phi_{\mathfrak{p}}(a_3) = \begin{pmatrix} u & v \\ w & -u \end{pmatrix}$ . Consider now the second equation in (19)

$$\phi_{\mathfrak{h}}(u)([a, b]_{\mathfrak{p}}) = [a, \phi_{\mathfrak{h}}(u)(b)]_{\mathfrak{p}} + [\phi_{\mathfrak{h}}(u)(a), b]_{\mathfrak{p}} + \phi_{\mathfrak{h}}(\phi_{\mathfrak{p}}(b)(u))(a) - \phi_{\mathfrak{h}}(\phi_{\mathfrak{p}}(a)(u))(b).$$

This equation is obviously true when  $u = e_1$  and  $(a, b) = (a_1, a_2)$ . For  $u = e_1$  and  $(a, b) = (a_1, a_3)$ , we get

$$\phi_{\mathfrak{h}}(\phi_{\mathfrak{p}}(a_3)(e_1))(a_1) = 0$$

and hence  $w = 0$ .

For  $u = e_2$  and  $(a, b) = (a_1, a_2)$ , we get  $\phi_{\mathfrak{h}}(e_2)([a_1, a_2]_{\mathfrak{p}}) = 0$  and hence  $[a_1, a_2]_{\mathfrak{p}} = \alpha a_3$ .

For  $u = e_2$  and  $(a, b) = (a_1, a_3)$  or  $(a, b) = (a_2, a_3)$ , we get

$$\phi_{\mathfrak{h}}(e_2)([a_1, a_3]_{\mathfrak{p}}) = \lambda[a_2, a_3]_{\mathfrak{p}} - \lambda u a_2 \quad \text{and} \quad \phi_{\mathfrak{h}}(e_2)([a_2, a_3]_{\mathfrak{p}}) = -\lambda[a_1, a_3]_{\mathfrak{p}} + \lambda u a_1.$$

This implies that  $[a_1, a_3]_{\mathfrak{p}}, [a_2, a_3]_{\mathfrak{p}} \in \text{span}\{a_1, a_2\}$  and hence

$$[a_1, a_3]_{\mathfrak{p}} = p a_1 + q a_2 \quad \text{and} \quad [a_2, a_3]_{\mathfrak{p}} = r a_1 + s a_2.$$

So

$$\begin{cases} \lambda(p a_2 - q a_1) = \lambda(r a_1 + s a_2 - u a_2), \\ \lambda(r a_2 - s a_1) = -\lambda(p a_1 + q a_2 - u a_1). \end{cases}$$

This is equivalent to

$$u = 0, p = s \quad \text{and} \quad r = -q.$$

To summarize, we get

$$[a_1, a_2]_{\mathfrak{p}} = \alpha a_3, [a_1, a_3]_{\mathfrak{p}} = p a_1 + q a_2, [a_2, a_3]_{\mathfrak{p}} = -q a_1 + p a_2 \quad \text{and}$$

$$\phi_{\mathfrak{p}}(a_i) = \begin{pmatrix} 0 & \mu_i \\ 0 & 0 \end{pmatrix}.$$

Let consider now the fourth equation in (19)

$$\phi_{\mathfrak{p}}([a, b]_{\mathfrak{p}})(u) = [\phi_{\mathfrak{p}}(a), \phi_{\mathfrak{p}}(b)](u) + [u, \mu(a, b)]_{\mathfrak{h}} - \mu(a, \phi_{\mathfrak{h}}(u)(b)) - \mu(\phi_{\mathfrak{h}}(u)(a), b).$$

This equation is obviously true for  $u = e_1$ .



For  $u = e_2$  and  $(a, b) = (a_1, a_2)$ ,  $(a, b) = (a_1, a_3)$  or  $(a, b) = (a_2, a_3)$ , we get

$$\begin{cases} \alpha\mu_3 = 0, \\ (p\mu_1 + q\mu_2)e_1 = -\lambda\mu(a_2, a_3), \\ (-q\mu_1 + p\mu_2)e_1 = \lambda\mu(a_1, a_3). \end{cases}$$

The last two equations are equivalent to

$$\phi_{\mathfrak{p}}(a_3)(\mu(a_1, a_2)) = -2p\mu(a_1, a_2) \quad \text{and} \quad p[a_1, a_2]_{\mathfrak{p}} = 0.$$

•  $p \neq 0$  then

$$\alpha = 0, \mu(a_1, a_2) = 0, \mu(a_2, a_3) = -\lambda^{-1}(p\mu_1 + q\mu_2)e_1 \quad \text{and} \\ \mu(a_1, a_3) = \lambda^{-1}(-q\mu_1 + p\mu_2)e_1.$$

•  $p = 0$  and  $\mu_3 \neq 0$  then  $\alpha = 0$  and

$$\mu(a_1, a_2) = ce_1, \mu(a_2, a_3) = -\lambda^{-1}(p\mu_1 + q\mu_2)e_1 \quad \text{and} \\ \mu(a_1, a_3) = \lambda^{-1}(-q\mu_1 + p\mu_2)e_1.$$

•  $p = 0$  and  $\mu_3 = 0$  then

$$\mu(a_1, a_2) = c_1e_1 + c_2e_2, \mu(a_2, a_3) = -\lambda^{-1}(p\mu_1 + q\mu_2)e_1 \quad \text{and} \\ \mu(a_1, a_3) = \lambda^{-1}(-q\mu_1 + p\mu_2)e_1.$$

□

By using Propositions 4.2–4.7, we can give all the Riemann-Poisson Lie algebras of dimension 3, 4 or 5.

Let  $(\mathfrak{g}, [\cdot, \cdot], \varrho, r)$  be a Riemann-Poisson Lie algebra of dimension less or equal to 5. According to what above then  $\mathfrak{g} = \mathfrak{h} \oplus \mathfrak{p}$  and the Lie bracket on  $\mathfrak{g}$  is given by (18) and  $((\mathfrak{h}, \omega, \varrho_{\mathfrak{h}}), (\mathfrak{p}, [\cdot, \cdot]_{\mathfrak{p}}, \varrho_{\mathfrak{p}}), \mu, \phi_{\mathfrak{h}}, \phi_{\mathfrak{p}})$  are solutions of Problem 1.

•  $\dim \mathfrak{g} = 3$ . In this case  $\dim \mathfrak{h} = 2$  and  $\dim \mathfrak{p} = 1$  and, by applying Proposition 4.2, the Lie bracket of  $\mathfrak{g}$ ,  $\varrho$  and  $r$  are given in Table 1, where  $e^{12} = e_1 \wedge e_2$ .

Non vanishing Lie brackets	Bivector $r$	Matrix of $\varrho$	Conditions
$[e_1, e_2] = ae_1, [e_3, e_2] = be_1$	$\alpha e^{12}$	$I_3$	$a \neq 0, \alpha \neq 0$
$[e_3, e_1] = -be_1 + ce_2, [e_3, e_2] = de_1 + be_2$	$\alpha e^{12}$	$I_3$	$\alpha \neq 0$

TAB. 1. Three dimensional Riemann-Poisson Lie algebras

•  $\dim \mathfrak{g} = 4$ . We have three cases:

(c41)  $\dim \mathfrak{h} = 2$ ,  $\dim \mathfrak{p} = 2$  and  $\mathfrak{h}$  is non abelian and we can apply Proposition 4.3 to get the Lie brackets on  $\mathfrak{g}$ ,  $\varrho$  and  $r$ . They are described in rows 1 and 2 in Table 2.

(c42)  $\dim \mathfrak{h} = 2$ ,  $\dim \mathfrak{p} = 2$  and  $\mathfrak{h}$  is abelian and we can apply Propositions 4.4 and 4.5 to get the Lie brackets on  $\mathfrak{g}$ ,  $\varrho$  and  $r$ . They are described in rows 3 and 8 in Table 2.

(c43)  $\dim \mathfrak{h} = 4$ . In this case  $\mathfrak{g}$  is a Kähler Lie algebra. We have used [12] to derive all four dimensional Kähler Lie algebra together with their symplectic derivations. The results are given in Table 3. The notation  $\text{Der}^s(\mathfrak{h})$  stands

for the vector spaces of derivations which are skew-symmetric with respect the symplectic form. The vector space  $\text{Der}^s(\mathfrak{h})$  is described by a family of generators and  $E_{ij}$  is the matrix with 1 in the  $i$  row and  $j$  column and 0 elsewhere.

Non vanishing Lie brackets	Bivector $r$	Matrix of $\varrho$	Conditions
$[e_1, e_2] = ae_1, [e_3, e_2] = be_1 + ce_4,$ $[e_4, e_2] = de_1 - ce_3$	$\alpha e^{12}$	$I_4$	$a \neq 0, \alpha \neq 0$
$[e_1, e_2] = ae_1, [e_3, e_2] = be_1,$ $[e_4, e_2] = de_1, [e_3, e_4] = ce_3 - a^{-1}cbe_1$	$\alpha e^{12}$	$I_4$	$\alpha ac \neq 0,$
$[e_3, e_4] = ae_1 + be_2$	$\alpha e^{12}$	$I_4$	$\alpha \neq 0$
$[e_3, e_4] = ae_1 + be_2 + ce_3, [e_4, e_1] = xe_1 + ye_2,$ $[e_4, e_2] = ze_1 - xe_2$	$\alpha e^{12}$	$I_4$	$\alpha \neq 0$
$[e_3, e_4] = ae_1 + be_2 + 2e_4, [e_3, e_1] = e_1,$ $[e_3, e_2] = -e_2, [e_4, e_2] = e_1$	$\alpha e^{12}$	$\text{Diag}\left(1, 1, \begin{pmatrix} \mu & \nu \\ \nu & \rho \end{pmatrix}\right)$	$\alpha \neq 0, \mu, \rho > 0$ $\mu\rho > \nu^2$
$[e_3, e_4] = ae_1 + be_2 - 2e_4, [e_3, e_1] = e_1,$ $[e_3, e_2] = -e_2, [e_4, e_1] = e_2$	$\alpha e^{12}$	$\text{Diag}\left(1, 1, \begin{pmatrix} \mu & \nu \\ \nu & \rho \end{pmatrix}\right)$	$\alpha \neq 0, \mu, \rho > 0$ $\mu\rho > \nu^2$
$[e_3, e_4] = ae_1 + be_2 - 2e_3, [e_3, e_1] = e_1 + xe_2,$ $[e_3, e_2] = -\frac{1}{x}e_1 - e_2, [e_4, e_1] = xe_2, [e_4, e_2] = \frac{1}{x}e_1$	$\alpha e^{12}$	$\text{Diag}\left(1, 1, \begin{pmatrix} \mu & \nu \\ \nu & \rho \end{pmatrix}\right)$	$\alpha \neq 0, \mu, \rho > 0$ $\mu\rho > \nu^2, x \neq 0$
$[e_3, e_4] = ae_1 + be_2, [e_3, e_2] = xe_1 + ye_4,$ $[e_4, e_2] = ze_1 - ye_3$	$\alpha e^{12}$	$I_4$	$\alpha y \neq 0$

TAB. 2. Four dimensional Riemann-Poisson Lie algebras of rank 2

Non vanishing Lie brackets	Bivector $r$	Matrix of $\varrho$	$\text{Der}^s(\mathfrak{h})$
$[e_1, e_2] = e_2,$	$\alpha e^{12} + \beta e^{34}$	$\text{Diag}(a, b, c, d)$	$\{E_{21}, E_{33} - E_{44}, E_{43}, E_{34}\}$
$[e_1, e_2] = -e_3, [e_1, e_3] = e_2,$	$\alpha e^{14} + \beta e^{23}$	$\text{Diag}(a, b, b, c)$	$\{E_{23} - E_{32}, E_{41}\}$
$[e_1, e_2] = e_2, [e_3, e_4] = e_4,$	$\alpha e^{12} + \beta e^{34}$	$\text{Diag}(a, b, c, d)$	$\{E_{21}, E_{43}\}$
$[e_4, e_1] = e_1, [e_4, e_2] = -\delta e_3,$ $[e_4, e_3] = \delta e_2$	$\alpha e^{14} + \beta e^{23}$	$\text{Diag}(a, b, b, c)$	$\{E_{14}, E_{23} - E_{32}\}$
$[e_1, e_2] = e_3, [e_4, e_3] = e_3,$ $[e_4, e_1] = \frac{1}{2}e_1, [e_4, e_2] = \frac{1}{2}e_2,$	$\alpha(e^{12} - e^{34})$	$\text{Diag}(a, \mu b, \mu a, b)$	$\{E_{34}, E_{22} - E_{11}, E_{12} + E_{21}\}$
$[e_1, e_2] = e_3, [e_4, e_3] = e_3,$ $[e_4, e_1] = 2e_1, [e_4, e_2] = -e_2,$	$\alpha(e^{23} + e^{14})$	$\text{Diag}(a, a, 2a, 2a)$	$\{2E_{14} - E_{32}\}$
$[e_1, e_2] = e_3, [e_4, e_3] = e_3,$ $[e_4, e_1] = \frac{1}{2}e_1 - e_2,$ $[e_4, e_2] = e_1 + \frac{1}{2}e_2,$	$\alpha(e^{12} - e^{34})$	$\text{Diag}(a, a, a, a)$	$\{E_{34}, E_{12} - E_{21}\}$

TAB. 3. Four-dimensional Kähler Lie algebras and their symplectic derivations,  $a, b, c, d > 0, \alpha\beta \neq 0$

•  $\dim \mathfrak{g} = 5$ . We have:

- (c51)  $\dim \mathfrak{h} = 4$  and  $\mathfrak{h}$  abelian and hence a symplectic vector space. We can apply Proposition 4.2 and  $\mathfrak{g}$  is semi-direct product.
- (c52)  $\dim \mathfrak{h} = 4$  and  $\mathfrak{h}$  non abelian. We can apply Proposition 4.2 and Table 3 to get the Lie brackets on  $\mathfrak{g}, \varrho$  and  $r$ . The result is summarized in Table 4.
- (c53)  $\dim \mathfrak{h} = 2$  and  $\mathfrak{h}$  non abelian. We apply Proposition 4.3. In this case  $(\mathfrak{p}, [\cdot, \cdot]_{\mathfrak{p}}, \varrho_{\mathfrak{p}})$  is a 3-dimensional Euclidean Lie algebra and one must compute  $\text{Der}(\mathfrak{p}) \cap \text{so}(\mathfrak{p})$  and solve (20). Three dimensional Euclidean Lie algebras were classified in [9]. For each of them we have computed  $\text{Der}(\mathfrak{p}) \cap \text{so}(\mathfrak{p})$

and solved (20) by using Maple. The result is summarized in Table 5 when  $\mathfrak{p}$  is unimodular and Table 6 when  $\mathfrak{p}$  is nonunimodular.

(c54)  $\dim \mathfrak{h} = 2$  and  $\mathfrak{h}$  abelian and  $\phi_{\mathfrak{h}} = 0$ . We apply Proposition 4.6 and we perform all the needed computations. We use the classification of 3-dimensional Euclidean Lie algebras given in [9]. The results are given in Tables 7-8.

(c55)  $\dim \mathfrak{h} = 2$  and  $\mathfrak{h}$  abelian and  $\phi_{\mathfrak{h}} \neq 0$ . We apply Proposition 4.7 and we perform all the needed computations. The results are given in Table 9.

Non vanishing Lie brackets	Bivector $r$	Matrix of $\rho$	Conditions
$[e_1, e_2] = e_2, [e_5, e_1] = xe_2,$ $[e_5, e_3] = ye_3 + te_4, [e_5, e_4] = ze_3 - ye_4$	$\alpha e^{12} + \beta e^{34}$	$\text{Diag}(a, b, c, d, e)$	$\alpha\beta \neq 0$ $a, b, c, d, e > 0$
$[e_1, e_2] = -e_3, [e_1, e_3] = e_2,$ $[e_5, e_1] = ye_4, [e_5, e_2] = -xe_3, [e_5, e_3] = xe_2$	$\alpha e^{14} + \beta e^{23}$	$\text{Diag}(a, b, b, c, d)$	$\alpha\beta \neq 0$ $a, b, c, d > 0$
$[e_1, e_2] = e_2, [e_3, e_4] = e_4,$ $[e_5, e_1] = xe_2, [e_5, e_3] = ye_4$	$\alpha e^{12} + \beta e^{34}$	$\text{Diag}(a, b, c, d, e)$	$\alpha\beta \neq 0$ $a, b, c, d, e > 0$
$[e_4, e_1] = e_1, [e_4, e_2] = -\delta e_3, [e_4, e_3] = \delta e_2$ $[e_5, e_2] = -ye_3, [e_5, e_3] = ye_2, [e_5, e_4] = xe_1$	$\alpha e^{14} + \beta e^{23}$	$\text{Diag}(a, b, b, c, d)$	$\alpha\beta \neq 0, \delta > 0$ $a, b, c, d > 0$
$[e_1, e_2] = e_3, [e_4, e_3] = e_3, [e_4, e_1] = \frac{1}{2}e_1$ $[e_4, e_2] = \frac{1}{2}e_2, [e_5, e_1] = xe_1 + ye_2,$ $[e_5, e_2] = ye_1 - xe_2, [e_5, e_4] = ze_3$	$\alpha(e^{12} - e^{34})$	$\text{Diag}(a, \mu b, \mu a, b, c)$	$\alpha \neq 0$ $a, b, c, \mu > 0$
$[e_1, e_2] = e_3, [e_4, e_3] = e_3, [e_4, e_1] = 2e_1$ $[e_4, e_2] = -e_2, [e_5, e_2] = xe_3, [e_5, e_4] = -2xe_1$	$\alpha(e^{23} + e^{14})$	$\text{Diag}(a, a, 2a, 2a, b)$	$\alpha \neq 0$ $a, b > 0$
$[e_1, e_2] = e_3, [e_4, e_3] = e_3, [e_4, e_1] = \frac{1}{2}e_1 - e_2$ $[e_4, e_2] = e_1 + \frac{1}{2}e_2, [e_5, e_1] = -xe_2, [e_5, e_2] = xe_1$ $[e_5, e_4] = ye_3$	$\alpha(e^{12} - e^{34})$	$\text{Diag}(a, a, a, a, b)$	$\alpha \neq 0$ $a, b > 0$

TAB. 4. Five-dimensional Riemann-Poisson Lie algebras of rank 4

Non vanishing Lie brackets	$r$	Matrix of $\rho$	Conditions
$[e_1, e_2] = e_1, [e_3, e_2] = b\mu e_1 - ce_4, [e_4, e_2] = d\mu e_1 + ce_3$ $[e_5, e_2] = fe_1, [e_3, e_4] = -fe_1 + e_5$	$\alpha e^{12}$	$\text{Diag}(1, \rho, \mu, \mu, 1)$	$c\alpha \neq 0$ $\mu, \rho > 0$
$[e_1, e_2] = e_1, [e_3, e_2] = be_1, [e_4, e_2] = ce_1$ $[e_5, e_2] = d\mu e_1, [e_3, e_5] = be_1 - e_2, [e_4, e_5] = -ce_1 + e_4$	$\alpha e^{12}$	$\text{Diag}(1, \rho, 1, 1, \mu)$	$\alpha \neq 0$ $\mu, \rho > 0$
$[e_1, e_2] = e_1, [e_3, e_2] = (b+c)e_1, [e_4, e_2] = (cx+b)e_1$ $[e_5, e_2] = d\mu e_1, [e_3, e_5] = (b+c)e_1 - e_3,$ $[e_4, e_5] = -(cx+b)e_1 + e_4$	$\alpha e^{12}$	$\text{Diag}(1, \rho, \begin{pmatrix} 1 & 1 \\ 1 & x \end{pmatrix}, \mu)$	$\alpha \neq 0$ $\mu, \rho > 0$
$[e_1, e_2] = e_1, [e_3, e_2] = be_1, [e_4, e_2] = c\mu e_1$ $[e_5, e_2] = d\nu e_1, [e_3, e_5] = -\mu ce_1 + e_4, [e_4, e_5] = be_1 - e_3$	$\alpha e^{12}$	$\text{Diag}(1, \rho, 1, \mu, \nu)$	$\alpha \neq 0$ $\mu, \nu, \rho > 0$
$[e_1, e_2] = e_1, [e_3, e_2] = b\mu e_1, [e_4, e_2] = c\nu e_1$ $[e_5, e_2] = d\rho e_1, [e_3, e_4] = -2\rho de_1 + 2e_5,$ $[e_3, e_5] = 2\nu ce_1 - 2e_4, [e_4, e_5] = 2\mu be_1 - 2e_3$	$\alpha e^{12}$	$\text{Diag}(1, \xi, \mu, \nu, \rho)$	$\alpha \neq 0, \nu \neq \rho$ $\mu, \nu, \rho, \xi > 0$ $\mu \neq \nu, \mu \neq \rho$
$[e_1, e_2] = e_1, [e_3, e_2] = b\mu e_1, [e_4, e_2] = c\nu e_1 - \lambda e_5$ $[e_5, e_2] = d\nu e_1 + \lambda e_4, [e_3, e_4] = -\frac{2\nu(\lambda c+d)}{1+\lambda^2}e_1 + 2e_5,$ $[e_3, e_5] = \frac{2\nu(c-\lambda d)}{1+\lambda^2}e_1 - 2e_4, [e_4, e_5] = 2\mu be_1 - 2e_3$	$\alpha e^{12}$	$\text{Diag}(1, \rho, \mu, \nu, \nu)$	$\lambda\alpha \neq 0$ $\mu, \nu, \rho > 0$
$[e_1, e_2] = e_1, [e_3, e_2] = b\mu e_1, [e_4, e_2] = c\nu e_1$ $[e_5, e_2] = d\rho e_1, [e_3, e_4] = -\rho de_1 + e_5,$ $[e_3, e_5] = \nu ce_1 - e_4, [e_4, e_5] = -\mu be_1 + e_3$	$\alpha e^{12}$	$\text{Diag}(1, \xi, \mu, \nu, \rho)$	$\alpha \neq 0, \nu \neq \rho$ $\mu, \nu, \rho, \xi > 0$ $\mu \neq \nu, \mu \neq \rho$
$[e_1, e_2] = e_1, [e_3, e_2] = b\mu e_1, [e_4, e_2] = c\nu e_1 - \lambda e_5$ $[e_5, e_2] = d\nu e_1 + \lambda e_4, [e_3, e_4] = -\frac{\nu(\lambda c+d)}{1+\lambda^2}e_1 + e_5,$ $[e_3, e_5] = \frac{\nu(c-\lambda d)}{1+\lambda^2}e_1 - e_4, [e_4, e_5] = -\mu be_1 + e_3$	$\alpha e^{12}$	$\text{Diag}(1, \rho, \mu, \nu, \nu)$	$\lambda\alpha \neq 0$ $\mu, \nu, \rho > 0$
$[e_1, e_2] = e_1, [e_3, e_2] = b\mu e_1 - ue_4 - ve_5,$ $[e_4, e_2] = c\mu e_1 + ue_3 - we_5, [e_5, e_2] = d\mu e_1 + ve_3 + we_4,$ $[e_3, e_4] = xe_1 + e_5, [e_3, e_5] = ye_1 - e_4, [e_4, e_5] = ze_1 + e_3$ $x = -\frac{\mu(buw-cv+du^2+bv+cw+d)}{1+u^2+v^2+w^2}$ $y = \frac{\mu(-bv+w+cv^2-duw+bu-dw+c)}{1+u^2+v^2+w^2}$ $z = -\frac{\mu(bw^2-cv+duw-cu-dv+b)}{1+u^2+v^2+w^2}$	$\alpha e^{12}$	$\text{Diag}(1, \rho, \mu, \mu, \mu)$	$\alpha \neq 0$ $\mu, \rho > 0$

TAB. 5. Five-dimensional Riemann-Poisson Lie algebras of rank 2 with non abelian Kähler subalgebra and unimodular complement

Non vanishing Lie brackets	$r$	Matrix of $\varrho$
$[e_1, e_2] = e_1, [e_3, e_2] = (f + c\lambda + f\lambda^2)e_1 - \lambda e_4, \lambda \neq 0,$ $[e_4, e_2] = ce_1 + \lambda e_3, [e_5, e_2] = d\mu e_1,$ $[e_3, e_5] = fe_1 - e_3, [e_4, e_5] = (\lambda f + c)e_1 - e_4,$	$\alpha e^{12}$	$\text{Diag}(1, \rho, 1, 1, \mu)$ $\mu, \rho > 0$
$[e_1, e_2] = e_1, [e_3, e_2] = be_1, [e_4, e_2] = c\mu e_1,$ $[e_5, e_2] = d\nu e_1, [e_3, e_5] = \mu ce_1 - e_4,$ $[e_4, e_5] = (-fb + 2\mu c)e_1 + fe_3 - 2e_4, f = 1 \text{ or } f \leq 0$	$\alpha e^{12}$	$\text{Diag}(1, \rho, 1, \mu, \nu)$ $0 < \mu <  f , \rho > 0$
$[e_1, e_2] = e_1, [e_3, e_2] = (b + c\mu)e_1, [e_4, e_2] = (c + b\mu)e_1,$ $[e_5, e_2] = d\nu e_1, [e_3, e_5] = (\mu b + c)e_1 - e_4,$ $[e_4, e_5] = ((2 - \mu)c + (2\mu - 1)b)e_1 + e_3 - 2e_4$	$\alpha e^{12}$	$\text{Diag}(1, \rho, \begin{pmatrix} 1 & \mu \\ \mu & 1 \end{pmatrix}, \nu)$ $\mu, \nu, \rho > 0$
$[e_1, e_2] = e_1, [e_3, e_2] = (b + c)e_1, [e_4, e_2] = (b + c\mu)e_1,$ $[e_5, e_2] = d\nu e_1, [e_3, e_5] = (b + c\mu)e_1 - e_4,$ $[e_4, e_5] = ((2 - f)b + (2\mu - f)c)e_1 + fe_3 - 2e_4$	$\alpha e^{12}$	$\text{Diag}(1, \rho, \begin{pmatrix} 1 & 1 \\ 1 & \mu \end{pmatrix}, \nu)$ $\nu, \rho > 0, c > \mu > 1$
$[e_1, e_2] = e_1, [e_3, e_2] = (b + \frac{1}{2}c)e_1, [e_4, e_2] = (c + \frac{1}{2}b)e_1,$ $[e_5, e_2] = d\nu e_1, [e_3, e_5] = (c + \frac{1}{2}b)e_1 - e_4,$ $[e_4, e_5] = (b + 2c)e_1 - 2e_4$	$\alpha e^{12}$	$\text{Diag}(1, \rho, \begin{pmatrix} 1 & \frac{1}{2} \\ \frac{1}{2} & 1 \end{pmatrix}, \nu)$ $\rho, \nu > 0$
$[e_1, e_2] = e_1, [e_3, e_2] = xe_1, [e_4, e_2] = ye_1,$ $[e_5, e_2] = d\nu e_1, [e_3, e_5] = ze_1 - e_4,$ $[e_4, e_5] = te_1 + fe_3 - 2e_4, 0 < f < 1,$ $x = \frac{((\mu+1)b+(\mu-1)c)f-2b}{2f^2(f-1)}, y = z = \frac{(\mu-1)(cf+b)}{2f(f-1)}$ $t = \frac{(1-\mu)cf+((f-2)\mu+f)b}{2f(1-f)}$	$\alpha e^{12}$	$A^t B A$ $A = \begin{pmatrix} \frac{1+s}{-2fs} & -\frac{1}{2s} & 0 \\ \frac{1-s}{2fs} & \frac{1}{2s} & 0 \\ 0 & 0 & 1 \end{pmatrix}$ $B = \text{Diag}(1, \rho, \begin{pmatrix} 1 & \mu \\ \mu & 1 \end{pmatrix}, \nu)$ $\nu, \rho > 0$ $s = \sqrt{1-f}, 0 \leq \mu < 1$

TABLE 6. Five-dimensional Riemann-Poisson Lie algebras of rank 2 with non abelian Kähler subalgebra and non unimodular complement. ( $\alpha \neq 0$ )

Non vanishing Lie brackets	Bivector $r$	Matrix of $\varrho$	Conditions
$[e_3, e_4] = ae_1 + be_2 + e_5, [e_3, e_5] = ce_1 + de_2$ $[e_4, e_5] = fe_1 + ge_2$	$\alpha e^{12}$	$\text{Diag}(1, 1, \mu, \mu, 1)$	$\alpha \neq 0$ $\mu > 0$
$[e_3, e_4] = ae_1 + be_2, [e_3, e_5] = ce_1 + de_2 - e_3$ $[e_4, e_5] = fe_1 + ge_2 + e_4$	$\alpha e^{12}$	$\text{Diag}(1, 1, 1, 1, \mu)$ $\text{Diag}(1, 1, \begin{pmatrix} 1 & 1 \\ 1 & x \end{pmatrix}, \mu)$	$\alpha \neq 0$ $\mu > 0$
$[e_3, e_4] = ae_1 + be_2, [e_3, e_5] = ce_1 + de_2 + e_4$ $[e_4, e_5] = fe_1 + ge_2 - e_3$	$\alpha e^{12}$	$\text{Diag}(1, 1, 1, \mu, \nu)$	$\alpha \neq 0$ $\mu, \nu > 0$
$[e_3, e_4] = ae_1 + be_2 + 2e_5, [e_3, e_5] = ce_1 + de_2 - 2e_4$ $[e_4, e_5] = fe_1 + ge_2 - 2e_3$	$\alpha e^{12}$	$\text{Diag}(1, 1, \mu, \nu, \rho)$	$\alpha \neq 0$ $\mu, \nu, \rho > 0$
$[e_3, e_4] = ae_1 + be_2 + e_5, [e_3, e_5] = ce_1 + de_2 - e_4$ $[e_4, e_5] = fe_1 + ge_2 + e_3$	$\alpha e^{12}$	$\text{Diag}(1, 1, \mu, \nu, \rho)$	$\alpha \neq 0$ $\mu, \nu, \rho > 0$
$[e_3, e_5] = ce_1 + de_2 - e_3$ $[e_4, e_5] = fe_1 + ge_2 - e_4$	$\alpha e^{12}$	$\text{Diag}(1, 1, 1, 1, \mu)$	$\alpha \neq 0$ $\mu > 0$
$[e_3, e_5] = ce_1 + de_2 - e_4$ $[e_4, e_5] = fe_1 + ge_2 + xe_3 - 2e_4$	$\alpha e^{12}$	There are many cases See [9]	$\alpha \neq 0$

TABLE 7. Five-dimensional Riemann-Poisson Lie algebras of rank 2 with abelian Kähler subalgebra

Non vanishing Lie brackets	$r$	Matrix of $\varrho$
$[e_3, e_1] = -e_2, [e_3, e_2] = e_1, [e_4, e_1] = e_2, [e_4, e_2] = e_1$ $[e_5, e_1] = e_1, [e_5, e_2] = -e_2,$ $[e_3, e_4] = 2e_5 + (l_{22} - l_{21} - 2l_{13})e_1 - (l_{12} + l_{11} + 2l_{23})e_2$ $[e_3, e_5] = -2e_4 + (l_{23} - l_{11} + 2l_{12})e_1 - (l_{13} - l_{21} - 2l_{22})e_2,$ $[e_4, e_5] = -2e_3 + (l_{23} - l_{12} + 2l_{11})e_1 + (l_{13} + l_{22} + 2l_{21})e_2$	$\alpha e^{12}$	Diag(1, 1, $\mu, \nu, \rho$ ) $\mu, \nu, \rho > 0$
$[e_4, e_2] = ue_1, [e_5, e_1] = -\frac{\alpha}{2}e_1, [e_5, e_2] = ve_1 + \frac{\alpha}{2}e_2,$ $[e_3, e_4] = xe_1 + ye_2, [e_3, e_5] = be_3 + ze_1 + te_2,$ $[e_4, e_5] = ce_3 + ae_4 + re_1 + se_2,$ $(a + 2b)x - 2tu + 2yv = 0, a \neq 0, b \neq 0, (3a + 2b)y = 0$	$\alpha e^{12}$	Diag(1, 1, 1, 1, 1)
$[e_4, e_2] = ue_1, [e_5, e_1] = -\frac{\alpha}{2}e_1, [e_5, e_2] = ve_1 + \frac{\alpha}{2}e_2,$ $[e_3, e_4] = xe_1, [e_3, e_5] = ze_1 + te_2,$ $[e_4, e_5] = ae_4 + re_1 + se_2, a \neq 0,$ $ax - 2tu = 0$	$\alpha e^{12}$	Diag(1, 1, $\begin{pmatrix} 1 & \mu \\ \mu & 1 \end{pmatrix}, 1)$
$[e_4, e_1] = ue_2, [e_5, e_1] = \frac{\alpha}{2}e_1 + ve_2, [e_5, e_2] = -\frac{\alpha}{2}e_2,$ $[e_3, e_4] = xe_1 + ye_2, [e_3, e_5] = be_3 + ze_1 + te_2,$ $[e_4, e_5] = ce_3 + ae_4 + re_1 + se_2,$ $(3a + 2b)x = 0, a \neq 0, b \neq 0$ $(a + 2b)y - 2zu + 2xv = 0$	$\alpha e^{12}$	Diag(1, 1, 1, 1, 1)
$[e_4, e_1] = ue_2, [e_5, e_1] = \frac{\alpha}{2}e_1 + ve_2, [e_5, e_2] = -\frac{\alpha}{2}e_2,$ $[e_3, e_4] = ye_2, [e_3, e_5] = ze_1 + te_2,$ $[e_4, e_5] = ae_4 + re_1 + se_2, a \neq 0, ay - 2zu = 0$	$\alpha e^{12}$	Diag(1, 1, $\begin{pmatrix} 1 & \mu \\ \mu & 1 \end{pmatrix}, 1)$
$[e_4, e_1] = ue_1 + upe_2, [e_4, e_2] = -\frac{u}{p}e_1 - ue_2,$ $[e_5, e_1] = ve_1 + \frac{(2v-a)p}{2}e_2, [e_5, e_2] = -\frac{(2v+a)}{2p}e_1 - ve_2$ $[e_3, e_4] = xe_1 + ye_2, [e_3, e_5] = be_3 + ze_1 + te_2, a \neq 0, b \neq 0$ $[e_4, e_5] = ce_3 + ae_4 + re_1 + se_2,$ $((2a + 2b + 2v)x - 2zu)p - ay + 2tu - 2yv = 0$ $(2xv - ax - 2zu)p + (2a + 2b - 2v)y + 2tu = 0$	$\alpha e^{12}$	Diag(1, 1, 1, 1, 1)
$[e_4, e_1] = ue_1 + upe_2, [e_4, e_2] = -\frac{u}{p}e_1 - ue_2,$ $[e_5, e_1] = ve_1 + \frac{(2v-a)p}{2}e_2, [e_5, e_2] = -\frac{(2v+a)}{2p}e_1 - ve_2$ $[e_3, e_4] = xe_1 + ye_2, [e_3, e_5] = ze_1 + te_2,$ $[e_4, e_5] = ae_4 + re_1 + se_2, a \neq 0, b \neq 0$ $((2a + 2v)x - 2zu)p - ay + 2tu - 2yv = 0$ $(2xv - ax - 2zu)p + (2a - 2v)y + 2tu = 0$	$\alpha e^{12}$	Diag(1, 1, $\begin{pmatrix} 1 & \mu \\ \mu & 1 \end{pmatrix}, 1)$
$[e_5, e_1] = ue_1 + ve_2, [e_5, e_2] = we_1 - ue_2,$ $[e_3, e_4] = xe_1 + ye_2, [e_3, e_5] = ae_3 + be_4 + ze_1 + te_2,$ $[e_4, e_5] = ce_3 + de_4 + re_1 + se_2,$ $(a + d + u)x + yw = 0, xv + (a + d - u)y = 0$	$\alpha e^{12}$	Diag(1, 1, 1, 1, 1)
$[e_5, e_1] = ue_1 + ve_2, [e_5, e_2] = we_1 - ue_2,$ $[e_3, e_4] = xe_1 + ye_2 + ae_4, [e_3, e_5] = be_4 + ze_1 + te_2,$ $[e_4, e_5] = ce_4 + re_1 + se_2, a \neq 0$ $(c + u)x - ar + yw = 0$ $(c - u)y - as + xv = 0$	$\alpha e^{12}$	Diag(1, 1, 1, 1, 1)

TAB. 8. Five-dimensional Riemann-Poisson Lie algebras of rank 2 with abelian Kähler subalgebra ( $\alpha \neq 0$ ) (Continued)

Non vanishing Lie brackets	Bivector $r$	Matrix of $\varrho$	Conditions
$[e_3, e_2] = xe_1 - ae_4, [e_4, e_2] = ye_1 + ae_3, [e_5, e_2] = ze_1$ $[e_3, e_5] = pe_3 + qe_4 + a^{-1}(-qx + py)e_1,$ $[e_3, e_5] = -qe_3 + pe_4 - a^{-1}(px + qy)e_1$	$\alpha e^{12}$	$\text{Diag}(1, 1, 1, 1, 1)$	$\alpha \neq 0$ $a \neq 0$
$[e_3, e_2] = xe_1 - ae_4, [e_4, e_2] = ye_1 + ae_3, [e_5, e_2] = ze_1$ $[e_3, e_4] = be_1$ $[e_3, e_5] = qe_4 - a^{-1}qx e_1,$ $[e_3, e_5] = -qe_3 - a^{-1}qye_1$	$\alpha e^{12}$	$\text{Diag}(1, 1, 1, 1, 1)$	$\alpha \neq 0$ $a \neq 0, z \neq 0$
$[e_3, e_2] = xe_1 - ae_4, [e_4, e_2] = ye_1 + ae_3,$ $[e_3, e_4] = be_1 + ce_2$ $[e_3, e_5] = qe_4 - a^{-1}qx e_1,$ $[e_3, e_5] = -qe_3 - a^{-1}qye_1$	$\alpha e^{12}$	$\text{Diag}(1, 1, 1, 1, 1)$	$\alpha \neq 0$ $a \neq 0$

TAB. 9. Five-dimensional Riemann-Poisson Lie algebras of rank 2 with abelian Kähler subalgebra (Continued)

This theorem unknown to our knowledge can be used to build examples of Riemann-Poisson Lie algebras.

**Theorem 4.1.** *Let  $(G, \langle , \rangle)$  be an even dimensional flat Riemannian Lie group. Then there exists a left invariant differential  $\Omega$  on  $G$  such that  $(G, \langle , \rangle, \Omega)$  is a Kähler Lie group.*

**Proof.** Let  $\mathfrak{g}$  be the Lie algebra of  $G$  and  $\varrho = \langle , \rangle(e)$ . According to Milnor’s Theorem [11, Theorem 1.5] and its improved version [1, Theorem 3.1] the flatness of the metric on  $G$  is equivalent to  $[\mathfrak{g}, \mathfrak{g}]$  is even dimensional abelian,  $[\mathfrak{g}, \mathfrak{g}]^\perp = \{u \in \mathfrak{g}, \text{ad}_u + \text{ad}_u^* = 0\}$  is also even dimensional abelian and  $\mathfrak{g} = [\mathfrak{g}, \mathfrak{g}] \oplus [\mathfrak{g}, \mathfrak{g}]^\perp$ . Moreover, the Levi-Civita product is given by

$$(21) \quad L_a = \begin{cases} \text{ad}_a & \text{if } a \in [\mathfrak{g}, \mathfrak{g}]^\perp, \\ 0 & \text{if } a \in [\mathfrak{g}, \mathfrak{g}] \end{cases}$$

and there exists a basis  $(e_1, f_1, \dots, e_r, f_r)$  of  $[\mathfrak{g}, \mathfrak{g}]$  and  $\lambda_1, \dots, \lambda_r \in [\mathfrak{g}, \mathfrak{g}]^\perp \setminus \{0\}$  such that for any  $a \in [\mathfrak{g}, \mathfrak{g}]^\perp$ ,

$$[a, e_i] = \lambda_i(a) f_i \quad \text{and} \quad [a, f_i] = -\lambda_i(a) e_i.$$

We consider a nondegenerate skew-symmetric 2-form  $\omega_0$  on  $[\mathfrak{g}, \mathfrak{g}]^\perp$  and  $\omega_1$  the nondegenerate skew-symmetric 2-form on  $[\mathfrak{g}, \mathfrak{g}]$  given by  $\omega_1 = \sum_{i=1}^r e_i^* \wedge f_i^*$ . One can see easily that  $\omega = \omega_0 \oplus \omega_1$  is a Kähler form on  $\mathfrak{g}$ .  $\square$

REFERENCES

- [1] Ait Haddou, M., Boucetta, M., Lebzioui, H., *Left-invariant Lorentzian flat metrics on Lie groups*, J. Lie Theory **22** (1) (2012), 269–289.
- [2] Boucetta, M., *Compatibilité des structures pseudo-riemanniennes et des structures de Poisson*, C.R. Acad. Sci. Paris Sér. I **333** (2001), 763–768.
- [3] Boucetta, M., *Riemann-poisson manifolds and kähler-riemann foliations*, C.R. Acad. Sci. Paris, Sér. I **336** (2003), 423–428.
- [4] Boucetta, M., *Poisson manifolds with compatible pseudo-metric and pseudo-Riemannian Lie algebras*, Differential Geom. Appl. **20** (2004), 279–291.
- [5] Boucetta, M., *On the Riemann-Lie algebras and Riemann-Poisson Lie groups*, J. Lie Theory **15** (1) (2005), 183–195.

- [6] Deninger, C., Singhof, W., *Real polarizable hodge structures arising from foliation*, Ann. Global Anal. Geom. **21** (2002), 377–399.
- [7] Dufour, J.P., Zung, N.T., *Poisson Structures and Their Normal Forms*, Progress in Mathematics, vol. 242, Birkhäuser Verlag, 2005.
- [8] Fernandes, R.L., *Connections in Poisson Geometry 1: Holonomy and invariants*, J. Differential Geom. **54** (2000), 303–366.
- [9] Ha, K.Y., Lee, J.B., *Left invariant metrics and curvatures on simply connected three dimensional Lie groups*, Math. Nachr. **282** (2009), 868–898.
- [10] Hawkin, E., *The structure of noncommutative deformations*, J. Differential Geom. **77** (2007), 385–424.
- [11] Milnor, J., *Curvatures of left invariant metrics on Lie Groups*, Adv. Math. **21** (1976), 293–329.
- [12] Ovando, G., *Invariant pseudo-Kähler metrics in dimension four*, J. Lie Theory **16** (2006), 371–391.
- [13] Vaisman, I., *Lectures on the Geometry of Poisson Manifolds*, Progress in Mathematics, vol. 118, Birkhäuser, Berlin, 1994.

<sup>a</sup>UNIVERSITÉ DE NOUAKCHOTT,  
FACULTÉ DES SCIENCES ET TECHNIQUES  
*E-mail*: pacha.ali86@gmail.com

<sup>b</sup>UNIVERSITÉ CADI-AYYAD,  
FACULTÉ DES SCIENCES ET TECHNIQUES,  
BP 549 MARRAKECH MAROC  
*E-mail*: m.boucetta@uca.ac.ma

<sup>c</sup>UNIVERSITÉ DE NOUAKCHOTT,  
FACULTÉ DES SCIENCES ET TECHNIQUES  
*E-mail*: lessiadahmed@gmail.com