

ON BOUNDED MODULE MAPS BETWEEN HILBERT MODULES OVER LOCALLY C^* -ALGEBRAS

M. JOIȚA

ABSTRACT. Let A be a locally C^* -algebra and let E be a Hilbert A -module. We show that the algebra $B_A(E)$ of all bounded A -module maps on E is a locally m -convex algebra which is algebraically and topologically isomorphic to $LM(K_A(E))$, the algebra of all left multipliers of $K_A(E)$, where $K_A(E)$ is the locally C^* -algebra of all "compact" A -module maps on E . Also we show that $b(B_A(E))$, the algebra of all bounded elements in $B_A(E)$, is a Banach algebra which is isometrically isomorphic to $B_{b(A)}(b(E))$.

1. INTRODUCTION

A locally C^* -algebra is a complete Hausdorff complex topological $*$ -algebra A whose topology is determined by its continuous C^* -seminorms in the sense that the net $\{a_i\}_i$ converges to 0 if and only if the net $\{p(a_i)\}_i$ converges to 0 for every continuous C^* -seminorm p on A . In fact a locally C^* -algebra is an inverse limit of C^* -algebras.

Hilbert modules over locally C^* -algebras generalize the notion of Hilbert C^* -modules by allowing the inner product to take values in a locally C^* -algebra. In [9], Phillips showed that many results about multipliers of a C^* -algebra are valid for multipliers of a locally C^* -algebra. Thus, he proved that $M(A)$, the multiplier algebra of a locally C^* -algebra A , is a locally C^* -algebra in the topology of seminorm [9, Theorem 3.14]. In this note we show that any left multiplier of a locally C^* -algebra A is automatically continuous (Proposition 3.4) and $LM(A)$, the algebra of left multipliers of A , is a complete locally m -convex algebra in the topology of seminorm (Theorem 3.5). Also, Phillips shows that if E is a Hilbert module over a locally C^* -algebra A , then the locally C^* -algebra $L_A(E)$

Received May 16, 2003.

2000 *Mathematics Subject Classification.* Primary 46L08, 46L05, 46A13.

Key words and phrases. Hilbert modules over locally C^* -algebras, bounded module maps, locally m -convex algebras.

of all adjointable maps on E is isomorphic to $M(K_A(E))$, where $K_A(E)$ is the locally C^* -algebra of all "compact" A -module maps on E [9, Theorem 4.2]. This result is a generalization of Theorem 1 of [5] for Hilbert module over locally C^* -algebras. We show that the locally m -convex algebra $B_A(E)$ of all bounded A -module maps is isomorphic to $LM(K_A(E))$ (Theorem 3.6). This result generalizes Theorem 1.5 of [6] in the context of Hilbert modules over locally C^* -algebras. Finally we prove that if E and F are Hilbert modules over a locally C^* -algebra A , then $b(B_A(E, F))$, the set of all bounded elements in $B_A(E, F)$, is a Banach space in the norm $\|\cdot\|_\infty$ which is isometrically isomorphic to $B_{b(A)}(b(E), b(F))$, the Banach space of all bounded $b(A)$ -module maps from $b(E)$ to $b(F)$ (Theorem 3.7). In particular, $b(B_A(E))$ is a Banach algebra which is isometrically isomorphic to $B_{b(A)}(b(E))$ and $b(L_A(E))$ is a C^* -algebra which is isomorphic to $L_{b(A)}(b(E))$.

2. PRELIMINARIES

If A is a locally C^* -algebra and $S(A)$ is the set of all continuous C^* -seminorms on A , then for each $p \in S(A)$, $A_p = A / \ker(p)$ is a C^* -algebra in the norm induced by p and $A = \lim_{p \leftarrow} A_p$ (see, for example, [9]). The canonical map from A onto A_p , $p \in S(A)$ is denoted by π_p and the image of a in A under π_p by a_p . The connecting maps of the inverse system $\{A_p\}_{p \in S(A)}$ are denoted by π_{pq} , $q, p \in S(A)$, with $p \geq q$.

Now we recall some facts about Hilbert modules over locally C^* -algebras from [9].

Definition 2.1. A pre-Hilbert A -module is a complex vector space E which is also a right A -module, compatible with the complex algebra structure, equipped with an A -valued inner product $\langle \cdot, \cdot \rangle : E \times E \rightarrow A$ which is \mathbb{C} - and A -linear in its second variable and satisfies the following relations:

- (i) $\langle x, y \rangle^* = \langle y, x \rangle$ for every $x, y \in E$;
- (ii) $\langle x, x \rangle \geq 0$ for every $x \in E$;
- (iii) $\langle x, x \rangle = 0$ if and only if $x = 0$.

We say that E is a Hilbert A -module if E is complete with respect to the topology determined by the family of seminorms $\bar{p}_E(x) = \sqrt{p(\langle x, x \rangle)}$, $x \in E$, $p \in S(A)$.

Given a Hilbert A -module E , for each $p \in S(A)$, $N_p^E = \ker(\bar{p}_E)$ is a closed submodule of E and $E_p = E/N_p^E$ is a Hilbert A_p -module with $(x + N_p^E)\pi_p(a) = xa + N_p^E$ and $\langle x + N_p^E, y + N_p^E \rangle = \pi_p(\langle x, y \rangle)$. The canonical map from E onto E_p is denoted by σ_p^E , and the image of x in E under σ_p^E by x_p , $p \in S(A)$.

For each $p, q \in S(A)$ with $p \geq q$ there is a canonical surjective linear map $\sigma_{pq}^E : E_p \rightarrow E_q$ such that $\sigma_{pq}^E(x_p) = x_q$, $x \in E$. Then $\{E_p; A_p; \sigma_{pq}^E, p \geq q, p, q \in S(A)\}$ is an inverse system of Hilbert C^* -modules in the following sense:

- $\sigma_{pq}^E(x_p a_p) = \sigma_{pq}^E(x_p)\pi_{pq}(a_p)$ for every $x_p \in E_p$ and for every $a_p \in A_p$;
- $\langle \sigma_{pq}^E(x_p), \sigma_{pq}^E(y_p) \rangle = \pi_{pq}(\langle x_p, y_p \rangle)$ for every $x_p, y_p \in E_p$;
- $\sigma_{qr}^E \circ \sigma_{pq}^E = \sigma_{pr}^E$, $p \geq q \geq r$;
- $\sigma_{pp}^E = \text{id}_{E_p}$;

and $\lim_{p \leftarrow} E_p$ is a Hilbert A -module with $((x_p)_p)((a_p)_p) = (x_p a_p)_p$ and $\langle (x_p)_p, (y_p)_p \rangle = \langle \langle x_p, y_p \rangle \rangle_p$. Moreover, $\lim_{p \leftarrow} E_p$ can be identified with E .

We recall that an element a in A respectively x in E is bounded if

$$\|a\|_\infty = \sup\{p(a); p \in S(A)\} < \infty$$

respectively

$$\|x\|_\infty = \sup\{\bar{p}_E(x); p \in S(A)\} < \infty$$

The set of all bounded elements in A respectively in E will be denoted by $b(A)$ respectively $b(E)$. We know that $b(A)$ is a C^* -algebra in the C^* -norm $\|\cdot\|_\infty$, and $b(E)$ is a Hilbert $b(A)$ -module.

3. BOUNDED MODULES MAPS

Let A be a locally C^* -algebra and let E and F be two Hilbert A -modules. An A -module map $T : E \rightarrow F$ is said to be bounded if for each $p \in S(A)$, there is $K_p > 0$ such that $\bar{p}_F(Tx) \leq K_p \bar{p}_E(x)$ for all $x \in E$. The set of all bounded A -module maps from E to F is denoted by $B_A(E, F)$ and we write $B_A(E)$ for $B_A(E, E)$.

Clearly, for each $p \in S(A)$, the map \tilde{p} defined by

$$\tilde{p}(T) = \sup \{ \bar{p}_F(Tx); x \in E \text{ and } \bar{p}_E(x) \leq 1 \}, \quad T \in B_A(E, F)$$

is a seminorm on $B_A(E, F)$.

Proposition 3.1. *Let A be a locally C^* -algebra and let E and F be two Hilbert A -modules. Then we have:*

1. $B_A(E, F)$ with the topology determined by the family of seminorms $\{\tilde{p}\}_{p \in S(A)}$ is a complete locally convex space.
2. $B_A(E)$ with the topology determined by the family of seminorms $\{\tilde{p}\}_{p \in S(A)}$ is a complete locally m -convex algebra.

Proof. (1): Let $p, q \in S(A)$ with $p \geq q$ and let $S \in B_{A_p}(E_p, F_p)$. Since

$$\begin{aligned} \langle \sigma_{pq}^F(S(\sigma_p^E(x))), \sigma_{pq}^F(S(\sigma_p^E(x))) \rangle &= \pi_{pq}(\langle S(\sigma_p^E(x)), S(\sigma_p^E(x)) \rangle) \\ &\leq \|S\|_p \pi_{pq}(\langle \sigma_p^E(x), \sigma_p^E(x) \rangle) \text{ cf. [7, 2.8]} \\ &= \|S\|_p \langle \sigma_q^E(x), \sigma_q^E(x) \rangle \end{aligned}$$

for all $x \in E$, where $\|\cdot\|_p$ is the norm on $B_{A_p}(E_p, F_p)$, we can define $(\pi_{pq})_*(S) : E_q \rightarrow F_q$ by $(\pi_{pq})_*(S)(\sigma_q^E(x)) = \sigma_{pq}^F(S(\sigma_p^E(x)))$. It is easy to see that $(\pi_{pq})_*(S)$ is a bounded A_q -module map from E_q to F_q . Thus we have obtained a map $(\pi_{pq})_*$ from $B_{A_p}(E_p, F_p)$ to $B_{A_q}(E_q, F_q)$. Also it is easy to see that $\{B_{A_p}(E_p, F_p); (\pi_{pq})_*, p \geq q, p, q \in S(A)\}$ is an inverse system of Banach spaces.

We will show that the locally convex spaces $B_A(E, F)$ and $\lim_{p \leftarrow} B_{A_p}(E_p, F_p)$ are isomorphic.

Let $p \in S(A)$ and let $T \in B_A(E, F)$. Since $T(N_p^E) \subseteq N_p^F$ there is a unique linear map $T_p : E_p \rightarrow F_p$ such that $\sigma_p^F \circ T = T_p \circ \sigma_p^E$. Moreover, T_p is a bounded A_p -module map. Thus we can define a map $(\pi_p)_* : B_A(E, F) \rightarrow B_{A_p}(E_p, F_p)$ by $(\pi_p)_*(T) = T_p$, where $\sigma_p^F \circ T = T_p \circ \sigma_p^E$. Clearly $(\pi_p)_*$ is a continuous linear map and $(\pi_{pq})_* \circ (\pi_p)_* = (\pi_q)_*$ for all $p, q \in S(A)$ with $p \geq q$. Therefore we can define a map Φ from $B_A(E, F)$ to $\lim_{p \leftarrow} B_{A_p}(E_p, F_p)$ by $\Phi(T) = ((\pi_p)_*(T))_p$. It is not difficult to check that Φ is linear and $\|\Phi(T)\|_p = \tilde{p}(T)$ for all $T \in B_A(E, F)$. To show that Φ is surjective, let $(T_p)_p \in \lim_{p \leftarrow} B_{A_p}(E_p, F_p)$. Define $T : E \rightarrow F$ by $T(x) = (T_p(\sigma_p^E(x)))_p$. Since $\sigma_{pq}^F(T_p(\sigma_p^E(x))) = (\pi_{pq})_*(T_p)(\sigma_p^E(x)) = T_q(\sigma_q^E(x))$ for all $p, q \in S(A)$ with $p \geq q$, T is well-defined. It is not difficult to check that T is a bounded A -module map and $\Phi(T) = (T_p)_p$. Hence Φ is surjective.

Thus we showed that the topological spaces $B_A(E, F)$ and $\lim_{p \leftarrow} B_{A_p}(E_p, F_p)$ are isomorphic, and since $\lim_{p \leftarrow} B_{A_p}(E_p, F_p)$ is complete, $B_A(E, F)$ is complete.

(2): It is not difficult to check that \tilde{p} is a submultiplicative seminorm on $B_A(E)$ for all $p \in S(A)$ and $\{B_p(E_p); (\pi_{pq})_*, p \geq q, p, q \in S(A)\}$ is an inverse system of Banach algebras. Also it is easy to check that the map $\tilde{\Phi}$ from $B_A(E)$ to $\lim_{p \leftarrow} B_{A_p}(E_p)$ defined by $\tilde{\Phi}(T) = ((\pi_p)_*(T))_p$ is an isomorphism of topological algebras, and since $\lim_{p \leftarrow} B_{A_p}(E_p)$ is complete, the assertion is proved. \square

Remark 3.2. *If A is a locally C^* -algebra and E and F are Hilbert A -modules, then the locally convex spaces $B_A(E, F)$ and $\lim_{p \leftarrow} B_{A_p}(E_p, F_p)$ as well as the locally m -convex algebras $B_A(E)$ and $\lim_{p \leftarrow} B_{A_p}(E_p)$ can be identified.*

A map T from E to F is adjointable if there is a map T^* from F to E such that $\langle T(x), y \rangle = \langle x, T^*(y) \rangle$ for all x in E and for all y in F . Any adjointable map from E into F is a bounded A -module map (cf. [11]). The set of all adjointable maps from E into F is denoted by $L_A(E, F)$, and we write $L_A(E)$ for $L_A(E, E)$. For x in E and for y in F the map $\theta_{y,x} : E \rightarrow F$ defined by $\theta_{y,x}(z) = y \langle x, z \rangle$ is adjointable. The closed subspace of $L_A(E, F)$

generated by $\{\theta_{y,x}; x \in E, y \in F\}$ is denoted by $K_A(E, F)$, and we write $K_A(E)$ for $K_A(E, E)$. It is easy to verify that $(\pi_{pq})_*(L_{A_p}(E_p, F_p)) \subseteq L_{A_q}(E_q, F_q)$ and $(\pi_{pq})_*(K_{A_p}(E_p, F_p)) \subseteq K_{A_q}(E_q, F_q)$ for all $p, q \in S(A)$ with $p \geq q$. Then the restriction of Φ on $L_A(E, F)$ is exactly the same map as defined in Proposition 4.7 of [9]. Therefore the restriction of Φ on $L_A(E, F)$ is an isomorphism between the locally convex spaces $L_A(E, F)$ and $\lim_{p \leftarrow} L_{A_p}(E_p, F_p)$, and the restriction of Φ on $K_A(E, F)$ is an isomorphism between the locally convex spaces $K_A(E, F)$ and $\lim_{p \leftarrow} K_{A_p}(E_p, F_p)$ [9, Proposition 4.7]. Also the restriction of $\tilde{\Phi}$ on $L_A(E)$ is an isomorphism between the locally C^* -algebras $L_A(E)$ and $\lim_{p \leftarrow} L_{A_p}(E_p)$, and the restriction of $\tilde{\Phi}$ on $K_A(E)$ is an isomorphism between the locally C^* -algebras $K_A(E)$ and $\lim_{p \leftarrow} K_{A_p}(E_p)$ [9, Theorem 4.2].

In [9, Theorem 4.2], Phillips shows that the locally C^* -algebras $L_A(E)$ and $M(K_A(E))$, the multiplier algebra of $K_A(E)$, are isomorphic. We will prove here that the locally m -convex algebras $B_A(E)$ and $LM(K_A(E))$, the algebra of left multipliers of $K_A(E)$, are isomorphic.

If A is a locally C^* -algebra, we recall that a left multiplier of A is a linear map $l : A \rightarrow A$ such that $l(ab) = l(a)b$ for all a and b in A . We know that any left multiplier of a C^* -algebra is automatically continuous. We will show that this result is still valid for left multipliers of a locally C^* -algebra. Recall that in [11], Weinder showed that the multipliers of a locally C^* -algebra are automatically continuous.

Lemma 3.3. *Let a be an element of a locally C^* -algebra A . If $0 < \alpha < 1$, then there is an element u in A such that $a = u|a|^\alpha$, where $|a|^2 = aa^*$.*

Proof. We know that for each p in $S(A)$, there is an element u_p in A_p such that $\pi_p(a) = u_p|\pi_p(a)|^\alpha$. Moreover, $u_p = \lim_n \pi_p(a) \left(\frac{1}{n} + |\pi_p(a)|^2\right)^{\frac{-1}{2}} |\pi_p(a)|^{1-\alpha}$ (see, for example, [8, 1.4.6]).

To show that $(u_p)_p$ is a coherent sequence in A_p , $p \in S(A)$, let $p, q \in S(A)$ with $p \geq q$. Since π_{pq} preserves spectral functions, we have

$$\pi_{pq}(u_p) = \lim_n \pi_{pq} \left(\pi_p(a) \left(\frac{1}{n} + |\pi_p(a)|^2 \right)^{\frac{-1}{2}} |\pi_p(a)|^{1-\alpha} \right) = \lim_n \pi_q(a) \left(\frac{1}{n} + |\pi_q(a)|^2 \right)^{\frac{-1}{2}} |\pi_q(a)|^{1-\alpha} = u_q.$$

Hence $(u_p)_p$ is a coherent sequence in A_p , $p \in S(A)$. Let u in A be such that $\pi_p(u) = u_p$ for all $p \in S(A)$. Then, since $\pi_p(|a|^\alpha) = |\pi_p(a)|^\alpha$ for all $p \in S(A)$ (see [9] or [2]), we have $a = u|a|^\alpha$. \square

Proposition 3.4. *Any left multiplier of a locally C^* -algebra A is automatically continuous.*

Proof. Let l be a left multiplier of A , let $p \in S(A)$ and $a \in \ker(p)$. By Lemma 3.3, there is $u \in A$ such that $a = u|a|^{\frac{1}{2}}$, and then

$$p(l(a)) = p(l(u)|a|^{\frac{1}{2}}) \leq p(l(u))p(a)^{\frac{1}{2}}$$

whence we conclude that $l(a) \in \ker(p)$. Hence there is a unique linear map $l_p : A_p \rightarrow A_p$ such that $\pi_p \circ l = l_p \circ \pi_p$. Moreover, l_p is a left multiplier of A_p and so it is continuous (see, for example, [8, 3.12.2]). From these facts we conclude that l is continuous and the proposition is proved. \square

We consider on $LM(A)$, the set of all left multipliers of A , the seminorm topology (that is the topology determined by that family of seminorms $\{\tilde{p}\}_{p \in S(A)}$, where $\tilde{p}(l) = \sup\{p(l(a)), a \in A \text{ and } p(a) \leq 1\}$).

Theorem 3.5. *Let A be a locally C^* -algebra. Then we have:*

- (1) $LM(A)$ is a complete locally m -convex algebra.
- (2) If $A = \lim_{\lambda \in \Lambda \leftarrow} A_\lambda$ and the canonical maps $\pi_\lambda : A \rightarrow A_\lambda$ are all surjective, then $LM(A)$ and $\lim_{\lambda \in \Lambda \leftarrow} LM(A_\lambda)$ are isomorphic. the locally m -convex algebras

Proof. To prove this theorem we use the same arguments as in the proof of Theorem 3.14 of [9].

(1): Let $p, q \in S(A)$ with $p \geq q$. Since π_{pq} is surjective, there is a unique morphism $\pi''_{pq} : A''_p \rightarrow A''_q$ which extends π_{pq} and $\pi''_{pq}(LM(A_p)) \subseteq LM(A_q)$ (see, for example, [8, 3.7.7 and 3.12]). Then $\{LM(A_p); \pi''_{pq}|_{LM(A_p)}, p \geq q, p, q \in S(A)\}$ is an inverse system of Banach algebras. It is not difficult to check that the map $\Psi : LM(A) \rightarrow \lim_{p \leftarrow} LM(A_p)$ defined by $\Psi(l) = (l_p)_p$, where $\pi_p \circ l = l_p \circ \pi_p$ for all $p \in S(A)$, is an isomorphism of locally m -convex algebras.

(2): Exactly as in the proof of Theorem 3.14 of [9] we show that the inverse systems $\{LM(A_\lambda)\}_{\lambda \in \Lambda}$ and $\{LM(A_p)\}_{p \in S(A)}$ have the same inverse limit and thus the assertion is proved. \square

The following theorem is a generalization of Theorem 1.5 of [6] in the context of Hilbert modules over locally C^* -algebras.

Theorem 3.6. *Let A be a locally C^* -algebra and let E be a Hilbert A -module. Then the locally m -convex algebras $B_A(E)$ and $LM(K_A(E))$ are isomorphic.*

Proof. Let $p, q \in S(A)$ with $p \geq q$. Since $(\pi_{pq})_* (\theta_{y,x}) = \theta_{\sigma_{pq}(y), \sigma_{pq}(x)}$ for all $x, y \in E_p$, and since the map σ_{pq} from E_p to E_q is surjective, the morphism $(\pi_{pq})_*$ from $K_{A_p}(E_p)$ to $K_{A_q}(E_q)$ is surjective. Then according to Theorem 3.5 (2), the locally m -convex algebras $LM(K_A(E))$ and $\lim_{p \leftarrow} LM(K_{A_p}(E_p))$ are isomorphic.

For each $p \in S(A)$, the map $\Phi_p : B_{A_p}(E_p) \rightarrow LM(K_{A_p}(E_p))$ defined by $\Phi_p(T_p)(S_p) = T_p \circ S_p$ is an isometric isomorphism of Banach algebras [6, Theorem 1.5]. It is easy to check that $(\Phi_p)_p$ is an inverse system of isometric isomorphisms of Banach algebras. Then $\lim_{p \leftarrow} \Phi_p$ is an isomorphism of locally m -convex algebras from $\lim_{p \leftarrow} B_{A_p}(E_p)$ onto $\lim_{p \leftarrow} LM(K_{A_p}(E_p))$ and the theorem is proved. \square

We say that an element T of $B_A(E, F)$ is bounded in $B_A(E, F)$ if there is $M > 0$ such that $\tilde{p}(T) \leq M$ for all $p \in S(A)$ and denote by $b(B_A(E, F))$ the set of all bounded elements in $B_A(E, F)$. It is clear that the map

$\|\cdot\|_\infty : b(B_A(E, F)) \rightarrow [0, \infty)$ defined by

$$\|T\|_\infty = \sup\{\tilde{p}(T); p \in S(A)\}$$

is a norm on $b(B_A(E, F))$.

Theorem 3.7. *If E and F are Hilbert A -modules, then $b(B_A(E, F))$ is a Banach space in the norm $\|\cdot\|_\infty$. Moreover, $b(B_A(E, F))$ is isometrically isomorphic to $B_{b(A)}(b(E), b(F))$.*

Proof. Let $T \in b(B_A(E, F))$. Then, since

$$\bar{p}_F(Tx) \leq \|T\|_\infty \|x\|_\infty$$

for every $x \in b(E)$ and for every $p \in S(A)$, $T(b(E)) \subseteq b(F)$ and it is easy to see that the restriction $T|_{b(E)}$ of T on $b(E)$ is an element in $B_{b(A)}(b(E), b(F))$. Moreover, $\|T|_{b(E)}\| \leq \|T\|_\infty$. On the other hand, since $b(E)$ is dense in E [4, Proposition 3.1], and since

$$\langle T|_{b(E)}x, T|_{b(E)}x \rangle \leq \|T|_{b(E)}\|^2 \langle x, x \rangle$$

for every $x \in b(E)$ (cf. [7, 2.8]), we have $\|T\|_\infty \leq \|T|_{b(E)}\|$. Hence $\|T\|_\infty = \|T|_{b(E)}\|$. Define $\Psi : b(B_A(E, F)) \rightarrow B_{b(A)}(b(E), b(F))$ by

$$\Psi(T) = T|_{b(E)}.$$

Clearly Ψ is an isometric morphism from $b(B_A(E, F))$ to $B_{b(A)}(b(E), b(F))$. To show that Ψ is surjective, let $S \in L(b(E), b(F))$. Since

$$\langle Sx, Sx \rangle \leq \|S\|^2 \langle x, x \rangle$$

for all $x \in b(E)$ (cf. [7, 2.8]) and $b(E)$ is dense in E , S can be extended to a bounded A -module map \tilde{S} from E to F . Moreover, since $\tilde{p}(\tilde{S}) \leq \|S\|$ for all $p \in S(A)$, \tilde{S} is a bounded element in $B_A(E, F)$. Hence Ψ is surjective.

Thus we showed that $b(B_A(E, F))$ is isometrically isomorphic to $B_{b(A)}(b(E), b(F))$, and so $b(B_A(E, F))$ is a Banach space. \square

It is easy to check that an element T in $b(B_A(E, F))$ is adjointable if and only if $T|_{b(E)}$ is adjointable.

Remark 3.8. *The restriction of Ψ on $b(L_A(E, F))$ is an isometric isomorphism from $b(L_A(E, F))$ onto $L_{b(A)}(b(E), b(F))$.*

Knowing that for each $p \in S(A)$, \tilde{p} is a submultiplicative seminorm on $B_A(E)$ and $\tilde{p}|_{L_A(E)}$ is a C^* -seminorm on $L_A(E)$, it is easy to see that $\|\cdot\|_\infty$ is a submultiplicative norm on $b(B_A(E))$ and a C^* -norm on $b(L_A(E))$.

Corollary 3.9. *Let A be a locally C^* -algebra and let E be a Hilbert A -module. Then we have:*

- (1) $b(B_A(E))$ with the norm $\|\cdot\|_\infty$ is a Banach algebra which is isometrically isomorphic to $B_{b(A)}(b(E))$.
- (2) $b(L_A(E))$ with the norm $\|\cdot\|_\infty$ is a C^* -algebra which is isomorphic to $L_{b(A)}(b(E))$ [4, Theorem 3.3].

Proof. Putting $F = E$ in Theorem 3.7, it is easy to verify that Ψ is an isometric isomorphism from $b(B_A(E))$ onto $B_{b(A)}(b(E))$ and the restriction Ψ on $b(L_A(E))$ is an isomorphism from $b(L_A(E))$ onto $L_{b(A)}(b(E))$. \square

Remark 3.10. Let E and F be two Hilbert A -modules. In general, $b(K_A(E, F))$ is not isomorphic to $K_{b(A)}(b(E), b(F))$.

Example. Let $A = C(\mathbb{Z}^+)$, the $*$ -algebra of all complex valued functions on \mathbb{Z}^+ . It is not difficult to see that A is just $\prod_{n=1}^{\infty} \mathbb{C}$. Also it is not difficult to check that A with the topology determined by the family of C^* -seminorms $\{p_n\}_n$, where $p_n((a_n)_n) = \sup\{|a_k|; 1 \leq k \leq n\}$, is a locally C^* -algebra, and A_{p_n} can be identified with the product of the first n factors of A for each n .

Let $E = \prod_{n=1}^{\infty} \mathbb{C}^n$. We make E into a Hilbert A -module via $(\xi_n)_n (a_n)_n = (\xi_n a_n)_n$ and $\langle (\xi_n)_n, (\eta_n)_n \rangle = \langle (\xi_n, \eta_n)_n \rangle_n$, where $\langle \cdot, \cdot \rangle_n$ denotes the usual \mathbb{C} -inner product on \mathbb{C}^n . Clearly E is not finitely generated as Hilbert A -module. Moreover, E_{p_n} can be identified with the product of the first n factors of E for each n . Therefore, $L_{A_{p_n}}(E_{p_n}) = K_{A_{p_n}}(E_{p_n})$ for each n . This implies that $L_A(E) = K_A(E)$ [9, Example 4.9], and by Corollary 3.9, $b(K_A(E))$ is isomorphic with $L_{b(A)}(b(E))$.

Suppose that $b(K_A(E))$ is isomorphic with $K_{b(A)}(b(E))$. Then the C^* -algebras $K_{b(A)}(b(E))$ and $L_{b(A)}(b(E))$ are isomorphic. This implies that $b(E)$ is finitely generated as Hilbert $b(A)$ -module [10] and so E is finitely generated as Hilbert A -module, a contradiction. Therefore $b(K_A(E))$ is not isomorphic with $K_{b(A)}(b(E))$.

Remark 3.11. If A is a locally C^* -algebra then A is a Hilbert A -module with $\langle a, b \rangle = a^*b$, $a, b \in A$ and the locally C^* -algebras $L_A(A)$ and $M(A)$, where $M(A)$ is the set of all multipliers of A , are isomorphic [9]. Putting $E = A$ in Corollary 3.9, we deduce that the C^* -algebras $M(b(A))$ and $b(M(A))$ are isomorphic, a result obtained independently by Bhatt and J. Karia [1, Theorem 5.1] and the author [3, Theorem 2].

1. Bhatt S. J. and Karia, D. J. *Complete positivity, tensor products and C^* -nuclearity for inverse limits of C^* -algebras*, Proc. Indian Acad. Sci. (Math. Sci.) **101** (1991), 149–167.
2. Inoue A., *Locally C^* -algebras*, Mem. Faculty Sci. Kyushu Univ. Ser. A **25** (1971), 197–235.
3. Joița M. *Multipliers of locally C^* -algebras*. An. Univ. Bucuresti, Mat., **48**(1) (1999), 17–24.
4. ———, *On the bounded part of a Hilbert module over a locally C^* -algebra*, Period. Math. Hungar. **45**(1–2) (2002), 81–85.
5. Kasparov G. G., *Hilbert C^* -modules: theorems of Stinespring and Voiculescu*, J. Operator Theory **4** (1980), 133–150.
6. Lin H., *Bounded module maps and pure completely positive maps*, J. Operator Theory **26** (1991), 121–139.
7. Paschke W. L., *Inner product modules over B^* -algebras*, Trans. Amer. Math. Soc. **182** (1973), 443–468.
8. Pedersen G. K., *C^* -algebras and their automorphism groups*, Academic Press, London, New York, San Francisco, 1979.
9. Phillips N. C., *Inverse limits of C^* -algebras*, J. Operator Theory **19** (1988), 159–195.
10. Rieffel M. A., *Morita equivalence for operator algebras*, Proc. Symp. Pure Math. Amer. Math. Soc. **38**(1) (1982), 285–298.
11. Weinder J., *Topological invariants for generalized operator algebras*, Ph. D. Thesis Heidelberg, 1987.

M. Joița, Department of Mathematics, Faculty of Chemistry, University of Bucharest, Bd. Regina Elisabeta nr. 4–12, Bucharest, Romania, e-mail: mjoita@fmi.unibuc.ro