

ON (k, l) -RADIUS OF RANDOM GRAPHS

M. HORVÁTHOVÁ

ABSTRACT. We introduce the concept of (k, l) -radius of a graph and prove that for any fixed pair k, l the (k, l) -radius is equal to $2\binom{k}{2} - \binom{l}{2}$ for almost all graphs. Since for $k = 2$ and $l = 0$ the (k, l) -radius is equal to the diameter, our result is a generalization of the known fact that almost all graphs have diameter two.

All graphs in this note are finite, undirected and simple. As usual, by *distance between two vertices* in a graph we mean the minimum length of a path connecting them. Then the *diameter* is the maximum distance between two vertices. The *transmission* of the graph, also called a *distance of the graph*, is defined as the sum of distances between all pairs of vertices (for general properties of the distance see [4]). The concepts of diameter and distance were generalized by Goddard, Swart and Swart in [3] by introducing the *k-diameter* as follows. The distance of k vertices $d_k(v_1, v_2, \dots, v_k)$ is the sum of distances between all pairs of vertices from $\{v_1, v_2, \dots, v_k\}$. The *k-diameter* is the maximum distance of a set of k vertices. Hence the *2-diameter* is the usual diameter and if n is the order of the graph, the *n-diameter* is the distance of the graph.

In this note we use the definition of distance of a set of k vertices to define (k, l) -eccentricity and (k, l) -radius. We study (k, l) -radius of random graphs and determine the value of this parameter for almost all graphs in a probability space. We also discuss the relationship between the (k, l) -radius and the *k-diameter* of a graph.

Let S be a set of l vertices, $0 \leq l \leq k$. We define (k, l) -eccentricity of S , $e_{k,l}(S)$, as the maximum distance of k vertices u_1, u_2, \dots, u_k , such that $S \subseteq \{u_1, u_2, \dots, u_k\}$. In symbols,

$$e_{k,l}(S) = \max_T \{d_k(T), |T| = k, S \subseteq T \subseteq V(G)\}.$$

The (k, l) -radius, $rad_{k,l}(G)$, is the minimum (k, l) -eccentricity of a set of l vertices in G , that is

$$rad_{k,l}(G) = \min_S (e_{k,l}(S)) = \min_S \left(\max_{S \subseteq T \subseteq V(G)} d_k(T) \right)$$

where $|S| = l$, $|T| = k$.

Received October 4, 2004.

2000 *Mathematics Subject Classification*. Primary 05C12, 05C80, 05C99.

Key words and phrases. random graphs, distance in graphs.

We recall that the eccentricity $e(v)$ of a vertex v is the maximum distance to another vertex, the radius $\text{rad}(G)$ is the minimum eccentricity, whereas the diameter $\text{diam}(G)$ is the maximum eccentricity. From the definition of (k, l) -radius it follows that $\text{rad}_{2,1}(G)$ is the usual radius and $\text{rad}_{k,0}(G)$ is the k -diameter.

Now, consider the probability space in the following sense. Let p be a real number, $0 < p < 1$, and let n be an integer. By $G(n, p)$ we denote a class of labelled random graphs on n vertices, in which the probability of an edge equals p . More precisely, for every $u, v \in V(G)$, we have $P[uv \in E(G)] = p$. Hence $G(n, p)$ is a probability space the elements of which are the $2^{\binom{n}{2}}$ differently labelled graphs. We say that almost all graphs have property A if

$$\lim_{n \rightarrow \infty} P[G \in G(n, p) \text{ has property } A] = 1.$$

The space of random graphs is one of the random structures studied in connection with the 0-1 law. This law states that for many properties. The probability that a random structure satisfies the property is guaranteed to approach either 0 or 1. The 0-1 law for graphs was proved by Glebskij [2] and later on by Fagin [1]. Fagin's method is based on considering the following properties.

Let r and s be nonnegative integers. By $A_{r,s}$ we denote the property that for any disjoint sets of vertices X and Y , such that $|X| = r$ and $|Y| = s$, there exists a vertex z , $z \notin X \cup Y$ such that z is adjacent to every vertex of X and to no vertex of Y .

The following statements are well-known and their proofs can be found in the excellent survey by Winkler [5].

Theorem 1. [5] *For any fixed nonnegative integers r and s and a real number p , $0 < p < 1$ we have*

$$\lim_{n \rightarrow \infty} P[G \in G(n, p) \text{ has property } A_{r,s}] = 1.$$

Theorem 2. [5] *Let be $T = \{A_{r_1, s_1}, A_{r_2, s_2}, \dots, A_{r_k, s_k}\}$ for some $k \geq 0$. Then almost all graphs have all the properties of T .*

From the fact that almost every graph has property $A_{2,0}$ (the distance of every pair of vertices is at most 2) and $A_{0,1}$ (the graph is not complete) we have:

Corollary 3. *For any fixed real p , $0 < p < 1$, almost all graphs are connected and have diameter 2.*

Now we are in a position to prove the main statement of this note.

Theorem 4. *Let k, l be nonnegative integers, $l \leq k$. For any fixed real p , $0 < p < 1$, almost all graphs G have*

$$\text{rad}_{k,l}(G) = 2 \binom{k}{2} - \binom{l}{2}.$$

Proof. Let L be a set of l vertices in a graph $G \in G(n, p)$. Let p_n denote the probability $P[G \in G(n, p) \text{ has diameter } 2]$. Then with the same probability p_n it

holds

$$(1) \quad e_{k,l}(L) \leq d_l(L) + 2l(k-l) + 2\binom{k-l}{2}.$$

By Corollary 3 $\lim_{n \rightarrow \infty} p_n = 1$, so that (1) holds for almost all graphs $G \in G(n, p)$. Now we prove that for almost all graphs

$$(2) \quad e_{k,l}(L) \geq d_l(L) + 2l(k-l) + 2\binom{k-l}{2}.$$

To do this, it suffices to prove that for almost all graphs there exist $k-l$ vertices from $V(G) \setminus L$ that are mutually nonadjacent and that are adjacent to no vertex of L . Let $T = \{A_{0,l}, A_{0,l+1}, \dots, A_{0,k-1}\}$. By Theorem 2, $\lim_{n \rightarrow \infty} P[G \in G(n, p) \text{ has all properties of } T] = 1$, i.e. almost all graphs G have all properties of T .

1. Let $L_l = L$. Property $A_{0,l}$ says that there exists a vertex $z_{l+1} \in V(G) \setminus L$ that is adjacent to no vertex of L .
2. For $i = l+1, l+2, \dots, k-1$ we define L_i inductively by $L_i = L_{i-1} \cup z_i$. Then property $A_{0,i}$ implies that there exists a vertex z_{i+1} that is adjacent to no vertex of L_i .

Hence, we have $k-l$ vertices $z_{l+1}, z_{l+2}, \dots, z_k$ that are mutually nonadjacent and are adjacent to no vertex of L_l , which proves (2). Thus, from (1) and (2) we have that for almost all graphs

$$(3) \quad e_{k,l}(L) = d_l(L) + 2l(k-l) + 2\binom{k-l}{2}.$$

Since $\text{rad}_{k,l}(G) = \min_L e_{k,l}(L)$, the radius is minimal whenever $d_l(L)$ is minimal, (see (3)). We show that in almost all graphs $G \in G(n, p)$ there exists a set L' of l vertices, such that $d_l(L') = \binom{l}{2}$. In other words, we show that there is a set L' of l mutually adjacent vertices. Let $T' = \{A_{1,0}, A_{2,0}, \dots, A_{l-1,0}\}$. Then almost all graphs have all properties of T' , since by Theorem 2 $\lim_{n \rightarrow \infty} P[G \in G(n, p) \text{ has all properties of } T'] = 1$.

1. Let L'_1 be a set containing a single vertex of G , say $L'_1 = \{z'_1\}$. Then $|L'_1| = 1$ and $A_{1,0}$ says that there exists a vertex z'_2 that is adjacent to z'_1 .
2. For $i = 2, 3, \dots, l-1$ let L'_i be a set of vertices, such that $L'_i = L'_{i-1} \cup z'_i$. Then $|L'_i| = i$ and $A_{i,0}$ implies that there exists a vertex z'_{i+1} that is adjacent to all vertices of L'_i .

In this way we obtain a set $L' = L'_l$ of l vertices that are mutually adjacent, so that $d_l(L') = \binom{l}{2}$. Since $d_l(L)$ cannot be less than $\binom{l}{2}$ for any set of l vertices, we have

$$\text{rad}_{k,l} = \binom{l}{2} + 2l(k-l) + 2\binom{k-l}{2} = 2\binom{k}{2} - \binom{l}{2}$$

for almost all graphs $G \in G(n, p)$, as required. □

Setting $l = 0$ in Theorem 2 we obtain:

Corollary 5. *For any $k \geq 0$ and for almost all graphs G we have*

$$\text{diam}_k(G) = k(k - 1).$$

It is obvious that Corollary 5 generalizes Corollary 3. Further, setting $k = 2$ and $l = 1$ we obtain:

Corollary 6. *For almost all graphs G we have $\text{rad}(G) = 2$.*

REFERENCES

1. Fagin R., *Probabilities on finite models*, J. Symbolic Logic **41**(1) (1976), 50–58.
2. Glebskij Y. V., Kogan D. I., Liogon'kii M. I. and Talanov V. A., *Range and degree of reliability of formulas in the restricted predicate calculus*, Kibernetika **2** (1969), 17–28.
3. Goddard W., Swart Ch. S. and Swart H. C., *On the graphs with maximum distance on k -diameter*, (preprint).
4. Šoltés L., *Transmission in graphs: a bound and vertex removing*, Math. Slovaca **41** (1991), 1–16.
5. Winkler P., *Random structures and zero-one laws*, Finite and Infinite Combinatorics of Sets and Logic, in: Sauer N., Woodrow R. and Sands B. eds., NATO Advanced Science Institutes Series, Kluwer Academic Publishers, Kluwer Academic, Dordrecht 1993, 283–288.

M. Horváthová, Department of Mathematics Slovak University of Technology, Radlinského 11, 813 68 Bratislava, Slovak Republic, *e-mail*: mhorvath@math.sk