

# A NOTE ON SOME NEW FRACTIONAL RESULTS INVOLVING CONVEX FUNCTIONS

Z. DAHMANI

**ABSTRACT.** In this paper, we establish some new integral inequalities for convex functions by using the Riemann-Liouville operator of non integer order. For our results some classical integral inequalities can be deduced as some special cases.

## 1. INTRODUCTION

The integral inequalities play a fundamental role in the theory of differential equations. Much significant development in this area has been established for the last two decades. For details we refer to [10, 12, 14, 15] and the references therein. Moreover, the study of fractional type inequalities is also of a great importance. For further information and applications we refer the reader to [1, 13]. Let us introduce now some results that have inspired our work. We begin by the paper of Ngo et al. [11], in which the authors proved that

$$(1) \quad \int_0^1 f^{\delta+1}(\tau) d\tau \geq \int_0^1 \tau^\delta f(\tau) d\tau$$

---

Received February 6, 2012; revised July 19, 2012.

2010 *Mathematics Subject Classification.* Primary 26D10, 26A33.

*Key words and phrases.* Convex function; Integral inequalities; Riemann-Liouville operator; Qi inequality.



Go back

Full Screen

Close

Quit

and

$$(2) \quad \int_0^1 f^{\delta+1}(\tau) d\tau \geq \int_0^1 \tau f^\delta(\tau) d\tau,$$

where  $\delta > 0$  and  $f$  is a positive continuous function on  $[0, 1]$  such that

$$\int_x^1 f(\tau) d\tau \geq \int_x^1 \tau d\tau, x \in [0, 1].$$

Then, in [8], W. J. Liu, G. S. Cheng and C. C. Li established the following result

$$(3) \quad \int_a^b f^{\alpha+\beta}(\tau) d\tau \geq \int_a^b (\tau - a)^\alpha f^\beta(\tau) d\tau,$$

provided that  $\alpha > 0$ ,  $\beta > 0$  and  $f$  is a positive continuous function on  $[a, b]$  satisfying

$$\int_x^b f^\gamma(\tau) d\tau \geq \int_x^b (\tau - a)^\gamma d\tau; \gamma := \min(1, \beta), x \in [a, b].$$

In [9], the following two theorems were proved.

**Theorem 1.1.** *Let  $f$  and  $h$  be two positive continuous functions on  $[a, b]$  with  $f \leq h$  on  $[a, b]$  such that  $\frac{f}{h}$  is decreasing and  $f$  is increasing. Assume that  $\phi$  is a convex function  $\phi; \phi(0) = 0$ . Then the inequality*

$$(4) \quad \frac{\int_a^b f(\tau) d\tau}{\int_a^b h(\tau) d\tau} \geq \frac{\int_a^b \phi(f(\tau)) d\tau}{\int_a^b \phi(h(\tau)) d\tau}$$

holds.

And




Go back

Full Screen

Close

Quit



**Theorem 1.2.** Let  $f, g$  and  $h$  be three positive continuous functions on  $[a, b]$  with  $f \leq h$  on  $[a, b]$  such that  $\frac{f}{h}$  is decreasing and  $f$  and  $g$  are increasing. Assume that  $\phi$  is a convex function  $\phi; \phi(0) = 0$ . Then the inequality

$$(5) \quad \frac{\int_a^b f(\tau) d\tau}{\int_a^b h(\tau) d\tau} \geq \frac{\int_a^b \phi(f(\tau))g(\tau) d\tau}{\int_a^b \phi(h(\tau))g(\tau) d\tau}$$

holds.

Many researchers have given considerable attention to (1), (2) and (3) and a number of extensions, generalizations and variants have appeared in the literature, (e.g. [2, 3, 4, 5, 7, 14]).

The purpose of this paper is to generalize some classical integral inequalities of [9] using the Riemann-Liouville integral operator. For our results Theorem 1.1 and Theorem 1.2 can be deduced as some special cases.

## 2. PRELIMINARIES

Let us introduce some definitions and properties concerning the Riemann-Liouville fractional integral operator.

**Definition 1.** The Riemann-Liouville fractional integral operator of order  $\alpha \geq 0$ , for a continuous function  $f$  on  $[a, b]$ , is defined as

$$(6) \quad \begin{aligned} J^\alpha[f(t)] &= \frac{1}{\Gamma(\alpha)} \int_a^t (t - \tau)^{\alpha-1} f(\tau) d\tau; & \alpha > 0, \quad a < t \leq b, \\ J^0[f(t)] &= f(t), \end{aligned}$$

where  $\Gamma(\alpha) := \int_0^\infty e^{-u} u^{\alpha-1} du$ .




Go back

Full Screen

Close

Quit



For the convenience of establishing the results we give the semigroup property

$$(7) \quad J^\alpha J^\beta[f(t)] = J^{\alpha+\beta}[f(t)], \quad \alpha \geq 0, \quad \beta \geq 0,$$

which implies the commutative property

$$(8) \quad J^\alpha J^\beta[f(t)] = J^\beta J^\alpha[f(t)].$$

For more details one can consult [6, 13].

### 3. MAIN RESULTS

**Theorem 3.1.** *Let  $f$  and  $h$  be two positive continuous functions on  $[a, b]$  and  $f \leq h$  on  $[a, b]$ . If  $\frac{f}{h}$  is decreasing and  $f$  is increasing on  $[a, b]$ , then for any convex function  $\phi$ ;  $\phi(0) = 0$ , the inequality*

$$(9) \quad \frac{J^\alpha[f(t)]}{J^\alpha[h(t)]} \geq \frac{J^\alpha[\phi(f(t))]}{J^\alpha[\phi(h(t))]}, \quad a < t \leq b, \quad \alpha > 0$$

is valid.

*Proof.* The function  $\phi$  is convex with  $\phi(0) = 0$ . Then the function  $\frac{\phi(x)}{x}$  is increasing. Since  $f$  is increasing, then  $\frac{\phi(f(x))}{f(x)}$  is also increasing. This and the fact that  $\frac{f(x)}{h(x)}$  is decreasing yield

$$(10) \quad \frac{\phi(f(\tau))}{f(\tau)} \frac{f(\rho)}{h(\rho)} + \frac{\phi(f(\rho))}{f(\rho)} \frac{f(\tau)}{h(\tau)} - \frac{\phi(f(\rho))}{f(\rho)} \frac{f(\rho)}{h(\rho)} - \frac{\phi(f(\tau))}{f(\tau)} \frac{f(\tau)}{h(\tau)} \geq 0$$

for all  $\tau, \rho \in [a, t]$ ,  $a < t \leq b$ .



Go back

Full Screen

Close

Quit



Hence, we can write

$$(11) \quad \begin{aligned} & \frac{\phi(f(\tau))}{f(\tau)} f(\rho) h(\tau) + \frac{\phi(f(\rho))}{f(\rho)} f(\tau) h(\rho) \\ & - \frac{\phi(f(\rho))}{f(\rho)} f(\rho) h(\tau) - \frac{\phi(f(\tau))}{f(\tau)} f(\tau) h(\rho) \geq 0 \end{aligned}$$

for all  $\tau, \rho \in [a, t]$ ,  $a < t \leq b$ .

Now, multiplying both sides of (11) by  $\frac{(t-\tau)^{\alpha-1}}{\Gamma(\alpha)}$ , then integrating the resulting inequality with respect to  $\tau$  over  $[a, t]$ ,  $a < t \leq b$ , we get

$$(12) \quad \begin{aligned} & f(\rho) J^\alpha \left[ \frac{\phi(f(t))}{f(t)} h(t) \right] + \frac{\phi(f(\rho))}{f(\rho)} h(\rho) J^\alpha [f(t)] \\ & - \frac{\phi(f(\rho))}{f(\rho)} f(\rho) J^\alpha [h(t)] - h(\rho) J^\alpha \left[ \frac{\phi(f(t))}{f(t)} f(t) \right] \geq 0. \end{aligned}$$

With the same argument as before, we obtain

$$(13) \quad J^\alpha [f(t)] J^\alpha \left[ \frac{\phi(f(t))}{f(t)} h(t) \right] - J^\alpha [h(t)] J^\alpha \left[ \frac{\phi(f(t))}{f(t)} f(t) \right] \geq 0.$$

Since  $f \leq h$  on  $[a, b]$ , then using the fact that the function  $\frac{\phi(x)}{x}$  is increasing, we can write

$$(14) \quad \frac{\phi(f(\tau))}{f(\tau)} \leq \frac{\phi(h(\tau))}{h(\tau)}, \quad \tau \in [a, t], \quad a < t \leq b.$$

This implies that

$$(15) \quad \frac{(t-\tau)^{\alpha-1}}{\Gamma(\alpha)} h(\tau) \frac{\phi(f(\tau))}{f(\tau)} \leq \frac{(t-\tau)^{\alpha-1}}{\Gamma(\alpha)} h(\tau) \frac{\phi(h(\tau))}{h(\tau)},$$




Go back

Full Screen

Close

Quit



where  $\tau \in [a, t]$ ,  $a < t \leq b$ .

Integrating both sides of (15) with respect to  $\tau$  over  $[a, t]$ ,  $a < t \leq b$ , yields

$$(16) \quad J^\alpha \left[ \frac{\phi(f(t))}{f(t)} h(t) \right] \leq J^\alpha \left[ \frac{\phi(h(t))}{h(t)} h(t) \right].$$

Hence, thanks to (13) and (16), we obtain (9). □

**Remark 3.2.** Applying Theorem 3.1 for  $\alpha = 1$ ,  $t = b$ , we obtain Theorem 1.1.

We further have the following theorem.

**Theorem 3.3.** *Let  $f$  and  $h$  be two positive continuous functions on  $[a, b]$  and  $f \leq h$  on  $[a, b]$ . If  $\frac{f}{h}$  is decreasing and  $f$  is increasing on  $[a, b]$ , then for any convex function  $\phi$ ;  $\phi(0) = 0$ , we have*

$$(17) \quad \frac{J^\alpha[f(t)]J^\omega[\phi(h(t))] + J^\omega[f(t)]J^\alpha[\phi(h(t))]}{J^\alpha[h(t)]J^\omega[\phi(f(t))] + J^\omega[h(t)]J^\alpha[\phi(f(t))]} \geq 1,$$

where  $\alpha > 0$ ,  $\omega > 0$ ,  $a < t \leq b$ .

*Proof.* The relation (12) allows us to obtain

$$(18) \quad J^\omega[f(t)]J^\alpha \left[ \frac{\phi(f(t))}{f(t)} h(t) \right] + J^\omega \left[ \frac{\phi(f(t))}{f(t)} h(t) \right] J^\alpha[f(t)] - J^\omega \left[ \frac{\phi(f(t))}{f(t)} f(t) \right] J^\alpha[h(t)] - J^\omega[h(t)]J^\alpha \left[ \frac{\phi(f(t))}{f(t)} f(t) \right] \geq 0.$$

Since  $f \leq h$  on  $[a, b]$  and using the fact that the function  $\frac{\phi(x)}{x}$  is increasing, then thanks to (14), we obtain

$$(19) \quad \frac{(t - \tau)^{\omega-1}}{\Gamma(\omega)} h(\tau) \frac{\phi(f(\tau))}{f(\tau)} \leq \frac{(t - \tau)^{\omega-1}}{\Gamma(\omega)} h(\tau) \frac{\phi(h(\tau))}{h(\tau)},$$



Go back

Full Screen

Close

Quit

where  $\tau \in [a, t]$ ,  $a < t \leq b$ . And then,

$$(20) \quad J^\omega \left[ \frac{\phi(f(t))}{f(t)} h(t) \right] \leq J^\omega \left[ \frac{\phi(h(t))}{h(t)} h(t) \right].$$

Hence, thanks to (16), (18) and (20), we get (17). □

**Remark 3.4.** (i) Applying Theorem 3.3 for  $\alpha = \omega$ , we obtain Theorem 3.1.

(ii) Applying Theorem 3.3 for  $\alpha = \omega = 1$ ,  $t = b$ , we obtain Theorem 1.1.

Another result which generalizes Theorem 1.2 is described in the following theorem.

**Theorem 3.5.** *Let  $f$ ,  $h$  and  $g$  be three positive continuous functions and  $f \leq h$  on  $[a, b]$ . Suppose that  $\frac{f}{h}$  is decreasing,  $f$  and  $g$  are increasing on  $[a, b]$  and  $\phi$  is a convex function,  $\phi(0) = 0$ . Then, for any  $\alpha > 0$ ,  $a < t \leq b$ , we have*

$$(21) \quad \frac{J^\alpha[f(t)]}{J^\alpha[h(t)]} \geq \frac{J^\alpha[\phi(f(t))g(t)]}{J^\alpha[\phi(h(t))g(t)]}.$$

*Proof.* Let  $\tau, \rho \in [a, t]$ ,  $a < t \leq b$ . We have

$$(22) \quad \begin{aligned} & \frac{\phi(f(\tau))g(\tau)}{f(\tau)} f(\rho)h(\tau) + \frac{\phi(f(\rho))g(\rho)}{f(\rho)} f(\tau)h(\rho) \\ & - \frac{\phi(f(\rho))g(\rho)}{f(\rho)} f(\rho)h(\tau) - \frac{\phi(f(\tau))g(\tau)}{f(\tau)} f(\tau)h(\rho) \geq 0. \end{aligned}$$

Hence we can write

$$(23) \quad \begin{aligned} & f(\rho)J^\alpha \left[ \frac{\phi(f(t))g(t)}{f(t)} h(t) \right] + \frac{\phi(f(\rho))g(\rho)}{f(\rho)} h(\rho)J^\alpha[f(t)] \\ & - \frac{\phi(f(\rho))g(\rho)}{f(\rho)} f(\rho)J^\alpha[h(t)] - h(\rho)J^\alpha \left[ \frac{\phi(f(t))g(t)}{f(t)} f(t) \right] \geq 0. \end{aligned}$$



Go back

Full Screen

Close

Quit



Therefore,

$$(24) \quad J^\alpha[f(t)]J^\alpha\left[\frac{\phi(f(t))g(t)}{f(t)}h(t)\right] - J^\alpha[h(t)]J^\alpha[\phi(f(t))g(t)] \geq 0.$$

On the other hand, we have

$$(25) \quad \frac{(t-\tau)^{\alpha-1}}{\Gamma(\alpha)}h(\tau)\frac{\phi(f(\tau))g(\tau)}{f(\tau)} \leq \frac{(t-\tau)^{\alpha-1}}{\Gamma(\alpha)}h(\tau)\frac{\phi(h(\tau))g(\tau)}{h(\tau)},$$

where  $\tau \in [a, t]$ ,  $a < t \leq b$ . Consequently,

$$(26) \quad J^\alpha\left[\frac{\phi(f(t))g(t)}{f(t)}h(t)\right] \leq J^\alpha[\phi(h(t))g(t)],$$

and so,

$$(27) \quad J^\alpha[f(t)]J^\alpha\left[\frac{\phi(f(t))g(t)}{f(t)}h(t)\right] \leq J^\alpha[f(t)]J^\alpha[\phi(h(t))g(t)].$$

Hence, thanks to (24) and (27) we obtain (21). □

**Remark 3.6.** It is clear that Theorem 1.2 would follow as a special case of Theorem 3.5 when  $\alpha = 1$  and  $t = b$ .

Another result which generalizes Theorem 3.5 is described in the following theorem.

**Theorem 3.7.** Let  $f, h$  and  $g$  be three positive continuous functions and  $f \leq h$  on  $[a, b]$ . Suppose that  $\frac{f}{h}$  is decreasing,  $f$  and  $g$  are increasing on  $[a, b]$  and  $\phi$  is a convex function,  $\phi(0) = 0$ . Then, for any  $\alpha > 0, \omega > 0, a < t \leq b$ , we have

$$(28) \quad \frac{J^\alpha[f(t)]J^\omega[\phi(h(t))g(t)] + J^\omega[f(t)]J^\alpha[\phi(h(t))g(t)]}{J^\alpha[h(t)]J^\omega[\phi(f(t))g(t)] + J^\omega[h(t)]J^\alpha[\phi(f(t))g(t)]} \geq 1.$$



Go back

Full Screen

Close

Quit



*Proof.* Using (23), we can write

$$(29) \quad \begin{aligned} & J^\omega [f(t)] J^\alpha \left[ \frac{\phi(f(t))g(t)}{f(t)} h(t) \right] + J^\omega \left[ \frac{\phi(f(t))g(t)}{f(t)} h(t) \right] J^\alpha [f(t)] \\ & - J^\omega \left[ \frac{\phi(f(t))g(t)}{f(t)} f(t) \right] J^\alpha [h(t)] - J^\omega [h(t)] J^\alpha \left[ \frac{\phi(f(t))g(t)}{f(t)} f(t) \right] \geq 0. \end{aligned}$$

Then, using the fact that the function  $\frac{\phi(x)g(x)}{x}$  is increasing and the hypothesis  $f \leq h$  on  $[a, b]$ , we obtain

$$(30) \quad J^k \left[ \frac{\phi(f(t))g(t)}{f(t)} h(t) \right] \leq J^k \left[ \frac{\phi(h(t))g(t)}{h(t)} h(t) \right], \quad k = \alpha, \omega.$$

Hence, thanks to (29) and (30), we get (28).  $\square$

**Remark 3.8.** It is clear that Theorem 3.5 would follow as a special case of Theorem 3.7 when  $\alpha = \beta$ .

**Acknowledgment.** This paper is supported by l'ANDRU, Agence Nationale pour le Developpement de la Recherche Universitaire and UMAB University of Mostaganem; (PNR Project 2011-2013).

1. Anastassiou G. A., *Fractional differentiation inequalities*, Springer, 2009.
2. Bougoufa L., *An integral inequality similar to Qi inequality*, JIPAM. J. Inequal. Pure Appl. Math., **6(1)** (2005), Art. 27.
3. Boukerrioua K. and Guezane Lakoud A., *On an open question regarding an integral inequality*, JIPAM. J. Inequal. Pure Appl. Math., **8(3)** (2007), Art. 77.
4. Dahmani Z. and Bedjaoui N., *Some generalized integral inequalities*, J. Advan. Resea. Appl. Math., **3(4)** (2011), 58–66.




Go back

Full Screen

Close

Quit



5. Dahmani Z., Metakkel Elard H., *Generalizations of some integral inequalities using Riemann-Liouville operator*, Int. J. Open Problems Compt. Math., **4(4)** (2011), 40–46.
6. Gorenflo R. and Mainardi F., *Fractional calculus: integral and differential equations of fractional order*, Springer Verlag, Wien (1997), 223–276.
7. Liu W. J., Li C. C. and Dong J. W., *On an open problem concerning an integral inequality*, JIPAM. J. Inequal. Pure Appl. Math., **8(3)** (2007), Art. 74.
8. Liu W. J., Cheng G. S. and Li C. C., *Further development of an open problem concerning an integral inequality*, JIPAM. J. Inequal. Pure Appl. Math., **9(1)** (2008), Art. 14.
9. Liu W. J., Ngo Q. A. and Huy V. N., *Several interesting integral inequalities*, Journal of Math. Inequal., **3(2)** (2009), 201–212.
10. Mitrinovic D. S., Pecaric J. E. and Fink A. M., *Classical and new inequalities in Analysis*, Kluwer Academic Publishers, Dordrecht, 1993.
11. Ngo Q. A., Thang D. D., Dat T. T., and Tuan D. A., *Notes on an integral inequality*, JIPAM. J. Inequal. Pure Appl. Math., **7(4)** (2006), Art. 120.
12. Pachpatte B. G., *Mathematical inequalities*, North Holland Mathematical Library, Vol. 67 (2005).
13. Podlubný I., *Fractional Differential Equations*, Academic Press, San Diego, 1999.
14. Qi F., *Several integral inequalities*, JIPAM. J. Inequal. Pure Appl. Math., **1(2)** (2000), Art. 19.
15. Sarikaya M. Z., Yildirim H. and Saglam A., *On Hardy type integral inequality associated with the generalized translation*, Int. J. Contemp. Math. Sci., **1(7)** (2006), 333–340.

Z. Dahmani, Laboratory LPAM, FSEI, UAMB, University of Mostaganem, Algeria,

e-mail: [zzdahmani@yahoo.fr](mailto:zzdahmani@yahoo.fr)



Go back

Full Screen

Close

Quit