

Hyperelliptic Plane Curves of Type $(d, d - 2)$

Fumio Sakai Mohammad Salem* Keita Tono

*Department of Mathematics, Graduate School of Science and Engineering
Saitama University, Shimo-Okubo 255, Sakura-ku, Saitama 338-8570, Japan
e-mail: fsakai@rimath.saitama-u.ac.jp*

*Department of Mathematics, Faculty of Science
Sohag University, Sohag 82524, Egypt
e-mail: abuelhassan@yahoo.com*

*Department of Mathematics, Graduate School of Science and Engineering
Saitama University, Shimo-Okubo 255, Sakura-ku, Saitama 338-8570, Japan
e-mail: ktono@rimath.saitama-u.ac.jp*

Abstract. In [7], we classified and constructed all rational plane curves of type $(d, d - 2)$. In this paper, we generalize these results to irreducible plane curves of type $(d, d - 2)$ with positive genus.

MSC 2000: 14H50, 14E07

1. Introduction

Let $C \subset \mathbf{P}^2 = \mathbf{P}^2(\mathbf{C})$ be a plane curve of degree d . We call C a *plane curve of type (d, ν)* if the maximal multiplicity of singular points on C is equal to ν . A unibranch singularity is called a *cuspidal*. Rational cuspidal plane curves of type $(d, d - 2)$ and $(d, d - 3)$ were classified by Flenner-Zaidenberg [5], [6] (see also [4], [8] for some cases). In [7], we classified rational plane curves of type $(d, d - 2)$ with arbitrary singularities. In order to describe a multibranch singularity P , we introduced the notion of the system of the multiplicity sequences $\underline{m}_P(C)$ (see

*partially supported by Post Doctoral Fellowship for Foreign Researchers, 1705292, JSPS

Section 2). We denote by $\text{Data}(C)$, the collection of such systems of multiplicity sequences. The purpose of this paper is to complete the classification of irreducible plane curves of type $(d, d - 2)$ with positive genus g . We remark that if $g \geq 2$, then C is a hyperelliptic curve, for the projection from the singular point of multiplicity $d - 2$ induces a double covering of C over \mathbf{P}^1 .

Theorem 1. *Let C be a plane curve of type $(d, d - 2)$ with genus g . Let $Q \in C$ be the singular point of multiplicity $d - 2$. Then, we have*

$$(i) \text{Data}(C) = \left[\underline{m}_Q(C), \binom{1}{1}_{b_1}, \dots, \binom{1}{1}_{b_n}, (2_{b_{n+1}}), \dots, (2_{b_{n+n'}}) \right], \text{ where}$$

$$\underline{m}_Q(C) = \left\{ \left(\begin{array}{c} k_1 \\ k'_1 \\ \vdots \\ k_s \\ k'_s \\ k_{s+1} \\ \vdots \\ k_N \end{array} \right) \left(\begin{array}{c} 1 \\ 1 \\ \vdots \\ 1 \\ 1 \\ 2_{a_{s+1}} \\ \vdots \\ 2_{a_N} \end{array} \right) \right\}$$

and the following conditions are satisfied:

- (1) $\sum_{h=1}^N k_h + \sum_{h'=1}^s k'_{h'} = d - 2$ and $\sum_{i=1}^N a_i + \sum_{j=1}^{n+n'} b_j = d - g - 2$, where $a_i \geq 0$
($a_i > 0$ for $i = 1, \dots, s$), $b_j > 0$,
- (2) we have $n, n', s \geq 0$ and $n' + s' \leq 2g + 2$, where $s' = \#\{j | a_{s+j} > 0\}$,
- (3) for $i = 1, 2, \dots, s$, if $k'_i = k_i$, then $a_i \geq k_i$ and if $k'_i > k_i$, then $a_i = k_i$,
- (4) for $i = s + 1, \dots, N$, if $a_i > 0$, then either k_i is even and $a_i \geq k_i/2$ or k_i is odd and $a_i = (k_i - 1)/2$.

Note that the N is the number of the different tangent lines to C at Q .

- (ii) $\text{Data}(C)$ can be derived from Degtyarev's 2-formula $T(C)$ defined for the defining equation of C (see Proposition 10 for details).

Corollary. *Let C be an irreducible plane curve of type $(d, d - 2)$ with genus g .*

- (i) *If C has only cusps, then C has the following data ($b_i > 0, k > 0, j \geq 0$):*

Class	$\text{Data}(C)$	
(a)	$[(k), (2_{b_1}), \dots, (2_{b_{n'}})]$	$(k = g + \sum_{i=1}^{n'} b_i)$ $(n' \leq 2g + 2)$
(b)	$[(2k + 1, 2_k), (2_{b_1}), \dots, (2_{b_{n'}})]$	$(k + 1 = g + \sum_{i=1}^{n'} b_i)$ $(n' \leq 2g + 1)$
(c)	$[(2k, 2_{k+j}), (2_{b_1}), \dots, (2_{b_{n'}})]$	$(k = g + j + \sum_{i=1}^{n'} b_i)$ $(n' \leq 2g + 1)$

(ii) If C has only bibranched singularities, then C has the following data $(b_i > 0, k > 0, r > 0, j \geq 0, l \geq 0)$:

Class	Data(C)
(e)	$\left[\binom{k}{k} \binom{1}{1}_{k+j}, \binom{1}{1}_{b_1}, \dots, \binom{1}{1}_{b_n} \right]$ $(k = g + j + \sum_{i=1}^n b_i)$
(f)	$\left[\binom{k}{k+r} \binom{1}{1}_k, \binom{1}{1}_{b_1}, \dots, \binom{1}{1}_{b_n} \right]$ $(k + r = g + \sum_{i=1}^n b_i)$
(aa)	$\left[\binom{k}{r}, \binom{1}{1}_{b_1}, \dots, \binom{1}{1}_{b_n} \right]$ $(k + r = g + \sum_{i=1}^n b_i)$
(ab)	$\left[\left\{ \binom{2k+1}{r} \binom{2k}{r} \right\}, \binom{1}{1}_{b_1}, \dots, \binom{1}{1}_{b_n} \right]$ $(k + r + 1 = g + \sum_{i=1}^n b_i)$
(ac)	$\left[\left\{ \binom{2k}{r} \binom{2k+j}{r} \right\}, \binom{1}{1}_{b_1}, \dots, \binom{1}{1}_{b_n} \right]$ $(k + r = g + j + \sum_{i=1}^n b_i)$
(bb)	$\left[\left\{ \binom{2k+1}{2r+1} \binom{2k}{2r} \right\}, \binom{1}{1}_{b_1}, \dots, \binom{1}{1}_{b_n} \right]$ $(k + r + 2 = g + \sum_{i=1}^n b_i)$
(bc)	$\left[\left\{ \binom{2k+1}{2r} \binom{2k}{2r+l} \right\}, \binom{1}{1}_{b_1}, \dots, \binom{1}{1}_{b_n} \right]$ $(k + r + 1 = g + l + \sum_{i=1}^n b_i)$
(cc)	$\left[\left\{ \binom{2k}{2r} \binom{2k+j}{2r+l} \right\}, \binom{1}{1}_{b_1}, \dots, \binom{1}{1}_{b_n} \right]$ $(k + r = g + j + l + \sum_{i=1}^n b_i)$

Theorem 2. (cf. Coble [1], Coolidge [2]) *Let C be an irreducible plane curve of type $(d, d - 2)$ with genus g . Then, there exists a Cremona transformation which transforms C into a plane curve:*

$$\Gamma : y^2 = \prod_{i=1}^{2g+2} (x - \lambda_i),$$

with some distinct λ_i 's.

Conversely, given a plane curve Γ as above and a collection of systems of multiplicity sequences M satisfying the conditions (1)–(4) in Theorem 1, (i) for $d \geq g + 2$, then we can find an irreducible plane curve C of type $(d, d - 2)$ such that

- (a) $\text{Data}(C) = M$,
- (b) C is Cremona birational to Γ .

Remark 3. For the first half of Theorem 2, we refer to Coble [1], p. 125 and Coolidge [2], Book III, Chapter V. As for the second half of Theorem 2, the particular cases in which $M = [(g)]$, $[\binom{g}{g} \binom{1}{1}_g]$ were discussed in [2], Book III, Chapter V, Theorems 8, 10.

In Section 2, we review the system of the multiplicity sequences, the 2-formula and quadratic Cremona transformations. In Section 3 (resp. Section 4), we will prove Theorem 1 (resp. Theorem 2). In Section 5, we discuss the defining equations for those curves given in Corollary.

2. Preliminaries

A cusp P can be described by its multiplicity sequence $\underline{m}_P = (m_0, m_1, m_2, \dots)$. For a multibranch singular point P on C , we introduced the system of the multiplicity sequences of P .

Definition 4. ([7]) *Let $P \in C$ be a multibranch singular point, having r local branches $\gamma_1, \dots, \gamma_r$. Let $\underline{m}(\gamma_i) = (m_{i0}, m_{i1}, m_{i2}, \dots)$ denote the multiplicity sequences of the branches γ_i , respectively. We define the system of the multiplicity sequences, which will be denoted by the same symbol $\underline{m}_P(C)$, to be the combination of $\underline{m}(\gamma_i)$ with brackets indicating the coincidence of the centers of the infinitely near points of the branches γ_i . For instance, for the case in which $r = 3$, we write it in the following form:*

$$\left\{ \begin{array}{l} \begin{pmatrix} m_{1,0} \\ m_{2,0} \\ m_{3,0} \end{pmatrix} \cdots \begin{pmatrix} m_{1,\rho} \\ m_{2,\rho} \\ m_{3,\rho} \end{pmatrix} \begin{pmatrix} m_{1,\rho+1} \\ m_{2,\rho+1} \\ m_{3,\rho+1} \end{pmatrix} \cdots \begin{pmatrix} m_{1,\rho'} \\ m_{2,\rho'} \\ m_{3,\rho'} \end{pmatrix} \begin{matrix} m_{1,\rho'+1}, \dots, m_{1,s_1} \\ m_{2,\rho'+1}, \dots, m_{2,s_2} \\ m_{3,\rho'+1}, \dots, m_{3,s_3} \end{matrix} \end{array} \right\}.$$

We also use some simplifications such as

$$(2_a) = \overbrace{(2, \dots, 2, 1, 1)}^a, \quad (2_0) = (1), \quad \begin{pmatrix} 1 \\ 1 \end{pmatrix}_a = \overbrace{\begin{pmatrix} 1 \\ 1 \end{pmatrix} \cdots \begin{pmatrix} 1 \\ 1 \end{pmatrix}}^a.$$

Example 5. We examine our notations for ADE singularities.

P	A_{2n}	A_{2n-1}	D_{2n-1}	D_{2n}	E_6	E_7	E_8
$\underline{m}_P(C)$	(2_n)	$\begin{pmatrix} 1 \\ 1 \end{pmatrix}_n$	$\left\{ \begin{pmatrix} 2 \\ 1 \end{pmatrix}^{2n-3} \right\}$	$\left\{ \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}_{n-2} \right\}$	(3)	$\begin{pmatrix} 2 \\ 1 \end{pmatrix} \begin{pmatrix} 1 \\ 1 \end{pmatrix}$	$(3, 2)$

Example 6. The hyperelliptic curve $y^2 = \prod_{i=1}^{2g+2} (x - \lambda_i)$ has one singularity Q on the line at infinity with $\underline{m}_Q = \begin{pmatrix} g \\ g \end{pmatrix} \begin{pmatrix} 1 \\ 1 \end{pmatrix}_g$.

Let C be an irreducible plane curve of type $(d, d - 2)$. Let $Q \in C$ be the singular point with multiplicity $d - 2$. Choosing homogeneous coordinates (x, y, z) so that $Q = (0, 0, 1)$, the curve C is defined by an equation:

$$F(x, y)z^2 + 2G(x, y)z + H(x, y) = 0,$$

where F , G and H are homogeneous polynomials of degree $d - 2$, $d - 1$ and d , respectively. Set $\Delta = G^2 - FH$. Let $t_1, \dots, t_l \in \mathbf{P}^1$ be all the distinct roots of the equation $F(t)\Delta(t) = 0$. For each i , let $(p_i, q_i) = (\text{ord}_{t_i}(F), \text{ord}_{t_i}(\Delta))$, where $\text{ord}_{t_i}(F)$ (resp. $\text{ord}_{t_i}(\Delta)$) is the multiplicity of the root t_i of the equation $F(t) = 0$ (resp. $\Delta(t) = 0$). Set $T(C) = \{(p_1, q_1), \dots, (p_l, q_l)\}$. This unordered l -tuple $T(C)$ is called the 2-formula of C (Degtyarev [3]). We remark that $T(C)$ does not depend on the choice of the coordinates (x, y, z) with $Q = (0, 0, 1)$.

Lemma 7. *The 2-formula $T(C)$ satisfies the following properties:*

- (i) $\sum_{i=1}^l q_i = 2 \sum_{i=1}^l p_i + 2,$
- (ii) $p_i = q_i$ or $\min\{p_i, q_i\}$ is even for each $i,$
- (iii) there exists a pair (p_i, q_i) such that q_i is an odd number.

Proof. (i) By definition, $\sum_{i=1}^l p_i = d - 2$ and $\sum_{i=1}^l q_i = 2d - 2.$

(ii) Suppose that $p_i \neq q_i.$ We may assume $t_i = (0, 1).$ We can write F, G and Δ as $F = x^{p_i} F_0, \Delta = x^{q_i} \Delta_0$ and $G = x^m G_0,$ respectively, where $m = \text{ord}_{t_i}(G).$ If $p_i > q_i,$ then $x^{2m} G_0^2 = x^{q_i} (\Delta_0 + x^{p_i - q_i} F_0 H).$ Thus $q_i = 2m.$ If $0 < p_i < q_i,$ then we get $x^{2m} G_0^2 = x^{p_i} (x^{q_i - p_i} \Delta_0 + F_0 H),$ which implies that $2m \geq p_i > 0.$ We have $x \nmid H,$ since C is irreducible. Hence $p_i = 2m.$

(iii) Suppose that all q_i 's are even. Then we can write as $\Delta = \Delta_0^2.$ We have $F(Fz^2 + 2Gz + H) = (Fz + G + \Delta_0)(Fz + G - \Delta_0).$ Since $\deg(Fz + G \pm \Delta_0) = d - 1,$ we infer that $Fz^2 + 2Gz + H$ is reducible. This is a contradiction. \square

Remark 8. We note that $P(x, y, z) = Fz^2 + 2Gz + H$ is irreducible if (a) $\text{GCD}(F, G, H) = 1,$ and if (b) the property (iii) holds. Indeed, under the assumption (a), if P is reducible, then $P = (Az + B)(Cz + D)$ with $A, B, C, D \in \mathbf{C}[x, y].$ But, in this case, $4\Delta = (AD - BC)^2,$ which contradicts the property (iii).

Example 9. Let C be the quartic curve $x^2 y^2 + y^2 z^2 + z^2 x^2 - 2xyz(x + y + z) = 0.$ We have $T(C) = \{(2, 0), (0, 3), (0, 3)\}.$

The (degenerate) quadratic Cremona transformation

$$\varphi_c : (x, y, z) \longrightarrow (xy, y^2, x(z - cx)) \quad (c \in \mathbf{C})$$

played an important role in [6, 7]. We find that $\varphi_c^{-1}(x, y, z) = (x^2, xy, yz + cx^2).$ We use the notations:

$$l : x = 0, \quad t : y = 0, \quad O = (0, 0, 1), \quad A = (1, 0, c), \quad B = (0, 1, 0).$$

Note that $\varphi_c(l \setminus \{O\}) = B$ and $\varphi_c(t \setminus \{O, A\}) = O.$

Let C be an irreducible plane curve of type $(d, d - 2)$ with $d \geq 4.$ Suppose the singular point $Q \in C$ of multiplicity $d - 2$ has coordinates $O.$ We have seen in [7] that the strict transform $C' = \varphi_c(C)$ is an irreducible plane curve of type $(d', d' - 2)$ for some $d'.$ In [7, 8], by analyzing how a local branch γ at $P \in \text{Sing}(C)$ is transformed by $\varphi_c,$ we described $\text{Data}[C']$ from $\text{Data}[C].$

3. Proof of Theorem 1

(i) We easily see that $P \in \text{Sing}(C) \setminus \{Q\}$ is a double point, because $LC = (d - 2)Q + 2P,$ where L is the line passing through $P, Q.$ Let $\pi : \tilde{\mathbf{P}}^2 \rightarrow \mathbf{P}^2$ be the blowing-up at $Q.$ Let E denote the exceptional curve. Take a line L passing through $Q.$ Let C' (resp. L') be the strict transform of C (resp. $L).$ We have

$C'L' = 2$. It follows that $P \in \text{Sing}(C') \cap E$ is also a double point. Thus, $\text{Data}(C)$ has the shape as in Theorem 1. Clearly, $\sum k_h + \sum k'_h = \text{mult}_Q(C) = d - 2$. The second part of the condition (1) follows from the genus formula. The condition (2) follows from the Hurwitz formula applied to the double covering $\tilde{C} \rightarrow \mathbf{P}^1$, which corresponds to the projection of C from Q , where the \tilde{C} is the non-singular model of C . For the proof of the conditions (3), (4), we refer to [7]. We will give an alternative, direct proof in Proposition 10.

(ii) Let $F(x, y)z^2 + 2G(x, y)z + H(x, y) = 0$ be the defining equation of C as in Section 2. Let $T(C)$ be the 2-formula of C . Setting

$$T'(C) = \{(p, q) \in T(C) \mid p > 0 \text{ or } q \geq 2\},$$

we renumber the pairs $(p_i, q_i) \in T'(C)$ in the following way:

- (1) $p_i > 0, q_i > 0$ and q_i is even for $i = 1, \dots, s$,
- (2) either $p_i > 0, q_i > 0$ and q_i is odd, or $p_i > 0, q_i = 0$ for $i = s + 1, \dots, N$,
- (3) $p_i = 0, q_i > 0$ and q_i is even, for $i = N + 1, \dots, N + n$,
- (4) $p_i = 0, q_i \geq 3$ and q_i is odd, for $i = N + n + 1, \dots, N + n + n'$.

Proposition 10. *Set*

- (1) for $i = 1, \dots, s$,

$$\begin{cases} k_i = k'_i = p_i/2, a_i = q_i/2 & \text{if } p_i \leq q_i, \\ k_i = q_i/2, k'_i = p_i - q_i/2, a_i = q_i/2 & \text{if } p_i > q_i, \end{cases}$$
- (2) for $i = s + 1, \dots, N$,

$$\begin{cases} k_i = p_i, a_i = (q_i - 1)/2 & \text{if } q_i > 0, \\ k_i = p_i, a_i = 0 & \text{if } q_i = 0, \end{cases}$$
- (3) $b_j = q_{N+j}/2$, for $j = 1, \dots, n$,
- (4) $b_j = (q_{N+j} - 1)/2$, for $j = n + 1, \dots, n + n'$.

Then $\text{Data}(C)$ is given as in Theorem 1, (i).

Proof. Take $(p_i, q_i) \in T'(C)$. Write t_i as $t_i = (\alpha_i, \beta_i)$. Let L_i be the line $\beta_i x = \alpha_i y$. By arranging the coordinates, we may assume $(\alpha_i, \beta_i) = (0, 1)$. Write F, G and Δ as $F = x^{p_i} F_0, G = x^{q_i} G_0$ and $\Delta = x^{q_i} \Delta_0$, where $m = \text{ord}_{t_i}(G)$.

We first consider the case in which $p_i = 0$. Since $\Delta(t_i) = 0$, we have

$$F(t_i)z^2 + 2G(t_i)z + H(t_i) = F(t_i)(z + G(t_i)/F(t_i))^2.$$

It follows that $CL_i = (d - 2)Q + 2P$, where $P = (0, 1, -G(t_i)/F(t_i))$. Let U be a neighbourhood of P such that $y \neq 0$ and $F(x, y) \neq 0$ for all $(x, y, z) \in U$. We use the affine coordinates $(\bar{x}, \bar{z}) = (x/y, z/y)$. We have

$$\begin{aligned} F(x, y)(F(x, y)z^2 + 2G(x, y)z + H(x, y)) \\ = y^{2d-2}((F(\bar{x}, 1)\bar{z} + G(\bar{x}, 1))^2 - \Delta(\bar{x}, 1)). \end{aligned}$$

Thus C is defined by the equation $(F(\bar{x}, 1)\bar{z} + G(\bar{x}, 1))^2 = \Delta(\bar{x}, 1)$ on U . Letting $u = F(\bar{x}, 1)\bar{z} + G(\bar{x}, 1)$ and $v = (\sqrt[q_i]{\Delta_0(\bar{x}, 1)})\bar{x}$, C is defined by the equation $u^2 = v^{q_i}$ around P . Thus $P \in \text{Sing}(C) \setminus \{Q\}$ if $q_i \geq 2$. In this case, we have

$$\underline{m}_P(C) = \begin{cases} \binom{1}{1}_{q_i/2} & \text{if } q_i \text{ is even,} \\ \binom{1}{2}_{(q_i-1)/2} & \text{if } q_i \text{ is odd,} \end{cases}$$

which gives the assertions (3), (4).

Conversely, take $P \in \text{Sing}(C) \setminus \{Q\}$. Let L be the line passing through P, Q . Write $L : \beta x = \alpha y$. Since $CL = (d - 2)Q + 2P$, we have $F(\alpha, \beta) \neq 0$ and $\Delta(\alpha, \beta) = 0$. For $(\alpha, \beta) \in \mathbf{P}^1$, we find a pair $(0, q) \in T(C)$. We see from the above argument that C is defined by the equation $u^2 = v^q$ near P . Thus $q \geq 2$.

We now consider the case in which $p_i > 0$. Let $\pi : \tilde{\mathbf{P}}^2 \rightarrow \mathbf{P}^2$ be the blowing-up at Q and E the exceptional curve of π . We use the affine coordinates $(\bar{x}, \bar{y}) = (x/z, y/z)$ of $U := \{(x, y, z) \in \mathbf{P}^2 \mid z \neq 0\}$. Put $V = \pi^{-1}(U)$. There exist an open cover $V = V_1 \cup V_2$ ($V_j \cong \mathbf{C}^2$) with standard coordinates (u_j, v_j) of V_j such that $\pi|_{V_1} : V_1 \ni (u_1, v_1) \mapsto (u_1 v_1, u_1)$ and $\pi|_{V_2} : V_2 \ni (u_2, v_2) \mapsto (u_2, u_2 v_2)$. Note that E is defined by $u_j = 0$ on V_j . The strict transform L'_i of L_i is defined by $v_1 = 0$ on V_1 . Let P be the unique point $E \cap L'_i$. We have $P = (0, 0)$ on V_1 . The strict transform C' of C is defined by the equation: $F(v_1, 1) + 2G(v_1, 1)u_1 + H(v_1, 1)u_1^2 = 0$ on V_1 . By the definition of p_i and m , the curve C' is defined by the equation: $F_0 v_1^{p_i} + 2G_0 v_1^m u_1 + H u_1^2 = 0$. In particular, we have $(C'E)_P = p_i$. If $q_i = 0$, then we must have $m = 0$ (see the proof of Lemma 7). Hence C' is smooth at P . If $q_i > 0$, then we have $m > 0$ (cf. the proof of Lemma 7). Since C is irreducible, we see that $H(t_i) \neq 0$. We have $H(F + 2G_0 v_1^m u_1 + H u_1^2) = (H u_1 + G_0 v_1^m)^2 - \Delta$. This means that C' is defined by the equation:

$$(H(v_1, 1)u_1 + G_0(v_1, 1)v_1^m)^2 - \Delta(v_1, 1) = 0$$

in a neighborhood of P .

Letting $u = H(v_1, 1)u_1 + G_0(v_1, 1)v_1^m$ and $v = (\sqrt[q_i]{\Delta_0(v_1, 1)})v_1$, C' is defined by the equation $u^2 = v^{q_i}$ around P . We have

$$\underline{m}_P(C') = \begin{cases} \binom{1}{1}_{q_i/2} & \text{if } q_i \text{ is even,} \\ \binom{1}{2}_{(q_i-1)/2} & \text{if } q_i \text{ is odd,} \\ \binom{1}{1} & \text{if } q_i = 0, \end{cases}$$

which gives the values of a_i in (1), (2). We prove the remaining assertions in (1). If q_i is even, then C' has two branches γ_+, γ_- at P defined by

$$H(v_1, 1)u_1 + G_0(v_1, 1)v_1^m \pm v_1^{q_i/2} \sqrt{\Delta_0(v_1, 1)} = 0.$$

In case $p_i > q_i$, we have $m = q_i/2$ (see the proof of Lemma 7). We infer that one of the intersection numbers $(E\gamma_+)_P$ and $(E\gamma_-)_P$ is equal to $q_i/2$. The other one must be equal to $p_i - q_i/2$, because $(EC')_P = p_i$. In case $p_i \leq q_i$, we have $m \geq p_i/2$ (cf. the proof of Lemma 7). Thus $(E\gamma_{\pm})_P \geq p_i/2$, hence $(E\gamma_{\pm})_P = p_i/2$. Consequently, we obtain the pair (k_i, k'_i) .

Conversely, take $P \in C' \cap E$. We assume $P \in V_1$. Write the coordinates of P as $P = (0, \beta)$. The equation $F(\beta, 1) + 2G(\beta, 1)u_1 + H(\beta, 1)u_1^2 = 0$ has the solution $u_1 = 0$ as C' passes through P . Thus $F(\beta, 1) = 0$. For $(\beta, 1) \in \mathbf{P}^1$, we find a pair $(p, q) \in T(C)$ with $p > 0$. \square

Remark 11. For $i = s+1, \dots, N$, if $(k_i, a_i) = (1, 0)$, then we have either $(p_i, q_i) = (1, 1)$ or $(1, 0)$. The case $(p_i, q_i) = (1, 1)$ occurs if and only if the line L_i is a flex-tangent line to the corresponding branch at Q .

4. Proof of Theorem 2

Let C be given by the equation (see Section 2):

$$F(x, y)z^2 + 2G(x, y)z + H(x, y) = 0.$$

Put $\Delta = G^2 - FH$. Via linear coordinates change of x and y , we may assume that $y \nmid F\Delta$. We then define a Cremona transformation (cf. [2], Book III, Chapter V):

$$\Phi(x, y, z) = (xy^{d-2}, y^{d-1}, Fz + G).$$

We find that $\Phi^{-1}(x, y, z) = (xF, yF, y^{d-2}z - G)$. We see easily that the strict transform $C' = \Phi(C)$ is defined by the equation:

$$y^{2(d-2)}z^2 = \Delta.$$

Write $\Delta = \prod_{i=1}^k (x - \lambda_i y)^{q_i}$, where the λ_i 's are distinct. Renumber q_i 's so that q_i 's are odd for $i = 1, \dots, l$ and q_i 's are even for $i = l + 1, \dots, k$. Letting $s_i = [q_i/2]$ for $i = 1, \dots, k$, we put $S = \prod_{i=1}^k (x - \lambda_i y)^{s_i}$ and $s = \sum_{i=1}^k s_i$. Note that $2d - 2 = \sum_{i=1}^k q_i = 2s + l$. We next define a Cremona transformation:

$$\Psi(x, y, z) = (xS, yS, y^s z).$$

We find that $\Psi^{-1}(x, y, z) = (xy^s, y^{s+1}, Sz)$. We see that $\Gamma' = \Psi(C')$ is defined by the equation:

$$y^{2(d-2)}z^2 = \prod_{i=1}^l (x - \lambda_i y).$$

We see that $l = 2g + 2$ and $g = d - s - 2$. Take a projective transformation: $\iota : (x, y, z) \rightarrow (x, z, y)$. Finally, the image $\Gamma = \iota(\Gamma')$ has the affine equation:

$$y^2 = \prod_{i=1}^{2g+2} (x - \lambda_i).$$

We now prove the second half of Theorem 2. We start with the curve Γ and a collection of systems of multiplicity sequences:

$$M = \left[m, \binom{1}{1}_{b_1}, \dots, \binom{1}{1}_{b_n}, (2b_{n+1}), \dots, (2b_{n+n'}) \right],$$

where the m is the system of the multiplicity sequences of the singular point with multiplicity $d - 2$. Let $r(M)$, $N(M)$ denote the number of the branches and the number of the different tangent lines of m . We have to construct an irreducible plane curve of type $(d, d - 2)$ with $\text{Data}(C) = M$. In [7], we considered the case in which $g = 0$. We here assume that $g \geq 1$. We follow the arguments in [7].

First we deal with the cuspidal case given in Corollary of Theorem 1. See also Proposition 13.

Case (a): $M = [(k), (2_{b_1}), \dots, (2_{b_{n'}})]$, where $k = g + \sum b_i$. We use the induction on n' . (i) $M = [(g)]$. Interchanging coordinates, we start with the curve:

$$\Gamma_0 : x^{2g}z^2 = \prod_{i=1}^{2g+2} (y - \lambda_i x).$$

After a linear change of coordinates, we may assume that $c = \prod_{i=1}^{2g+2} (-\lambda_i) \neq 0$. Letting $c_1 = \sqrt{c}$, we have $\Gamma_0 t = (2g)O + A_1 + A'_1$, where $A_1 = (1, 0, c_1)$, $A'_1 = (1, 0, -c_1)$. Let Γ_1 be the strict transform of Γ_0 via φ_{c_1} . Using Lemma 1, (a) and Lemma 2, (e)* in [7], we see that $\Gamma_1 t = (2g - 1)O + A_2$. Write $A_2 = (1, 0, c_2)$. Let Γ_2 be the strict transform of Γ_1 via φ_{c_2} . In this way, we successively choose c_1, \dots, c_g . It turns out that $\text{Data}(\Gamma_g) = [(g)]$.

(ii) Suppose we have constructed C_0 with $\text{Data}(C_0) = [(k_0), (2_{b_1}), \dots, (2_{b_{n'-1}})]$, where $k_0 = g + \sum_{i=1}^{n'-1} b_i$. After a suitable change of coordinates, we may assume $C_0 l = k_0 O + 2B_1$ and $C_0 t = (k_0 + 1)O + A_1$. Note that the double covering $\tilde{C} \rightarrow \mathbf{P}^1$ defined through the projection from O to a line, must have $2g + 2$ branch points. Since $n' - 1 < 2g + 2$, we see that a line passing through O is tangent to C_0 at a smooth point B_1 . Write $A_1 = (1, 0, c_1)$. Let C_1 be the strict transform of C_0 via φ_{c_1} . We have $C_1 l = (k_0 + 1)O + 2B$ and $C_1 t = (k_0 + 2)O + A_2$. Write $A_2 = (1, 0, c_2)$. Let C_2 be the strict transform of C_1 via φ_{c_2} . We have again $C_2 t = (k_0 + 3)O + A_3$. Repeating in this way, we successively choose $c_1, \dots, c_{b_{n'}}$ and define $C_1, \dots, C_{b_{n'}}$. Then, the curve $C = C_{b_{n'}}$ has the desired property.

Case (b): $M = [(2k + 1, 2_k), (2_{b_1}), \dots, (2_{b_{n'}})]$, where $k + 1 = g + \sum b_i$. As in Case (a), we can similarly prove this case. For the first step: $M = [(2g - 1, 2_{g-1})]$, it suffices to arrange coordinates so that $\Gamma_0 t = gO + 2A_1$ with $A_1 = (1, 0, 0)$. Put $c_1 = 0$ and choose c_2, \dots, c_g arbitrarily. Then we obtain $\text{Data}(\Gamma_g) = M$ (cf. Lemma 1, (b) and Lemma 2, (e)* in [7]).

Case (c): $M = [(2k, 2_{k+j}), (2_{b_1}), \dots, (2_{b_{n'}})]$, where $k = g + j + \sum b_i$. We also use the induction on n' as in Case (a). For the first step: $M = [(2(g + j), 2_{g+2j})]$, we start with a curve C_0 with $\text{Data}(C_0) = [(g + j), (2_j)]$ constructed in Case (a). We again arrange coordinates so that $C_0 t = (g + j)O + 2R$, where $\underline{m}_R(C_0) = (2_j)$ and $R = (1, 0, a)$. Choose $c_1 \neq a$ and c_2, \dots, c_{g+j} arbitrarily. Then we have $\text{Data}(C_{g+j}) = M$ (cf. Lemma 1, (a)*, (c) in [7]).

Starting with the cuspidal case, we can prove the general case in a similar manner to that in [7]. We have three subcases:

- I. $N(M) = r(M) = 1$, II. $N(M) = 1, r(M) = 2$, III. $N(M) \geq 2$.

Here, we only give a proof for $M = [(k), \binom{1}{1}_{b_1}, \dots, \binom{1}{1}_{b_n}, (2b_{n+1}), \dots, (2b_{n+n'})]$, where $k = g + \sum_{j=1}^{n+n'} b_j$, which is one of the remaining cases in I. We use the induction on n .

- (i) We constructed a cuspidal curve C with $\text{Data}(C) = [(k), (2b_{n+1}), \dots, (2b_{n+n'})]$.
(ii) Suppose we have already constructed C_0 with

$$\text{Data}(C_0) = [(k_0), \binom{1}{1}_{b_1}, \dots, \binom{1}{1}_{b_{n-1}}, (2b_{n+1}), \dots, (2b_{n+n'})],$$

where $k_0 = g + \sum_{j=1}^{n-1} b_j + \sum_{j=n+1}^{n+n'} b_j$. By arranging coordinates, we have $C_0 l = k_0 O + B_1 + B'_1$ and $C_0 t = (k_0 + 1)O + A_1$. Letting $A_1 = (1, 0, c_1)$, the strict transform C_1 of C_0 via φ_{c_1} has the property $C_0 t = (k_0 + 2)O + A_2$. Write $A_2 = (1, 0, c_2)$. We successively choose c_2, \dots, c_{b_n} in this way. Then the strict transform C of C_0 via $\varphi_{c_{b_n}} \circ \dots \circ \varphi_{c_1}$ has the desired property (cf. Lemma 1, (d) and Lemma 2, (tn) in [7]). In particular, C contains a tacnode $\binom{1}{1}_{b_n}$ at $B = (0, 1, 0)$.

5. Defining equations

We now describe the defining equations for those curves listed in Corollary. In [6, 7, 8], the defining equations were computed step by step by using quadratic Cremona transformations. But, for some cases, we encountered a difficulty to evaluate points in some special positions. We here employ the method used by Degtyarev in [3].

Lemma 12. *Consider two polynomials*

$$g(t) = \sum_{i=0}^d c_i t^i, \quad \delta(t) = \sum_{i=0}^{2d} d_i t^i \in \mathbb{C}[t].$$

Suppose $\delta(0) = d_0 \neq 0$. For $k \leq d$, we have $t^k \mid (g^2 - \delta)$ if and only if

- (1) $c_0 = \pm \sqrt{d_0}$,
- (2) $c_j = (d_j - \sum_{i=1}^{j-1} c_i c_{j-i}) / (2c_0)$ for $j = 1, \dots, k-1$.

Proof. Write $g(t)^2 = \sum_{j=0}^{2d} b_j t^j$. We see that $b_j = \sum_{i=0}^j c_i c_{j-i}$ for $j \leq d$. □

Proposition 13. *The defining equations of irreducible plane curves of type $(d, d-2)$ with genus g having only cusps are the following (up to projective equivalence, the λ_i 's are distinct).*

- (a) $y^k z^2 + 2Gz + \{G^2 - \Delta\} / y^k = 0$, where

$$\Delta(x, y) = \prod_{i=1}^{n'} (x - \lambda_i y)^{2b_i+1} \prod_{i=n'+1}^{2g+2} (x - \lambda_i y).$$

Letting $G(x, y) = \sum_{h=0}^{k+1} c_h x^{k+1-h} y^h$, the coefficients c_0, \dots, c_{k-1} are determined by the condition $y^k \mid (G(1, y)^2 - \Delta(1, y))$ (see Lemma 12).

$$(b) \quad (y^k z + \sum_{h=0}^{k+1} c_h x^{k+1-h} y^h)^2 y - \prod_{i=1}^{n'} (x - \lambda_i y)^{2b_i+1} \prod_{i=n'+1}^{2g+1} (x - \lambda_i y) = 0.$$

$$(c) \quad (y^k z + \sum_{h=0}^{k+1} c_h x^{k+1-h} y^h)^2 - y^{2j+1} \prod_{i=1}^{n'} (x - \lambda_i y)^{2b_i+1} \prod_{i=n'+1}^{2g+1} (x - \lambda_i y) = 0,$$

where $c_0 \neq 0$.

Proof. Class (a). In this case, in view of the argument in the proof of Theorem 1, (ii), we have $T(C) = \{(k, 0), (0, 2b_1 + 1), \dots, (0, 2b_{n'} + 1), (0, 1), \dots, (0, 1)\}$. Thus, we can write $F = y^k$ and Δ as above. We must have $y^k | (G^2 - \Delta)$. In view of Lemma 12, the coefficients c_0, \dots, c_{k-1} are uniquely determined. In particular, $c_0 = \pm 1$. So by Remark 8, the defining equation is irreducible.

Class (b): We have $T(C) = \{(2k + 1, 2k + 1), (0, 2b_1 + 1), \dots, (0, 2b_{n'+1}), (0, 1), \dots, (0, 1)\}$. We can arrange coordinates as

$$F = y^{2k+1}, \quad \Delta = y^{2k+1} \prod_{i=1}^{n'} (x - \lambda_i y)^{2b_i+1} \prod_{i=n'+1}^{2g+1} (x - \lambda_i y).$$

We infer that $G = y^{k+1} G_0$ for some G_0 .

Class (c): We have $T(C) = \{(2k, 2k + 2j + 1), (0, 2b_1 + 1), \dots, (0, 2b_{n'+1}), (0, 1), \dots, (0, 1)\}$. We can arrange coordinates as

$$F = y^{2k}, \quad \Delta = y^{2k+2j+1} \prod_{i=1}^{n'} (x - \lambda_i y)^{2b_i+1} \prod_{i=n'+1}^{2g+1} (x - \lambda_i y).$$

It follows that $G = y^k G_0$ for some G_0 . If we write $G_0 = \sum_{h=0}^{k+1} c_h x^{k+1-h} y^h$, then we must have $c_0 \neq 0$, for otherwise the defining equation becomes reducible (see Remark 8). \square

Example 14. We give the defining equation of a cuspidal septic curve C with $\text{Data}(C) = [(5), (2), (2), (2), (2)]$ which are birational to the elliptic curve $y^2 = (x^2 - 1)(x^2 - \lambda^2)$, $(\lambda \neq \pm 1, 0)$.

$$\begin{aligned} & y^5 z^2 + \{2x^4 - 3(\lambda^2 + 1)x^2 y^2 + \frac{3}{4}(\lambda^4 + 6\lambda^2 + 1)y^4\} x^2 z \\ & - \frac{1}{8}(\lambda^2 + 1)(\lambda^4 - 10\lambda^2 + 1)x^6 y + \frac{3}{64}\{3\lambda^8 - 28\lambda^6 - 78\lambda^4 - 28\lambda^2 + 3\} x^4 y^3 \\ & + 3\lambda^4(\lambda^2 + 1)x^2 y^5 - \lambda^6 y^7 = 0 \end{aligned}$$

Proposition 15. *The defining equations of irreducible plane curves of type $(d, d - 2)$ with genus g having only bibranch singularities are the following (up to projective equivalence, the λ_i 's are distinct).*

$$(e) \quad (y^k z + \sum_{h=0}^{k+1} c_h x^{k+1-h} y^h)^2 - y^{2j} \prod_{i=1}^n (x - \lambda_i y)^{2b_i} \prod_{i=n+1}^{n+2g+2} (x - \lambda_i y) = 0,$$

where $c_0 \neq 0$.

$$(f) \quad y^{2k+r}z^2 + 2y^kG_0z + \left\{G_0^2 - \Delta_0\right\}/y^r = 0,$$

where

$$\Delta_0(x, y) = \prod_{i=1}^n (x - \lambda_i y)^{2b_i} \prod_{i=n+1}^{n+2g+2} (x - \lambda_i y)$$

and the coefficients c_0, \dots, c_{r-1} of $G_0(x, y) = \sum_{h=0}^{k+r+1} c_h x^{k+r+1-h} y^h$ are determined by the condition $y^r \mid (G_0(1, y)^2 - \Delta_0(1, y))$ (cf. Lemma 12) and c_r is chosen so that $y^{r+1} \nmid (G_0(1, y)^2 - \Delta_0(1, y))$.

$$(aa) \quad x^r y^k z^2 + 2Gz + \left\{G^2 - \Delta\right\}/(x^r y^k) = 0,$$

where

$$\Delta(x, y) = \prod_{i=1}^n (x - \lambda_i y)^{2b_i} \prod_{i=n+1}^{n+2g+2} (x - \lambda_i y) \quad (\lambda_i \neq 0 \text{ for all } i).$$

Write $G(x, y) = \sum_{h=0}^{k+r+1} c_h x^{k+r+1-h} y^h$. The coefficients $c_0, \dots, c_{k-1}, c_{k+2}, \dots, c_{k+r+1}$ are determined by the conditions $y^k \mid (G(1, y)^2 - \Delta(1, y))$ and $x^r \mid (G(x, 1)^2 - \Delta(x, 1))$ (cf. Lemma 12).

$$(aa+) \quad x(y^k z + 2G_0)z + \left\{xG_0^2 - \Delta_0\right\}/y^k = 0,$$

where

$$\Delta_0(x, y) = \prod_{i=1}^n (x - \lambda_i y)^{2b_i} \prod_{i=n+1}^{n+2g+1} (x - \lambda_i y) \quad (\lambda_i \neq 0 \text{ for all } i).$$

Write $G_0(x, y) = \sum_{h=0}^{k+1} c_h x^{k+1-h} y^h$. The coefficients c_0, \dots, c_{k-1} are determined by the condition $y^k \mid (G_0(1, y)^2 - \Delta_0(1, y))$.

$$(aa1) \quad xyz^2 - (x - \lambda y)^4 = 0 \quad (\lambda \neq 0, g = 0).$$

$$(aa2) \quad xyz^2 - (x - \lambda_1 y)^2 (x - \lambda_2 y)^2 = 0 \quad (\lambda_1 \lambda_2 \neq 0, g = 0).$$

$$(aa3) \quad xyz^2 - (x - \lambda_1 y)^2 (x - \lambda_2 y) (x - \lambda_3 y) = 0 \quad (\lambda_1 \lambda_2 \lambda_3 \neq 0, g = 1).$$

$$(aa4) \quad xyz^2 - \prod_{i=1}^4 (x - \lambda_i y) = 0 \quad (\lambda_i \neq 0 \text{ for all } i, g = 2).$$

$$(ab) \quad x^r y^{2k+1} z^2 + 2y^{k+1} G_0 z + \left\{yG_0^2 - \Delta_0\right\}/x^r = 0,$$

where

$$\Delta_0(x, y) = \prod_{i=1}^n (x - \lambda_i y)^{2b_i} \prod_{i=n+1}^{n+2g+1} (x - \lambda_i y) \quad (\lambda_i \neq 0 \text{ for all } i)$$

and the coefficients $c_{k+2}, \dots, c_{k+r+1}$ of $G_0(x, y) = \sum_{h=0}^{k+r+1} c_h x^{k+r+1-h} y^h$ are determined by the condition $x^r \mid (G_0(x, 1)^2 - \Delta_0(x, 1))$.

$$(ab+) \quad (y^k z + \sum_{h=0}^{k+1} c_h x^{k+1-h} y^h)^2 xy - \prod_{i=1}^n (x - \lambda_i y)^{2b_i} \prod_{i=n+1}^{n+2g} (x - \lambda_i y) = 0,$$

where $\lambda_i \neq 0$ for all i .

$$(ac) \quad x^r y^{2k} z^2 + 2y^k G_0 z + \{G_0^2 - \Delta_0\} / x^r = 0,$$

where

$$\Delta_0(x, y) = y^{2j+1} \prod_{i=1}^n (x - \lambda_i y)^{2b_i} \prod_{i=n+1}^{n+2g+1} (x - \lambda_i y) \quad (\lambda_i \neq 0 \text{ for all } i)$$

and the coefficients $c_{k+2}, \dots, c_{k+r+1}$ of $G_0(x, y) = \sum_{h=0}^{k+r+1} c_h x^{k+r+1-h} y^h$ are determined by the condition $x^r \mid (G_0(x, 1)^2 - \Delta_0(x, 1))$ and $c_0 \neq 0$, which is required for the irreducibility of the defining equation (Remark 8).

$$(ac+) \quad (y^k z + \sum_{h=0}^{k+1} c_h x^{k+1-h} y^h)^2 x - y^{2j+1} \prod_{i=1}^n (x - \lambda_i y)^{2b_i} \prod_{i=n+1}^{n+2g} (x - \lambda_i y) = 0,$$

where $\lambda_i \neq 0$ for all i and $c_0 \neq 0$.

$$(bb) \quad (x^r y^k z + \sum_{h=0}^{k+r+1} c_h x^{k+r+1-h} y^h)^2 x y - \prod_{i=1}^n (x - \lambda_i y)^{2b_i} \prod_{i=n+1}^{n+2g} (x - \lambda_i y) = 0,$$

where $\lambda_i \neq 0$ for all i .

$$(bc) \quad (x^r y^k z + \sum_{h=0}^{k+r+1} c_h x^{k+r+1-h} y^h)^2 y - x^{2l+1} \prod_{i=1}^n (x - \lambda_i y)^{2b_i} \prod_{i=n+1}^{n+2g} (x - \lambda_i y) = 0,$$

where $\lambda_i \neq 0$ for all i and $c_{k+r+1} \neq 0$.

$$(cc) \quad (x^r y^k z + \sum_{h=0}^{k+r+1} c_h x^{k+r+1-h} y^h)^2 - x^{2l+1} y^{2j+1} \prod_{i=1}^n (x - \lambda_i y)^{2b_i} \prod_{i=n+1}^{n+2g} (x - \lambda_i y) = 0,$$

where $\lambda_i \neq 0$ for all i and $c_0 c_{k+r+1} \neq 0$.

Proof. Class (e): In this case, we have $T(C) = \{(2k, 2k + 2j), (0, 2b_1), \dots, (0, 2b_n), (0, 1), \dots, (0, 1)\}$. We can arrange coordinates as

$$F = y^{2k}, \quad \Delta = y^{2k+2j} \prod_{i=1}^n (x - \lambda_i y)^{2b_i} \prod_{i=n+1}^{n+2g+2} (x - \lambda_i y).$$

We infer that $G = y^k G_0$ for some G_0 .

Class (f): We have $T(C) = \{(2k + r, 2k), (0, 2b_1), \dots, (0, 2b_n), (0, 1), \dots, (0, 1)\}$. We can arrange coordinates as

$$F = y^{2k+r}, \quad \Delta = y^{2k} \prod_{i=1}^n (x - \lambda_i y)^{2b_i} \prod_{i=n+1}^{n+2g+2} (x - \lambda_i y).$$

We infer that $G = y^k G_0$ for some G_0 . Write $\Delta = y^{2k} \Delta_0$. Furthermore, we must have $y^r \mid (G_0^2 - \Delta_0)$.

Class (aa): We may assume $k \geq r$. We have the case in which $T(C) = \{(k, 0), (r, 0), (0, 2b_1), \dots, (0, 2b_n), (0, 1), \dots, (0, 1)\}$. We can then arrange coordinates as

$$F = x^r y^k \quad \Delta = \prod_{i=1}^n (x - \lambda_i y)^{2b_i} \prod_{i=n+1}^{n+2g+2} (x - \lambda_i y) \quad (\lambda_i \neq 0 \text{ for all } i).$$

We infer that $y^k|(G^2 - \Delta)$ and $x^r|(G^2 - \Delta)$.

In case $r = 1$, we also have the case in which $T(C) = \{(k, 0), (1, 1), (0, 2b_1), \dots, (0, 2b_n), (0, 1), \dots, (0, 1)\}$. We obtain Class (aa+). If $d = 4$, then we have four more classes:

Class	$T(C)$	g
(aa1)	$\{(1, 1), (1, 1), (0, 4)\}$	0
(aa2)	$\{(1, 1), (1, 1), (0, 2), (0, 2)\}$	0
(aa3)	$\{(1, 1), (1, 1), (0, 2), (0, 1), (0, 1)\}$	1
(aa4)	$\{(1, 1), (1, 1), (0, 1), (0, 1), (0, 1), (0, 1)\}$	2

For the remaining classes, we omit the details. □

Acknowledgement. The first author would like to thank Prof. I. V. Dolgachev for comments on the book [1].

References

- [1] Coble, A. B.: *Algebraic Geometry and Theta Functions*. Colloq. Publ. **10**, Amer. Math. Soc., Providence, RI, 1929. [JFM 55.0808.02](#)
- [2] Coolidge, J. L.: *A Treatise on Algebraic Plane Curves*. Oxford Univ. Press. 1931 (reprinted 1959). [JFM 57.0820.06](#) and [Zbl 0085.36403](#)
- [3] Degtyarev, A.: *Isotopy classification of complex plane projective curves of degree five*. Leningr. Math. J. **1** (1990), 881–904. [Zbl 0725.14025](#)
- [4] Fenske, T.: *Rational 1- and 2-cuspidal plane curves*. Beitr. Algebra Geom. **40**(2) (1999), 309–329. [Zbl 0959.14012](#)
- [5] Flenner, H.; Zaidenberg, M.: *On a class of rational cuspidal plane curves*. Manuscr. Math. **89** (1996), 439–460. [Zbl 0868.14014](#)
- [6] Flenner, H.; Zaidenberg, M.: *Rational cuspidal plane curves of type $(d, d - 3)$* . Math. Nachr. **210** (2000), 93–110. [Zbl 0948.14020](#)
- [7] Sakai, F.; Saleem, M.: *Rational plane curves of type $(d, d - 2)$* . Saitama Math. J. **22** (2004), 11–34. [Zbl 1079.14042](#)
- [8] Sakai, F.; Tono, K.: *Rational cuspidal curves of type $(d, d - 2)$ with one or two cusps*. Osaka J. Math. **37** (2000), 405–415. [Zbl 0969.14020](#)

Received January 7, 2008