



Banach J. Math. Anal. 3 (2009), no. 1, 111–121

**BANACH JOURNAL OF MATHEMATICAL ANALYSIS**

ISSN: 1735-8787 (electronic)

<http://www.math-analysis.org>

## NEIGHBORHOODS OF A CERTAIN CLASS OF ANALYTIC FUNCTIONS WITH NEGATIVE COEFFICIENTS

ADRIANA CĂTAȘ<sup>1</sup>

Communicated by K. S. Berenhaut

**ABSTRACT.** A certain subclass of analytic functions in the open unit disc with negative coefficients is introduced. The new class is defined by means of multiplier transformations. By making use of the familiar concept of neighborhoods of analytic function, the author proves coefficient inequalities, distortion theorems and associated inclusion relations for the  $(n, \delta)$ -neighborhoods of functions belonging to the new class, which satisfy a certain nonhomogeneous Cauchy-Euler differential equation.

<sup>1</sup> FACULTY OF SCIENCES, DEPARTMENT OF MATHEMATICS, 1 UNIVERSITY STREET, 410087, ORADEA, ROMANIA.

*E-mail address:* [acatas@gmail.com](mailto:acatas@gmail.com), [acatas@uoradea.ro](mailto:acatas@uoradea.ro)

---

*Date:* Received: 12 June 2008; Accepted: 19 November 2008.

*2000 Mathematics Subject Classification.* Primary 30C45; Secondary 30C50.

*Key words and phrases.* Analytic function, multiplier transformation, coefficient inequality, distortion bound, inclusion relationship.