



Banach J. Math. Anal. 8 (2014), no. 2, 124–130

BANACH **J**OURNAL OF **M**ATHEMATICAL **A**NALYSIS

ISSN: 1735-8787 (electronic)

www.emis.de/journals/BJMA/

GENERALIZATION OF SOME RESULTS ON $p\alpha$ -DUALS

IVANA DJOLOVIĆ^{1*} AND EBERHARD MALKOWSKY²

Communicated by D. Werner

ABSTRACT. We will find the $p\alpha$ -dual for X_T , where X is one of the spaces c , c_0 , ℓ_∞ and T is a triangle matrix. This will be achieved in two ways: firstly, under some conditions for the inverse matrix S of T and secondly, for arbitrary triangles T .

¹ TECHNICAL FACULTY IN BOR, UNIVERSITY OF BELGRADE, VJ 12, 19210 BOR, SERBIA.

E-mail address: zucko@open.telekom.rs

² DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE, FATIH UNIVERSITY, BÜYÜKÇEKMECE 34500, ISTANBUL, TURKEY.

E-mail address: Eberhard.Malkowsky@math.uni-giessen.de; ema@Bankerinter.net

Date: Received: Oct. 19, 2013; Accepted: Nov. 6, 2013.

* Corresponding author.

2010 *Mathematics Subject Classification.* Primary: 40H05; Secondary: 46H05.

Key words and phrases. $p\alpha$ -dual, sequence spaces, matrix domain of triangle.