



**APPROXIMATION OF COMMON RANDOM FIXED POINTS
OF FINITE FAMILIES OF N-UNIFORMLY L_i -LIPSCHITZIAN
ASYMPTOTICALLY HEMICONTRACTIVE RANDOM MAPS IN
BANACH SPACES.**

CHIKA MOORE¹, C. P. NNANWA^{2*} AND B. C. UGWU³

Communicated by K. Ciesielski

ABSTRACT. Let (Ω, Σ, μ) be a complete probability measure space, E be a real separable Banach space, K a nonempty closed convex subset of E . Let $T : \Omega \times K \rightarrow K$, such that $\{T_i\}_{i=1}^N$, be N -uniformly L_i -Lipschitzian asymptotically hemicontractive random maps of K with $F = \bigcap_{i=1}^N F(T_i) \neq \emptyset$. We construct an explicit iteration scheme and prove necessary and sufficient conditions for approximating common fixed points of finite family of asymptotically hemicontractive random maps.

1. INTRODUCTION AND PRELIMINARIES

Let E be a real normed linear space, E^* its dual and let the map $J : E \rightarrow 2^{E^*}$ denote the generalized duality mapping define for each $x \in E$ by

$$J(x) = \{f^* \in E^* : \langle x, f^* \rangle = \|x\|^2 = \|f^*\|^2\}$$

where \langle, \rangle denotes the duality pairing between elements of E and E^* . It is well known that if E is smooth, then J is single-valued. In the sequel we shall denote the single-valued normalized duality map by j .

Date: Received: 7 October 2008; Revised: 15 May 2009; Accepted: 3 June 2009.

* Corresponding author.

2000 *Mathematics Subject Classification.* Primary 47H06; Secondary 47H09, 47J25.

Key words and phrases. N -uniformly L_i -Lipschitzian, finite family, asymptotically hemicontractive map, explicit iteration, Banach space.

Let (Ω, Σ, μ) be a complete probability measure space with Σ , a σ -algebra of subset of Ω and μ a probability measure on Σ . Let E be a (separable) normed linear space. A map $\xi : \Omega \rightarrow E$ is measurable if $\xi^{-1}(K) \in \Sigma$ for each open subset K of E ; alternatively, $\xi^{-1} \in \Sigma$ for each open ball B in E . A map $T : \Omega \times E \rightarrow E$ is said to be a random map if for fixed $\xi \in E$ the map $T(\omega)\xi(\omega) : \Omega \rightarrow E$ is measurable. A measurable map $\xi : \Omega \rightarrow E$ is called a random fixed point of the random map $T : \Omega \times E \rightarrow E$ if $\mu(\{\omega \in \Omega : T(\omega)\xi(\omega) = \xi(\omega)\}) = 1$; that is $T\xi = \xi$ almost surely (a.s.) in Ω . The n th iterate, $n \in \mathbb{N}$ of the map $T : \Omega \times E \rightarrow E$ is given by $T^n(\omega) = T(\omega)T^{n-1}(\omega)$; that is, $T^n(\omega)\xi(\omega) = T^n(\omega)(T^{n-1}(\omega)\xi(\omega))$. Let $\xi, \eta : \Omega \rightarrow E$ be measurable maps.

A random map $T : \Omega \times E \rightarrow E$ is said to be nonexpansive if

$$\|T(\omega)\xi(\omega) - T(\omega)\eta(\omega)\| \leq \|\xi(\omega) - \eta(\omega)\| \quad (\omega \in \Omega)$$

and is L-Lipschitzian if for all $\omega \in \Omega$ there exists $L(\omega) \geq 0$ such that

$$\|T(\omega)\xi(\omega) - T(\omega)\eta(\omega)\| \leq L(\omega)\|\xi(\omega) - \eta(\omega)\|$$

where $L(\omega) \leq L$ a.s. in Ω that is $\mu(\{\omega \in \Omega : L(\omega) \leq L\}) = 1$. The map T is said to be uniformly L-Lipschitzian if for all $\omega \in \Omega$, there exists $L(\omega) \geq 0$, such that $L(\omega) \leq L$ a.s. a constant such that for all $\xi(\omega), \eta(\omega) \in E, \omega \in \Omega, n \in \mathbb{N}$,

$$\|T^n(\omega)\xi(\omega) - T^n(\omega)\eta(\omega)\| \leq L(\omega)\|\xi(\omega) - \eta(\omega)\|$$

A map T is said to be *asymptotically nonexpansive* if for all $\omega \in \Omega$, there exists $\{k_n(\omega)\}_{n \geq 0} \subset [1, +\infty)$ with $\lim_{n \rightarrow \infty} k_n(\omega) = 1$ a.s. such that

$$\|T^n(\omega)\xi(\omega) - T^n(\omega)\eta(\omega)\| \leq k_n(\omega)\|\xi(\omega) - \eta(\omega)\| \quad (n \in \mathbb{N})$$

and T is said to be *asymptotically pseudocontractive* if for all $\omega \in \Omega$, there exists $\{k_n(\omega)\}_{n \geq 0} \subset [1, +\infty)$ with $\lim_{n \rightarrow \infty} k_n(\omega) = 1$ a.s. and for all $\xi(\omega), \eta(\omega) \in E$, there exists $j(\xi(\omega) - \eta(\omega)) \in J(\xi(\omega) - \eta(\omega))$ such that

$$\langle T^n(\omega)\xi(\omega) - T^n(\omega)\eta(\omega), j(\xi(\omega) - \eta(\omega)) \rangle \leq k_n(\omega)\|\xi(\omega) - \eta(\omega)\|^2 \quad (n \in \mathbb{N}). \quad (1.1)$$

T is said to be asymptotically hemicontractive if $F(T) = \{\xi(\omega) \in D(T) : T(\omega)\xi(\omega) = \xi(\omega)\} \neq \emptyset$ and (1.1) is satisfied for all $\xi(\omega) \in D(T)$ and $\eta(\omega) = \xi^*(\omega) \in F(T)$, $k_n(\omega) = a_n(\omega)$

and there exists $j(\xi(\omega) - \xi^*(\omega)) \in J(\xi(\omega) - \xi^*(\omega))$ such that

$$\langle T^n(\omega)\xi(\omega) - \xi^*(\omega), j(\xi(\omega) - \xi^*(\omega)) \rangle \leq a_n(\omega)\|\xi(\omega) - \xi^*(\omega)\|^2 \quad (n \in \mathbb{N}).$$

In late 50's Spacek [12] and Hans [7] initiated works on random operator theory or probabilistic analysis. since then, it has been an area for active research, a host of other researchers have done several work on random (probabilistic) fixed point theorems and applications (see e.g., Beg [1], Beg and Shahzad [2, 3], Benavides et.al [4], Bharucha-Reid [5, 6], Itoh [8, 9], Lin [10], Tan and Yuan [13], Xu [14, 15],)

In recent time, some authors have obtained solutions to real life problems using the deterministic model (see e.g., Bharuch-Reid [5, 6]).

Moore and Ofoedu [11] extended results of Beg [1] from the class of asymptotically nonexpansive random maps to more general class of asymptotically hemicontractive random maps.

In this paper, it is our purpose to construct a random explicit iteration scheme for approximation of common fixed points of finite families of N -uniformly L_i -Lipschitzian asymptotically hemicontractive maps.

Our theorems extended that of Moore and Ofoedu [11] from a single operator to a finite families of the operator and a host of others.

2. PRELIMINARIES

We shall make use of the following lemmas.

Lemma 2.1. *Let $\{\beta_n\}_{n=0}^\infty$ and $\{b_n\}_{n=0}^\infty$ be sequences of nonnegative real numbers satisfying the inequality*

$$\beta_{n+1} \leq \beta_n + b_n, n \geq 0$$

if $\sum_{n=0}^\infty b_n < \infty$ then $\lim_{n \rightarrow \infty} \beta_n$ exists.

Lemma 2.2. *Let E be a real normed linear space. Then for all $\xi(\omega), \eta(\omega) \in E$ and $j(\xi(\omega) + \eta(\omega)) \in J(\xi(\omega) + \eta(\omega))$ the following inequality holds.*

$$\|\xi(\omega) + \eta(\omega)\|^2 \leq \|\xi(\omega)\|^2 + 2\langle \eta(\omega), j(\xi(\omega) + \eta(\omega)) \rangle$$

3. MAIN RESULTS

If K is a nonempty closed convex subset of E and $\{T_i\}_{i=1}^N$ is a family of N uniformly L_i -Lipschitzian asymptotically hemicontractive self mappings of K , then $\xi_0(\omega) \in K$ and $\{\alpha_n\}_{n \geq 0} \subset (0, 1)$, the iteration process is generated as follows

$$\begin{aligned} \xi_1(\omega) &= (1 - \alpha_0)\xi_0(\omega) + \alpha_0 T_1(\omega)\xi_0(\omega), \\ \xi_2(\omega) &= (1 - \alpha_1)\xi_1(\omega) + \alpha_1 T_2(\omega)\xi_1(\omega), \\ &\vdots \\ \xi_N(\omega) &= (1 - \alpha_{N-1})\xi_{N-1}(\omega) + \alpha_{N-1} T_N(\omega)\xi_{N-1}(\omega), \\ \xi_{N+1}(\omega) &= (1 - \alpha_N)\xi_N(\omega) + \alpha_N T_1^2(\omega)\xi_N(\omega), \\ &\vdots \\ \xi_{2N}(\omega) &= (1 - \alpha_{2N-1})\xi_{2N-1}(\omega) + \alpha_{2N-1} T_N^2(\omega)\xi_{2N-1}(\omega), \\ \xi_{2N+1}(\omega) &= (1 - \alpha_{2N})\xi_{2N}(\omega) + \alpha_{2N} T_1^3(\omega)\xi_{2N}(\omega), \\ &\vdots \end{aligned}$$

Then the compact form of the iteration process is

$$\xi_{n+1}(\omega) = (1 - \alpha_n)\xi_n(\omega) + \alpha_n T_i^k(\omega)\xi_n(\omega), \quad n \geq 0, \omega \in \Omega \quad (3.1)$$

where $k = \{\frac{n-i}{N}\} + 1$.

Theorem 3.1. *Let E be a real Banach space and K , a nonempty closed convex subset of E . Let $\{T_i\}_{i=1}^N$ be N uniformly L_i -Lipschitzian asymptotically hemicontractive self mappings of K such that $F = \bigcap_{i=1}^N F(T_i) \neq \emptyset$. Let $\{\alpha_n\}_{n \geq 0}$ be a sequence in $(0,1)$ satisfying the conditions*

$$\begin{aligned} (i) & \sum_{n \geq 0} \alpha_n = \infty; \\ (ii) & \sum_{n \geq 0} \alpha_n^2 < \infty; \\ (iii) & \sum_{n \geq 0} \alpha_n (a_{in}(\omega) - 1) < \infty. \end{aligned}$$

Then the explicit iterative sequence $\{\xi_n(\omega)\}_{n \geq 0}$ generated from an arbitrary $\xi_0(\omega) \in K$ by (3.1) converges strongly surely to a random common fixed point of the family $\{T_i\}_{i=1}^N$ if and only if $\liminf_{n \rightarrow \infty} d(\xi_n(\omega), F) = 0$ almost surely in Ω .

Proof. We have that

$$\|\xi_{n+1}(\omega) - \xi^*(\omega)\|^2 = \|(1 - \alpha_n)(\xi_n(\omega) - \xi^*(\omega)) + \alpha_n(T_i^k(\omega)\xi_n(\omega) - \xi^*(\omega))\|^2$$

and

$$\begin{aligned} & \|\xi_{n+1}(\omega) - \xi^*(\omega)\|^2 \\ & \leq \|(1 - \alpha_n)(\xi_n(\omega) - \xi^*(\omega))\|^2 \\ & \quad + 2\alpha_n \langle (T_i^k(\omega)\xi_n(\omega) - \xi^*(\omega)), j(\xi_{n+1}(\omega) - \xi^*(\omega)) \rangle \\ & \leq (1 - \alpha_n)^2 \|\xi_n(\omega) - \xi^*(\omega)\|^2 \\ & \quad - 2\alpha_n \langle \xi_{n+1}(\omega) - T_i^k(\omega)\xi_{n+1}(\omega), j(\xi_{n+1}(\omega) - \xi^*(\omega)) \rangle \\ & \quad + 2\alpha_n \langle T_i^k(\omega)\xi_n(\omega) - T_i^k(\omega)\xi_{n+1}(\omega), j(\xi_{n+1}(\omega) - \xi^*(\omega)) \rangle \\ & \quad + 2\alpha_n \langle \xi_{n+1}(\omega) - \xi^*(\omega), j(\xi_{n+1}(\omega) - \xi^*(\omega)) \rangle \\ & = (1 - \alpha_n)^2 \|\xi_n(\omega) - \xi^*(\omega)\|^2 + 2\alpha_n \{ (a_{in}(\omega) - 1) \|\xi_{n+1}(\omega) - \xi^*(\omega)\|^2 \} \\ & \quad + 2\alpha_n \|T_i^k(\omega)\xi_n(\omega) - T_i^k(\omega)\xi_{n+1}(\omega)\| \|\xi_{n+1}(\omega) - \xi^*(\omega)\| \\ & \quad + 2\alpha_n \|\xi_{n+1}(\omega) - \xi^*(\omega)\|^2. \end{aligned} \quad (3.2)$$

Moreover

$$\|T_i^k(\omega)\xi_n(\omega) - T_i^k(\omega)\xi_{n+1}(\omega)\| \leq L_i(1 + L_i)\alpha_n \|\xi_n(\omega) - \xi^*(\omega)\|$$

Also,

$$\|\xi_{n+1}(\omega) - \xi^*(\omega)\| = [1 + (1 + L_i)\alpha_n] \|\xi_n(\omega) - \xi^*(\omega)\|$$

Therefore, (3.2) gives

$$\begin{aligned}
 \|\xi_{n+1}(\omega) - \xi^*(\omega)\|^2 &\leq (1 - \alpha_n)^2 \|\xi_n(\omega) - \xi^*(\omega)\|^2 \\
 &\quad + 2\alpha_n(a_{in}(\omega) - 1)(\alpha_n + \alpha_n L_i + 1)^2 \|\xi_n(\omega) - \xi^*(\omega)\|^2 \\
 &\quad + 2\alpha_n^2(1 + L_i)L_i[\alpha_n(L_i + 1) + 1] \|\xi_n(\omega) - \xi^*(\omega)\|^2 \\
 &\quad + 2\alpha_n[\alpha_n^2(L_i + 1)^2 + 2\alpha_n(1 + L_i)] \|\xi_n(\omega) - \xi^*(\omega)\|^2 \\
 &\quad + 2\alpha_n \|\xi_n(\omega) - \xi^*(\omega)\|^2 \\
 &= (1 + \alpha_n^2) \|\xi_n(\omega) - \xi^*(\omega)\|^2 \\
 &\quad + 2\alpha_n(a_{in}(\omega) - 1)[\alpha_n(L_i + 1) + 1]^2 \|\xi_n(\omega) - \xi^*(\omega)\|^2 \\
 &\quad + 2\alpha_n^2(1 + L_i)L_i[\alpha_n(L_i + 1) + 1] \|\xi_n(\omega) - \xi^*(\omega)\|^2 \\
 &\quad + 2\alpha_n[\alpha_n^2(L_i + 1)^2 + 2\alpha_n(1 + L_i)] \|\xi_n(\omega) - \xi^*(\omega)\|^2 \\
 &= (1 + \gamma_{in}) \|\xi_n(\omega) - \xi^*(\omega)\|^2, \tag{3.3}
 \end{aligned}$$

where

$$\begin{aligned}
 \gamma_{in}(\omega) &= \{\alpha_n^2 + 2\alpha_n(a_{in}(\omega) - 1)[\alpha_n(1 + L_i) + 1]^2 \\
 &\quad + 2\alpha_n^2(L_i + 1)L_i[\alpha_n(1 + L_i) + 1] + 2\alpha_n[\alpha_n^2(1 + L_i)^2 + 2\alpha_n(1 + L_i)]\}
 \end{aligned}$$

We observe that

$$\sum_{n \geq 0}^{\infty} \gamma_{in}(\omega) < \infty \text{ almost surely in } \Omega$$

therefore, from (3.3) we have

$$\begin{aligned}
 \|\xi_{n+1}(\omega) - \xi^*(\omega)\|^2 &\leq \prod_{j=0}^n (1 + \gamma_{ij}(\omega)) \|\xi_0(\omega) - \xi^*(\omega)\|^2 \\
 &\leq e^{\sum_{j=0}^{\infty} \gamma_{ij}(\omega)} \|\xi_0(\omega) - \xi^*(\omega)\|^2
 \end{aligned}$$

therefore,

$$\|\xi_{n+1}(\omega) - \xi^*(\omega)\| \leq M \quad (n \in \mathbb{N})$$

since $\|\xi_{n+1}(\omega) - \xi^*(\omega)\| \leq M$ for some $M > 0$, now, we observe that if we set

$$\beta_n = \|\xi_n(\omega) - \xi^*(\omega)\|^2 \text{ and } b_n = \gamma_{in}(\omega)M^2.$$

Then by Lemma 2.1

$$\lim_{n \rightarrow \infty} \|\xi_n(\omega) - \xi^*(\omega)\| \text{ exists almost surely in } \Omega \tag{3.4}$$

If from (3.3), we have that

$$\|\xi_{n+1}(\omega) - \xi^*(\omega)\|^2 \leq [1 + \alpha_n^2 + \lambda_{in}(\omega)] \|\xi_n(\omega) - \xi^*(\omega)\|^2$$

where $\lambda_{in}(\omega) = \gamma_{in}(\omega) - \alpha_n^2$, i.e. $\alpha_n^2 + \lambda_{in}(\omega) = \gamma_{in}(\omega)$ then

$$\begin{aligned}
\|\xi_{n+1}(\omega) - \xi^*(\omega)\| &\leq [1 + \alpha_n^2 + \lambda_{in}(\omega)]^{\frac{1}{2}} \|\xi_n(\omega) - \xi^*(\omega)\| \\
&\leq (1 + \alpha_n^2) \|\xi_n(\omega) - \xi^*(\omega)\| + \lambda_{in}(\omega) \|\xi_n(\omega) - \xi^*(\omega)\| \\
&\leq (1 + \alpha_n^2) \|\xi_n(\omega) - \xi^*(\omega)\| + \mu_{in}(\omega)
\end{aligned}$$

where $\mu_{in}(\omega) = \lambda_{in}(\omega)M = (\gamma_{in}(\omega) - \alpha_n^2)M$ so we observe that

$$\sum_{n \geq 0}^{\infty} \mu_{in}(\omega) < \infty \quad \text{almost surely in } \Omega$$

And for all $n, m \in \mathbb{N}$ we have

$$\begin{aligned}
\|\xi_{n+m}(\omega) - \xi^*(\omega)\| &\leq (1 + \alpha_{n+m-1}^2) \|\xi_{n+m+1}(\omega) - \xi^*(\omega)\| + \mu_{in+m-1}(\omega) \\
&\leq (1 + \alpha_{n+m-1}^2)(1 + \alpha_{n+m-2}^2) \|\xi_{n+m+2}(\omega) - \xi^*(\omega)\| \\
&\quad + (1 + \alpha_{n+m-1}^2) \mu_{in+m-2}(\omega) + \mu_{in+m-1}(\omega) \\
&= \prod_{j=n}^{n+m-1} (1 + \alpha_j^2) \|\xi_n(\omega) - \xi^*(\omega)\| + \prod_{j=n}^{n+m-1} (1 + \alpha_j^2) \sum_{j=n}^{n+m-1} \mu_{ij}(\omega) \\
&\leq e^{\sum_{j=n}^{n+m-1} \alpha_j^2} \|\xi_n(\omega) - \xi^*(\omega)\| + e^{\sum_{j=n}^{n+m-1} \alpha_j^2} \sum_{j=n}^{n+m-1} \mu_{ij}(\omega) \\
&= D \|\xi_n(\omega) - \xi^*(\omega)\| + D \sum_{j=n}^{n+m-1} \mu_{ij}(\omega) < \infty \tag{3.5}
\end{aligned}$$

where $D = \exp\left(\sum_{j=1}^{\infty} \alpha_j^2\right)$.

Thus, taking infimum over $\xi^*(\omega) \in F(\omega)$, we obtain

$$d(\xi_{n+1}(\omega), F(\omega)) \leq (1 + \alpha_n^2) d(\xi_n(\omega), F(\omega)) + \mu_{in}(\omega)$$

since the $\liminf_{n \rightarrow \infty} d(\xi_n(\omega), F(\omega)) = 0$ almost surely in Ω .

Thus, we have from (3.4), that $\lim_{n \rightarrow \infty} d(\xi_n(\omega), F(\omega)) = 0$ almost surely in Ω . That is

$$\mu\left(\{\omega \in \Omega : \liminf_{n \rightarrow \infty} d(\xi(\omega), F(\omega)) = 0\}\right) = 1$$

implies

$$\mu\left(\{\omega \in \Omega : \lim_{n \rightarrow \infty} d(\xi(\omega), F(\omega)) = 0\}\right) = 1$$

It suffices to show that $\{\xi_n(\omega)\}_{n \geq 0}$ is Cauchy.

Let $\epsilon > 0$ be given, since $\lim_{n \rightarrow \infty} d(\xi_n(\omega), F(\omega)) = 0$ almost surely in Ω and

$\sum_{i=1}^{\infty} \delta_i(\omega) < \infty$ there exists a positive integer N_1 such that for all $n \geq N_1$,

$$d(\xi_n(\omega), F(\omega)) < \frac{\epsilon}{3D}$$

and

$$\sum_{i=1}^{\infty} \delta_i(\omega) < \frac{\epsilon}{6D}.$$

In particular there exists $\xi^*(\omega) \in F(\omega)$ such that $d(\xi_{N_1}(\omega), \xi^*(\omega)) < \frac{\epsilon}{3D}$.

Now from (3.5), we have that for all $n \geq N_1$,

$$\begin{aligned} \|\xi_{n+m}(\omega) - \xi_n(\omega)\| &\leq \|\xi_{n+m}(\omega) - \xi^*(\omega)\| + \|\xi_n(\omega) - \xi^*(\omega)\| \\ &\leq D\|\xi_{N_1}(\omega) - \xi^*(\omega)\| + D \sum_{i=N_1}^{N_1+m-1} \delta_i(\omega) \\ &\quad + D\|\xi_{N_1}(\omega) - \xi^*(\omega)\| + D \sum_{i=N_1}^{N_1+m-1} \delta_i(\omega) \\ &< \epsilon \end{aligned}$$

Hence, $\lim_{n \rightarrow \infty} \xi_n(\omega)$ exists almost surely in Ω (Since E is complete).

Suppose that $\lim_{n \rightarrow \infty} \xi_n(\omega) = \xi^*(\omega)$ we show that $\xi^*(\omega) \in F(\omega)$. But given $\epsilon_2 > 0$ there exists a positive $N_2 \geq N_1$ such that for all $n \geq N_2$

$$\begin{aligned} &\mu\left(\{\omega \in \Omega : \|\xi_n(\omega) - \xi^*(\omega)\| \right. \\ &\quad \left. < \frac{\epsilon_2}{2(1+L)}\} \cap \{\omega \in \Omega : d(\xi_n(\omega), F(\omega)) < \frac{\epsilon_2}{2(1+3L)}\}\right) = 1 \end{aligned}$$

Thus, there exists $\eta^*(\omega) \in F(\omega)$ such that

$$\begin{aligned} &\mu\left(\{\omega \in \Omega : \|\xi_{N_2}(\omega) - \eta^*(\omega)\| \right. \\ &\quad \left. = d(\xi_{N_2}(\omega), \eta^*(\omega))\} \cap \{\omega \in \Omega : d(\xi_{N_2}(\omega), \eta^*(\omega)) < \frac{\epsilon_2}{2(1+3L)}\}\right) = 1 \end{aligned}$$

with the following estimates

$$\begin{aligned} \|T(\omega)\xi^*(\omega) - \xi^*(\omega)\| &\leq \|T(\omega)\xi^*(\omega) - \eta^*(\omega)\| + 2\|T(\omega)\xi_{N_2}(\omega) - \eta^*(\omega)\| \\ &\quad + \|\xi_{N_2}(\omega) - \eta^*(\omega)\| + \|\xi_{N_2}(\omega) - \xi^*(\omega)\| \\ &\leq L\|\xi^*(\omega) - \eta^*(\omega)\| + 2L\|\xi_{N_2}(\omega) - \eta^*(\omega)\| \\ &\quad + \|\xi_{N_2}(\omega) - \eta^*(\omega)\| + \|\xi_{N_2}(\omega) - \xi^*(\omega)\| \\ &\leq (1+L)\|\xi_{N_2}(\omega) - \xi^*(\omega)\| + (1+3L)\|\xi_{N_2}(\omega) - \eta^*(\omega)\| \\ &< \epsilon_2 \end{aligned}$$

Since $\epsilon_2 > 0$ is arbitrary we have that

$$\mu(\{\omega \in \Omega : T(\omega)\xi^*(\omega) = \xi^*(\omega)\}) = 1$$

□

Theorem 3.2. *Let E be a real Banach space and K , a nonempty closed convex subset of E . Let $\{T_i\}_{i=1}^N$ be N uniformly L_i -Lipschitzian asymptotically hemicontractive self mappings of K such that $F = \bigcap_{i=1}^N F(T_i) \neq \emptyset$. Let $\{\alpha_n\}_{n \geq 0}$ be a*

sequence in $(0,1)$ satisfying the conditions (i) $\sum_{n \geq 0} \alpha_n = \infty$ (ii) $\sum_{n \geq 0} \alpha_n^2 < \infty$ (iii)

$\sum_{n \geq 0} \alpha_n (a_{in}(\omega) - 1) < \infty$. Then the explicit iterative sequence $\{\xi_n(\omega)\}_{n \geq 0}$ generated from an arbitrary $\xi_0(\omega) \in K$, $\omega \in \Omega$ by (3.1) converges strongly to a common fixed point of the family $\{T_i\}_{i=1}^N$ if and only if there exists an infinite subsequence of $\{\xi_n(\omega)\}_{n \geq 0}$ which converges strongly to a random common fixed point of the family $\{T_i\}_{i=1}^N$.

Proof. Let $\xi^*(\omega) \in F(\omega)$ and $\{\xi_{n_j}(\omega)\}_{j \geq 0}$ be a subsequence of $\{\xi_n(\omega)\}_{n \geq 0}$ such that $\lim_{j \rightarrow \infty} \|\xi_{n_j}(\omega) - \xi^*(\omega)\| = 0$ almost surely, since $\lim_{n \rightarrow \infty} \|\xi_n(\omega) - \xi^*(\omega)\|$ exists almost surely, then, $\lim_{n \rightarrow \infty} \|\xi_n(\omega) - \xi^*(\omega)\| = 0$ almost surely. \square

Remark 3.3. Our theorems unify, extend and generalize the corresponding results of Beg [1], Beg and Shahzad [2], Moore and Ofoedu [11], Xu [14] and host of other results recently announced, to more general class of finite families of N -uniformly L_i -Lipschitzian asymptotically hemicontractive Random maps.

REFERENCES

1. I.Beg, *Fixed points of random operators satisfying semicontractivity conditions*, Math. Japan **46(1)** (1997), 151–155.
2. I.Beg and N.Shahzad, *Random fixed point theorems for nonexpansive and contractive type random operators on Banach spaces*, J. Math. Stochas Anal. **7(4)** (1994), 566–580.
3. I.Beg and N.Shahzad, *Some random approximation theorem with applications*, Nonlinear Analysis, **35** (1999), 609–616.
4. T.D. Benavides, G.L.Acedo and H.K.Xu, *Random fixed points set-valued operators*, Proc. Amer. Math. Soc. **121(3)** (1996), 831–838.
5. A.T.Bharucha-Reid, *Random integral equations*, academic press, New York, 1972.
6. A.T.Bharucha-Reid, *Fixed points theorems in probabilistic analysis*, Bull. Amer. Math. Soc., **82** (1976), 641–657
7. O. Hans, *Random operator equations*, In Proc. Berkely symposium on mathematical, statistics and probability Vol.11, part 1, 185–200, University of California Press, Berkely 1961.
8. S.Itoh, *A random fixed point theorem for a multivalued contraction mappings*, Pacific J.Math., **68** (1977), 85–90.
9. S.Itoh, *Random fixed point theorems with application to random dif and only ifferential equations in Banach spaces*, J. Math. Anal. Appl., **68** (1979), 261–373.
10. T.C. Lin, *Random approximations and random fixed point theorems for a non-self maps*, Proc. Amer. Mth. Soc., **103** (1998), 1129–1135.
11. C. Moore and E.U. Ofoedu, *Convergence of an iterative process to random fixed points of uniformly l -Lipschitzian asymptotically hemicontractive random maps in Banach Spaces*, J. Nig. Math Soc. **25** (2006), 87–94.
12. A.Spacek, *Zufallige gleichugen*, Czechoslovak Math. J. **5** (1955), 162–466.
13. K.K. Tan, X.Z.Yaun, *Random fixed point theorems and approximations*, Stochas. Anal.Appl., **15(1)** (1997), 103–123.
14. H.K. Xu, *Some random fixed point theorems for condensing and nonexpansive operators* Proc. Amer. Math. Soc., **110** (1990), 395–400.
15. H.K. Xu, *A random fixed point theorem for multivalued nonexpansive operators in uniformly convex Banach Spaces* Proc. Amer. Math. Soc., **117** (1993), 1089–1092.

^{1,3} DEPARTMENT OF MATHEMATICS, NNAMDI AZIKIWE UNIVERSITY, AWKA, P. M. B. 5025, AWKA, ANAMBRA STATE, NIGERIA.

E-mail address: drchikamoore@yahoo.com and debosky2002@yahoo.com

² DEPARTMENT OF MATHEMATICS, NNAMDI AZIKIWE UNIVERSITY, AWKA, P. M. B. 5025, AWKA, ANAMBRA STATE, NIGERIA.

NON-LINEAR ANALYSIS RESEARCH GROUP, UNIZIK AWKA, NIGERIA

E-mail address: chimexnnanwa@yahoo.com