



## A FIXED POINT THEOREM ON CONE METRIC SPACES WITH NEW TYPE CONTRACTIVITY

ISHAK ALTUN<sup>1\*</sup>, MUJAHID ABBAS<sup>2</sup> AND HAKAN SIMSEK<sup>3</sup>

Communicated by M. A. Japón Pineda

**ABSTRACT.** In the present work, a common fixed point theorem for self maps on cone metric spaces is proved. Also two examples, which shows that our main theorem is generalized version of main theorems of [A. Branciari, Int. J. Math. Math. Sci., 29 (2002), no. 9, 531–536] and [L.G. Huang and X. Zhang, J. Math. Anal. Appl. 332 (2007), no. 2, 1468–1476], are given.

### 1. INTRODUCTION

There are a lot of generalizations of Banach fixed point principle in the literature. One of the most interesting generalization of it has been given by Branciari [10]. Branciari has made this by taking integral type contraction instead of ordinary contraction. He has given an example showing that the integral type contraction is more general than ordinary contraction. This interesting fixed point result is as follows:

**Theorem 1.1** ([10]). *Let  $(X, d)$  be a complete metric space,  $k \in (0, 1)$  and  $T : X \rightarrow X$  be mapping such that for each  $x, y \in X$  one has*

$$\int_0^{d(Tx, Ty)} f(t) dt \leq k \int_0^{d(x, y)} f(t) dt \quad (1.1)$$

where  $f : [0, \infty) \rightarrow [0, \infty]$  is a Lebesgue integrable mapping which is finite integral on each compact subset of  $[0, \infty)$ , non-negative and such that for each  $t > 0$ ,

---

*Date:* Received: 1 July 2010; Revised: 13 October 2010; Accepted: 13 November 2010.

\* Corresponding author.

2010 *Mathematics Subject Classification.* Primary 54H25; Secondary 47H10.

*Key words and phrases.* Fixed point, cone metric space, integral type contraction.

$\int_0^t f(s)ds > 0$ . Then  $T$  has a unique fixed point  $z \in X$  such that for each  $x \in X$ ,  $\lim_{n \rightarrow \infty} T^n x = z$ .

Since then many fixed point theorists have given some fixed point and common fixed point results using integral type contractions (see for example [3, 4, 5, 18, 23, 24, 27]).

On the other hand, existence of fixed point in cone metric space has been considered recently in [14]. Before giving this result, we recall some definitions and properties of cone metric spaces.

Let  $E$  be a real Banach space and  $P$  be a subset of  $E$ . By  $\theta$  we denote the zero element of  $E$  and by  $IntP$  the interior of  $P$ . The subset  $P$  is called a cone if and only if:

- (i)  $P$  is closed, nonempty and  $P \neq \{\theta\}$ ,
- (ii)  $a, b \in \mathbb{R}$ ,  $a, b \geq 0$ ,  $x, y \in P \implies ax + by \in P$ ,
- (iii)  $x \in P$  and  $-x \in P \implies x = \theta$ .

A cone  $P$  is called solid if it contains interior points, that is, if  $IntP \neq \emptyset$ . Given a cone  $P \subset E$ , we define a partial ordering  $\preceq$  with respect to  $P$  by  $x \preceq y$  if and only if  $y - x \in P$ . We shall write  $x \prec y$  if  $x \preceq y$  and  $x \neq y$ , we shall write  $x \ll y$  if  $y - x \in IntP$ . The cone  $P$  is called normal if there is a number  $K > 0$  such that for all  $x, y \in E$ ,  $\theta \preceq x \preceq y$  implies  $\|x\| \leq K \|y\|$ . The least positive number satisfying the above is called the normal constant of  $P$ .

The following example shows that there are non-normal cones.

**Example 1.2** ([22]). Let  $E = C_R^2([0, 1])$  with the norm  $\|f\| = \|f\|_\infty + \|f'\|_\infty$  and consider the cone  $P = \{f \in E : f \geq 0\}$ . For each  $K > 1$ , put  $f(x) = x$  and  $g(x) = x^{2K}$ . Then  $0 \leq g \leq f$ ,  $\|f\| = 2$  and  $\|g\| = 2K + 1$ . Since  $K \|f\| < \|g\|$ ,  $K$  is not normal constant of  $P$ . Therefore,  $P$  is a non-normal cone.

In the following we always suppose that  $E$  is a real Banach space,  $P$  is a cone in  $E$  with  $IntP \neq \emptyset$  and  $\preceq$  a partial ordering with respect to  $P$ .

**Definition 1.3** ([14]). Let  $X$  be a nonempty set. Suppose the mapping  $d : X \times X \rightarrow E$  satisfies

- (d<sub>1</sub>)  $\theta \prec d(x, y)$  for all  $x, y \in X$  with  $x \neq y$  and  $d(x, y) = \theta$  if and only if  $x = y$ ,
- (d<sub>2</sub>)  $d(x, y) = d(y, x)$  for all  $x, y \in X$ ,
- (d<sub>3</sub>)  $d(x, y) \preceq d(x, z) + d(z, y)$  for all  $x, y, z \in X$ .

Then  $d$  is called a cone metric on  $X$  and  $(X, d)$  is called a cone metric space.

It is obvious that the cone metric spaces generalize metric spaces.

**Example 1.4** ([14]). Let  $E = \mathbb{R}^2$ ,  $P = \{(x, y) \in E \mid x, y \geq 0\}$ ,  $X = \mathbb{R}$  and  $d : X \times X \rightarrow E$  such that  $d(x, y) = (|x - y|, \alpha |x - y|)$ , where  $\alpha \geq 0$  is a constant. Then  $(X, d)$  is a cone metric space.

**Definition 1.5** ([14]). Let  $(X, d)$  be a cone metric space. Let  $\{x_n\}$  be a sequence in  $X$  and  $x \in X$ . If for every  $c \in E$  with  $\theta \ll c$  there is  $N$  such that for all  $n > N$ ,  $d(x_n, x) \ll c$ , then  $\{x_n\}$  is said to be convergent and  $\{x_n\}$  converges to  $x$  and  $x$  is the limit of  $\{x_n\}$ . We denote this by  $\lim_{n \rightarrow \infty} x_n = x$  or  $x_n \rightarrow x$  as  $n \rightarrow \infty$ . If

for every  $c \in E$  with  $\theta \ll c$  there is  $N$  such that for all  $n, m > N$ ,  $d(x_n, x_m) \ll c$ , then  $\{x_n\}$  is called a Cauchy sequence in  $X$ .  $(X, d)$  is a complete cone metric space if every Cauchy sequence is convergent.

**Lemma 1.6** ([14]). *Let  $(X, d)$  be a cone metric space,  $P$  be a normal cone and let  $\{x_n\}$  be a sequence in  $X$ . Then*

- (i)  $\{x_n\}$  converges to  $x$  if and only if  $d(x_n, x) \rightarrow \theta$  ( $n \rightarrow \infty$ ),
- (ii)  $\{x_n\}$  is a Cauchy sequence if and only if  $d(x_n, x_m) \rightarrow \theta$  ( $n, m \rightarrow \infty$ ).

Let  $(X, d)$  be a cone metric space,  $T : X \rightarrow X$  and  $x_0 \in X$ . Then the function  $T$  is continuous at  $x_0$  if for any sequence  $x_n \rightarrow x_0$  we have  $Tx_n \rightarrow Tx_0$  [15].

Now we give the result of Huang and Zhang.

**Theorem 1.7** ([14]). *Let  $(X, d)$  be a complete cone metric space,  $P$  be a normal cone with normal constant  $K$ . Suppose the mapping  $f : X \rightarrow X$  satisfies the contractive condition*

$$d(fx, fy) \preceq kd(x, y) \text{ for all } x, y \in X, \quad (1.2)$$

where  $k \in [0, 1)$  is a constant. Then  $f$  has a unique fixed point in  $X$ .

Also, Huang and Zhang [14] gave an example showing that Theorem 1.7 is a generalization of Banach fixed point principle.

After the definition of the concept of cone metric space, fixed point theory on these spaces has been developing (see for example [1, 2, 6, 7, 8, 9, 11, 13, 16, 17, 19, 20, 21, 22, 25, 26]).

The aim of this work is to give a fixed point result using the ideas of Branciari [10] (or Zhang [27]) and Huang and Zhang [14]. Our result a generalized version of both Theorem 1.1 and Theorem 1.7 in cone metric spaces.

Let  $(X, d)$  be a cone metric space with the real Banach space  $E$  and the cone  $P$ . Let  $Q_a = \{u \in P : \|u\| < a, a \in (0, \infty]\}$  and let  $F : Q_a \rightarrow P$  satisfy that

- (F<sub>1</sub>)  $F(\theta) = \theta$  and  $\theta \prec F(u)$  for each  $u \in Q_a \setminus \{\theta\}$ ,
- (F<sub>2</sub>)  $F$  is nondecreasing on  $Q_a$ , that is,  $u, v \in Q_a$  and  $u \preceq v$  implies  $F(u) \preceq F(v)$ ,
- (F<sub>3</sub>)  $F$  is continuous.
- (F<sub>4</sub>)  $F(u_n) \rightarrow \theta$  implies  $u_n \rightarrow \theta$  as  $n \rightarrow \infty$ .

Define  $\mathcal{F}(E, P, Q_a) = \{F \mid F : Q_a \rightarrow P \text{ satisfies (F}_1\text{)-(F}_4\text{)}\}$ .

**Example 1.8.**

- (a) Let  $E$  be a real Banach space and  $P$  be a cone in  $E$ . Define  $F : Q_a \rightarrow P$  with  $F(u) = u$ , then  $F \in \mathcal{F}(E, P, Q_a)$  for all  $a \in (0, \infty]$ .
- (b) Let  $E = \mathbb{R}$  be real numbers,  $P_1 = [0, \infty)$ . Suppose that  $\phi$  is nonnegative, Lebesgue integrable on  $[0, \infty)$  and satisfies

$$\int_0^\varepsilon \phi(s) ds > 0 \text{ for each } \varepsilon > 0.$$

Let  $F(u) = \int_0^u \phi(s) ds$ . Then  $F \in \mathcal{F}(\mathbb{R}, P_1, Q_a)$  for all  $a \in (0, \infty]$ .

- (c) Let  $P_2 = \{(x, y) \in \mathbb{R}^2 : x, y \geq 0\}$  and  $F_1(u) = (\|u\|, \alpha \|u\|)$ . Then  $F_1 \in \mathcal{F}(\mathbb{R}^2, P_2, Q_a)$  for all  $a \in (0, \infty]$ , where  $\alpha \geq 0$ . Similarly, let

$$F_2(u) = \left( \int_0^{\|u\|} \phi(s) ds, \alpha \int_0^{\|u\|} \phi(s) ds \right).$$

Then  $F_2 \in \mathcal{F}(\mathbb{R}^2, P_2, Q_a)$  for all  $a \in (0, \infty]$ , where  $\alpha$  and  $\phi$  as above.

- (d) Let  $E = C_{\mathbb{R}}[0, 1]$  with the supremum norm,  $P_3 = \{f \in E : f \geq 0\}$  and  $F(u) = \frac{\|u\|}{1 + \|u\|} \psi$ . Then  $F \in \mathcal{F}(E, P_3, Q_a)$  for all  $a \in (0, \infty]$ , where  $\psi \in \text{Int}P_3$ .

- (e) Let  $E = l^1$ ,  $P_4 = \{\{x_n\} \in E : x_n \geq 0, \text{ for all } n\}$  and  $F(u) = \frac{\|u\|}{2^n}$ . Then  $F \in \mathcal{F}(E, P_4, Q_a)$  for all  $a \in (0, \infty]$ .

**Definition 1.9** ([11]). Let  $P$  be a cone on a Banach space  $E$ . A nondecreasing function  $\varphi : P \rightarrow P$  is called  $\varphi$ -map if

- ( $\varphi_1$ )  $\varphi(\theta) = \theta$  and  $\theta \prec \varphi(\omega) \prec \omega$  for  $\omega \in P \setminus \{\theta\}$ ,
- ( $\varphi_2$ )  $\omega \in \text{Int}P$  implies  $\omega - \varphi(\omega) \in \text{Int}P$ ,
- ( $\varphi_3$ )  $\lim_{n \rightarrow \infty} \varphi^n(\omega) = \theta$  for every  $\omega \in P \setminus \{\theta\}$ .

There is an example of  $\varphi$ -map in [12].

## 2. MAIN RESULTS

Let  $E$  be a real Banach space,  $P$  be a cone in  $E$  and  $\preceq$  is partial ordering with respect to  $P$ . We consider the following condition for convenience:

$$\begin{cases} \text{If } \{u_n\} \text{ is a sequence in } E \text{ such that } \|u_n\| \text{ is convergent,} \\ \text{then } \{u_n\} \text{ has an upper limit in } E \text{ w.r.t } \preceq. \end{cases} \quad (2.1)$$

It is clear that  $\mathbb{R}^2$  is satisfied the condition (2.1) with the cone  $P_2$ . In the following theorem we assume that  $E$  satisfies the condition (2.1).

**Theorem 2.1.** *Let  $(X, d)$  be a complete cone metric space with the normal cone  $P$  on a Banach space  $E$ . Let  $D = \sup\{\|d(x, y)\| : x, y \in X\}$  and set  $a = D$  if  $D = \infty$  and  $a > D$  if  $D < \infty$ . Suppose that  $T, S : X \rightarrow X$ ,  $F \in \mathcal{F}(E, P, Q_a)$  satisfy*

$$F(d(Tx, Sy)) \preceq \varphi(F(d(x, y))), \quad (2.2)$$

for all  $x, y \in X$ , where  $\varphi$  is a continuous  $\varphi$ -map. Then  $T$  and  $S$  have a unique common fixed point in  $X$ .

*Proof.* First we prove that any fixed point of  $T$  is also a fixed point of  $S$ , and conversely. If  $Tx = x$  but  $Sx \neq x$ , then we have from (2.2)

$$F(d(x, Sx)) = F(d(Tx, Sx)) \preceq \varphi(F(\theta)) = \theta,$$

which is a contradiction. So  $Sx = x$ . Similarly, if  $Sx = x$ , then  $Tx = x$ .

Now we show that if  $T$  and  $S$  have a common fixed point, then the fixed point is unique. Let  $Tx = Sx = x$  and  $Ty = Sy = y$ . If  $x \neq y$ , then we have from (2.2)

$$F(d(x, y)) = F(d(Tx, Sy)) \preceq \varphi(F(d(x, y))),$$

which is a contradiction. Thus  $x = y$ .

Let  $x_0 \in X$  and let  $x_{2n+1} = Tx_{2n}$ ,  $x_{2n+2} = Sx_{2n+1}$  for all  $n = 0, 1, 2, \dots$ . Suppose that for any  $n$ ,  $x_{n+1} \neq x_n$ . Otherwise  $T$  or  $S$  has a fixed point and the proof is complete. Now we show that

$$\begin{aligned} F(d(x_{n+1}, x_n)) &\preceq \varphi(F(d(x_n, x_{n-1}))) \\ &\prec F(d(x_n, x_{n-1})) \text{ for } n \geq 1. \end{aligned} \quad (2.3)$$

We have that

$$\begin{aligned} F(d(x_{2n+1}, x_{2n})) &= F(d(Tx_{2n}, Sx_{2n-1})) \\ &\preceq \varphi(F(d(x_{2n}, x_{2n-1}))) \\ &\prec F(d(x_{2n}, x_{2n-1})). \end{aligned} \quad (2.4)$$

Similarly we have

$$\begin{aligned} F(d(x_{2n+2}, x_{2n+1})) &= F(d(Sx_{2n+1}, Tx_{2n})) \\ &\preceq \varphi(F(d(x_{2n+1}, x_{2n}))) \\ &\prec F(d(x_{2n+1}, x_{2n})). \end{aligned} \quad (2.5)$$

Therefore from (2.4) and (2.5) we have (2.3). Then from (2.3)

$$\begin{aligned} F(d(x_{n+1}, x_n)) &\preceq \varphi(F(d(x_n, x_{n-1}))) \\ &\preceq \varphi^2(F(d(x_{n-1}, x_{n-2}))) \\ &\quad \vdots \\ &\preceq \varphi^n(F(d(x_1, x_0))). \end{aligned}$$

Fix  $\theta \ll c$  and we choose a positive real number  $\delta$  such that  $c + N_\delta(\theta) \subset \text{Int}P$ , where  $N_\delta(\theta) = \{x \in P : \|x\| < \delta\}$ . Also, choose a natural number  $n_0$  such that  $\varphi^n(F(d(x_1, x_0))) \in N_\delta(\theta)$ , for all  $n \geq n_0$ . Then

$$\varphi^n(F(d(x_1, x_0))) \ll c,$$

for all  $n \geq n_0$ . Consequently  $F(d(x_{n+1}, x_n)) \ll c$ , for all  $n \geq n_0$ . From (F<sub>4</sub>), we have, for all  $\theta \ll c$ , there exists a natural number  $n_0$  such that for all  $n \geq n_0$

$$d(x_{n+1}, x_n) \ll c.$$

Now we claim that  $\{x_n\}$  is a Cauchy sequence in  $X$ . If not, there exists  $\varepsilon > 0$  for which we can find subsequences  $\{x_{m_k}\}$  and  $\{x_{n_k}\}$  of  $\{x_n\}$  with  $n_k > m_k > k$  such that

$$\varepsilon \leq \|d(x_{m_k}, x_{n_k})\|.$$

Further, corresponding to odd numbers  $m_k$ , we can choose even numbers  $n_k$  in such a way that it is the smallest integer with  $n_k > m_k$  such that  $\|d(x_{m_k}, x_{n_k-2})\| < \varepsilon$  and  $\varepsilon \leq \|d(x_{m_k}, x_{n_k})\|$ . From

$$\begin{aligned} \varepsilon &\leq \|d(x_{m_k}, x_{n_k})\| \\ &\leq \|d(x_{m_k}, x_{n_k-2})\| + \|d(x_{n_k-2}, x_{n_k})\| < \varepsilon + \|d(x_{n_k-2}, x_{n_k})\|, \end{aligned}$$

it follows

$$\lim_{k \rightarrow \infty} \|d(x_{m_k}, x_{n_k})\| = \varepsilon.$$

Now

$$\|d(x_{n_{k-1}}, x_{m_{k-1}})\| \leq \|d(x_{n_{k-1}}, x_{n_k})\| + \|d(x_{n_k}, x_{m_k})\| + \|d(x_{m_k}, x_{m_{k-1}})\|$$

and

$$\|d(x_{n_k}, x_{m_k})\| \leq \|d(x_{n_k}, x_{n_{k-1}})\| + \|d(x_{n_{k-1}}, x_{m_{k-1}})\| + \|d(x_{m_{k-1}}, x_{m_k})\|$$

gives

$$\lim_{k \rightarrow \infty} \|d(x_{m_{k-1}}, x_{n_{k-1}})\| = \varepsilon.$$

Therefore from the condition (2.1) there exists  $\theta \ll c$  such that

$$\limsup_{k \rightarrow \infty} d(x_{m_k}, x_{n_k}) = \limsup_{k \rightarrow \infty} d(x_{m_{k-1}}, x_{n_{k-1}}) = c.$$

Setting  $x = x_{m_{k-1}}$  and  $y = x_{n_{k-1}}$  in (2.2), we obtain

$$\begin{aligned} F(d(x_{m_k}, x_{n_k})) &= F(d(Tx_{m_{k-1}}, Sx_{n_{k-1}})) \\ &\preceq \varphi(F(d(x_{m_{k-1}}, x_{n_{k-1}}))) \end{aligned}$$

which on take upper limit

$$F(c) \preceq \varphi(F(c)) \prec F(c)$$

a contradiction which shows that  $\{x_n\}$  is a Cauchy sequence in  $X$ , and so  $x_n \rightarrow u$ , for some  $u$  in  $X$ . Take  $x = x_{2n}$  and  $y = u$  in (2.2), we have

$$F(d(Tx_{2n}, Su)) \preceq \varphi(F(d(x_{2n}, u)))$$

and

$$F(d(u, Su)) = \varphi(F(\theta)) = \theta$$

implies that  $d(u, Su) = \theta$ , and  $u = Su$ .  $\square$

**Corollary 2.2.** *Let  $(X, d)$  be a complete cone metric space with the normal cone  $P$  on a Banach space  $E$ . Let  $D = \sup\{\|d(x, y)\| : x, y \in X\}$  and set  $a = D$  if  $D = \infty$  and  $a > D$  if  $D < \infty$ . Suppose that  $T : X \rightarrow X$ ,  $F \in \mathcal{F}(E, P, Q_a)$  satisfy*

$$F(d(Tx, Ty)) \preceq \varphi(F(d(x, y))), \quad (2.6)$$

for all  $x, y \in X$ , where  $\varphi$  is a continuous  $\varphi$ -map. Then  $T$  has a unique fixed point in  $X$ .

*Remark 2.3.* Corollary 2.2 is a generalized version of Theorem 1.1. Indeed, if we take  $E = \mathbb{R}$ ,  $P_1 = [0, \infty)$ ,  $F \in \mathcal{F}(\mathbb{R}, P_1, Q_a)$  as in Example 1.8 (b) and  $\varphi(\omega) = k\omega$  ( $k \in [0, 1)$ ) in Corollary 2.2, we have Theorem 1.1.

The following example supports this remark.

**Example 2.4.** Let  $E = \mathbb{R}^2$  with the Euclidean norm, and  $P_2 = \{(x, y) \in \mathbb{R}^2 : x, y \geq 0\}$  a cone in  $E$ . Let

$$X = \{(x, 0) \in \mathbb{R}^2 : 0 \leq x \leq 1\} \cup \{(0, x) \in \mathbb{R}^2 : 0 \leq x \leq 1\}$$

and let  $d : X \times X \rightarrow E$  defined by (as in [14])

$$\begin{aligned} d((x, 0), (y, 0)) &= \left(\frac{4}{3}|x - y|, |x - y|\right), \\ d((0, x), (0, y)) &= \left(|x - y|, \frac{2}{3}|x - y|\right) \\ d((x, 0), (0, y)) &= d((0, y), (x, 0)) = \left(\frac{4}{3}x + y, x + \frac{2}{3}y\right). \end{aligned}$$

Then  $(X, d)$  is a complete cone metric space.

Let mapping  $T : X \rightarrow X$  with  $T(x, 0) = (0, x)$  and  $T(0, x) = \left(\frac{x}{2}, 0\right)$ . It is easy to see that  $T$  satisfies the condition (2.6) of Corollary 2.2 with  $F(u) = u$  and  $\varphi(\omega) = \frac{3}{4}\omega$ . Therefore, we can apply Corollary 2.2 to this example. Note that, the conditions of Theorem 1.1 are not satisfied if  $d$  is the usual metric in  $\mathbb{R}^2$ . For example, if  $d$  is the usual metric in  $\mathbb{R}^2$ , then there are not a function  $f$  and a constant  $k \in (0, 1)$  satisfying the condition (1.1) for  $x = (1, 0)$  and  $y = (0, 0)$  in  $X$ .

*Remark 2.5.* Theorem 2.1 is a generalized version of Theorem 1.7. Indeed, if we take  $F \in \mathcal{F}(E, P, Q_a)$  defined by  $F(u) = u$  and  $\varphi(\omega) = k\omega$  ( $k \in [0, 1)$ ) in Theorem 2.1, we have Theorem 1.7.

The following example supports this remark.

**Example 2.6.** Let  $E = \mathbb{R}^2$  with the maximum norm, and  $P_2 = \{(x, y) \in \mathbb{R}^2 : x, y \geq 0\}$  a cone in  $E$ . Let  $X = \{\frac{1}{n} : n = 1, 2, \dots\} \cup \{0\}$  and let  $d : X \times X \rightarrow E$  defined by  $d(x, y) = (|x - y|, |x - y|)$ . Then  $(X, d)$  is a complete cone metric space and  $D = 1$ . Let  $F(\theta) = \theta$  and  $F(u) = \left(\|u\|^{\frac{1}{\|u\|}}, \alpha \|u\|^{\frac{1}{\|u\|}}\right)$ , for  $u \in Q_e \setminus \{\theta\}$ . Then  $F \in \mathcal{F}(\mathbb{R}^2, P_2, Q_e)$  and let  $\varphi(\omega) = \frac{1}{2}\omega$ . Suppose that  $T$  and  $S$  are mappings on  $X$  defined by

$$Tx = Sx = \begin{cases} 0 & , \quad x = 0 \\ \frac{1}{n+1} & , \quad x = \frac{1}{n}. \end{cases}$$

Now we prove that, for each  $x, y \in X$ ,

$$F(d(Tx, Sy)) \preceq \varphi(F(d(x, y))).$$

For each  $x, y \in X$  we have

$$\begin{aligned} F(d(Tx, Sy)) &= \left( \left( \|d(Tx, Sy)\| \frac{1}{\|d(Tx, Sy)\|}, \alpha \|d(Tx, Sy)\| \frac{1}{\|d(Tx, Sy)\|} \right) \right) \\ &= \left( \left( |Tx - Sy| \frac{1}{|Tx - Sy|}, \alpha |Tx - Sy| \frac{1}{|Tx - Sy|} \right) \right), \end{aligned}$$

since  $\|d(Tx, Sy)\| = \|(|Tx - Sy|, |Tx - Sy|)\| = |Tx - Sy|$ . On the other hand, for each  $x, y \in X$  we have

$$\begin{aligned} \varphi(F(d(x, y))) &= \frac{1}{2}F(d(x, y)) \\ &= \frac{1}{2} \left( \|d(x, y)\| \frac{1}{\|d(x, y)\|}, \alpha \|d(x, y)\| \frac{1}{\|d(x, y)\|} \right) \\ &= \frac{1}{2} \left( |x - y| \frac{1}{|x - y|}, \alpha |x - y| \frac{1}{|x - y|} \right). \end{aligned}$$

Now using Example 2 of [27] for each  $x, y \in X$  we have

$$|Tx - Sy| \frac{1}{|Tx - Sy|} \leq \frac{1}{2} \left( |x - y| \frac{1}{|x - y|} \right)$$

and therefore we have

$$\begin{aligned} F(d(Tx, Sy)) &= \left( |Tx - Sy| \frac{1}{|Tx - Sy|}, \alpha |Tx - Sy| \frac{1}{|Tx - Sy|} \right) \\ &\preceq \frac{1}{2} \left( |x - y| \frac{1}{|x - y|}, \alpha |x - y| \frac{1}{|x - y|} \right) \\ &= \varphi(F(d(x, y))). \end{aligned}$$

This shows that the condition (2.2) of Theorem 2.1 is satisfied. Now suppose there exists  $k \in [0, 1)$  such that, for all  $x, y \in X$ ,

$$d(Tx, Sy) \preceq kd(x, y).$$

This implies, for all  $x, y \in X$ ,

$$(|Tx - Ty|, |Tx - Ty|) \preceq k(|x - y|, |x - y|)$$

or

$$|Tx - Ty| \leq k|x - y|. \quad (2.7)$$

But we can not get the inequality (2.7). To see this, let  $x = \frac{1}{n}$  and  $y = \frac{1}{n+1}$ .

Then we have

$$|Tx - Sy| = \frac{1}{(n+1)(n+2)}, \quad |x - y| = \frac{1}{n(n+1)}$$

and so

$$\sup_{x, y \in X, x \neq y} \frac{|Tx - Sy|}{|x - y|} \geq \sup_{n \in \mathbb{N}} \frac{|T\frac{1}{n} - S\frac{1}{n+1}|}{|\frac{1}{n} - \frac{1}{n+1}|} = 1.$$

Therefore the condition (1.2) of Theorem 1.7 is not satisfied.



**Acknowledgement.** The authors thank the referees for their appreciation, valuable comments and suggestions.

## REFERENCES

- [1] M. Abbas and G. Jungck, *Common fixed point results for noncommuting mappings without continuity in cone metric spaces*, J. Math. Anal. Appl. **341** (2008), 416–420.
- [2] M. Abbas and B.E. Rhoades, *Fixed and periodic point results in cone metric spaces*, Appl. Math. Lett. **22** (2008), 511–515.
- [3] M. Abbas and B.E. Rhoades, *Common fixed point theorems for occasionally weakly compatible mappings satisfying a generalized contractive condition*, Math. Commun. **13** (2) (2008), 295–301.
- [4] A. Aliouche, *A common fixed point theorem for weakly compatible mappings in symmetric spaces satisfying a contractive condition of integral type*, J. Math. Anal. Appl. **322** (2)(2006), 796–802.
- [5] I. Altun, D. Turkoglu and B.E. Rhoades, *Fixed points of weakly compatible maps satisfying a general contractive condition of integral type*, Fixed Point Theory and Appl. **2007** (2007), Article ID 17301.
- [6] I. Altun, B. Damjanović and D. Djorić, *Fixed point and common fixed point theorems on ordered cone metric spaces*, Appl. Math. Lett. **23** (2010), 310–316.
- [7] I. Altun and G. Durmaz, *Some fixed point theorems on ordered cone metric spaces*, Rend. Circ. Mat. Palermo **58** (2009), 319–325.
- [8] M. Arshad, A. Azam and P. Vetro, *Some common fixed point results in cone metric spaces*, Fixed Point Theory Appl. **2009** (2009), Article ID 493965.
- [9] A. Azam, M. Arshad and I. Beg, *Common fixed points of two maps in cone metric spaces*, Rend. Circ. Mat. Palermo **57** (2008), 433–441.
- [10] A. Branciari, *A fixed point theorem for mappings satisfying a general contractive condition of integral type*, Int. J. Math. Math. Sci. **29** (9) (2002), 531–536.
- [11] C. Di Bari and P. Vetro,  *$\varphi$ -pairs and common fixed points in cone metric spaces*, Rend. Circ. Mat. Palermo **57** (2008), 279–285.
- [12] C. Di Bari and P. Vetro, *Weakly  $\varphi$ -pairs and common fixed points in cone metric spaces*, Rend. Circ. Mat. Palermo **58** (2009), 125–132.
- [13] R.H. Haghi and Sh. Rezapour, *Fixed points of multifunctions on regular cone metric spaces*, Expo. Math. **28** (2010), 71–77.
- [14] L.G. Huang and X. Zhang, *Cone metric spaces and fixed point theorems of contractive mappings*, J. Math. Anal. Appl. **332** (2007), no. 2, 1468–1476.
- [15] D. Ilić and V. Rakočević, *Common fixed points for maps on cone metric space*, J. Math. Anal. Appl. **341** (2008) 876–882.
- [16] G. Jungck, S. Radenović, S. Radojević and V. Rakočević, *Common fixed point theorems for weakly compatible pairs on cone metric spaces*, Fixed Point Theory Appl. **2009** (2009), Article ID 643840.
- [17] D. Klim and D. Wardowski, *Dynamic processes and fixed points of set-valued nonlinear contractions in cone metric spaces*, Nonlinear Anal. **71** (2009), 5170–5175.
- [18] J. Matkowski, *Remarks on Lipschitzian mappings and some fixed point theorems*, Banach J. Math. Anal. **1** (2) (2007), 237–244.
- [19] S. Radenović, *Common fixed points under contractive conditions in cone metric spaces*, Comput. Math. Appl. **58** (6) (2009), 1273–1278.
- [20] P. Raja and S.M. Vaezpour, *Some extensions of Banach’s contraction principle in complete cone metric spaces*, Fixed Point Theory Appl. **2008** (2008), Article ID 768294.
- [21] Sh. Rezapour and R. H. Hagli, *Fixed point of multifunctions on cone metric spaces*, Numer. Funct. Anal. Opt. **30** (7-8) (2009), 1–8.
- [22] Sh. Rezapour and R. Hamlbarani, *Some notes on the paper “Cone metric spaces and fixed point theorems of contractive mappings*, J. Math. Anal. Appl. **345** (2008) 719–724.

- [23] B.E. Rhoades, *Two fixed point theorems for mappings satisfying a general contractive condition of integral type*, Int. J. Math. Math. Sci. **2003** (63) (2003), 4007–4013.
- [24] D. Turkoglu and I. Altun, *A common fixed point theorem for weakly compatible mappings in symmetric spaces satisfying an implicit relation*, Bol. Soc. Mat. Mexicana **13** (1) (2007), 195–205.
- [25] P. Vetro, *Common fixed points in cone metric spaces*, Rend. Circ. Math. Palermo **56** (2007), 464–468.
- [26] P. Vetro, A. Azam and M. Arshad, *Fixed point results in cone metric spaces*, Int.J. Modern Math. **5** (1) (2010), 101–108.
- [27] X. Zhang, *Common fixed point theorems for new generalized contractive type mappings*, J. Math. Anal. Appl. **333** (2007), 780–786.

<sup>1,3</sup> DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE AND ARTS, KIRIKKALE UNIVERSITY, 71450 YAHSIHAN, KIRIKKALE, TURKEY.

*E-mail address:* [ishakaltun@yahoo.com](mailto:ishakaltun@yahoo.com)

*E-mail address:* [hsimsek@kku.edu.tr](mailto:hsimsek@kku.edu.tr)

<sup>2</sup> DEPARTMENT OF MATHEMATICS, LAHORE UNIVERSITY OF MANAGEMENT SCIENCES, 54792-LAHORE, PAKISTAN.

*E-mail address:* [mujahid@lums.edu.pk](mailto:mujahid@lums.edu.pk)