

NOTE ON LAPLACIAN ENERGY OF GRAPHS

G. H. FATH-TABAR, A. R. ASHRAFI, I. GUTMAN

(Presented at the 7th Meeting, held on October 26, 2007)

A b s t r a c t. Let G be an (n, m) -graph and $\mu_1, \mu_2, \dots, \mu_n$ its Laplacian eigenvalues. The Laplacian energy LE of G is defined as $\sum_{i=1}^n |\mu_i - 2m/n|$. Some new bounds for LE are presented, and some results from the paper B. Zhou, I. Gutman, *Bull. Acad. Serbe Sci. Arts (Cl. Math. Natur.)* **134** (2007) 1–11 are improved and extended.

AMS Mathematics Subject Classification (2000): 05C50

Key Words: Laplacian spectrum (of graph), Laplacian energy (of graph)

1. *Introduction*

Throughout this paper we are concerned with finite graphs. Let G be a graph of order (= number of vertices) n and size (= number of edges) m . We say that G is an (n, m) -graph. In some cases, the number of vertices of the graph G will be denoted as $|G|$.

As usual, the vertex and edge sets of G are denoted by $V(G)$ and $E(G)$, respectively. Let $V(G) = \{v_1, v_2, \dots, v_n\}$. The adjacency matrix $\mathbf{A}(G) = [a_{ij}]$ of G is a square matrix of order n whose (i, j) -entry is equal

to the number of edges between the vertices v_i and v_j . In this paper, we consider only simple graphs, i.e., graphs without multiple edges and loops. In this case, the adjacency matrix $\mathbf{A}(G)$ is a (0,1)-matrix. The spectrum of the graph G is the set of eigenvalues of $\mathbf{A}(G)$, together with their multiplicities [1]. The energy of G is defined as the sum of absolute values of the eigenvalues of G . This quantity, introduced long time ago by one of the present authors [4], has noteworthy chemical applications [6, 9] and interesting mathematical properties [5].

Let $\mathbf{D}(G) = [d_{ij}]$ be the diagonal matrix associated with the graph G , defined so that d_{ii} is the degree of the vertex v_i and $d_{ij} = 0$ if $i \neq j$. Then $\mathbf{L}(G) = \mathbf{D}(G) - \mathbf{A}(G)$ is the *Laplacian matrix* and its eigenvalues $\mu_1, \mu_2, \dots, \mu_n$ the *Laplacian eigenvalues* of the graph G , forming its *Laplacian spectrum*. For details of the theory of Laplacian spectra see [15, 16, 17].

The Laplacian polynomial $\psi(G, \lambda)$ of the graph G is the characteristic polynomial of its Laplacian matrix. Then the Laplacian eigenvalues are the zeros of $\psi(G, \lambda)$.

In what follows we assume that $\mu_1 \geq \mu_2 \geq \dots \geq \mu_n$.

It is well known that for all graphs, $\mu_n = 0$, and that the multiplicity of 0 in the Laplacian spectrum of G is equal to the number of (connected) components of G .

The Laplacian energy is a recently conceived graph invariant [10], defined as

$$LE = LE(G) = \sum_{i=1}^n \left| \mu_i - \frac{2m}{n} \right|.$$

The Laplacian energy is a proper extension of the graph-energy concept. The few, hitherto communicated, results on LE are found in the papers [2, 8, 10, 14, 20, 21].

The complement of the graph G is denoted by \overline{G} .

Suppose that G_1 and G_2 are two graphs with disjoint vertex and edge sets. Their *disjoint union* $G_1 \cup G_2$ has the vertex set $V(G_1) \cup V(G_2)$ and edge set $E(G_1) \cup E(G_2)$. Their *Cartesian product* $G_1 \times G_2$ has the vertex set $V(G_1) \times V(G_2)$ and $(a, x)(b, y)$ is an edge of $G_1 \times G_2$ if $a = b$ and $xy \in E(G_2)$, or $ab \in E(G_1)$ and $x = y$. The product $G_1 \times G_2 \times \dots \times G_k$ is defined analogously and in what follows will be denoted by $\prod_{i=1}^k G_i$.

The *join* $G_1 + G_2$ of the graphs G_1 and G_2 is obtained from $G_1 \cup G_2$, by connecting all vertices of G_1 with all vertices of G_2 . If $G = G_1 + G_2 + \dots + G_k$,

then we write $G = \sum_{i=1}^k G_i$. In the case that $G_1 \cong G_2 \cong \dots \cong G_k \cong H$, we denote G by kH .

2. Bounds for the Laplacian energy

Proposition 1. *If G is an (n, m) -graph and $\psi(G, \lambda)$ its Laplacian characteristic polynomial, then*

$$LE(G) \geq n \left| \psi \left(G, \frac{2m}{n} \right) \right|^{1/n} \quad (1)$$

with equality if and only if G is a disjoint union of the empty graph $\overline{K_p}$ and $(n-p)/2$ copies of K_2 , $0 \leq p \leq n$.

Proof. By the inequality between the geometric and arithmetic means,

$$\frac{LE(G)}{n} = \frac{1}{n} \sum_{i=1}^n \left| \mu_i - \frac{2m}{n} \right| \geq \left(\prod_{i=1}^n \left| \mu_i - \frac{2m}{n} \right| \right)^{1/n} = \left| \prod_{i=1}^n \left(\frac{2m}{n} - \mu_i \right) \right|^{1/n}$$

and (1) follows from the fact that

$$\psi(G, \lambda) = \prod_{i=1}^n (\lambda - \mu_i).$$

Equality in (1) occurs if and only if for every i, j , $1 \leq i, j \leq n$, the equality $|\mu_i - 2m/n| = |\mu_j - 2m/n|$ is obeyed. This requires that either $\mu_i = 0$ or $\mu_i = 4m/n$, which happens if and only if the degree of every vertex of G is not greater than unity. \square

Proposition 1 can be somewhat enhanced. Consider $LE(G)^2$:

$$\begin{aligned} LE(G)^2 &= \sum_{i=1}^n \sum_{j=1}^n \left| \mu_i - \frac{2m}{n} \right| \cdot \left| \mu_j - \frac{2m}{n} \right| \\ &= \sum_{i=1}^n \left(\mu_i - \frac{2m}{n} \right)^2 + \sum_{i \neq j} \left| \mu_i - \frac{2m}{n} \right| \cdot \left| \mu_j - \frac{2m}{n} \right|. \end{aligned}$$

By direct calculation we obtain

$$\sum_{i=1}^n \left(\mu_i - \frac{2m}{n} \right)^2 = Zg - 2m \left(\frac{2m}{n} - 1 \right)$$

where Zg is the sum of squares of vertex degrees (often referred to as the *Zagreb index* [7, 18, 19]). By the geometric–arithmetic mean inequality,

$$\begin{aligned} \sum_{i \neq j} \left| \mu_i - \frac{2m}{n} \right| \cdot \left| \mu_j - \frac{2m}{n} \right| &\geq n(n-1) \left[\prod_{i \neq j} \left| \mu_i - \frac{2m}{n} \right| \cdot \left| \mu_j - \frac{2m}{n} \right| \right]^{1/n(n-1)} \\ &= n(n-1) \left[\prod_{i=1}^n \left| \mu_i - \frac{2m}{n} \right|^{2(n-1)} \right]^{1/n(n-1)} = \left| \prod_{i=1}^n \left(\mu_i - \frac{2m}{n} \right) \right|^{2/n} \\ &= \left| \prod_{i=1}^n \left(\frac{2m}{n} - \mu_i \right) \right|^{2/n} = \left| \psi \left(G, \frac{2m}{n} \right) \right|^{2/n}. \end{aligned}$$

Combining these two results we get

$$LE(G)^2 \geq Zg - 2m \left(\frac{2m}{n} - 1 \right) + \left| \psi \left(G, \frac{2m}{n} \right) \right|^{2/n}$$

i.e.,

$$LE(G) \geq \sqrt{Zg - 2m \left(\frac{2m}{n} - 1 \right) + \left| \psi \left(G, \frac{2m}{n} \right) \right|^{2/n}}.$$

Proposition 2. *If G is an (n, m) -graph, and \overline{G} is its complement, then*

$$LE(G) - \frac{4m}{n} < LE(\overline{G}) \leq LE(G) + 2(n-1) - \frac{4m}{n}. \quad (2)$$

Moreover, equality on the right-hand side of (2) is attained if and only if G is the empty graph.

P r o o f. Clearly, $\mathbf{L}(G) + \mathbf{L}(\overline{G}) = n \mathbf{I}_n - \mathbf{J}_n$, where \mathbf{J}_n is the square matrix of order n whose all entries are equal to unity. Because $\mathbf{L}(G) \mathbf{J}_n = \mathbf{0} = \mathbf{J}_n \mathbf{L}(G)$, the matrix $\mathbf{L}(G)$ commutes with $\mathbf{L}(\overline{G})$. It follows that if $\mu_1(G) \geq \mu_2(G) \geq \dots \geq \mu_n(G)$ then $\mu_n(\overline{G}) = 0$ and $\mu_{n-i}(\overline{G}) = n - \mu_i(G)$, $1 \leq i \leq n-1$. Thus $\overline{m} + m = n(n-1)/2$ and so $2\overline{m}/n = n-1 - 2m/n$. Therefore,

$$\begin{aligned} LE(\overline{G}) &= \sum_{i=1}^n \left| \mu_i(\overline{G}) - \frac{2\overline{m}}{n} \right| \\ &= \sum_{i=1}^{n-1} \left| n - \mu_i(G) - \left(n - 1 - \frac{2m}{n} \right) \right| + \left| 0 - \left(n - 1 - \frac{2m}{n} \right) \right| \end{aligned}$$

$$\begin{aligned}
&= \sum_{i=1}^{n-1} \left| \mu_i(G) - \frac{2m}{n} - 1 \right| + \left| n - 1 - \frac{2m}{n} \right| \\
&\leq \sum_{i=1}^{n-1} \left(\left| \mu_i(G) - \frac{2m}{n} \right| + 1 \right) + n - 1 - \frac{2m}{n} \\
&= \sum_{i=1}^{n-1} \left| \mu_i(G) - \frac{2m}{n} \right| + 2(n-1) - \frac{2m}{n} \\
&= LE(G) - \frac{2m}{n} + 2(n-1) - \frac{2m}{n}
\end{aligned}$$

which directly implies the right-hand side inequality in (2).

In order to prove the left-hand side inequality in (2), notice that

$$\begin{aligned}
LE(\overline{G}) &= \sum_{i=1}^{n-1} \left| n - \mu_i(G) - \frac{2m}{n} \right| + \frac{2m}{n} \\
&= \sum_{i=1}^{n-1} \left| n - \mu_i(G) - n + 1 + \frac{2m}{n} \right| + n - 1 - \frac{2m}{n} \\
&= \sum_{i=1}^{n-1} \left| \mu_i(G) - \frac{2m}{n} - 1 \right| + n - 1 - \frac{2m}{n} \\
&\geq \sum_{i=1}^{n-1} \left(\left| \mu_i(G) - \frac{2m}{n} \right| - 1 \right) + n - 1 - \frac{2m}{n} \\
&= \sum_{i=1}^{n-1} \left| \mu_i(G) - \frac{2m}{n} \right| - (n-1) + n - 1 - \frac{2m}{n} \\
&= \sum_{i=1}^n \left| \mu_i(G) - \frac{2m}{n} \right| - \frac{2m}{n} - \frac{2m}{n} \\
&= LE(G) - \frac{4m}{n}.
\end{aligned}$$

In order to complete the proof, assume that G is an empty graph. Then $LE(G) = 0$ and so $LE(\overline{G}) = 2(n-1) = LE(G) + 2(n-1) - 4m/n$.

We now suppose that $LE(\overline{G}) = LE(G) + 2(n-1) - 4m/n$. Then in the inequality (2) it must be

$$\sum_{i=1}^{n-1} \left| \mu_i(G) - \frac{2m}{n} - 1 \right| + \left| n-1 - \frac{2m}{n} \right| = \sum_{i=1}^{n-1} \left(\left| \mu_i(G) - \frac{2m}{n} \right| + 1 \right) + n-1 - \frac{2m}{n}.$$

Hence,

$$\sum_{i=1}^{n-1} \left| \mu_i(G) - \frac{2m}{n} - 1 \right| = \sum_{i=1}^{n-1} \left(\left| \mu_i(G) - \frac{2m}{n} \right| + 1 \right)$$

which implies that

$$\left| \mu_i(G) - \frac{2m}{n} - 1 \right| = \left| \mu_i(G) - \frac{2m}{n} \right| + 1$$

holds for every i . Therefore $\mu_i(G) \leq 2m/n$ and so G is an empty graph. \square

The inequalities (2) can be written also as

$$\frac{4m}{n} - 2(n-1) \leq LE(G) - LE(\overline{G}) < \frac{4m}{n}$$

which should be compared with the inequalities obtained in [21]:

$$2(n-1) \leq LE(G) + LE(\overline{G}) < n\sqrt{n^2-1}.$$

By Proposition 2, one can see that

$$\left| LE(\overline{G}) - \frac{2\overline{m}}{n} \right| - \left| LE(G) - \frac{2m}{n} \right| \leq n-1.$$

Proposition 3. *Let G be a regular (n, m) -graph of degree r , $r \geq 2$, and $L(G)$ its line graph. Then $LE(G) \leq LE(L(G)) < LE(G) + 2n(r-2)$.*

Proof. By a result of Kel'mans [13], $\psi(L(G), x) = (x-2r)^{m-n} \psi(G, x)$. Therefore the Laplacian eigenvalues of $L(G)$ are $2r$ ($m-n$ times) and $\mu_i(G)$, $1, 2, \dots, n$. Clearly, $|\mu_i(G) - (2r-2)| \leq |\mu_i(G) - r| + |r-2|$ and so $LE(G) \leq LE(L(G)) \leq LE(G) + 2n(r-2)$. In order to complete the proof, we must show that $LE(L(G)) \neq LE(G) + 2n(r-2)$. Otherwise, $|\mu_i - (2r-2)| = |\mu_i - r - (r-2)| = |\mu_i - r| + |r-2|$ and so $\mu_i \leq r$, which is impossible. \square

In the case $r = 2$, Proposition 3 yields $LE(L(G)) = LE(G)$, which is trivially true since then $L(G) \cong G$.

3. Laplacian energy of graph products

In [21] it was shown that for the Cartesian product $G_1 \times G_2$ of two graphs G_1 and G_2 of the same size n ,

$$LE(G_1 \times G_2) \leq n LE(G_1) + n LE(G_2) .$$

We now obtain a generalization of this result:

Proposition 4. *Let G_1, G_2, \dots, G_k be graphs with disjoint vertex sets. Then*

$$LE\left(\prod_{i=1}^k G_i\right) \leq \left(\prod_{i=1}^k |G_i|\right) \sum_{i=1}^k \frac{LE(G_i)}{|G_i|} \quad (3)$$

with equality if and only if at most one of the graphs G_i is non-empty.

P r o o f. Suppose that G_i is an (n_i, m_i) -graph, $i = 1, 2, \dots, k$, and that $G = \prod_{i=1}^k G_i = G_1 \times G_2 \times \dots \times G_k$ is an (n, m) -graph. Then it is easy to see that $2m/n = \sum_{i=1}^k 2m_i/n_i$. On the other hand, by a result of Fiedler [3], the Laplacian eigenvalues of $\prod_{i=1}^k G_i$ are of the form $\sum_{i=1}^k \mu_{j_i}(G_i)$, $1 \leq j_i \leq n_i$. Therefore,

$$\begin{aligned} LE(G) &= \sum_{j_1, j_2, \dots, j_k} \left| \sum_{i=1}^k \mu_{j_i}(G_i) - \frac{2m}{n} \right| = \sum_{j_1, j_2, \dots, j_k} \left| \sum_{i=1}^k \mu_{j_i}(G_i) - \sum_{i=1}^k \frac{2m_i}{n_i} \right| \\ &= \sum_{j_1, j_2, \dots, j_k} \left| \sum_{i=1}^k \left(\mu_{j_i}(G_i) - \frac{2m_i}{n_i} \right) \right| \leq \sum_{j_1, j_2, \dots, j_k} \sum_{i=1}^k \left| \mu_{j_i}(G_i) - \frac{2m_i}{n_i} \right| \\ &= \sum_{i=1}^k \left(\prod_{j=1, j \neq i}^k |G_j| \right) LE(G_i) . \end{aligned}$$

This implies (3).

For the second part of the proposition, we notice that $LE = 0$ for an empty graph, and so if all G_i 's are empty, then the equality in (3) holds. On the other hand, if one of the graphs G_i is non-empty and all other graphs

are empty, then

$$LE\left(\prod_{j=1}^n G_j\right) = LE\left(G_i \times \prod_{j \neq i} G_j\right) = \frac{n}{n_i} LE(G_i) = \left(\prod_{i=1}^k n_i\right) \sum_{i=1}^k \frac{LE(G_i)}{n_i}$$

and (3) holds again.

We now assume that the equality is satisfied. If so, then

$$\left|\sum_{i=1}^k \mu_{j_i} - \frac{2m_i}{n_i}\right| = \sum_{i=1}^k \left|\mu_{j_i} - \frac{2m_i}{n_i}\right|$$

and if $\mu_{j_t} = 0$, then $\mu_{j_i} - 2m_i/n_i \leq 0$ for $i \neq t$. Therefore, $\mu_{j_i} \leq 2m_i/n_i \leq 0$ and G_i is an empty graph for $i \neq t$.

By this the proof has been completed. \square

In [21] it was shown that if G_1 and G_2 are both (n, m) -graphs, and $G_1 + G_2$ is their joint, then

$$LE(G_1 + G_2) = LE(G_1) + LE(G_2) + 2n - \frac{4m}{n}.$$

In what follows we state, without proof, the straightforward extension of this result:

Proposition 5. *If all the graphs G_1, G_2, \dots, G_k are (n, m) -graphs, then*

$$LE\left(\sum_{i=1}^k G_i\right) = \sum_{i=1}^k LE(G_i) + 2(k-1)\left(n - \frac{2m}{n}\right).$$

Corollary 5.1. $LE(kG) = kLE(G) + 2(k-1)(n - 2m/n)$.

Corollary 5.2. $LE(K_n) = 2(n-1)$ and $LE(\underbrace{K_{n, n, \dots, n}}_{k \text{ times}}) = 2n(k-1)$.

Acknowledgment. One of us (A. R. A.) was partially supported by a grant from IUT (CEAMA). This work was also partially supported by the Serbian Ministry of Science and Environmental Protection through Grant no. 144015G.

REFERENCES

- [1] D. Cvetković, M. Doob, H. Sachs, *Spectra of Graphs – Theory and Application*, Barth, Heidelberg, 1995.
- [2] N. M. M. de Abreu, C. T. M. Vinagre, A. S. Bonifácio, I. Gutman, *The Laplacian energy of some Laplacian integral graphs*, MATCH Commun. Math. Comput. Chem. **60** (2008), 447–460.
- [3] M. Fiedler, *Algebraic connectivity of graphs*, Czech. Math. J. **23** (1973), 298–305.
- [4] I. Gutman, *The energy of a graph*, Ber. Math.-Statist. Sect. Forschungsz. Graz. **103** (1978), 1–22.
- [5] I. Gutman, *The energy of a graph: old and new results*, in: A. Betten, A. Kohnert, R. Laue, A. Wassermann (Eds.), *Algebraic Combinatorics and Applications*, Springer-Verlag, Berlin, 2001, pp. 196–211.
- [6] I. Gutman, *Topology and stability of conjugated hydrocarbons. The dependence of total π -electron energy on molecular topology*, J. Serb. Chem. Soc. **70** (2005), 441–456.
- [7] I. Gutman, K. C. Das, *The first Zagreb index 30 years after*, MATCH Commun. Math. Comput. Chem. **50** (2004), 83–92.
- [8] I. Gutman, N. M. M. de Abreu, C. T. M. Vinagre, A. S. Bonifácio, S. Radenković, *Relation between energy and Laplacian energy*, MATCH Commun. Math. Comput. Chem. **59** (2008), 343–354.
- [9] I. Gutman, O. E. Polansky, *Mathematical Concepts in Organic Chemistry*, Springer-Verlag, Berlin, 1986, chapter 8.
- [10] I. Gutman, B. Zhou, *Laplacian energy of a graph*, Lin. Algebra Appl. **414** (2006), 29–37.
- [11] A. K. Kelmans, *The number of trees in a graph I*, Automat. Remote Control **26** (1965), 2118–2129.
- [12] A. K. Kelmans, *The number of trees in a graph II*, Automat. Remote Control **27** (1966), 233–241.
- [13] A. K. Kelmans, *Properties of the characteristic polynomial of a graph*, Kibernetiky na sluzhbu komunizmu **4** (1967) 27–41 (in Russian).
- [14] J. Liu, B. Liu, *A Laplacian-energy-like invariant of a graph*, MATCH Commun. Math. Comput. Chem. **59** (2008), 355–372.
- [15] R. Merris, *Laplacian matrices of graphs: A survey*, Lin. Algebra Appl. **197–198** (1994), 143–176.
- [16] R. Merris, *A survey of graph Laplacians*, Lin. Multilin. Algebra **39** (1995), 19–31.
- [17] B. Mohar, *The Laplacian spectrum of graphs*, in: Y. Alavi, G. Chartrand, O. R. Oellermann, A. J. Schwenk (Eds.), *Graph Theory, Combinatorics, and Applications*, Wiley, New York, 1991, pp. 871–898.
- [18] S. Nikolić, G. Kovačević, A. Miličević, N. Trinajstić, *The Zagreb indices 30 years after*, Croat. Chem. Acta **76** (2003), 113–124.
- [19] B. Zhou, *Zagreb indices*, MATCH Commun. Math. Comput. Chem. **52** (2004), 113–118.

- [20] B. Z h o u, I. G u t m a n, *On Laplacian energy of graphs*, MATCH Commun. Math. Comput. Chem. **57** (2007), 211–220.
- [21] B. Z h o u, I. G u t m a n, *Nordhaus–Gaddum–type relations for the energy and Laplacian energy of graphs*, Bull. Acad. Serbe Sci. Arts (Cl. Math. Natur.) **134** (2007), 1–11.

Department of Mathematics
Faculty of Science
University of Kashan
Kashan 87317–51167
I. R. Iran

Faculty of Science
University of Kragujevac
P. O. Box 60
34000 Kragujevac
Serbia