

HODGE-WITT COHOMOLOGY
AND WITT-RATIONAL SINGULARITIES

ANDRE CHATZISTAMATIOU AND KAY RÜLLING¹

Received: September 30, 2011

Communicated by Takeshi Saito

ABSTRACT. We prove the vanishing modulo torsion of the higher direct images of the sheaf of Witt vectors (and the Witt canonical sheaf) for a purely inseparable projective alteration between normal finite quotients over a perfect field. For this, we show that the relative Hodge-Witt cohomology admits an action of correspondences. As an application we define Witt-rational singularities which form a broader class than rational singularities. In particular, finite quotients have Witt-rational singularities. In addition, we prove that the torsion part of the Witt vector cohomology of a smooth, proper scheme is a birational invariant.

2010 Mathematics Subject Classification: 14J17, 14C25, 14F30

Keywords and Phrases: De Rham-Witt complex, Ekedahl duality, correspondences, singularities, Witt-vector cohomology

CONTENTS

Introduction	664
1. De Rham-Witt systems after Ekedahl	671
1.1. Witt schemes	671
1.2. De Rham-Witt systems	672
1.3. Direct image, inverse image and inverse limit	674
1.4. Global sections with support	675
1.5. Derived functors	676
1.6. Witt-dualizing systems	681
1.7. Residual complexes and traces	684
1.8. Witt residual complexes	690
1.9. The dualizing functor	694
1.10. Ekedahl's results	696

¹This work has been supported by the SFB/TR 45 “Periods, moduli spaces and arithmetic of algebraic varieties”

1.11. The trace morphism for a regular closed immersion	699
2. Pushforward and pullback for Hodge-Witt cohomology with supports	703
2.1. Relative Hodge-Witt cohomology with supports	704
2.2. Pullback	705
2.3. Pushforward	705
2.4. Compatibility of pushforward and pullback	707
3. Correspondences and Hodge-Witt cohomology	714
3.1. Exterior product	714
3.2. The cycle class of Gros	718
3.3. Hodge-Witt cohomology as weak cohomology theory with supports	720
3.4. Cycle classes	726
3.5. Correspondence action on relative Hodge-Witt cohomology	731
3.6. Vanishing results	736
3.7. De Rham-Witt systems and modules modulo torsion	739
4. Witt-rational singularities	742
4.1. The Witt canonical system	743
4.2. Topological finite quotients	747
4.3. Quasi-resolutions and relative Hodge-Witt cohomology	750
4.4. Rational and Witt-rational singularities	753
4.5. Complexes and sheaves attached to singularities of schemes	758
4.6. Isolated singularities	762
4.7. Cones and Witt-rational singularities	765
4.8. Morphisms with rationally connected generic fibre	769
5. Further applications	772
5.1. Results on torsion	772
References	778

INTRODUCTION

An important class of singularities over fields of characteristic zero are the rational singularities. For example, quotient singularities and log terminal singularities are rational singularities (see e.g. [KM98]). Over fields with positive characteristic the situation is more subtle. The definition of rational singularities requires resolution of singularities which is not yet available in all dimensions. Moreover, quotient singularities are only rational singularities under a further tameness condition, but in general fail to be rational singularities.

The purpose of this paper is to define a broader class of singularities in positive characteristic, which we call Witt-rational singularities. The main idea is that we replace the structure sheaf \mathcal{O}_X and the canonical sheaf ω_X by the Witt sheaves $W\mathcal{O}_{X,\mathbb{Q}}$ and $W\omega_{X,\mathbb{Q}}$. One important difference is that multiplication with p is invertible in $W\mathcal{O}_{X,\mathbb{Q}}$ and $W\omega_{X,\mathbb{Q}}$. Instead of resolution of singularities we can use alterations.

Witt-rational singularities have been first introduced by Blickle and Esnault [BE08]. In this paper we use a slightly different and more restrictive definition, which seems to be more accessible. Conjecturally, our definition agrees with the one of Blickle-Esnault by using a Grauert-Riemenschneider vanishing theorem for the Witt canonical sheaf $W\omega_{X,\mathbb{Q}}$. We hope to say more about this in the future.

0.1. Let k be a perfect field of positive characteristic. We denote by $W = W(k)$ the ring of Witt vectors and by $K_0 = \text{Frac}(W)$ the field of fractions. For a smooth proper k -scheme the crystalline cohomology $H_{\text{crys}}^*(X/W)$ has, by the work of Bloch and Deligne-Illusie, a natural interpretation as hypercohomology of the de Rham-Witt complex $W\Omega_X^\bullet$,

$$H_{\text{crys}}^*(X/W) \cong H^*(X, W\Omega_X^\bullet).$$

After inverting p , the slope spectral sequence degenerates which yields a decomposition

$$H^n(X/K_0) = \bigoplus_{i+j=n} H^j(X, W\Omega_X^i) \otimes_W K_0.$$

The de Rham-Witt complex is the limit of a pro-complex $(W_n\Omega_X^\bullet)_n$, and for us $W_n\mathcal{O}_X$ and $W_n\omega_X = W_n\Omega_X^{\dim X}$ will be most important. The sheaf $W_n\mathcal{O}_X$ is the sheaf of Witt vectors of length n , and defines a scheme structure W_nX on the topological space X . The structure map $\pi : X \rightarrow \text{Spec}(k)$ induces a morphism $W_n(\pi) : W_nX \rightarrow \text{Spec } W_n(k)$, but $W_n(\pi)$ is almost never flat. By the work of Ekedahl (see [Eke84]) $W_n\omega_X$ equals $W_n(\pi)^!W_n[-\dim X]$, hence $W_n\omega_X$ is a dualizing sheaf for W_nX .

The main technical problem in order to define Witt-rational singularities is to prove the independence of the chosen alteration. Our approach is to use the action of algebraic cycles in a similar way as in [CR09]. For this, we have to extend the work of Gros [Gro85] on the de Rham-Witt complex in Theorem 1 below.

For a k -scheme S we denote by \mathcal{C}_S the category whose objects are S -schemes which are smooth and quasi-projective over k . For two objects $f : X \rightarrow S$ and $g : Y \rightarrow S$ in \mathcal{C}_S , the morphisms $\text{Hom}_S(f : X \rightarrow S, g : Y \rightarrow S)$ are defined by $\varinjlim_Z \text{CH}(Z)$, where the limit is taken over all proper correspondences over S between X and Y , i.e. closed subschemes $Z \subset X \times_S Y$ such that the projection to Y is proper ($\text{CH}(Z) = \bigoplus_{i=0}^{\dim Z} \text{CH}_i(Z)$ denotes the Chow group). The composition of two morphisms is defined using Fulton's refined intersection product. The following theorem on the action of proper correspondences on relative Hodge-Witt cohomology is the main technical tool of the article.

THEOREM 1 (cf. Proposition 3.5.4). *There is a functor*

$$\hat{\mathcal{H}}(?/S) : \mathcal{C}_S \rightarrow (W\mathcal{O}_S\text{-modules}),$$

$$\hat{\mathcal{H}}(f : X \rightarrow S/S) = \bigoplus_{i,j} R^i f_* W\Omega_X^j,$$

with the following properties.

If $h : X \rightarrow Y$ is an S -morphism between two smooth k -schemes and $\Gamma_h^t \subset Y \times_S X$ denotes the transpose of its graph, then $\hat{\mathcal{H}}([\Gamma_h^t]/S)$ is the natural pull-back. If in addition h is projective then $\hat{\mathcal{H}}([\Gamma_h]/S)$ is the pushforward defined by Gros in [Gro85] using Ekedahl's duality theory [Eke84].

For a morphism $\alpha \in \text{Hom}_{\mathcal{C}_S}(X/S, Y/S)$ in \mathcal{C}_S , the map $\hat{\mathcal{H}}(\alpha/S)$ is compatible with Frobenius, Verschiebung and the differential.

0.2. We say that an integral normal k -scheme X is a *finite quotient* if there exists a finite and surjective morphism from a smooth k -scheme $Y \rightarrow X$ (e.g. $X = Y/G$ for some finite group G acting on Y .) We say that a normal integral scheme X is a *topological finite quotient* if there exists a finite, surjective and purely inseparable morphism $u : X \rightarrow X'$, where X' is a finite quotient. The morphism u is in fact a universal homeomorphism. Finally we say that a morphism $f : X \rightarrow Y$ between two integral k -schemes is a *quasi-resolution of Y* if X is a topologically finite quotient and the morphism f is surjective, projective, generically finite and purely inseparable. (In characteristic zero these conditions imply that X is a finite quotient and f is projective and birational.) By a result of de Jong (see [dJ96], [dJ97]) quasi-resolutions always exist. When working with \mathbb{Q} -coefficients the notion of quasi-resolutions suffices to define an analog of rational singularities. This follows from the following theorem.

THEOREM 2 (Theorem 4.3.3). *Let Y be a topological finite quotient and $f : X \rightarrow Y$ a quasi-resolution. Then*

$$Rf_*W\mathcal{O}_{X,\mathbb{Q}} \cong W\mathcal{O}_{Y,\mathbb{Q}}.$$

If X and Y are smooth and f is birational, this is a direct consequence of Theorem 1 and the vanishing Lemmas 3.6.1 and 3.6.2. Indeed, in $\text{CH}(X \times_Y X)$ the diagonal $\Delta_X \subset X \times_Y X$ can be written as $[\Gamma_f^t] \circ [\Gamma_f] + E$, where E is a cycle whose projections to X have at least codimension ≥ 1 . Thus E acts as zero on the $W\mathcal{O}$ part and hence $[\Gamma_f^t] \circ [\Gamma_f]$ acts as the identity on $R^i f_* W\mathcal{O}_{X,\mathbb{Q}}$, but it factors over 0 for $i > 0$; this will prove the theorem in case X and Y is smooth. Because the Frobenius is invertible when working with \mathbb{Q} -coefficients we can neglect all purely inseparable phenomena. Therefore the main point in the general case is to realize the higher direct images of $R^i f_* W\mathcal{O}_{X,\mathbb{Q}}$ (and also for Y) as certain direct factors in the relative cohomology of smooth schemes, which is possible since X and Y are topological finite quotients.

0.3. Before explaining our definition of Witt-rational singularities we need to introduce some notations. If X is a k -scheme of pure dimension d and with structure map $\pi : X \rightarrow \text{Spec } k$, then we define the Witt canonical sheaf of length n by

$$W_n \omega_X := H^{-d}(W_n(\pi)^! W_n).$$

It follows from the duality theory developed by Ekedahl in [Eke84], that these sheaves form a projective system $W_\bullet \omega_X$ with Frobenius, Verschiebung and

Cartier morphisms. Further properties are (the first two are due to Ekedahl, see [Eke84] and Proposition 4.1.4)

- (1) If X is smooth, then $W_{\bullet}\omega_X \cong W_{\bullet}\Omega_X^d$.
- (2) If X is Cohen-Macaulay, then $W_n\omega_X[d] \cong W_n(\pi)^!W_n$, in particular $W_n\omega_X$ is dualizing.
- (3) If $f : X \rightarrow Y$ is a proper morphism between k -schemes of the same pure dimension, then there is a $W_{\bullet}\mathcal{O}_Y$ -linear morphism

$$f_* : f_*W_{\bullet}\omega_X \rightarrow W_{\bullet}\omega_Y,$$

which is compatible with composition and localization.

We define $W\omega_X := \varprojlim W_{\bullet}\omega_X$.

We say that an integral k -scheme S has *Witt-rational singularities* (Definition 4.4.4) if for any quasi-resolution $f : X \rightarrow S$ the following conditions are satisfied:

- (1) $f^* : W\mathcal{O}_{S,\mathbb{Q}} \xrightarrow{\cong} f_*W\mathcal{O}_{X,\mathbb{Q}}$ is an isomorphism,
- (2) $R^i f_*W\mathcal{O}_{X,\mathbb{Q}} = 0$, for all $i \geq 1$,
- (3) $R^i f_*W\omega_{X,\mathbb{Q}} = 0$, for all $i \geq 1$.

In case only the first two properties are satisfied we say that S has *WO-rational singularities*. Condition (1) is satisfied provided that S is normal.

Our main example for varieties with Witt-rational singularities are topologically finite quotients, because the vanishing property in Theorem 2 also holds for $W\omega$.

THEOREM 3 (Corollary 4.4.7). *Topological finite quotient have Witt-rational singularities.*

A particular case are normalizations of smooth schemes X in a purely inseparable finite field extension of the function field of X . More generally, if $u : Y \rightarrow X$ is a universal homeomorphism between normal schemes then Y has Witt-rational singularities if and only if X has Witt-rational singularities (Proposition 4.4.9).

Every scheme with rational singularities has Witt-rational singularities, but varieties with Witt-rational singularities form a broader class. For example, finite quotients may fail to be Cohen-Macaulay and thus are in general not rational singularities.

A different definition of Witt-rational singularities has been introduced by Blickle and Esnault as follows. Let S be an integral k -scheme and $f : X \rightarrow S$ a generically étale alteration with X a smooth k -scheme. We say that S has *BE-Witt-rational singularities* if the natural morphism

$$W\mathcal{O}_{S,\mathbb{Q}} \rightarrow Rf_*W\mathcal{O}_{X,\mathbb{Q}}$$

admits a splitting in the derived category of sheaves of abelian groups on X . A scheme with Witt-rational singularities in our sense has BE-Witt-rational singularities (Proposition 4.4.17). We conjecture that the converse is also true. The existence of quasi-resolutions implies the following corollary.

COROLLARY 1 (Corollary 4.4.11). *Let S be a k -scheme and X and Y two integral S -schemes. Suppose that there exists a commutative diagram*

$$\begin{array}{ccc} & Z & \\ \pi_X \swarrow & & \searrow \pi_Y \\ X & & Y \\ f \searrow & & \swarrow g \\ & S & \end{array}$$

with π_X and π_Y quasi-resolutions. Suppose that X, Y have Witt-rational singularities. Then we get induced isomorphisms in $D^b(W\mathcal{O}_S)$

$$(1) \quad Rf_*W\mathcal{O}_{X,\mathbb{Q}} \cong Rg_*W\mathcal{O}_{Y,\mathbb{Q}}, \quad Rf_*W\omega_{X,\mathbb{Q}} \cong Rg_*W\omega_{Y,\mathbb{Q}}.$$

The isomorphisms are compatible with the action of the Frobenius and the Verschiebung.

If $f : X \rightarrow S$ and $g : Y \rightarrow S$ are quasi-resolutions then the isomorphisms in (1) are independent of the choice of Z (Corollary 4.5.1). In this way we obtain natural complexes

$$W\mathcal{S}_{0,S} := Rf_*W\mathcal{O}_{X,\mathbb{Q}}, \quad W\mathcal{S}_{\dim(S),S} := Rf_*W\omega_{S,\mathbb{Q}},$$

(Definition 4.5.2).

0.4. By using the work of Berthelot-Bloch-Esnault Corollary 1 yields congruences for the number of rational points over finite fields.

COROLLARY 2 (Corollary 4.4.16). *Let $S = \text{Spec } k$ be a finite field. Let X and Y be as in Corollary 1, and suppose that X, Y are proper. Then for any finite field extension k' of k we have*

$$|X(k')| \equiv |Y(k')| \pmod{|k'|}.$$

If X, Y are smooth this is a theorem due to Ekedahl [Eke83].

For a normal integral scheme S with an isolated singularity $s \in S$ we can give a criterion for the $W\mathcal{O}$ -rationality of S , provided that a resolution of singularities $f : X \rightarrow S$ exists such that $f : f^{-1}(S \setminus \{s\}) \rightarrow S \setminus \{s\}$ is an isomorphism; we denote by $E := f^{-1}(s)$ the fibre over s . Then S has $W\mathcal{O}$ -rational singularities if and only if

$$(2) \quad H^i(E, W\mathcal{O}_{E,\mathbb{Q}}) = 0 \quad \text{for all } i > 0,$$

(Corollary 4.6.4). This implies that a normal surface has $W\mathcal{O}$ -rational singularities if and only if the exceptional divisor is a tree of smooth rational curves. For cones C of smooth projective schemes X , we obtain that C has $W\mathcal{O}$ -rational singularities if and only if $H^i(X, W\mathcal{O}_{X,\mathbb{Q}}) = 0$ for $i > 0$. We can show that C has Witt-rational singularities provided that Kodaira vanishing holds for X (Section 4.7). We expect that this assumption can be dropped; in general, a Grauert-Riemenschneider type vanishing theorem for $W\omega$ should imply that $W\mathcal{O}$ -rationality is equivalent to Witt-rationality.

Over a finite field k we use a weight argument to refine the criterion (2) if E is a strict normal crossing divisor. Let E_i be the irreducible components of E , via the restriction maps we obtain for all $t \geq 0$ a complex $C_t(E)$:

$$\underbrace{\bigoplus_{i_0} H^t(E_{i_0}, W\mathcal{O}_{E_{i_0}, \mathbb{Q}})}_{\text{deg}=0} \rightarrow \bigoplus_{i_0 < i_1} H^t(E_{i_0} \cap E_{i_1}, W\mathcal{O}_{E_{i_0} \cap E_{i_1}, \mathbb{Q}}) \rightarrow \dots$$

THEOREM 4 (Theorem 4.6.7). *Let k be a finite field. In the above situation, S has $W\mathcal{O}$ -rational singularities if and only if*

$$H^i(C_t(E)) = 0 \quad \text{for all } (i, t) \neq (0, 0).$$

Theorem 4 is inspired by the results of Kerz-Saito [Sai10, Theorem 8.2] on the weight homology of the exceptional divisor.

For morphisms with generically smooth fibre with trivial Chow group of zero cycles we can show the following vanishing theorem.

THEOREM 5 (Theorem 4.8.1). *Let X be an integral scheme with Witt-rational singularities. Let $f : X \rightarrow Y$ be a projective morphism to an integral, normal and quasi-projective scheme Y . We denote by η the generic point of Y , and X_η denotes the generic fibre of f . Suppose that X_η is smooth and for every field extension $L \supset k(\eta)$ the degree map*

$$\text{CH}_0(X_\eta \times_{k(\eta)} L) \otimes_{\mathbb{Z}} \mathbb{Q} \rightarrow \mathbb{Q}$$

is an isomorphism. Then, for all $i > 0$,

$$R^i f_* W\mathcal{O}_{X, \mathbb{Q}} \cong H^i(\mathcal{WS}_{0, Y}), \quad R^i f_* W\omega_{X, \mathbb{Q}} \cong H^i(\mathcal{WS}_{\dim(Y), Y}).$$

In particular, if Y has Witt-rational singularities then

$$R^i f_* W\mathcal{O}_{X, \mathbb{Q}} = 0, \quad R^i f_* W\omega_{X, \mathbb{Q}} = 0, \quad \text{for all } i > 0.$$

0.5. For smooth schemes we can show the following result which takes the torsion into account.

THEOREM 6 (Theorem 5.1.10). *Let S be a k -scheme. Let $f : X \rightarrow S$ and $g : Y \rightarrow S$ be two S -schemes which are integral and smooth over k and have dimension N . Assume X and Y are properly birational over S , i.e. there exists a closed integral subscheme $Z \subset X \times_S Y$, such that the projections $Z \rightarrow X$ and $Z \rightarrow Y$ are proper and birational. There are isomorphisms in $D^b(S, W(k))$:*

$$Rf_* W\mathcal{O}_X \cong Rg_* W\mathcal{O}_Y, \quad Rf_* W\Omega_X^N \cong Rg_* W\Omega_Y^N.$$

Taking cohomology we obtain isomorphisms of $W\mathcal{O}_S$ -modules which are compatible with Frobenius and Verschiebung:

$$R^i f_* W\mathcal{O}_X \cong R^i g_* W\mathcal{O}_Y, \quad R^i f_* W\Omega_X^N \cong R^i g_* W\Omega_Y^N, \quad \text{for all } i \geq 0.$$

If X and Y are tame finite quotients and there exists a proper and birational morphism $h : X \rightarrow Y$ then a similar statement holds (see Theorem 5.1.13).

If X and Y are two smooth and proper k -schemes, which are birational and of pure dimension N . Then we obtain isomorphisms of $W(k)[F, V]$ -modules

$$H^i(X, W\mathcal{O}_X) \cong H^i(Y, W\mathcal{O}_Y), \quad H^i(X, W\Omega_X^N) \cong H^i(Y, W\Omega_Y^N), \quad \text{for all } i \geq 0.$$

Modulo torsion the statement for $W\mathcal{O}$ is a theorem due to Ekedahl (see [Eke83]).

0.6. We give a brief overview of the content of each section. In Section 1 we introduce the category dRW_X of de Rham-Witt systems on a k -scheme X . In the language of Ekedahl [Eke84] an object in dRW_X is both, a direct and an inverse de Rham-Witt system at the same time. Furthermore, we introduce the derived pushforward, derived cohomology with supports and $R\varprojlim$ on $D^b(\mathrm{dRW}_X)$. We recall the definition of Witt-dualizing systems from [Eke84] in 1.6, and some facts about residual complexes in 1.7. In particular, we observe that if $f : X \rightarrow Y$ is a morphism between k -schemes, which is proper along a family of supports Φ on X , then for any residual complex K on Y the trace morphism $f_* f^\Delta K \rightarrow K$, which always exists as a map of graded sheaves, induces a morphism of complexes $f_* \Gamma_\Phi f^\Delta K \rightarrow K$. In 1.8 we show that for any $\pi : X \rightarrow \mathrm{Spec} k$ the residual complexes $W_n \pi^\Delta W_n(k)$ form a projective system K_X , which is term-wise a Witt-dualizing system. In 1.9 we define the functor $D_X = \mathcal{H}om(-, K_X)$ on $D(\mathrm{dRW}_{X, \mathrm{qc}})^\circ$. (It is only defined on complexes of quasi-coherent de Rham-Witt systems.) In 1.10 we recall the results of Ekedahl in the smooth case relating K_X to $W_\bullet \Omega_X^{\dim X}$, and in 1.11 we calculate the trace morphism for a regular closed immersion. A similar description is given in [Gro85], but it refers to work in progress by Ekedahl, which we could not find in the literature, therefore we give another argument.

In Section 2 we introduce relative Hodge-Witt cohomology with supports on smooth and quasi-projective k -schemes, which are defined over some base scheme S . We define a pullback for arbitrary morphisms and using the trace map from Section 1 also a pushforward for morphisms which are proper along a family of supports. Then in 2.4 we give an explicit description of the pushforward in the case of a regular closed immersion and also for the projection $\mathbb{P}_X^n \rightarrow X$, where X is a smooth scheme X . From this description we deduce the expected compatibility between pushforward and pullback with respect to maps in a certain cartesian diagram.

In Section 3 we collect and prove the remaining facts, which we need to show that $(X, \Phi) \mapsto \bigoplus_{i,j} H_\Phi^i(X, W\Omega_X^j)$ is a weak cohomology theory with supports in the sense of [CR09]. In particular, we need the cycle class constructed by Gros in [Gro85]. From this we deduce Theorem 1 above. In 3.6 we prove the two vanishing Lemmas, which give a criterion for certain correspondences to act as zero on certain parts of the Hodge-Witt cohomology. In 3.7 we introduce the notation $\mathrm{dRW}_{X, \mathbb{Q}}$, which is the \mathbb{Q} -linearization of dRW_X . In general, for $M \in \mathrm{dRW}_X$ the notation $M_{\mathbb{Q}}$ means the image of M in $\mathrm{dRW}_{X, \mathbb{Q}}$ (which is not the same as $M \otimes_{\mathbb{Z}} \mathbb{Q}$).

In Section 4 we introduce the Witt canonical system $W_\bullet\omega_X$ for a pure-dimensional k -scheme X and prove some of its properties. Moreover we show in 4.2 that the cohomology of $W\mathcal{O}$ and $W\omega$ for a topological finite quotient is a direct summand of the Hodge-Witt cohomology of a certain smooth scheme. Then we prove Theorem 2 and define Witt rational singularities. It follows some elaboration on this notion, in particular the Theorems 3, 4, 5.

Finally in Section 5 we prove some results on torsion, as in Theorem 6. In order to do this, we show that a correspondence actually gives rise to a morphism in the derived category of modules over the Cartier-Dieudonné-Raynaud ring and then use Ekedahl’s Nakayama Lemma to deduce the statement from [CR09]. We advise the reader who is mostly interested in the geometric application to start for a first time reading with Section 1.1 and 1.2 to get some basic notations and then jump directly to Section 4.

0.7. NOTATION AND GENERAL CONVENTIONS. We are working over a perfect ground field k of characteristic $p > 0$. We denote by $W_n = W_n(k)$ the ring of Witt vectors of length n over k and by $W = W(k)$ the ring of infinite Witt vectors. By a k -scheme we always mean a scheme X , which is separated and of finite type over k . If X and Y are k -schemes, then a morphism $X \rightarrow Y$ is always assumed to be a k -morphism.

1. DE RHAM-WITT SYSTEMS AFTER EKEDAHL

1.1. WITT SCHEMES. For the following facts see e.g. [Ill79, 0.1.5], [LZ04, Appendix A]. Let X be a k -scheme. For $n \geq 1$, we denote

$$W_n X = (|X|, W_n \mathcal{O}_X) = \text{Spec } W_n \mathcal{O}_X,$$

where $W_n \mathcal{O}_X$ is the sheaf of rings of Witt vectors of length n . This construction yields a functor from the category of k -schemes to the category of separated, finite type W_n -schemes. If $f : X \rightarrow Y$ is a separated (resp. finite type, proper or étale) morphism of k -schemes, then $W_n f : W_n X \rightarrow W_n Y$ is a separated (resp. finite type, proper or étale) morphism of W_n -schemes. If f is an open (resp. closed) immersion, so is $W_n f$. We denote by $i_n : W_{n-1} X \hookrightarrow W_n X$ (or sometimes by $i_{n,X}$) the nilimmersion induced by the restriction $W_n \mathcal{O}_X \rightarrow W_{n-1} \mathcal{O}_X$. We will write $\pi : W_n \mathcal{O}_X \rightarrow i_{n*} W_{n-1} \mathcal{O}_X$ instead of i_n^* . The absolute Frobenius on X is denoted by $F_X : X \rightarrow X$. The morphism $W_n(F_X) : W_n X \rightarrow W_n X$ is finite for all n . With this notation the Frobenius and Verschiebung morphisms on the Witt vectors become morphisms of $W_n \mathcal{O}_X$ -modules

$$F = W_n(F_X)^* \circ \pi : W_n \mathcal{O}_X \rightarrow (W_n(F_X)i_n)_* W_{n-1} \mathcal{O}_X,$$

$$V : (W_n(F_X)i_n)_* W_{n-1} \mathcal{O}_X \rightarrow W_n \mathcal{O}_X.$$

Further “lift and multiply by p ” induces a morphism of $W_n \mathcal{O}_X$ -modules

$$\underline{p} : i_{n*} W_{n-1} \mathcal{O}_X \rightarrow W_n \mathcal{O}_X.$$

If $f : X \rightarrow Y$ is a morphism of k -schemes, then we have $W_n(f)i_{n,X} = i_{n,Y}W_{n-1}(f)$ and $W_n(f)W_n(F_X) = W_n(F_Y)W_n(f)$. If $f : X \rightarrow Y$ is étale,

then the following diagrams are *cartesian*:

$$(1.1.1) \quad \begin{array}{ccc} W_{n-1}X & \xrightarrow{i_n} & W_nX \\ W_{n-1}f \downarrow & & \downarrow W_n f \\ W_{n-1}Y & \xrightarrow{i_n} & W_nY, \end{array} \quad \begin{array}{ccc} W_nX & \xrightarrow{W_n(F_X)} & W_nX \\ W_n f \downarrow & & \downarrow W_n f \\ W_nY & \xrightarrow{W_n(F_Y)} & W_nY. \end{array}$$

1.2. DE RHAM-WITT SYSTEMS.

DEFINITION 1.2.1. For an integer $n \geq 1$ we denote by \mathcal{C}_n the category of \mathbb{Z} -graded $W_n\mathcal{O}_X$ -modules on X . We define

$$\mathcal{C}_{\mathbb{N}} := \prod_{n \in \mathbb{Z}, n \geq 1} \mathcal{C}_n.$$

For an object $M \in \mathcal{C}_{\mathbb{N}}$ and $n \geq 1$ we denote by M_n the n -th component. An object M in $\mathcal{C}_{\mathbb{N}}$ is (quasi-)coherent, if all M_n are (quasi-)coherent $W_n(\mathcal{O}_X)$ -modules. We denote by $\mathcal{C}_{\mathbb{N},qc}$ (resp. $\mathcal{C}_{\mathbb{N},c}$) the full subcategory of (quasi-)coherent objects of $\mathcal{C}_{\mathbb{N}}$. There are two natural endo-functors:

$$\begin{aligned} i_* : \mathcal{C}_{\mathbb{N}} &\rightarrow \mathcal{C}_{\mathbb{N}} \\ (i_*M)_n &:= \begin{cases} i_{n*}M_{n-1} & \text{if } n > 1, \\ 0 & \text{if } n = 1, \end{cases} \\ \sigma : \mathcal{C}_{\mathbb{N}} &\rightarrow \mathcal{C}_{\mathbb{N}} \\ (\sigma_*M)_n &:= W_n(F_X)_*M_n \end{aligned}$$

The two functors commute

$$(1.2.1) \quad \sigma_*i_* = i_*\sigma_*,$$

since $W_n(F_X)_*i_{n*} = i_{n*}W_{n-1}(F_X)_*$.

We will also need the following functor:

$$\begin{aligned} \Sigma_* : \mathcal{C}_{\mathbb{N}} &\rightarrow \mathcal{C}_{\mathbb{N}} \\ (\Sigma_*M)_n &:= W_n(F_X)_*^n M_n. \end{aligned}$$

We have the equalities

$$(1.2.2) \quad \sigma_*i_*\Sigma_* = \Sigma_*i_*, \quad \sigma_*\Sigma_* = \Sigma_*\sigma_*.$$

Furthermore, since the components of $M \in \mathcal{C}_{\mathbb{N}}$ are \mathbb{Z} -graded we can define for all $i \in \mathbb{Z}$ the shift functor

$$(1.2.3) \quad M(i)_n := M_n(i).$$

The shift functor commutes in an obvious way with i_*, σ_*, Σ_* .

DEFINITION 1.2.2. A *graded Witt system* $(M, F, V, \pi, \underline{p})$ on X is an object M in $\mathcal{C}_{\mathbb{N}}$ equipped with morphisms in $\mathcal{C}_{\mathbb{N}}$:

$$F : M \rightarrow \sigma_*i_*M, \quad V : \sigma_*i_*M \rightarrow M, \quad \pi : M \rightarrow i_*M, \quad \underline{p} : i_*M \rightarrow M,$$

such that

- (a) $V \circ F$ is multiplication with p ,

- (b) $F \circ V$ is multiplication with p ,
- (c) $\sigma_* i_*(\pi) \circ F = i_*(F) \circ \pi$,
- (d) $\pi \circ V = i_*(V \circ \sigma_*(\pi))$,
- (e) $i_*(\sigma_*(\underline{p}) \circ F) = F \circ \underline{p}$,
- (f) $V \circ \sigma_* i_*(\underline{p}) = \underline{p} \circ i_*(V)$,
- (g) $i_*(\underline{p} \circ \pi) = \pi \circ \underline{p}$.

Graded Witt systems form in the obvious way a category which we denote by W_X . It is straightforward to check that W_X is abelian.

We have an obvious forgetful functor $W_X \rightarrow \mathcal{C}_{\mathbb{N}}$. We say that $(M, F, V, \pi, \underline{p})$ is *(quasi-)coherent* if M_n is (quasi-)coherent for every n .

Remark 1.2.3. One should memorise (c) as “ $\pi \circ F = F \circ \pi$ ”, (d) as “ $\pi \circ V = V \circ \pi$ ”, (e) as “ $\underline{p} \circ F = F \circ \underline{p}$ ”, (f) as “ $V \circ \underline{p} = \underline{p} \circ V$ ”, and (g) as “ $\underline{p} \circ \pi = \pi \circ \underline{p}$ ”.

DEFINITION 1.2.4. A *de Rham-Witt system* (M, d) is a graded Witt system M together with a morphism in W_X :

$$d : \Sigma_* M \rightarrow \Sigma_* M(1),$$

such that the following conditions are satisfied:

- (a) $\Sigma_* F(1) \circ d \circ \Sigma_* V = \sigma_*^2 i_* d$ (we used 1.2.2),
- (b) $\Sigma_* \pi(1) \circ d = \sigma_* i_* d \circ \Sigma_*(\pi)$ (we used 1.2.2),
- (c) $d \circ \Sigma_*(\underline{p}) = \Sigma(\underline{p}) \circ \sigma_* i_* d$ (again, we used 1.2.2).
- (d) $d(1) \circ d = 0$.

De Rham Witt systems form in the obvious way a category which we denote by dRW_X . We say that a de Rham-Witt system is *(quasi-)coherent* if the underlying graded Witt system is. We denote the category of (quasi-)coherent de Rham-Witt systems by $dRW_{X, qc}$ (resp. $dRW_{X, c}$). It is straightforward to check that dRW_X , $dRW_{X, qc}$ and $dRW_{X, c}$ are abelian. We denote by $D^+(dRW_X)$, $D^+(dRW_{X, qc})$ and $D^+(dRW_{X, c})$ the corresponding derived categories of bounded below complexes.

Remark 1.2.5. One should memorise (a) as “ $F \circ d \circ V = d$ ”, (b) as “ $\pi \circ d = d \circ \pi$ ”, and (c) as “ $d \circ \underline{p} = \underline{p} \circ d$ ”.

DEFINITION 1.2.6. A *de Rham-Witt module* (M, F, V, d) is a graded $W\mathcal{O}_X$ -module M together with morphisms of $W\mathcal{O}_X$ -modules

$$F : M \rightarrow W(F_X)_* M, \quad V : W(F_X)_* M \rightarrow M$$

and a morphism of $W(k)$ -modules

$$d : M \rightarrow M(1)$$

such that

- (a) $F \circ V$ is multiplication with p ,
- (b) $V \circ F$ is multiplication with p ,
- (c) $F \circ d \circ V = d$,
- (d) $d(1) \circ d = 0$.

De Rham-Witt modules form in the obvious way a category which we denote by $\widehat{\mathrm{dRW}}_X$. It is straightforward to check that $\widehat{\mathrm{dRW}}_X$ is abelian. We denote by $D^+(\widehat{\mathrm{dRW}}_X)$ the derived category of bounded below complexes of de Rham-Witt modules.

EXAMPLE 1.2.7. Let X be a k -scheme

- (1) The sheaves of Witt vectors of finite length on X define a coherent graded Witt system

$$W_\bullet \mathcal{O}_X = (\{W_n \mathcal{O}_X\}_{n \geq 1}, \pi, F, V, \underline{p}),$$

which is concentrated in degree 0. If $X = \mathrm{Spec} k$, we simply write W_\bullet instead of $W_\bullet k$.

- (2) The de Rham Witt complex of Bloch-Deligne-Illusie $W_\bullet \Omega_X$ is a coherent de Rham-Witt system (see [Ill79]) and $W \Omega_X = \varprojlim_n W_n \Omega_X$ is a de Rham-Witt module.
- (3) Let M be a de Rham-Witt system on X and $i \in \mathbb{Z}$. Then we define

$$M(i) := (\{M_n(i)\}_{n \geq 1}, \pi_M, F_M, V_M, (-1)^i d_M, \underline{p}_M) \in \mathrm{dRW}_X.$$

1.3. DIRECT IMAGE, INVERSE IMAGE AND INVERSE LIMIT.

1.3.1. Let $f : X \rightarrow Y$ be a morphism between k -schemes. We get an induced functor

$$f_* : \mathcal{C}_{\mathbb{N}, X} \rightarrow \mathcal{C}_{\mathbb{N}, Y}, \quad (M_n) \mapsto (W_n(f)_* M_n)$$

which commutes in the obvious way with i_*, σ, Σ_* . We thus obtain a functor

$$f_* : \mathrm{dRW}_X \rightarrow \mathrm{dRW}_Y.$$

1.3.2. Let $f : X \rightarrow Y$ be an *étale* morphism between k -schemes. We get an induced functor

$$f^* : \mathcal{C}_{\mathbb{N}, Y} \rightarrow \mathcal{C}_{\mathbb{N}, X}, \quad (M_n) \mapsto (W_n(f)^* M_n)$$

which by (1.1.1) commutes with i_*, σ, Σ_* . We thus obtain a functor

$$f^* : \mathrm{dRW}_Y \rightarrow \mathrm{dRW}_X.$$

1.3.3. Let $(M, F, V, \pi, \underline{p}, d)$ be a de Rham Witt system. Then (M, π) forms naturally a projective system of $W \mathcal{O}_X$ -modules, F and V induce morphisms of projective systems of $W \mathcal{O}_X$ -modules $F : (M, \pi) \rightarrow (W(F_X)_* M, W(F_X)_* \pi)$, $V : (W(F_X)_* M, W(F_X)_* \pi) \rightarrow (M, \pi)$ and induces a morphism of projective systems of $W(k)$ -modules $d : (M, \pi) \rightarrow (M(1), \pi(1))$. We thus obtain a functor

$$\varprojlim : \mathrm{dRW}_X \rightarrow \widehat{\mathrm{dRW}}_X.$$

1.4. GLOBAL SECTIONS WITH SUPPORT.

DEFINITION 1.4.1. A family of supports Φ on X is a non-empty set of closed subsets of X such that the following holds:

- (i) The union of two elements in Φ is contained in Φ .
- (ii) Every closed subset of an element in Φ is contained in Φ .

Let A be any set of closed subsets of X . The smallest family of supports Φ_A which contains A is given by

$$(1.4.1) \quad \Phi_A := \left\{ \bigcup_{i=1}^n Z'_i; Z'_i \underset{\text{closed}}{\subset} Z_i \in A \right\}.$$

For a closed subset $Z \subset X$ we write Φ_Z for $\Phi_{\{Z\}}$.

NOTATION 1.4.2. Let $f : X \rightarrow Y$ be a morphism of schemes and Φ resp. Ψ a family of supports of X resp. Y .

- (1) We denote by $f^{-1}(\Psi)$ the smallest family of supports on X which contains $\{f^{-1}(Z); Z \in \Psi\}$.
- (2) We say that $f \mid \Phi$ is proper if $f \mid Z$ is proper for every $Z \in \Phi$. If $f \mid \Phi$ is proper then $f(\Phi)$ is a family of supports on Y .
- (3) If Φ_1, Φ_2 are two families of supports then $\Phi_1 \cap \Phi_2$ is a family of supports.
- (4) If Φ resp. Ψ is a family of supports of X resp. Y then we denote by $\Phi \times \Psi$ the smallest family of supports on $X \times_k Y$ which contains $\{Z_1 \times Z_2; Z_1 \in \Phi, Z_2 \in \Psi\}$.

1.4.3. Let Φ be a family of supports on X . We consider the sections-with-support-in- Φ functor (see e.g. [Har66, IV, §1])

$$\underline{\Gamma}_\Phi : \mathcal{C}_{\mathbb{N}, X} \rightarrow \mathcal{C}_{\mathbb{N}, X}, \quad (M_n) \mapsto (\underline{\Gamma}_\Phi(M_n)).$$

Since $\underline{\Gamma}_\Phi$ commutes in the obvious way with i_*, σ, Σ_* we obtain

$$\underline{\Gamma}_\Phi : \text{dRW}_X \rightarrow \text{dRW}_X.$$

For a closed subset $Z \subset X$ we also write $\underline{\Gamma}_Z$ instead of $\underline{\Gamma}_{\Phi_Z}$.

If $f : X \rightarrow Y$ is a morphism and Ψ a family of supports on Y , then

$$(1.4.2) \quad \underline{\Gamma}_\Psi f_* = f_* \underline{\Gamma}_{f^{-1}(\Psi)}.$$

If $f : X \rightarrow Y$ is a morphism and Φ is a family of supports on X , then we define

$$(1.4.3) \quad f_\Phi := f_* \circ \underline{\Gamma}_\Phi : \text{dRW}_X \rightarrow \text{dRW}_Y,$$

$$(1.4.4) \quad \hat{f}_\Phi := \varprojlim \circ f_\Phi : \text{dRW}_X \rightarrow \widehat{\text{dRW}}_Y.$$

Notice that if $\Phi = \Phi_Z$, with Z a closed subset of X , then

$$(1.4.5) \quad \hat{f}_{\Phi_Z} = f_* \circ \Gamma_{\Phi_Z} \circ \varprojlim.$$

This relation does not hold for arbitrary families of support on X .

1.5. DERIVED FUNCTORS.

LEMMA 1.5.1. *Let (X, \mathcal{O}_X) be a ringed space and $E = (E_n)$ a projective system of \mathcal{O}_X -modules (indexed by integers $n \geq 1$). Let \mathcal{B} be a basis of the topology of X . We consider the following two conditions:*

- a) *For all $U \in \mathcal{B}$, $H^i(U, E_n) = 0$ for all $i, n \geq 1$.*
- b) *For all $U \in \mathcal{B}$, the projective system $(H^0(U, E_n))_{n \geq 1}$ satisfies the Mittag-Leffler condition.*

Then

- (1) *If E satisfies condition a), then $R^i \varprojlim_n E_n = 0$, for all $i \geq 2$.*
- (2) *If E satisfies the conditions a) and b), then $R^i \varprojlim_n E_n = 0$, for all $i \geq 1$, i.e. E is \varprojlim -acyclic.*

Proof. It is a basic fact that there are sufficiently many injective \mathcal{O}_X -modules. Notice that a projective system of \mathcal{O}_X -modules $I = (I_n)$ is injective if and only if each I_n is an injective \mathcal{O}_X -module and the transition maps $I_{n+1} \rightarrow I_n$ are split surjective. (The “if” direction is easy, as well as I injective implies each I_n is injective. If I is injective, let J be the projective system with $J_n = I_1 \oplus \dots \oplus I_n$ and projections as transition maps. We have an obvious inclusion of projective systems $I \hookrightarrow J$, hence a surjection $\text{Hom}(J, I) \rightarrow \text{Hom}(I, I)$. Now a lift of the identity on I together with the split surjectivity of the transition maps of J gives the splitting of the transition maps of I .)

Now let $E \rightarrow I^\bullet$ be an injective resolution (which always exist). The transition maps of the projective system (of abelian groups) $(\Gamma(U, I_n^q))_n$ are surjective (since split) for all $q \geq 0$ and all open subsets $U \subset X$. Hence they satisfy the Mittag-Leffler condition and are \varprojlim -acyclic.

On the other hand, $\varprojlim_n I_n^q$ is an injective \mathcal{O}_X -module for every q . Indeed, since the transition maps $I_{n+1}^q \rightarrow I_n^q$ are surjective and split, we may write $I_n^q \cong \bigoplus_{i=1}^n I'_i$ for $I'_i = \ker(I_i^q \rightarrow I_{i-1}^q)$. The \mathcal{O}_X -modules I'_i are injective for all n , and the transition maps

$$\bigoplus_{i=1}^{n+1} I'_i \cong I_{n+1}^q \rightarrow I_n^q \cong \bigoplus_{i=1}^n I'_i$$

are the obvious projections. Thus $\varprojlim_n I_n^q = \prod_{i \geq 1} I'_i$ is injective.

By using $\varprojlim \circ \Gamma_U = \Gamma_U \circ \varprojlim$ we obtain a spectral sequence

$$R^i \varprojlim H^j(U, E_n) \implies H^{i+j}(U, R \varprojlim E_n),$$

where $R \varprojlim E_n = \varprojlim_n I_n^\bullet$. If $U \in \mathcal{B}$, condition a) implies $R^i \varprojlim H^0(U, E_n) = H^i(U, R \varprojlim E_n) = H^i(\varprojlim I^\bullet(U))$. We know that $R^i \varprojlim H^0(U, E_n)$ is zero for all $i \geq 2$ (see e.g. [Wei, Cor. 3.5.4]) and - in case condition b) is satisfied - also for all $i \geq 1$. Now the assertion follows from

$$\varprojlim_{U \in \mathcal{B}, U \ni x} H^i(U, R \varprojlim E_n) = \varprojlim_{U \in \mathcal{B}, U \ni x} H^i(\varprojlim I^\bullet(U)) = (R^i \varprojlim E_n)_x,$$

for all $x \in X$. □

LEMMA 1.5.2. *Let A be a sheaf of abelian groups on a noetherian topological space X . If A is flasque, so is $\underline{\Gamma}_\Phi(A)$ for all families of supports Φ on X .*

Proof. Let Y and Z be two closed subsets of X . Since $\underline{\Gamma}_Z(I)$ is injective if I is ([SGA2, Exp. I, Cor. 1.4]), there exists a spectral sequence $H_Y^i(X, \mathcal{H}_Z^j(A)) \implies H_{Y \cap Z}^{i+j}(X, A)$. Now assume A is flasque, then $\mathcal{H}_Z^j(A) = 0$ for $j \neq 0$. In particular $H_Y^1(X, \underline{\Gamma}_Z(A)) = H_{Y \cap Z}^1(X, A) = 0$. Thus $\underline{\Gamma}_Z(A)$ is flasque. The space X is noetherian and therefore $\underline{\Gamma}_\Phi(A) = \varinjlim_{Z \in \Phi} \underline{\Gamma}_Z(A)$ is also flasque. \square

DEFINITION 1.5.3. We say that a de Rham-Witt system M on a k -scheme X is *flasque*, if for all n

$$0 \rightarrow K_n \rightarrow M_n \xrightarrow{\pi} M_{n-1} \rightarrow 0$$

is an exact sequence of flasque abelian sheaves on X , where $K_n = \text{Ker}(\pi : M_n \rightarrow M_{n-1})$.

LEMMA 1.5.4. *Let X be a k -scheme.*

- (1) *Let $0 \rightarrow M' \rightarrow M \rightarrow M'' \rightarrow 0$ be a short exact sequence of de Rham-Witt systems on X and assume that M' is flasque. Then M is flasque iff M'' is.*
- (2) *Let Φ be a family of supports on X . Then $\underline{\Gamma}_\Phi$ restricts to an exact endo-functor on the full subcategory of flasque de Rham-Witt systems.*
- (3) *Let $f : X \rightarrow Y$ be a morphism. Then f_* restricts to an exact functor between the full subcategories of flasque de Rham-Witt systems on X and Y .*
- (4) *The functor $\varprojlim : \text{dRW}_X \rightarrow \widehat{\text{dRW}}_X$ restricts to an exact functor from the full subcategory of flasque de Rham-Witt systems to the full subcategory of flasque de Rham-Witt modules (i.e. de Rham-Witt modules, which are flasque as abelian sheaves on X).*

Proof. The proof of (1) is straightforward. (2) follows from Lemma 1.5.2. (3) is clear. Finally (4). It follows directly from the definition, that the transition maps on the sections over any open $U \subset X$ of a flasque de Rham-Witt systems are surjective. The exactness of \varprojlim on the category of flasque de Rham-Witt systems, thus follows from Lemma 1.5.1, (2). Now let M be a flasque de Rham-Witt system. It remains to show that $\varprojlim M$ is flasque again. For this let $U \subset X$ be open and define $L_n = \text{Ker}(\Gamma(\widehat{X}, M_n) \rightarrow \Gamma(U, M_n))$. Thus we have an exact sequence

$$(1.5.1) \quad \varprojlim \Gamma(X, M) \rightarrow \varprojlim \Gamma(U, M) \rightarrow R^1 \varprojlim_n L_n.$$

Consider the following diagram:

$$\begin{array}{ccccccc}
 & & 0 & & 0 & & \\
 & & \downarrow & & \downarrow & & \\
 & & \Gamma(X, K_n) & \longrightarrow & \Gamma(U, K_n) & \longrightarrow & 0 \\
 & & \downarrow & & \downarrow & & \\
 0 & \longrightarrow & L_n & \longrightarrow & \Gamma(X, M_n) & \longrightarrow & \Gamma(U, M_n) \longrightarrow 0 \\
 & & \downarrow a & & \downarrow & & \downarrow \\
 0 & \longrightarrow & L_{n-1} & \longrightarrow & \Gamma(X, M_{n-1}) & \longrightarrow & \Gamma(U, M_{n-1}) \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \\
 & & 0 & & 0 & &
 \end{array}$$

All rows and columns are exact, since M is flasque. Now it follows from an easy diagram chase that a is surjective. Therefore $R^1 \varprojlim_n L_n = 0$ and the flasqueness of $\varprojlim M$ follows from (1.5.1). \square

LEMMA 1.5.5. *The categories dRW_X and $\widehat{\text{dRW}}_X$ have enough flasque objects, i.e. any M in dRW_X (or in $\widehat{\text{dRW}}_X$) admits an injection into a flasque object.*

Proof. For the de Rham-Witt modules this is just the usual Godement construction. For the de Rham-Witt systems this has to be refined as follows: Let M be a de Rham-Witt system. Denote by $G(M_n)$ the $W_n \mathcal{O}_X$ -module given by

$$G(M_n)(U) = \prod_{x \in U} M_{n,x}, \quad U \subset X \text{ open,}$$

with the restriction maps given by projection. These sheaves fit together to form a de Rham-Witt system $G(M) = \{G(M_n)\}_{n \geq 1}$, such that the natural map $M \rightarrow G(M)$ is a morphism of de Rham-Witt systems.

For $m < n$ we denote by $i_{m,n} : W_m X \hookrightarrow W_n X$ the closed immersion induced by the restriction $W_n \mathcal{O}_X \rightarrow W_m \mathcal{O}_X$, in particular $i_{n-1,n} = i_n$. We set

$$\tilde{G}_n(M) := i_{1,n*} G(M_1) \oplus \dots \oplus i_{n-1,n*} G(M_{n-1}) \oplus G(M_n).$$

Then $\tilde{G}_n(M)$ is a graded $W_n \mathcal{O}_X$ -module. We define $W_n \mathcal{O}_X$ -linear maps $\pi, F, d, V, \underline{p}$, as follows

$$\begin{array}{ll}
 \pi : \tilde{G}_n \rightarrow i_{n*} \tilde{G}_{n-1}, & (m_1, \dots, m_n) \mapsto (m_1, \dots, m_{n-1}), \\
 F : \tilde{G}_n \rightarrow (W_n(F_X) i_n)_* \tilde{G}_{n-1}, & (m_1, \dots, m_n) \mapsto (F m_2, \dots, F m_n), \\
 d : W_n(F_X^n)_* \tilde{G}_n \rightarrow W_n(F_X^n)_* \tilde{G}_n(1), & (m_1, \dots, m_n) \mapsto (d m_1, \dots, d m_n), \\
 V : (W_n(F_X) i_n)_* \tilde{G}_{n-1} \rightarrow \tilde{G}_n, & (m_1, \dots, m_{n-1}) \mapsto (0, V m_1, \dots, V m_{n-1}), \\
 \underline{p} : i_{n*} \tilde{G}_{n-1} \rightarrow \tilde{G}_n, & (m_1, \dots, m_{n-1}) \mapsto (0, \underline{p} m_1, \dots, \underline{p} m_{n-1}).
 \end{array}$$

It is straightforward to check that $\tilde{G}(M) = (\{\tilde{G}_n(M)\}_{n \geq 1}, \pi, F, d, V, \underline{p})$ becomes a de Rham-Witt system and it is flasque by its definition. Also, the inclusion $M \hookrightarrow G(M)$ induces an inclusion

$$M_n \hookrightarrow \tilde{G}_n(M), \quad m \mapsto (\pi^{n-1}(m), \dots, \pi(m), m).$$

By definition this yields an inclusion of de Rham-Witt systems $M \hookrightarrow \tilde{G}(M)$ and we are done. \square

PROPOSITION 1.5.6. *Let $f : X \rightarrow Y$ be a morphism between k -schemes and Φ a family of supports on X . Then the right derived functors*

$$\begin{aligned} R\underline{\Gamma}_{\Phi} &: D^+(\mathrm{dRW}_X) \rightarrow D^+(\mathrm{dRW}_X), \\ Rf_* &: D^+(\mathrm{dRW}_X) \rightarrow D^+(\mathrm{dRW}_Y), \\ R\underline{\varprojlim} &: D^+(\mathrm{dRW}_X) \rightarrow D^+(\widehat{\mathrm{dRW}}_X) \\ Rf_{\Phi} &: D^+(\mathrm{dRW}_X) \rightarrow D^+(\mathrm{dRW}_Y), \\ R\hat{f}_{\Phi} &: D^+(\mathrm{dRW}_X) \rightarrow D^+(\widehat{\mathrm{dRW}}_Y), \end{aligned}$$

exist. Furthermore there are the following natural isomorphisms:

- (1) *Let $f : X \rightarrow Y$ and $g : Y \rightarrow Z$ morphisms, then $Rg_*Rf_* = R(g \circ f)_*$.*
- (2) *Let Φ and Ψ be two families of supports on X , then $R\underline{\Gamma}_{\Phi}R\underline{\Gamma}_{\Psi} = R\underline{\Gamma}_{\Phi \cap \Psi}$.*
- (3) *Let $f : X \rightarrow Y$ be a morphism and Ψ a family of supports on Y , then $R\underline{\Gamma}_{\Psi}Rf_* = Rf_*R\underline{\Gamma}_{f^{-1}(\Psi)}$.*
- (4) *Let $f : X \rightarrow Y$ be a morphism, then $R\underline{\varprojlim} Rf_* = Rf_*R\underline{\varprojlim}$.*
- (5) *Let $f : X \rightarrow Y$ be a morphism and Φ a family of supports on X . Then $Rf_{\Phi} = Rf_*R\underline{\Gamma}_{\Phi}$ and $R\hat{f}_{\Phi} = R\underline{\varprojlim} Rf_{\Phi}$. If Z is a closed subset of X and $\Phi = \Phi_Z$, then also $R\hat{f}_{\Phi_Z} = Rf_*R\underline{\Gamma}_Z R\underline{\varprojlim}$.*

Proof. The existence follows from [Har66, I, Cor. 5.3, β] (take P there to be the flasque objects) together with the Lemmas 1.5.5 and 1.5.4. The compatibility isomorphisms follow from [Har66, I, Cor 5.5] and Lemma 1.5.4, (2)-(4). \square

Remark 1.5.7. Let $f : X \rightarrow Y$ be an étale morphism between k -schemes. Then $W_n(f)$ is étale and thus $W_n(f)^*$ is exact on the category of $W_n(\mathcal{O}_Y)$ -modules. Therefore $f^* : \mathrm{dRW}_Y \rightarrow \mathrm{dRW}_X$ is exact and thus extends to

$$f^* : D^+(\mathrm{dRW}_Y) \rightarrow D^+(\mathrm{dRW}_X).$$

In case $j : U \hookrightarrow X$ is an open immersion we write $M|_U$ instead of j^*M for $M \in D^+(\mathrm{dRW}_X)$.

1.5.8. *Cousin-complex for de Rham-Witt systems.* Let X be a k -scheme and Z^\bullet the codimension filtration of X , i.e. Z^q is the family of supports on X consisting of all closed subsets of X whose codimension is at least q . Let M be a de Rham-Witt system on X . Take a complex of flasque de Rham-Witt systems G on X , which is a resolution of M , i.e. there is a quasi-isomorphism $M[0] \rightarrow G$. The filtration of complexes of de Rham-Witt systems

$$G \supset \underline{\Gamma}_{Z^1}(G) \supset \dots \supset \underline{\Gamma}_{Z^q}(G) \supset \dots$$

defines a spectral sequence of de Rham-Witt systems

$$E_1^{i,j} = \mathcal{H}_{Z^i/Z^{i+1}}^{i+j}(M) \implies \mathcal{H}^{i+j}(M),$$

where we put $\mathcal{H}_{Z^i/Z^{i+1}}^{i+j}(M) = H^{i+j}(\Gamma_{Z^i}(G)/\Gamma_{Z^{i+1}}(G))$. We define the *Cousin complex* of M (with respect to the codimension filtration) $E(M)$ to be the complex $E_1^{\bullet,0}$ coming from this spectral sequence, i.e. it is the complex of de Rham-Witt systems

$$E(M) : \mathcal{H}_{Z^0/Z^1}^0(M) \xrightarrow{d_1^{0,0}} \mathcal{H}_{Z^1/Z^0}^1(M) \xrightarrow{d_1^{1,0}} \dots \rightarrow \mathcal{H}_{Z^i/Z^{i+1}}^i(M) \xrightarrow{d_1^{i,0}} \dots$$

It satisfies the following properties:

- (1) $(E(M))_n = E(M_n)$ is the usual Cousin complex associated to M_n (see e.g. [Har66, IV, §2] or [Con00, p. 107-109]).
- (2)

$$E^i(M) = \mathcal{H}_{Z^i/Z^{i+1}}^i(M) = \bigoplus_{x \in X^{(i)}} i_{x*} H_x^i(M),$$

where $H_x^i(M) = (\varinjlim_{U \ni x} H_{\overline{\{x\}} \cap U}^i(U, M_n))_n$, which is a de Rham-Witt system on $\text{Spec } \mathcal{O}_{X,x}$ supported in the closed point x , $i_x : \text{Spec } \mathcal{O}_{X,x} \rightarrow X$ is the natural map and $X^{(i)}$ is the set of points x of codimension i in X (i.e. $\dim \mathcal{O}_{X,x} = i$).

- (3) The natural augmentation $M \rightarrow E(M)$ is a resolution of M if and only if $H_x^i(M_n) = 0$ for all $x \in X^{(j)}$ with $j \neq i$ and for all $n \geq 1$.

((1) holds since each G_n is a flasque resolution of M_n ; (2) follows from (1) and [Har66, IV, §1, Var. 8, Motif F]; (3) follows from (1) and [Har66, IV, Prop. 2.6, (iii) \iff (iv)] and [Har66, IV, §1, Var. 8, Motif F].)

LEMMA 1.5.9. *Let X be a smooth k -scheme. Then $E(W_\bullet \Omega_X)$ is a flasque resolution of quasi-coherent de Rham-Witt systems of the coherent de Rham-Witt system $W_\bullet \Omega_X$.*

Proof. By [Ill79, I, Cor. 3.9] the graded pieces of the standard filtration on $W_n \Omega_X^q$ are extensions of locally free \mathcal{O}_X -modules. Thus

$$(1.5.2) \quad H_x^i(W_n \Omega_X^q) = 0 \quad \text{for all } x \in X^{(j)}, \text{ with } j \neq i, \text{ and all } q, n \geq 1.$$

Thus $E(W_\bullet \Omega_X)$ is a quasi-coherent resolution of $W_\bullet \Omega_X$. Next we claim, that the transition morphisms

$$(1.5.3) \quad H_x^i(W_n \Omega_X^q) \rightarrow H_x^i(W_{n-1} \Omega_X^q)$$

are surjective for all $x \in X^{(i)}$ and $n \geq 2$. Indeed, for $x \in X^{(i)}$ we can always find an open affine neighborhood $U = \text{Spec } A$ of x and sections t_1, \dots, t_i such that $\overline{\{x\}} \cap U = V(t_1, \dots, t_i)$. This also implies for all $n \geq 1$, $W_n(\overline{\{x\}}) \cap W_n U = V([t_1], \dots, [t_i]) \subset W_n U$, where $[t] \in W_n A$ is the Teichmüller lift of $t \in A$. Then by [SGA2, Exp. II, Prop. 5]

$$H_{\overline{\{x\}} \cap U}^i(U, W_n \Omega_X) = \varinjlim_r \frac{\Gamma(U, W_n \Omega_X)}{([t_1]^r, \dots, [t_i]^r) \Gamma(U, W_n \Omega_X)}.$$

In particular the transition maps (1.5.3) are surjective. If we denote the kernel of the restriction morphism $W_n \Omega_X \rightarrow W_{n-1} \Omega_X$ by K_n , then this and (1.5.2)

implies, that the sequence

$$0 \rightarrow E^i(K_n) \rightarrow E^i(W_n \Omega_X) \rightarrow E^i(W_{n-1} \Omega_X) \rightarrow 0$$

is an exact sequence of flasque abelian sheaves on X and this proves the lemma. \square

1.6. WITT-DUALIZING SYSTEMS.

1.6.1. Let $f : X \rightarrow Y$ be a finite morphism between two finite dimensional noetherian schemes. Using the notation from [Har66, III, §6] we denote by $f^b : D^+(\mathcal{O}_Y) \rightarrow D^+(\mathcal{O}_X)$, the functor which sends a complex C to

$$f^b(C) = f^{-1} R\mathcal{H}om_{\mathcal{O}_Y}(f_* \mathcal{O}_X, C) \otimes_{f^{-1} f_* \mathcal{O}_X} \mathcal{O}_X.$$

Evaluation by 1 induces the finite trace morphism on $D_{qc}^+(\mathcal{O}_Y)$ (see [Har66, III, Prop. 6.5])

$$(1.6.1) \quad \text{Trf} : f_* f^b \rightarrow \text{id}_{D_{qc}^+(\mathcal{O}_Y)}$$

and composition with the natural map

$$(1.6.2) \quad \epsilon_f : f_* R\mathcal{H}om_X(-, -) \rightarrow R\mathcal{H}om_Y(f_*(-), f_*(-))$$

induces an isomorphism for any $A \in D_{qc}^-(\mathcal{O}_X)$, $B \in D_{qc}^+(\mathcal{O}_Y)$

$$(1.6.3) \quad \theta_f = \text{Trf}_f \circ \epsilon_f : f_* R\mathcal{H}om_X(A, f^b B) \xrightarrow{\cong} R\mathcal{H}om_Y(f_* A, B).$$

In particular, we see that for any morphism $\varphi : f_* A \rightarrow B$ in $D_{qc}(\mathcal{O}_Y)$, with A bounded above and B bounded below there exists a morphism ${}^a\varphi : A \rightarrow f^b B$ in $D_{qc}(\mathcal{O}_X)$, such that φ equals the composition

$$f_* A \xrightarrow{f_*({}^a\varphi)} f_* f^b B \xrightarrow{\text{Trf}_f} B.$$

We call ${}^a\varphi$ the adjoint of φ .

1.6.2. Let X be a k -scheme and denote by $D(\mathcal{C}_{\mathbb{N}, X}) = \prod_{n \geq 1} D(\mathcal{C}_{n, X})$ the derived category of $\mathcal{C}_{\mathbb{N}}$. Since the morphisms i_n and $W_n(F_X)$ are finite for all n , the functors i_* , σ_* , Σ_* are exact and extend to functors on $D(\mathcal{C}_{\mathbb{N}})$, which still satisfy the identities (1.2.1), (1.2.2). On $D_{qc}^+(\mathcal{C}_{\mathbb{N}})$ we define i^b , σ^b , Σ^b as follows:

$$(i^b M)_n := i_{n+1}^b M_{n+1}, \quad (\sigma^b M)_n := W_n(F_X)^b M_n, \quad (\Sigma^b M)_n = W_n(F_X^n)^b M_n.$$

There is an obvious way to define Trf_i , Trf_σ , Trf_Σ , ϵ_i , ϵ_σ , ϵ_Σ such that the compositions

$$i_* R\mathcal{H}om(M, i^b N) \xrightarrow{\epsilon_i} R\mathcal{H}om(i_* M, i_* i^b N) \xrightarrow{\text{Trf}_i} R\mathcal{H}om(i_* M, N),$$

$$\sigma_* R\mathcal{H}om(M, \sigma^b N) \xrightarrow{\epsilon_\sigma} R\mathcal{H}om(\sigma_* M, \sigma_* \sigma^b N) \xrightarrow{\text{Trf}_\sigma} R\mathcal{H}om(\sigma_* M, N),$$

$$\Sigma_* R\mathcal{H}om(M, \Sigma^b N) \xrightarrow{\epsilon_\Sigma} R\mathcal{H}om(\Sigma_* M, \Sigma_* \Sigma^b N) \xrightarrow{\text{Trf}_\Sigma} R\mathcal{H}om(\Sigma_* M, N)$$

are isomorphisms for $M \in D_{qc}^-(\mathcal{C}_{\mathbb{N}})$ and $N \in D_{qc}^+(\mathcal{C}_{\mathbb{N}})$.

DEFINITION 1.6.3 ([Eke84, III, Def. 2.2]). A *Witt quasi-dualizing system* on X is a collection (Q, \underline{p}, C, V) where Q is an object in $\mathcal{C}_{\mathbb{N}, \text{qc}}$ and

$$\underline{p} : i_*Q \rightarrow Q, \quad C : \Sigma_*Q \rightarrow Q, \quad V : \sigma_*i_*Q \rightarrow Q$$

are morphisms in $\mathcal{C}_{\mathbb{N}}$ such that the following holds:

- (a) $V \circ \sigma_*i_*C = C \circ \Sigma_*\underline{p}$,
- (b) $\underline{p} \circ i_*V = V \circ \sigma_*i_*\underline{p}$.

A *Witt dualizing system* is a Witt quasi-dualizing system, which has the additional property, that the adjoints

$$(1.6.4) \quad {}^a\underline{p} : Q \xrightarrow{\simeq} i^bQ, \quad {}^aC : Q \xrightarrow{\simeq} \Sigma^bQ, \quad {}^aV : Q \xrightarrow{\simeq} i^b\sigma^bQ$$

are quasi-isomorphisms.

A morphism φ between Witt (quasi-) dualizing systems is a morphism in $\mathcal{C}_{\mathbb{N}}$ commuting with \underline{p}, V , and C .

A Witt (quasi-) dualizing system (Q, \underline{p}, C, V) on X is called *coherent* if Q_n is coherent for all $n \geq 1$.

EXAMPLE 1.6.4. (1) The system

$$W_\bullet\omega := (\{W_n\}_{n \geq 1}, \underline{p}, C := \{W_n(F_{\text{Spec } k})^{-n}\}, V := \{W_n(F_{\text{Spec } k})^{-1}\underline{p}\})$$

is a Witt dualizing system on $\text{Spec } k$, where \underline{p} is the usual map “lift and multiply by p ”, which is concentrated degree 0. For this, first notice that W_n is an injective W_n -module for all $n \geq 1$. Then one easily checks that the following maps are isomorphisms and adjoint to \underline{p}, C and V respectively:

$$\begin{aligned} W_\bullet &\xrightarrow{\simeq} i^bW_\bullet = \text{Hom}_{W_\bullet}(i_*W_\bullet, W_\bullet), & a \mapsto (b \mapsto \underline{p}ab), \\ W_\bullet &\xrightarrow{\simeq} \Sigma^bW_\bullet = \text{Hom}_{W_\bullet}((\Sigma_*W_\bullet), W_\bullet), & a \mapsto (b \mapsto Cab), \\ W_\bullet &\xrightarrow{\simeq} i^b\sigma^bW_\bullet = \text{Hom}_{W_\bullet}(\sigma_*i_*W_\bullet, W_\bullet), & a \mapsto (b \mapsto Vab). \end{aligned}$$

(2) Let X be a smooth k -scheme of pure dimension N . Then

$$W_\bullet\omega_X := (\{W_n\Omega_X^N\}_{n \geq 1}, \underline{p}, C, V)$$

is a Witt dualizing system which by definition is concentrated in degree N . Here \underline{p} is “lift and multiply by p ” and V is the Verschiebung. On the n -th level C is defined as the composition:

$$C^n : (\Sigma_*W_\bullet\Omega_X^N)_n \rightarrow (\Sigma_*W_\bullet\Omega_X^N/d(W_\bullet\Omega_X^{N-1}))_n \xrightarrow{(C^{-n})^{-1}} W_n\Omega_X^N,$$

where $C^{-n} : W_n\Omega_X^N \xrightarrow{\simeq} W_n(F_X^n)_*W_n\Omega_X^N/d(W_n\Omega_X^{N-1})$ is the inverse Cartier isomorphism from [IR83, III, Prop. (1.4)]. One easily checks that \underline{p}, C^n and V satisfy the relations (a), (b) in Definition (1.6.3). The condition on the adjoints (1.6.4) is harder and follows from Ekedahl’s result $W_n\Omega_X^N = W_n(f)^!W_n$, with $f : X \rightarrow \text{Spec } k$ the structure map, see [Eke84, I and II, Ex. 2.2.1.]. Notice that $W_\bullet\omega_{\text{Spec } k} = W_\bullet\omega$.

1.6.5. Let (Q, \underline{p}, C, V) be a Witt dualizing system on X . We may express the equalities in (1.6.4) as

$$H^0({}^a\varphi) : Q \xrightarrow{\cong} H^0(f^b Q) \quad \text{and} \quad H^0(f^b Q)[0] \cong f^b Q,$$

where $(f, \varphi) \in \{(i, \underline{p}), (\sigma, V), (\Sigma, C)\}$. Therefore by the definition of the adjoints, \underline{p} , C and V factor as follows:

$$\begin{aligned} \underline{p} : i_* Q &\xrightarrow{\cong, H^0({}^a\underline{p})} i_* H^0(i^b Q) \xrightarrow{H^0(\text{Tr}_f)} Q, \\ C : \Sigma_* Q &\xrightarrow{\cong, H^0({}^a C)} \Sigma_* H^0(\Sigma^b Q) \xrightarrow{H^0(\text{Tr}_\Sigma)} Q, \\ V : \sigma_* i_* Q &\xrightarrow{\cong, H^0({}^a V)} \sigma_* i_* H^0(i^b \sigma^b Q) \xrightarrow{H^0(\text{Tr}_{\sigma i})} Q. \end{aligned}$$

Furthermore it follows from 1.6.2, that the natural transformations

$$\begin{aligned} \underline{p} \circ \epsilon_i : i_* \mathcal{H}om_{\mathcal{C}_{\mathbb{N}}}((-), Q) &\rightarrow \mathcal{H}om_{\mathcal{C}_{\mathbb{N}}}(i_*(-), Q), \\ C \circ \epsilon_\Sigma : \Sigma_* \mathcal{H}om_{\mathcal{C}_{\mathbb{N}}}((-), Q) &\rightarrow \mathcal{H}om_{\mathcal{C}_{\mathbb{N}}}(\Sigma_*(-), Q), \\ V \circ \epsilon_{\sigma i} : \sigma_* i_* \mathcal{H}om_{\mathcal{C}_{\mathbb{N}}}((-), Q) &\rightarrow \mathcal{H}om_{\mathcal{C}_{\mathbb{N}}}(\sigma_* i_*(-), Q) \end{aligned}$$

are isomorphisms when restricted to the category $\mathcal{C}_{\mathbb{N}, \text{qc}}$.

1.6.6. Let M be a quasi-coherent de Rham-Witt system and Q a Witt dualizing system on X . Then we may define maps π , F , V , d and \underline{p} on $\mathcal{H}om_{\mathcal{C}_{\mathbb{N}}}(M, Q) \in \mathcal{C}_{\mathbb{N}}$ as follows:

$$\begin{aligned} \pi : \mathcal{H}om(M, Q) &\xrightarrow{\circ \underline{p}} \mathcal{H}om(i_* M, Q) \xrightarrow{(\underline{p} \circ \epsilon_i)^{-1}} i_* \mathcal{H}om(M, Q), \\ F : \mathcal{H}om(M, Q) &\xrightarrow{\circ V} \mathcal{H}om(\sigma_* i_* M, Q) \xrightarrow{(V \circ \epsilon_{\sigma i})^{-1}} \sigma_* i_* \mathcal{H}om(M, Q), \\ V : \sigma_* i_* \mathcal{H}om(M, Q) &\xrightarrow{(V \circ \epsilon_{\sigma i})} \mathcal{H}om(\sigma_* i_* M, Q) \xrightarrow{\circ F} \mathcal{H}om(M, Q), \\ \underline{p} : i_* \mathcal{H}om(M, Q) &\xrightarrow{\underline{p} \circ \epsilon_i} \mathcal{H}om(i_* M, Q) \xrightarrow{\circ \pi} \mathcal{H}om(M, Q), \\ d : \Sigma_* \mathcal{H}om(M, Q) &\xrightarrow{(C \circ \epsilon_\Sigma)} \mathcal{H}om(\Sigma_* M, Q) \xrightarrow{\circ d} \mathcal{H}om(\Sigma_* M(-1), Q) \\ &\xrightarrow{\cong, \alpha} \mathcal{H}om(\Sigma_* M, Q)(1) \xrightarrow{(C \circ \epsilon_\Sigma)^{-1}} \Sigma_* \mathcal{H}om(M, Q)(1), \end{aligned}$$

where the isomorphism $\alpha : \mathcal{H}om(\Sigma_* M(-1), Q) \xrightarrow{\cong} \mathcal{H}om(\Sigma_* M, Q)(1)$ is given by multiplication with $(-1)^{q+1}$ in degree q .

PROPOSITION 1.6.7 ([Eke84, III, 2.]). *The above construction yields a functor*

$$\mathcal{H}om(-, Q) : (\text{dRW}_{X, \text{qc}})^o \longrightarrow \text{dRW}_X,$$

which has the following properties

- (1) A morphism of Witt dualizing systems $Q \rightarrow Q'$ induces a natural transformation of functors $\mathcal{H}om(-, Q) \rightarrow \mathcal{H}om(-, Q')$.
- (2) The functor $\mathcal{H}om(-, Q)$ restricts to a functor $(\text{dRW}_{X, c})^o \rightarrow \text{dRW}_{X, \text{qc}}$ and if Q is coherent, then also to $(\text{dRW}_{X, c})^o \rightarrow \text{dRW}_{X, c}$.

(3) For all $M \in \mathrm{dRW}_{X,\mathrm{qc}}$ and all $m \in \mathbb{Z}$ there is a natural isomorphism

$$\mathrm{Hom}(M(m), Q) \simeq \mathrm{Hom}(M, Q)(-m),$$

given by multiplication with $(-1)^{qm + \frac{m(m-1)}{2}}$ in degree q and a natural isomorphism

$$\mathrm{Hom}(M, Q(m)) \simeq \mathrm{Hom}(M, Q)(m)$$

given by the identity in each degree. (There is some freedom in defining these isomorphisms; our choice is compatible with the sign convention for complexes in [Con00].)

Proof. It is straightforward to check the relations in Definition 1.2.4. \square

1.7. RESIDUAL COMPLEXES AND TRACES. In this section A will always be a regular local ring, all schemes are of finite type over A and all morphisms will be A -morphisms. The results of this section will be applied in the next sections in the case $A = W$ and schemes of finite type over W_n , some n .

1.7.1. *Review of residual complexes.* The general references for residual complexes are [Har66, VI], [Con00, 3.2]. Let X be an A -scheme. A *residual complex* on X is a bounded complex K of quasi-coherent and injective \mathcal{O}_X -modules, which has *coherent cohomology* and such that there exists an isomorphism of \mathcal{O}_X -modules $\bigoplus_{q \in \mathbb{Z}} K^q \cong \bigoplus_{x \in X} i_{x*} J(x)$, where $i_x : \mathrm{Spec} \mathcal{O}_{X,x} \hookrightarrow X$ is the inclusion and $J(x)$ is an injective hull of $k(x)$ in $\mathcal{O}_{X,x}$ (i.e. it is an injective $\mathcal{O}_{X,x}$ -module which contains $k(x)$ and such that, for any $0 \neq a \in J(x)$ exists a $b \in \mathcal{O}_{X,x}$ with $0 \neq ba \in k(x)$). It follows that $i_{x*} J(x)$ is supported in $\overline{\{x\}}$. The *codimension function* on X associated to K is the unique function $d_K : |X| \rightarrow \mathbb{Z}$ such that $K^q = \bigoplus_{d_{K^\bullet}(x)=q} i_{x*} J(x)$ for all q . If x_0 is an immediate specialization of $x \in X$ (i.e. x_0 is a point of codimension 1 in $\overline{\{x\}}$), then $d_K(x_0) = d_K(x) + 1$. The *filtration* $\dots \subset Z_K^q \subset Z_K^{q-1} \subset \dots \subset X$ associated to K is defined by $Z_K^q := \{x \in X \mid d_K(x) \geq q\}$. On each irreducible component of X this filtration equals the shifted codimension filtration.

If $R \in D_c^b(X)$ is a dualizing complex with associated codimension filtration Z^\bullet (see [Con00, 3.1] for these notions), then the Cousin complex $E_{Z^\bullet}(R)$ of R with respect to Z^\bullet is a residual complex with associated filtration also Z^\bullet . In $D_c^b(X)$ we have $R \cong E_{Z^\bullet}(R)$ (since a dualizing complex is Gorenstein). Particular examples of dualizing complexes are W_n on $X = \mathrm{Spec} W_n$, $\mathcal{O}_X[0]$ in case X is regular, $\omega_X[N] = \Omega_{X/A}^N[N]$ in case X is smooth of pure dimension N and more general $f^! R \in D_c^b(X')$, where $f : X' \rightarrow X$ is a finite type morphism and R is a dualizing complex on X .

On the other hand any residual complex on X is a dualizing complex when viewed as a complex in $D_c^b(X)$. Furthermore, if K is a residual complex on Y with filtration Z^\bullet , then we have an equality of complexes $E_{Z^\bullet}(K) = K$.

1.7.2. $(-)^!$ for residual complexes. Let $f : X \rightarrow Y$ be a finite type morphism between A -schemes and K a residual complex on Y with associated filtration $Z_K^\bullet =: Z^\bullet$ (which exists by the above). Then there is a functor f^Δ from the category of residual complexes with filtration Z^\bullet on Y to the category of residual complexes on X with a fixed filtration denoted by $f^\Delta Z^\bullet$, having the following properties:

- (1) For two finite type morphisms $f : X \rightarrow Y$ and $g : Y \rightarrow Z$ of A -schemes, there is an isomorphism $c_{f,g} : (gf)^\Delta \xrightarrow{\cong} f^\Delta g^\Delta$, which is compatible with triple compositions and such that $c_{\text{id},f} = \text{id} = c_{f,\text{id}}$.
- (2) If $f : X \rightarrow Y$ is smooth and separated of relative dimension r , then there is an isomorphism

$$\varphi_f : f^\Delta K \xrightarrow{\cong} E_{f^{-1}Z^\bullet[r]}(\Omega_{X/Y}^r[r] \otimes f^*K).$$

Here $E_{f^{-1}Z^\bullet[r]}(\Omega_{X/Y}^r[r] \otimes f^*K)$ is the Cousin complex associated to the complex $\Omega_{X/Y}^r[r] \otimes f^*K$ and the filtration $f^{-1}Z^\bullet[r]$. If we have two smooth separated maps of some fixed relative dimension f and g as in (1), then $c_{f,g}$ is compatible with the natural map on the right.

- (3) If $f : X \rightarrow Y$ is finite, then there is an isomorphism

$$\psi_f : f^\Delta K \xrightarrow{\cong} E_{f^{-1}Z^\bullet}(\bar{f}^*R\mathcal{H}om_{\mathcal{O}_Y}(f_*\mathcal{O}_X, K)) = \bar{f}^*\mathcal{H}om_{\mathcal{O}_Y}(f_*\mathcal{O}_X, K),$$

where we set $\bar{f}^*(-) := f^{-1}(-) \otimes_{f^{-1}f_*\mathcal{O}_X} \mathcal{O}_X$ (which is an exact functor). If we have two finite maps f and g as in (1), then $c_{f,g}$ is compatible with the natural map on the right of the above isomorphism.

- (4) Let

$$\begin{array}{ccc} X_U & \xrightarrow{u'} & X \\ \downarrow f' & \searrow h & \downarrow f \\ U & \xrightarrow{u} & Y \end{array}$$

be a cartesian diagram of A -schemes with u étale. Then there is an isomorphism

$$d_{u,f} : f'^\Delta u^* \xrightarrow{\cong} u'^* f^\Delta,$$

which is compatible with compositions in u and f and with the isomorphisms in (2) and (3). Furthermore, by (2) we have $u^* \cong u^\Delta$ and $u'^* \cong u'^\Delta$ and the following diagram commutes

$$\begin{array}{ccc} f'^\Delta u^* & \xrightarrow{d_{u,f}} & u'^* f^\Delta \\ \downarrow \cong & & \downarrow \cong \\ f'^\Delta u^\Delta & \xleftarrow{c_{f',u}} h^\Delta \xrightarrow{c_{u',f}} & u'^\Delta f^\Delta. \end{array}$$

(To prove this commutativity one may assume that f factors as $X \xrightarrow{i} P \xrightarrow{\pi} Y$, with i a closed immersion and π smooth and then use (2)

and (3), cf. [Con00, (3.3.34)].) Finally if f and g are étale, then $c_{f,g} : (gf)^\Delta \cong f^\Delta g^\Delta$ corresponds to the natural isomorphism $(gf)^* \cong f^*g^*$.

In fact f^Δ is defined by locally factoring $f : X \rightarrow Y$ into a closed immersion followed by a smooth morphism and then use (2) and (3) and glue (and then show that what you obtain is independent of all the choices). In $D_c^b(X)$ we have $f^\Delta K \cong f^!K$. For details, as well as more compatibilities, see [Har66, VI] and [Con00, 3.2].

1.7.3. *The Trace for residual complexes.* The reference for this section is [Har66, VI, 4,5, VII, 2] and [Con00, 3.4]. Let $f : X \rightarrow Y$ be a finite type morphism between A -schemes. Let K be a residual complex on Y . Then there exists a morphism of *graded sheaves* (in general not of complexes, which we indicate by the dotted arrow)

$$\mathrm{Tr}_f : f_* f^\Delta K \cdots \cdots \dashrightarrow K,$$

which satisfies the following properties (and is also uniquely determined by the first three of them):

- (1) Tr_f is functorial with respect to maps between residual complexes with same associated filtration and $\mathrm{Tr}_{\mathrm{id}} = \mathrm{id}$.
- (2) If $g : Y \rightarrow Z$ is another morphism of finite type between A -schemes, then

$$\mathrm{Tr}_{gf} = \mathrm{Tr}_g \circ g_*(\mathrm{Tr}_f) \circ (gf)_* c_{f,g}.$$

- (3) If f is finite, then Tr_f is a morphism of complexes and equals the composition

$$\mathrm{Tr}_f : f_* f^\Delta K \xrightarrow{\simeq, \psi_f} \mathcal{H}om_{\mathcal{O}_Y}(f_* \mathcal{O}_X, K) \xrightarrow{\mathrm{ev. at 1}} K.$$

- (4) Tr_f is compatible with étale base change (using the maps $d_{u,f}$ from 1.7.2, (4)).
- (5) If $f : X \rightarrow Y$ is *proper*, then $\mathrm{Tr}_f : f_* f^\Delta K \rightarrow K$ is a morphism of complexes.
- (6) If $f : \mathbb{P}_X^n \rightarrow X$ is the projection, then Tr_f is the composition (in $D_c^b(X)$)

$$\mathrm{Tr}_f : f_* f^\Delta K \xrightarrow{\simeq, \varphi_f} Rf_*(\Omega_{\mathbb{P}_X^n/X}^n[n]) \otimes K \simeq K,$$

where for the first isomorphism we used 1.7.2, (2) and the projection formula, the second isomorphism is induced by base change from the isomorphism

$$\mathbb{Z} \xrightarrow{\simeq} H^n(\mathbb{P}_{\mathbb{Z}}^n, \Omega_{\mathbb{P}_{\mathbb{Z}}^n/\mathbb{Z}}^n) = \check{H}^n(\mathfrak{U}, \Omega_{\mathbb{P}_{\mathbb{Z}}^n/\mathbb{Z}}^n), \quad 1 \mapsto (-1)^{\frac{n(n+1)}{2}} \frac{dt_1 \wedge \dots \wedge dt_n}{t_1 \cdots t_n},$$

where $\mathfrak{U} = \{U_0, \dots, U_n\}$ is the standard covering of $\mathbb{P}_{\mathbb{Z}}^n$ and the t_i 's are the coordinate functions on U_0 .

- (7) (Grothendieck-Serre duality, special case) If $f : X \rightarrow Y$ is *proper*, then for any $C \in D_{\mathrm{qc}}^-(X)$ the composition

$$Rf_* R\mathcal{H}om_X(C, f^\Delta K) \xrightarrow{\epsilon_f} R\mathcal{H}om_Y(Rf_* C, f_* f^\Delta K) \xrightarrow{\mathrm{Tr}_f} R\mathcal{H}om_Y(Rf_* C, K)$$

is an isomorphism. It is compatible with étale base change.

DEFINITION 1.7.4. We denote by $\text{Sch}_{A,*}$ the category with objects given by pairs (X, Φ) , where X is a scheme of finite type over A and Φ is a family of supports on X , and the morphisms $f : (X, \Phi) \rightarrow (Y, \Psi)$ are given by separated A -morphisms $f : X \rightarrow Y$, whose restriction to Φ is *proper* and which satisfy $f(\Phi) \subset \Psi$.

Remark 1.7.5. Let (X, Φ) be an object in $\text{Sch}_{A,*}$ and K a residual complex on X . Then $\Gamma_{\Phi} K$ is a direct summand of K and is a complex of quasi-coherent and injective \mathcal{O}_X -modules. Indeed the isomorphism $\bigoplus_{q \in \mathbb{Z}} K^q \cong \bigoplus_{x \in X} i_{x*} J(x)$ (see 1.7.1) implies $\bigoplus_{q \in \mathbb{Z}} \Gamma_{\Phi} K^q \cong \bigoplus_{\overline{\{x\}} \in \Phi} i_{x*} J(x)$.

COROLLARY 1.7.6. Let $f : (X, \Phi) \rightarrow (Y, \Psi)$ be a morphism in $\text{Sch}_{A,*}$ and K a residual complex on Y , then there exists a morphism of complexes

$$\text{Tr}_f : f_{\Phi} f^{\Delta} K \rightarrow \Gamma_{\Psi}(K),$$

where we set $f_{\Phi} := f_* \circ \Gamma_{\Phi}$, which satisfies the following properties:

- (1) Tr_f is functorial with respect to maps between residual complexes with same associated filtration and $\text{Tr}_{\text{id}} = \text{id}$.
- (2) The following diagram commutes

$$\begin{array}{ccc} f_{\Phi} f^{\Delta} K & \xrightarrow{\text{Tr}_f} & \Gamma_{\Psi}(K) \\ \downarrow & & \downarrow \\ f_* f^{\Delta} K & \xrightarrow{\text{Tr}_f} & K, \end{array}$$

where the vertical maps are the natural ones and the lower horizontal map is the trace from 1.7.3, which is only a map of graded sheaves (visualized by the dotted arrow). By abuse of notation we write f for both, the morphism $(X, \Phi) \rightarrow (Y, \Psi)$ in $\text{Sch}_{A,*}$ and the underlying morphism of schemes $X \rightarrow Y$.

- (3) If $g : (Y, \Psi) \rightarrow (Z, \Xi)$ is another morphism in $\text{Sch}_{A,*}$, then

$$\text{Tr}_{gf} = \text{Tr}_g \circ g_*(\text{Tr}_f) \circ (gf)_* c_{f,g} : (gf)_{\Phi} (gf)^{\Delta} K \xrightarrow{c_{f,g}} g_{\Psi} f_{\Phi} f^{\Delta} g^{\Delta} K \xrightarrow{\text{Tr}_f} g_{\Psi} g^{\Delta} K \xrightarrow{\text{Tr}_g} \Gamma_{\Xi}(K).$$

- (4) Tr_f is compatible with étale base change in the following sense: Let

$$\begin{array}{ccc} X_U & \xrightarrow{u'} & X \\ f' \downarrow & & \downarrow f \\ U & \xrightarrow{u} & Y \end{array}$$

be a cartesian square of finite type A -schemes with u étale; let Φ, Ψ, Φ' and Ψ' be families of supports on X, Y, X_U and U , such that $f : (X, \Phi) \rightarrow (Y, \Psi)$ and $f' : (X_U, \Phi') \rightarrow (U, \Psi')$ are in $\text{Sch}_{A,*}$ and

additionally $u^{-1}(\Psi) \subset \Psi'$, $u'^{-1}(\Phi) \subset \Phi'$. Then $u^*K \simeq u^\Delta K$ is a residual complex and the following diagram commutes:

$$\begin{array}{ccc} u^* f_\Phi f^\Delta K & \xrightarrow{u^*(\text{Tr}_f)} & u^* \Gamma_\Psi K \\ \downarrow & & \downarrow \\ f'_{\Phi'} f'^\Delta u^* K & \xrightarrow{\text{Tr}_{f'}} & \Gamma_{\Psi'} u^* K. \end{array}$$

Here the vertical maps are given as follows: First, the composition $\Gamma_\Psi \rightarrow \Gamma_\Psi u_* u^* \simeq u_* \Gamma_{u^{-1}(\Psi)} u^* \rightarrow u_* \Gamma_{\Psi'} u^*$ gives by adjunction a map $u^* \Gamma_\Psi \rightarrow \Gamma_{\Psi'} u^*$, yielding the vertical map on the right in the diagram; similar we have a map $u'^* \Gamma_{\Phi'} \rightarrow \Gamma_{\Phi'} u'^*$ inducing a map $u^* f_\Phi \rightarrow f'_{\Phi'} u'^*$; the vertical map on the left in the diagram is then given by the composition $u^* f_\Phi f^\Delta \rightarrow f'_{\Phi'} u'^* f^\Delta \xrightarrow{d_{u,f}^{-1}} f'_{\Phi'} f'^\Delta u^*$, where $d_{u,f}$ is the map from 1.7.2, (4).

- (5) Let $j : U \rightarrow X$ be an open immersion such that Φ is contained in U . Then $j : (U, \Phi) \hookrightarrow (X, \Phi)$ is a morphism in $\text{Sch}_{A,*}$ and $\text{Tr}_j : j_\Phi j^\Delta K \rightarrow \Gamma_\Phi K$ is the excision isomorphism, more precisely: Tr_j is given by the composition

$$(1.7.1) \quad \text{Tr}_j : j_\Phi j^\Delta K = \Gamma_{(X,\Phi)} j_* j^\Delta K \simeq \Gamma_{(X,\Phi)} j_* j^* K \stackrel{\text{exc.}}{=} \Gamma_{(X,\Phi)} K.$$

Proof. We define $\text{Tr}_f : f_\Phi f^\Delta K \rightarrow \Gamma_\Psi K$ to be the following composition

$$(1.7.2) \quad f_\Phi f^\Delta K \rightarrow f_* \Gamma_{f^{-1}(\Psi)} f^\Delta K \simeq \Gamma_\Psi f_* f^\Delta K \xrightarrow{1.7.3} \Gamma_\Psi K.$$

A priori this is only a map of graded sheaves. But we already observe, that the properties 1)- 4) follow immediately from the definition and the corresponding properties in 1.7.3. If $j : (U, \Phi) \hookrightarrow (X, \Phi)$ is an open immersion as in (5), we may apply the excision identity $\Gamma_{(X,\Phi)} = \Gamma_{(X,\Phi)} j_* j^*$ to the map of graded sheaves $\text{Tr}_j : j_* j^\Delta(-) \dashrightarrow (-)$ to obtain a commutative diagram

$$\begin{array}{ccc} j_\Phi j^\Delta K = \Gamma_{(X,\Phi)} j_* j^\Delta K & \dashrightarrow^{\text{Tr}_j} & \Gamma_{(X,\Phi)} K \\ \parallel & & \parallel \\ \Gamma_{(X,\Phi)} j_* j^*(j_* j^\Delta K) & \dashrightarrow^{j_* j^*(\text{Tr}_j)} & \Gamma_{(X,\Phi)} j_* j^* K. \end{array}$$

Now using the compatibility with base change as in (4) (in the situation $j = u = f$) implies that going around the diagram from the top left corner to the top right corner counterclockwise is the isomorphism (of complexes) (1.7.1). This gives (5) and in particular Tr_j is a morphism of complexes. It remains to show that Tr_f as defined in (1.7.2) is in fact a morphism of complexes. For this we factor $f : X \rightarrow Y$ into an open immersion $j : X \hookrightarrow \bar{X}$ followed by a proper A -morphism $\bar{f} : \bar{X} \rightarrow Y$ (Nagata compactification). Since

the restriction of f to Φ is proper, it follows that Φ also defines a family of supports on \bar{X} . We consider the following diagram

$$\begin{array}{ccc}
 f_{\Phi} f^{\Delta} K & \longrightarrow & \Gamma_{\Psi} f_* f^{\Delta} K \cdots \xrightarrow{\text{Tr}_f} \Gamma_{\Psi}(K) \\
 \simeq \downarrow & & \downarrow \text{Tr}_j \circ c_{\bar{f}, j} \\
 \bar{f}_{\Phi} \bar{f}^{\Delta} K & \longrightarrow & \Gamma_{\Psi} \bar{f}_* \bar{f}^{\Delta} K,
 \end{array}$$

where the vertical isomorphism on the left is a morphism of complexes, which is given by

$$f_* \Gamma_{(X, \Phi)} f^{\Delta} K = \bar{f}_* \Gamma_{(\bar{X}, \Phi)} j_* f^{\Delta} K \xrightarrow{c_{\bar{f}, j}} \bar{f}_* \Gamma_{(\bar{X}, \Phi)} j_* j^{\Delta} \bar{f}^{\Delta} K \xrightarrow{\text{Tr}_j, (5)} \bar{f}_* \Gamma_{(\bar{X}, \Phi)} \bar{f}^{\Delta} K.$$

Further $\text{Tr}_{\bar{f}}$ is a morphism of complexes by (1.7.3, (5)). The diagram is commutative by 1.7.3, 2) and hence Tr_f as defined in (1.7.2) is a morphism of complexes. \square

1.7.7. Let X be a finite type A -scheme and $\emptyset = Z^r \subset \dots \subset Z^{q+1} \subset Z^q \subset \dots \subset Z^s = X$ be a filtration with $r > s \in \mathbb{Z}$ and such that Z^q is stable under specialization and any $y \in Z^q \setminus Z^{q+1}$ is not a specialization of any other point in Z^q . Recall that a *Cousin complex on X with respect to Z^{\bullet}* is a complex C^{\bullet} of quasi-coherent \mathcal{O}_X -modules, such that for all q the terms C^q are supported in the Z^q/Z^{q+1} -skeleton, i.e. $C^q \cong \bigoplus_{x \in Z^q \setminus Z^{q+1}} i_{x*} M_x$, where $i_x : \text{Spec } \mathcal{O}_{X,x} \hookrightarrow X$ is the inclusion and M_x is a quasi-coherent sheaf on $\text{Spec } \mathcal{O}_{X,x}$ supported only in the closed point x . Notice that $i_{x*} M_x$ is the extension by zero of the constant sheaf M_x on $\{x\}$. Any residual complex with associated filtration Z^{\bullet} is in particular a Cousin complex with respect to Z^{\bullet} . If G is any complex of quasi-coherent \mathcal{O}_X -modules, then $E_{Z^{\bullet}}(G)$ is a Cousin complex with respect to Z^{\bullet} .

LEMMA 1.7.8. *If $f : X \rightarrow Y$ is finite and D is a Cousin complex on X with respect to $f^{-1}Z^{\bullet}$ (Z^{\bullet} as above), then $f_* D$ is a Cousin complex on Y with respect to Z^{\bullet} .*

Proof. Write $D^q \cong \bigoplus_{x \in f^{-1}Z^q \setminus f^{-1}Z^{q+1}} i_{x*} N_x$ as above, in particular N_x is supported in x . Then

$$f_* D^q \cong \bigoplus_{y \in Z^q \setminus Z^{q+1}} i_{y*} M_y, \quad \text{with } M_y := f_{|X_{(y)}|*} \bigoplus_{x \in f^{-1}(y)} i_{x, f^{-1}(y)*} N_x,$$

where $i_{x, f^{-1}(y)} : \text{Spec } \mathcal{O}_{X,x} \hookrightarrow X_{(y)} = X \times_Y \text{Spec } \mathcal{O}_{Y,y}$ is the inclusion. M_y is supported in y and this gives the claim. \square

COROLLARY 1.7.9. *Let $f : X \rightarrow Y$ be a finite morphism between finite-type A -schemes and K a residual complex on Y with filtration Z^{\bullet} and C a Cousin complex on X with respect to $f^{-1}Z^{\bullet}$. Then the isomorphism of 1.7.3, (7) induces an isomorphism*

$$\text{Hom}_X(C, f^{\Delta} K) \cong \text{Hom}_Y(f_* C, K).$$

This isomorphism is compatible with étale base change and is concretely induced by sending a morphism of complexes $\varphi : C \rightarrow f^\Delta K$ to the composition $f_*C \xrightarrow{f_*(\varphi)} f_*f^\Delta K \xrightarrow{\text{Tr}_f} K$.

Proof. Since K and $f^\Delta K$ are complexes of injectives and f is finite, 1.7.3, (7) immediately gives (for all C)

$$\text{Hom}_X(C, f^\Delta K)/\text{homotopy} \cong \text{Hom}_Y(f_*C, K)/\text{homotopy}.$$

By Lemma 1.7.8 above f_*C^q is supported in the Z^q/Z^{q+1} -skeleton, which by definition also holds for K^q , for all q . Thus there is only the trivial homotopy between f_*C^\bullet and K . Similar with $f^\Delta K$ and C . \square

LEMMA 1.7.10. *Let $f : X \rightarrow Y$ be a finite morphism between finite type A -schemes and K a residual complex on Y with associated filtration Z^\bullet . Then for all $q \in \mathbb{Z}$ the following equality holds in $D_{\text{qc}}^b(X)$*

$$(f^\Delta K)^q \cong \bar{f}^* \mathcal{H}om_Y(f_*\mathcal{O}_X, K^q) \cong f^b(K^q) \cong \bigoplus_{x \in f^{-1}(Z^q) \setminus f^{-1}(Z^{q+1})} i_{x*} J(x),$$

where $\bar{f}^*(-) := f^{-1}(-) \otimes_{f^{-1}f_*\mathcal{O}_X} \mathcal{O}_X$, $i_x : \text{Spec } \mathcal{O}_{X,x} \hookrightarrow X$ is the inclusion and $J(y')$ is an injective hull of $k(y')$ in $\mathcal{O}_{Y',y'}$.

Proof. The first isomorphism follows from 1.7.2, (3), the second holds since K^q is injective and the third is [Har66, VI, Lem. 4.1]. \square

1.8. WITT RESIDUAL COMPLEXES. Let X be a k -scheme.

1.8.1. Let $C(\mathcal{C}_{\mathbb{N}})$ be the category of complexes of $\mathcal{C}_{\mathbb{N}}$. Recall that it is equipped with endo-functors i_*, σ_*, Σ_* .

NOTATION 1.8.2. We say that a complex K in $C(\mathcal{C}_{\mathbb{N}})$ is a *residual complex* if K_n is a residual complex on $W_n(X)$ for all n and the associated filtrations on $|X| = |W_n(X)|$ are all the same.

We say that a complex C in $C(\mathcal{C}_{\mathbb{N}})$ is a *Cousin complex* if C_n is a Cousin complex for all n with respect to the same filtration.

For a residual complex $K \in C(\mathcal{C}_{\mathbb{N}})$ we define

$$\begin{aligned} (i^\Delta K)_n &:= i_{n+1}^\Delta K_{n+1} \\ (\sigma^\Delta K)_n &:= W_n(F_X)^\Delta K_n \\ (\Sigma^\Delta K)_n &:= W_n(F_X^n)^\Delta K_n. \end{aligned}$$

This yields endo-functors on the full subcategory of residual complexes with some fixed filtration of $C(\mathcal{C}_{\mathbb{N}})$.

In view of Corollary 1.7.9 we get

$$\begin{aligned} \text{Hom}_{C(\mathcal{C}_{\mathbb{N}})}(i_*C, K) &= \text{Hom}_{C(\mathcal{C}_{\mathbb{N}})}(C, i^\Delta K) \\ \text{Hom}_{C(\mathcal{C}_{\mathbb{N}})}(\sigma_*C, K) &= \text{Hom}_{C(\mathcal{C}_{\mathbb{N}})}(C, \sigma^\Delta K) \\ \text{Hom}_{C(\mathcal{C}_{\mathbb{N}})}(\Sigma_*C, K) &= \text{Hom}_{C(\mathcal{C}_{\mathbb{N}})}(C, \Sigma^\Delta K) \end{aligned} \tag{1.8.1}$$

if C is a Cousin complex with respect to the filtration of K .

DEFINITION 1.8.3. Let X be a k -scheme. A *Witt residual complex* on X is a collection (K, \underline{p}, C, V) where $K \in C(\mathcal{C}_{\mathbb{N}})$ is a complex and

$$\underline{p} : i_*K \rightarrow K, C : \Sigma_*K \rightarrow K, V : \sigma_*i_*K \rightarrow K,$$

are morphisms of complexes such that:

- (a) K a residual complex,
- (b) the morphisms \underline{p}, C, V , satisfy the relations

$$V \circ \sigma_*i_*C = C \circ \Sigma_*\underline{p}, \quad \underline{p} \circ i_*V = V \circ \sigma_*i_*\underline{p},$$

and the adjoints of \underline{p}, C and V under (1.8.1)

$${}^a\underline{p} : K \xrightarrow{\cong} i^\Delta K, {}^aC : K \xrightarrow{\cong} \Sigma^\Delta K, {}^aV : K \xrightarrow{\cong} i^\Delta \sigma^\Delta K$$

are isomorphisms of complexes.

A morphism between Witt residual complexes is a morphism in $C(\mathcal{C}_{\mathbb{N}})$ which is compatible with \underline{p}, C, V in the obvious sense.

Remark 1.8.4. One should memorise the relations in (b) above as $VC = C\underline{p}$ and $V\underline{p} = \underline{p}V$.

Remark 1.8.5. It follows from Lemma 1.7.10, that if (K, \underline{p}, C, V) is a Witt residual complex on X , then for all $q \in \mathbb{Z}$ the systems $(\{K_n^q\}_{n \geq 1}, \underline{p}^q, C^q, V^q)$ are Witt dualizing systems. Furthermore, if Z^\bullet is the filtration of \bar{K}_1 , then

$$K_n^q = \bigoplus_{x \in Z^q \setminus Z^{q+1}} i_{W_n(x)*} J_n(x),$$

where $i_{W_n(x)} : \text{Spec } W_n \mathcal{O}_{X,x} \hookrightarrow W_n X$ is the inclusion and $J_n(x)$ is an injective hull of $k(x)$ in $W_n \mathcal{O}_{X,x}$.

1.8.6. By extending the notions from 1.7.2 and Corollary 1.7.6 term by term to $C(\mathcal{C}_{\mathbb{N}})$ we obtain: For any finite-type morphism $f : X \rightarrow Y$ between k -schemes a functor f^Δ of residual complexes in $C(\mathcal{C}_{\mathbb{N},Y})$ with associated filtration Z^\bullet to residual complexes in $C(\mathcal{C}_{\mathbb{N},X})$ with filtration $f^\Delta Z^\bullet$. For $g : Y \rightarrow Z$ another morphism a canonical isomorphism $c_{f,g} : (g \circ f)^\Delta \cong f^\Delta g^\Delta$ of natural transformations on the category of residual complexes on $C(\mathcal{C}_{\mathbb{N},Z})$. For $f : (X, \Phi) \rightarrow (Y, \Psi)$ a morphism in $\text{Sch}_{k,*}$ (see Definition 1.7.4) a natural transformation $\text{Tr}_f : f_\Phi f^\Delta \rightarrow \underline{\Gamma}_\Psi$ on the category of residual complexes with fixed filtration in $C(\mathcal{C}_{\mathbb{N},Y})$. And these data satisfies the compatibilities from 1.7.2 and Corollary 1.7.6.

1.8.7. Let $f : X \rightarrow Y$ be a finite-type morphism between k -schemes. Let K be a Witt residual complex. We use (1.8.1) to define morphisms of complexes

$$\underline{p} : i_*f^\Delta K \rightarrow f^\Delta K, \quad C : \Sigma_*f^\Delta K \rightarrow f^\Delta K, \quad V : \sigma_*i_*f^\Delta K \rightarrow f^\Delta K,$$

as adjoints of the compositions

$$f^\Delta K \xrightarrow{f^\Delta({}^a\underline{p}_K), \cong} f^\Delta i^\Delta K \cong i^\Delta f^\Delta K$$

$$f^\Delta K \xrightarrow{f^\Delta({}^aC_K), \cong} f^\Delta \Sigma^\Delta K \cong \Sigma^\Delta f^\Delta K,$$

$$f^\Delta K \xrightarrow{f^\Delta({}^aV_K), \simeq} f^\Delta i^\Delta \sigma^\Delta K \cong i^\Delta \sigma^\Delta f^\Delta K.$$

Here \underline{p}_K , C_K and V_K denote the corresponding morphisms for K and the isomorphisms on the right are induced by the isomorphisms $c_{W_n f, i_n, Y}$ etc. from 1.7.2, (1).

PROPOSITION 1.8.8. *Let $f : X \rightarrow Y$ be a finite-type morphism between k -schemes and K a Witt residual complex on Y . Then the system*

$$(f^\Delta K, \underline{p}, C, V)$$

defined above is a Witt residual complex on X . Furthermore, if $g : Y \rightarrow Z$ is another finite-type morphism of k -schemes, then the isomorphism $c_{f,g} : (g \circ f)^\Delta K \cong f^\Delta g^\Delta K$ from 1.8.6 defines an isomorphism of Witt residual complexes, which is compatible with triple compositions.

If $f : X \rightarrow Y$ is étale, then $f^ K$ is isomorphic to $f^\Delta K$. If $f : X \rightarrow Y$ and $g : Y \rightarrow Z$ are étale, then the isomorphism $(gf)^\Delta \cong f^\Delta g^\Delta$ is induced by the isomorphisms $(gf)^* \cong f^* g^*$.*

Proof. First notice that if K and L are two residual complexes, then their associated filtrations are the same iff their associated codimension functions (see 1.7.1) are the same. Since the codimension functions of the K_n 's are the same by assumption we obtain (using the formula of [Con00, (3.2.4)]) for any $x \in |X| = |W_n X|$

$$\begin{aligned} d_{(W_n f)^\Delta K_n}(x) &= d_{K_n}(W_n f(x)) - \text{trdeg}(k(x)/k(W_n f(x))) \\ &= d_{K_1}(f(x)) - \text{trdeg}(k(x)/k(f(x))) \\ &= d_{f^\Delta K_1}(x). \end{aligned}$$

Thus the $(W_n f)^\Delta K_n$ are residual complexes and all have the same associated filtration. The condition on the adjoints of \underline{p} , C and V holds by definition. It remains to check the relations $VC = C\underline{p}$ and $V\underline{p} = \underline{p}V$, which are equivalent to ${}^aV {}^aC = {}^aC {}^a\underline{p}$ and ${}^aV {}^a\underline{p} = {}^a\underline{p} {}^aV$. To prove the first equality consider the following diagram

$$\begin{array}{ccccc} f^\Delta K & \xrightarrow{{}^a\underline{p}_K} & f^\Delta i^\Delta K & \xrightarrow{\simeq} & i^\Delta f^\Delta K \\ {}^aC_K \downarrow & & \downarrow {}^aC_K & & \downarrow {}^aC_K \\ f^\Delta \Sigma^\Delta K & \xrightarrow{{}^aV_K} & f^\Delta i^\Delta \Sigma^\Delta K & \xrightarrow{\simeq} & i^\Delta f^\Delta (\Sigma)^\Delta K \\ \simeq \downarrow & & \downarrow \simeq & & \downarrow \simeq \\ \Sigma^\Delta f^\Delta K & \xrightarrow{{}^aV_K} & \Sigma^\Delta f^\Delta i^\Delta \sigma^\Delta K & \xrightarrow{\simeq} & i^\Delta \Sigma^\Delta f^\Delta K. \end{array}$$

Here all arrows labeled by \simeq are induced by compositions of the isomorphisms $c_{f,g}$ and their inverses with f, g appropriate and also in the two arrows labeled by aV_K there is such an isomorphism involved. (We also need the relation (1.2.2).) The square in the upper left commutes since K is a Witt residual complex, all other squares commute because of the functoriality of the $c_{f,g}$'s

and their compatibility with compositions. Thus going around the diagram from the top left corner to the lower right corner clockwise is the same as going around counter clockwise, which yields ${}^a C \underline{p} = {}^a V {}^a C$. The other relation is proved by drawing a similar diagram. Thus $f^\Delta K$ is a Witt residual complex. The second statement amounts to prove that $c_{g,f}$ is compatible with \underline{p} , C^n and V . This follows again from the functoriality of the $c_{(-,-)}$'s and their compatibility with compositions by drawing diagrams as above, which we omit. Finally, the statement about étale morphisms follows from 1.7.2, (4). \square

LEMMA 1.8.9. *Let $(X, \Phi) \rightarrow (Y, \Psi)$ be a morphism in $\text{Sch}_{k,*}$ and K a Witt residual complex on Y . Then the complex $f_\Phi f^\Delta K$, with $f_\Phi := f_* \circ \underline{\Gamma}_\Phi$, is a complex of Witt quasi-dualizing systems (see Definition 1.6.3) and each term $(f_\Phi f^\Delta K)_n$ is a bounded complex of quasi-coherent, flasque sheaves. Furthermore*

$$\text{Tr}_f : f_\Phi f^\Delta K \longrightarrow \underline{\Gamma}_\Psi K,$$

is a morphism of complexes of Witt quasi-dualizing systems, which is compatible with composition and étale base change, as in Corollary 1.7.6, (3) and (4), and it is functorial with respect to maps between Witt residual complexes with the same associated filtration (e.g. isomorphisms). Finally, if f is an open immersion and $\Psi = \Phi$, then Tr_f is the excision isomorphism as in Corollary 1.7.6, (5).

Proof. We define the maps \underline{p} , C and V on $f_\Phi f^\Delta K$ to be the following compositions

$$\underline{p} : i_* f_\Phi f^\Delta K \cong f_\Phi i_* f^\Delta K \xrightarrow{\underline{p}_{f^\Delta K}} f_\Phi f^\Delta K,$$

$$C : \Sigma_* f_\Phi f^\Delta K \cong f_\Phi \Sigma_* f^\Delta K \xrightarrow{C_{f^\Delta K}} f_\Phi f^\Delta K,$$

$$V : \sigma_* i_* f_\Phi f^\Delta K \cong f_\Phi \sigma_* i_* f^\Delta K \xrightarrow{V_{f^\Delta K}} f_\Phi f^\Delta K.$$

These maps obviously make $f_\Phi f^\Delta K$ into a complex of Witt quasi-dualizing systems. Also the $(f^\Delta K)_n$ are bounded complexes of quasi-coherent injective $W_n \mathcal{O}_X$ -modules (since the $(f^\Delta K)_n$ are residual complexes), thus all the $(f_\Phi f^\Delta K)_n$ are complexes of quasi-coherent and flasque sheaves. It remains to check, that the trace morphism commutes with \underline{p} , C^n and V . We will prove

$$(1.8.2) \quad \text{Tr}_f \circ \underline{p} = \underline{p} \circ i_*(\text{Tr}_f).$$

For this, we consider the following diagram:

$$\begin{array}{ccccc}
 & & i_* f_\Phi f^\Delta K & \xrightarrow{\cong} & f_\Phi i_* f^\Delta K \\
 & \swarrow \text{Tr}_f & \downarrow \underline{a}_p & & \downarrow \underline{a}_p \\
 i_* \underline{\Gamma}_\Psi K & & i_* f_\Phi f^\Delta i^\Delta K & \xrightarrow{\cong} & f_\Phi i_* f^\Delta i^\Delta K \\
 & \searrow \underline{a}_p & \downarrow \text{Tr}_f & & \downarrow \cong \\
 & & i_* \underline{\Gamma}_\Psi i^\Delta K & & f_\Phi i_* i^\Delta f^\Delta K \\
 & & \downarrow \text{Tr}_i & & \downarrow \text{Tr}_i \\
 & & \underline{\Gamma}_\Psi K & \xleftarrow{\text{Tr}_f} & f_\Phi f^\Delta K.
 \end{array}$$

The triangle in the upper left (which should be a square) commutes by the functoriality of Tr_f (Corollary 1.7.6, (1)), the upper square in the middle commutes by the functoriality of the isomorphism $i_* f_\Phi \cong f_\Phi i_*$ and the big square in the middle commutes by Corollary 1.7.6, (3). Thus going around the diagram from the upper left to the lower left corner clockwise, which is $\text{Tr}_f \circ \underline{a}_p$, is the same as going around counterclockwise, which is $\underline{a}_p \circ i_*(\text{Tr}_f)$. This gives (1.8.2). Similar diagrams prove the relations $\text{Tr}_f \circ C \cong C \circ \Sigma_*(\text{Tr}_f)$ and $V \circ \sigma_* i_*(\text{Tr}_f) \cong \text{Tr}_f \circ V$. The compatibility of Tr_f with composition and étale base change follows directly from Corollary 1.7.6, (3) and (4). The functoriality statement follows from the corresponding statement in 1.7.6, (1). If f is an open immersion, then so is $W_n f$ for all n and hence the last statement follows from 1.7.6, (5). \square

1.9. THE DUALIZING FUNCTOR.

1.9.1. Let X be a k -scheme and K a Witt residual complex on X . If M is a complex of *quasi-coherent* de Rham-Witt systems on X , then by Proposition 1.6.7 and Remark 1.8.5, we have de Rham-Witt systems $\mathcal{H}\text{om}(M^i, K^j) \in \text{dRW}_X$ for all i, j . Thus we obtain a complex of de Rham-Witt systems on X , $\mathcal{H}\text{om}(M, K) \in C(\text{dRW}_X)$ which is defined by

$$\mathcal{H}\text{om}^q(M, K) := \prod_{i \in \mathbb{Z}} \mathcal{H}\text{om}(M^i, K^{i+q}),$$

$$d_{\mathcal{H}\text{om}}^q((f^i)) := (d_K^{i+q} \circ f^i + (-1)^{q+1} f^{i+1} \circ d_M^i).$$

In this way we clearly obtain a functor

$$D_K : C(\text{dRW}_{X,\text{qc}})^\circ \rightarrow C(\text{dRW}_X), \quad M \mapsto \mathcal{H}\text{om}(M, K),$$

It restricts to a functor $D_K : C(\text{dRW}_{X,c})^\circ \rightarrow C(\text{dRW}_{X,\text{qc}})$. Since K_n is a bounded complex of injective $W_n \mathcal{O}_X$ -modules for each $n \geq 1$, $\mathcal{H}\text{om}(M, K)$ is acyclic, whenever M is ([Har66, II, Lem. 3.1]). Thus D_K preserves quasi-isomorphisms and therefore induces functors

$$D_K : D(\text{dRW}_{X,\text{qc}})^\circ \rightarrow D(\text{dRW}_X), \quad D_K : D(\text{dRW}_{X,c})^\circ \rightarrow D(\text{dRW}_{X,\text{qc}}).$$

A morphism $K \rightarrow L$ of Witt residual complexes clearly induces a natural transformation of functors

$$(1.9.1) \quad D_K \rightarrow D_L.$$

NOTATION 1.9.2. Let X be a k -scheme with structure map $\rho_X : X \rightarrow \text{Spec } k$. Then we denote by K_X the Witt residual complex $\rho_X^\Delta W_\bullet \omega$ and write

$$D_X = D_{K_X} = \mathcal{H}om(-, K_X).$$

Notice that K_X is concentrated in degree 0, since $W_\bullet \omega$ is.

Remark 1.9.3. Let X be a scheme, Φ a family of supports on X and K a Witt residual complex. Then we have the following equalities on $C(\mathcal{C}_{\mathbb{N}, \text{qc}})$

$$\Gamma_\Phi D_K(-) = \Gamma_\Phi \mathcal{H}om(-, K) = \mathcal{H}om(-, \Gamma_\Phi K).$$

Therefore, if M is a complex of quasi-coherent de Rham-Witt systems on X , then $\mathcal{H}om(M, \Gamma_\Phi K)$ inherits the structure of a complex of de Rham-Witt systems from $\Gamma_\Phi D_K(M)$. Furthermore $\Gamma_\Phi(K)_n$ is a complex of injectives (by Remark 1.7.5),

$$\mathcal{H}om(-, \Gamma_\Phi K) : D(\text{dRW}_{X, \text{qc}})^o \rightarrow D(\text{dRW}_X).$$

PROPOSITION 1.9.4. Let $f : (X, \Phi) \rightarrow (Y, \Psi)$ be a morphism in $\text{Sch}_{k, *}$, K a Witt residual complex on Y and M a bounded above complex of quasi-coherent de Rham-Witt systems on X . Let

$$\vartheta_f : f_\Phi D_{f^* K}(M) \rightarrow \Gamma_\Psi D_K(f_* M)$$

be the composition

$$f_\Phi \mathcal{H}om(M, f^\Delta K) \xrightarrow{\text{nat.}} \mathcal{H}om(f_* M, f_\Phi f^\Delta K) \xrightarrow{\text{Tr}_f} \mathcal{H}om(f_* M, \Gamma_\Psi K).$$

Then ϑ_f is a morphism of complexes of de Rham-Witt systems on Y and it has the following properties:

- (1) It defines a natural transformation between the bifunctors $f_\Phi \mathcal{H}om(-, f^\Delta -)$ and $\mathcal{H}om(f_* -, \Gamma_\Psi(-))$ defined on the product of the category $C^-(\text{dRW}_{X, \text{qc}})$ with the category of Witt residual complexes with the same associated filtration.
- (2) It is compatible with composition, i.e. if $g : (Y, \Psi) \rightarrow (Z, \Xi)$ is another morphism in $\text{Sch}_{k, *}$, then $\vartheta_{gf} \cong \vartheta_g \circ g_*(\vartheta_f) \circ c_{f, g}$.
- (3) It is compatible with étale base change as in Corollary 1.7.6, (4).
- (4) Let $j : U \hookrightarrow X$ be an open immersion, such that Φ is contained in U , then $\vartheta_j : j_\Phi \mathcal{H}om(-, j^\Delta K) \xrightarrow{\cong} \mathcal{H}om(j_*(-), \Gamma_\Phi K)$ is the excision isomorphism (cf. Corollary 1.7.6, (5)).

Proof. We have to show, that ϑ_f commutes with $\pi, F, d, V, \underline{p}$. Recall from 1.6.6, how the de Rham-Witt system structure on $\mathcal{H}om(M, Q)$ is defined, where M is a de Rham-Witt - and Q a Witt dualizing system. For example the map

π is uniquely determined by the property that it makes the following diagram commutative:

$$\begin{array}{ccc} \mathcal{H}om(M, Q) & \xrightarrow{\circ \underline{p}_M} & \mathcal{H}om(i_* M, Q) \\ \downarrow \pi & & \uparrow \underline{p}_Q \\ i_* \mathcal{H}om(M, Q) & \xrightarrow{\epsilon_i} & \mathcal{H}om(i_* M, i_* Q). \end{array}$$

The maps F, d, V, \underline{p} are defined uniquely by making similar diagrams commutative. Using these diagrams, the second part of 1.6.5 and Lemma 1.8.9 it is straightforward to check the compatibility of ϑ_f with the de Rham-Witt system structure. (Notice, that ϑ_f factors over $\mathcal{H}om(f_* M, f_\Phi f^\Delta K)$, which in general will not be a de Rham-Witt system, since $f_\Phi f^\Delta K$ will not be a complex of Witt dualizing systems. But it is a complex of Witt quasi-dualizing systems, which is sufficient to conclude.) Thus ϑ_f gives a morphism of complexes of de Rham-Witt systems. The functoriality of ϑ_f in M is clear and in K follows from the functoriality of $\text{Tr}_f : f_\Phi f^\Delta \rightarrow \text{id}$ (see Lemma 1.8.9). The properties (2)-(4) follow from the corresponding properties of Tr_f , see Lemma 1.8.9. □

1.10. EKEDAHL’S RESULTS.

THEOREM 1.10.1 ([Eke84, II, Thm 2.2 and III, Prop. 2.4]). *Let X be a smooth k -scheme of pure dimension N . Then the morphism induced by multiplication*

$$W_\bullet \Omega_X \xrightarrow{\cong} \mathcal{H}om(W_\bullet \Omega_X, W_\bullet \omega_X), \quad \alpha \mapsto (\beta \mapsto \alpha\beta).$$

is an isomorphism of de Rham-Witt systems. Furthermore

$$\mathcal{E}xt_{W_n \mathcal{O}_X}^i(W_n \Omega_X, W_n \omega_X) = 0, \quad \text{for all } i, n \geq 1.$$

Proof. This is all due to Ekedahl. Let us just point out that $W_\bullet \omega_X$ sits in degree N , hence the multiplication map is a graded morphism; and that $C^n d = 0$ gives the equality

$$C^n(xdy) = (-1)^{q+1} C^n(d(x)y), \quad \text{for all } x \in W_n \Omega_X^q, y \in W_n \Omega_X^{N-q}$$

and this together with the sign α introduced in the definition of d in 1.6.6 (which is missing in [Eke84]) gives the compatibility with d . □

LEMMA 1.10.2 ([BER10, Prop. 8.4, (ii)]). *Let X' be a smooth W_n -scheme and denote by X its reduction modulo p . Let $\varphi : W_n \Omega_X \xrightarrow{\cong} \sigma_* \mathcal{H}_{\text{DR}}^\bullet(X'/W_n)$, with $\sigma = W_n(F_{\text{Spec } k})$, be the canonical isomorphism of graded W_n -algebras from [IRS3, III, (1.5)]. Then φ is the unique morphism, which makes the following diagram commutative*

$$\begin{array}{ccc} W_{n+1} \Omega_{X'/W_n} & \xrightarrow{F^n} & Z \Omega_{X'/W_n} \\ \downarrow & & \downarrow \\ W_n \Omega_X & \xrightarrow{\varphi} & \mathcal{H}_{\text{DR}}^\bullet(X'/W_n). \end{array}$$

Here $W_{n+1}\Omega_{X'/W_n}$ is the relative de Rham-Witt complex defined in [LZ04] and the vertical maps are the natural surjections. In particular we have

$$(1.10.1) \quad \varphi\left(\sum_{i=0}^{n-1} V^i([a_i])\right) = \sum_{i=0}^{n-1} F^n V^i([\tilde{a}_i]) = \tilde{a}_0^{p^n} + p\tilde{a}_1^{p^{n-1}} + \dots + p^{n-1}\tilde{a}_{n-1}^p$$

and

$$(1.10.2) \quad \varphi\left(\sum_{i=0}^{n-1} dV^i([a_i])\right) = \sum_{i=0}^{n-1} F^n dV^i([\tilde{a}_i]) = \tilde{a}_0^{p^n-1} d\tilde{a}_0 + \tilde{a}_1^{p^{n-1}-1} d\tilde{a}_1 + \dots + \tilde{a}_{n-1}^{p-1} d\tilde{a}_{n-1},$$

where the \tilde{a}_i are any liftings of $a_i \in \mathcal{O}_X$ to $\mathcal{O}_{X'}$.

THEOREM 1.10.3 (Ekedahl). *Let X be a smooth k -scheme of pure dimension N with structure map $\rho_X : X \rightarrow \text{Spec } k$. Then $(W_n\rho_X)^!W_n \in D_c^b(W_nX)$ is concentrated in degree $-N$, for all $n \geq 1$ and there is a quasi-isomorphism of complexes of Witt dualizing systems (see Example 1.6.4, (1) and (2))*

$$\tau : W_\bullet\omega_X(N)[N] \xrightarrow{\text{qis}} K_X = \rho_X^\Delta W_\bullet\omega,$$

such that

- (1) The map $\tau_1 : \Omega_X^N[N] \xrightarrow{\text{qis}} \rho_X^\Delta k$ is the classical (ungraded) quasi-isomorphism, i.e. it is the composition of the natural quasi-isomorphism $\Omega_X^N[N] \rightarrow E_{Z^\bullet[N]}(\Omega_X^N[N])$ with the inverse of 1.7.2, (2). Here $Z^\bullet[N]$ is the shifted codimension filtration of X and $E_{Z^\bullet[N]}$ the Cousin functor.
- (2) τ is compatible with étale pullback, i.e. if $f : U \rightarrow X$ is étale, then the following diagram commutes

$$\begin{array}{ccc} f^*W_\bullet\omega_X(N)[N] & \xrightarrow{\tau_X} & f^*K_X = f^*\rho_X^\Delta W_\bullet\omega \\ \downarrow \simeq & & \downarrow \simeq \\ W_\bullet\omega_U(N)[N] & \xrightarrow{\tau_U} & K_U = \rho_U^\Delta W_\bullet\omega, \end{array}$$

where $\rho_U : U \rightarrow \text{Spec } k$ is the structure morphism of U , the vertical isomorphism on the right is induced by $f^*\rho_X^\Delta \cong f^*\rho_X^\Delta \cong \rho_U^\Delta$ and the isomorphism on the left is induced by $(W_n f)^*W_n\Omega_X^N = W_n\Omega_U^N$, $n \geq 1$ (see [Ill79, I, Prop. 1.14]).

Proof. We do not write the shift (N) in the following (the statement about the grading being obvious). In [Eke84, I,Thm. 4.1] it is proven that one has an isomorphism $\tau_n : W_n\Omega_X^N \xrightarrow{\simeq} (W_n\rho_X)^!W_n[-N]$ in $D_c^b(W_nX)$. (We give a sketch of the construction in case X admits a smooth lift over W_n in the remark below.) Let Z^\bullet be the filtration by codimension on W_nX . For C any complex let $E_{Z^\bullet}(C)$ be the associated Cousin complex. Then E_{Z^\bullet} induces an equivalence from the category of dualizing complexes with filtration Z^\bullet in $D_c^b(W_nX)$ and the category of residual complexes with associated filtration Z^\bullet

on $W_n X$ with quasi inverse the natural localization functor (see [Con00, Lem. 3.2.1]). Thus we obtain an isomorphism of residual complexes

$$E_{Z^\bullet}(W_n \Omega_X^N) \xrightarrow{\cong} E_{Z^\bullet}((W_n \rho_X)^! W_n[-N]) \cong (W_n \rho_X)^\Delta W_n[-N].$$

By Lemma 1.5.9 the natural map $W_n \Omega_X^N \rightarrow E_{Z^\bullet}(W_n \Omega_X^N)$ is a resolution, for all $n \geq 1$ (by [Har66, IV, Prop. 2.6.]). Therefore we obtain a quasi-isomorphism (by abuse of notation again denoted by τ_n)

$$\tau_n : W_n \Omega_X^N[N] \xrightarrow{qis} (W_n \rho)^\Delta W_n.$$

It follows from [Eke84, I, Lem 3.3 and II, Lem. 2.1], that the τ_n , $n \geq 1$, are compatible with \underline{p}, C^n, V and hence induce a morphism of complexes of Witt dualizing systems as in the statement. The property (1) is proved at the end of the proof of [Eke84, I, Thm. 4.1, p. 198] and (2) follows from the construction of τ_n in [Eke84, I, 2.] (see in particular the second paragraph on page 194). \square

Remark 1.10.4. Let us sketch for later purposes how the isomorphism $\tau_n : W_n \omega_X \rightarrow (W_n \rho_X)^! W_n[-N]$ is constructed in case X admits a smooth lifting $\rho_{X'} : X' \rightarrow \text{Spec } W_n$. For details see [Eke84, I, 2.]. Let

$$(1.10.3) \quad \varphi : W_n \Omega_X \xrightarrow{\cong} \sigma_*^n \mathcal{H}_{\text{DR}}^\bullet(X'/W_n)$$

be the canonical isomorphism of graded W_n -algebras of [IR83, III, (1.5)] (see Lemma 1.10.2). The composition $W_n \mathcal{O}_X \xrightarrow{\varphi^0} \sigma_*^n \mathcal{H}_{\text{DR}}^0(X'/W_n) \rightarrow \mathcal{O}_{X'}$ defines a finite morphism $\epsilon : X' \rightarrow W_n X$, which fits into a commutative diagram

$$\begin{array}{ccc} X' & \xrightarrow{\epsilon} & W_n X \\ \rho_{X'} \downarrow & & \downarrow W_n \rho_X \\ \text{Spec } W_n & \xrightarrow{\sigma^n} & \text{Spec } W_n. \end{array}$$

It follows that φ becomes an isomorphism $W_n \Omega_X \xrightarrow{\cong} \epsilon_* \mathcal{H}_{\text{DR}}^\bullet(X'/W_n)$. (Here we abuse the notation ϵ_* to indicate where the $W_n \mathcal{O}_X$ -module structure on the W_n -module $\mathcal{H}_{\text{DR}}^\bullet(X'/W_n)$ is coming from, which is of course not an $\mathcal{O}_{X'}$ -module.) Since $\rho_{X'}$ is smooth we have a canonical isomorphism $\tau_{X'} : \omega_{X'/W_n} \xrightarrow{\cong} \rho_{X'}^! W_n[-N]$. Now Ekedahl shows, that the composition

$$\begin{aligned} \epsilon_* \omega_{X'/W_n} &\xrightarrow{\tau_{X'}} \epsilon_* \rho_{X'}^! W_n[-N] \simeq \epsilon_* \rho_{X'}^! \sigma^{n!} W_n[-N] \\ &\simeq \epsilon_* \epsilon^! (W_n \rho_X)^! W_n[-N] \xrightarrow{\text{Tr}_\epsilon} (W_n \rho_X)^! W_n[-N] \end{aligned}$$

factors over $\epsilon_* \mathcal{H}_{\text{DR}}^N(X'/W_n)$. Then τ_n is defined as the composition

$$W_n \omega_X \xrightarrow{\varphi, \cong} \epsilon_* \mathcal{H}_{\text{DR}}^N(X'/W_n) \rightarrow (W_n \rho_X)^! W_n[-N].$$

COROLLARY 1.10.5. *Let X be a smooth k -scheme of pure dimension N . Denote by $E(W_\bullet \omega_X)$ the Cousin complex associated to $W_\bullet \omega_X$ with respect to the codimension filtration on X (cf. 1.5.8). Then the maps \underline{p}, C^n, V on $W_\bullet \omega_X$ induce morphisms of complexes on the system $\{E(W_n \omega_X)(N)[N]\}_n$; this defines a Witt residual complex $E(W_\bullet \omega_X)(N)[N]$, which is isomorphic as Witt*

residual complex to K_X (see Notation 1.9.2). This isomorphism is compatible with étale base change.

COROLLARY 1.10.6. *Let the notations be as above. Then there is an isomorphism in $C^b(\mathrm{dRW}_{X,\mathrm{qc}})$*

$$\mu : E(W_\bullet \Omega_X) \xrightarrow{\cong} D_X(W_\bullet \Omega_X)(-N)[-N],$$

given by the composition

$$E(W_\bullet \Omega_X) \xrightarrow{E(1.10.1), \cong} E(\mathcal{H}\mathrm{om}(W_\bullet \Omega_X, W_\bullet \omega_X)) \xrightarrow{\cong} \mathcal{H}\mathrm{om}(W_\bullet \Omega_X, E(W_\bullet \omega_X)) \xrightarrow{\text{Cor. 1.10.5}, \cong} \mathcal{H}\mathrm{om}(W_\bullet \Omega_X, K_X)(-N)[-N].$$

The map μ is compatible with étale base change.

1.11. THE TRACE MORPHISM FOR A REGULAR CLOSED IMMERSION.

1.11.1. *Local Cohomology.* Let $X = \mathrm{Spec} A$ be an affine Cohen-Macaulay scheme and $Z \subset X$ a closed subscheme of pure codimension c , defined by the ideal $I \subset A$. Let $t = t_1, \dots, t_c \in I$ be an A -regular sequence with $\sqrt{(t)} = \sqrt{I}$, here (t) denotes the ideal $(t_1, \dots, t_c) \subset A$. (After shrinking X such a sequence always exists.) We denote by $K^\bullet(t)$ the Koszul complex of the sequence t , i.e. $K^{-q}(t) = K_q(t) = \bigwedge^q A^c$, $q = 0, \dots, c$, and if $\{e_1, \dots, e_c\}$ is the standard basis of A^c and $e_{i_1, \dots, i_q} := e_{i_1} \wedge \dots \wedge e_{i_q}$, then the differential is given by

$$d_{K^\bullet}^{-q}(e_{i_1, \dots, i_q}) = d_q^{K^\bullet}(e_{i_1, \dots, i_q}) = \sum_{j=1}^q (-1)^{j+1} t_{i_j} e_{i_1, \dots, \widehat{i_j}, \dots, i_q}.$$

We define the complex

$$K^\bullet(t, M) := \mathrm{Hom}_A(K^{-\bullet}(t), M),$$

and denote its n -th cohomology by $H^n(t, M)$. The map

$$\mathrm{Hom}_A\left(\bigwedge^c A^c, M\right) \rightarrow M/(t)M, \quad \varphi \mapsto \varphi(e_{1, \dots, c})$$

induces a canonical isomorphism $H^c(t, M) \simeq M/(t)M$.

If t and t' are two sequences as above with $(t') \subset (t)$, then there exists a $c \times c$ -matrix T with coefficients in A such that $t' = Tt$ and T induces a morphism of complexes $K^\bullet(t') \rightarrow K^\bullet(t)$, which is the unique (up to homotopy) morphism lifting the natural map $A/(t') \rightarrow A/(t)$. Furthermore we observe that, for any pair of sequences t, t' as above there exists an $N \geq 0$ such that $(t^N) \subset (t')$, where t^N denotes the sequence t_1^N, \dots, t_c^N . Thus the sequences t form a directed set and $H^c(t, M) \rightarrow H^c(t', M)$, $(t') \subset (t)$, becomes a direct system. It follows from [SGA2, Exp. II, Prop. 5], that we have an isomorphism

$$(1.11.1) \quad \varinjlim_t M/(t)M = \varinjlim_t H^c(t, M) \cong H_Z^c(X, \widetilde{M}),$$

where the limit is over all A -regular sequences $t = t_1, \dots, t_c$ with $V((t)) = Z$ and \widetilde{M} is the sheaf associated to M . In fact it is enough to take the limit

over the powers $t^n = t_1^n, \dots, t_c^n$, $n \geq 1$, of just one regular sequence t with $V((t)) = Z$. We denote by

$$\begin{bmatrix} m \\ t \end{bmatrix}$$

the image of $m \in M$ under the composition

$$M \rightarrow M/(t)M \rightarrow H^c(t, M) \rightarrow H_Z^c(X, \widetilde{M}).$$

It is a consequence of the above explanations that we have the following properties:

- (1) Let t and t' be two sequences as above with $(t') \subset (t)$. Let T be a $c \times c$ -matrix with $t' = Tt$, then

$$\begin{bmatrix} \det(T) m \\ t' \end{bmatrix} = \begin{bmatrix} m \\ t \end{bmatrix}.$$

- (2)

$$\begin{bmatrix} m + m' \\ t \end{bmatrix} = \begin{bmatrix} m \\ t \end{bmatrix} + \begin{bmatrix} m' \\ t \end{bmatrix}, \quad \begin{bmatrix} t_i m \\ t \end{bmatrix} = 0 \quad \text{all } i.$$

- (3) If M is any A -module, then

$$H_Z^c(X, \mathcal{O}_X) \otimes_A M \xrightarrow{\cong} H_Z^c(X, \widetilde{M}), \quad \begin{bmatrix} a \\ t \end{bmatrix} \otimes m \mapsto \begin{bmatrix} am \\ t \end{bmatrix}$$

is an isomorphism.

Remark 1.11.2. Since for an A -regular sequence t as above $K^\bullet(t) \rightarrow A/(t)$ is a free resolution, we have an isomorphism

$$\mathrm{Ext}^n(A/(t), M) \simeq \mathrm{Hom}_A^\bullet(K^\bullet(t), M).$$

Notice that we also have an isomorphism

$$\mathrm{Hom}_A^\bullet(K^\bullet(t), M) \simeq K^\bullet(t, M),$$

which is multiplication with $(-1)^{n(n+1)/2}$ in degree n (see [Con00, (1.3.28)]). We obtain an isomorphism

$$\psi_{t,M} : \mathrm{Ext}^c(A/(t), M) \xrightarrow{\cong} H^c(t, M) = M/(t)M,$$

which has the sign $(-1)^{c(c+1)/2}$ in it. In particular under the composition

$$\mathrm{Ext}^c(A/(t), M) \xrightarrow{\psi_{t,M}} M/(t)M \rightarrow H_Z^c(X, \widetilde{M})$$

the class of a map $\varphi \in \mathrm{Hom}_A(\bigwedge^c A^c, M)$ is sent to

$$(-1)^{c(c+1)/2} \begin{bmatrix} \varphi(e_1, \dots, e_c) \\ t \end{bmatrix}.$$

Remark 1.11.3. If $X = \mathrm{Spec} A$ is a k -scheme and $t = t_1, \dots, t_c$ is a regular sequence of sections of \mathcal{O}_X , then for any n the sequence $[t] = [t_1], \dots, [t_c]$ of sections of $W_n \mathcal{O}_X$ is a regular sequence. Here $[t_i]$ denotes the Teichmüller lift of t_i . Indeed the sequence $[t]$ is regular iff its Koszul complex is a resolution

of $W_n A/([t])$ and thus the statement follows by induction from the short exact sequence

$$0 \rightarrow K_\bullet([t]^p, W_{n-1}A) \xrightarrow{V} K_\bullet([t], W_n A) \rightarrow K_\bullet(t, A) \rightarrow 0.$$

PROPOSITION 1.11.4. *Let $i : Z \hookrightarrow X$ be a closed immersion between smooth k -schemes of pure dimension N_Z, N_X and structure maps ρ_Z, ρ_X and let c be the codimension of Z in X , i.e. $c = N_X - N_Z$. Then we have the following isomorphism in $D_{qc}^b(W_n \mathcal{O}_X)$,*

$$(1.11.2) \quad R\Gamma_Z(W_n \omega_X) \cong \mathcal{H}_Z^c(W_n \omega_X)[-c]$$

Assume furthermore, that the ideal sheaf of Z is generated by a regular sequence $t = t_1, \dots, t_c$ of global sections of \mathcal{O}_X and define a morphism $\iota_{Z,n}$ by

$$\iota_{Z,n} : (W_n i)_* W_n \omega_Z \rightarrow \mathcal{H}_Z^c(W_n \omega_X), \quad \alpha \mapsto (-1)^c \begin{bmatrix} d[t] \tilde{\alpha} \\ [t] \end{bmatrix},$$

with $\tilde{\alpha} \in W_n \Omega_X^{N_Z}$ any lift of α , and $d[t] = d[t_1] \cdot \dots \cdot d[t_c]$. Then the following diagram in $D_{qc}^b(W_n \mathcal{O}_X)$ is commutative

$$\begin{array}{ccccc} (W_n i)_* (W_n \rho_Z)^\Delta W_n \xrightarrow{c_{W_n i, W_n \rho_X}} (W_n i)_* (W_n i)^\Delta (W_n \rho_X)^\Delta W_n & \xrightarrow{\text{Tr}_{W_n i}} & R\Gamma_Z(W_n \rho_X)^\Delta W_n & & \\ \tau_{Z,n} \uparrow \simeq & & \simeq \uparrow \tau_{X,n} & & \\ (W_n i)_* W_n \omega_Z[N_Z] & \xrightarrow{\iota_{Z,n}} & \mathcal{H}_Z^c(W_n \omega_X)[N_Z] & \xrightarrow{\simeq} & R\Gamma_Z(W_n \omega_X)[N_X] \end{array}$$

where $\text{Tr}_{W_n i}$ is the n -th level of the trace morphism from Lemma 1.8.9.

Proof. We write $i_n = W_n i$, $\rho_{X,n} = W_n \rho_X$ and $W_n \rho_Z = \rho_{Z,n}$. By (1.5.2) we have $\mathcal{H}_Z^i(W_n \omega_X) = 0$ for $i < c$. Furthermore $\mathcal{H}_Z^i(W_n \omega_X) = 0$ for $i > c$, by Čech considerations. (Locally the ideal of Z is generated by a regular sequence of length c and thus $U \setminus Z$ may locally be covered by c affine open subschemes.) This gives the isomorphism (1.11.2). To prove the commutativity of the diagram in the statement of the proposition we have to show that two elements in $\text{Hom}_{W_n \mathcal{O}_X}(i_n^* W_n \omega_Z, \mathcal{H}_Z^c(W_n \omega_X))$ are equal. This is a local question. We may thus assume, that the closed immersion i lifts to a closed immersion $i' : Z' \hookrightarrow X'$ between smooth W_n -schemes and that there exists a regular sequence $t' = t'_1, \dots, t'_c$ in $\mathcal{O}_{X'}$ which generates the ideal of Z' and reduces modulo p to t . As in Remark 1.10.4 we obtain a commutative diagram

$$\begin{array}{ccc} Z' & \xrightarrow{\epsilon_Z} & W_n Z \\ \downarrow i' & & \downarrow i_n \\ X' & \xrightarrow{\epsilon_X} & W_n X \\ \downarrow \rho_{X'} & & \downarrow \rho_{X,n} \\ \text{Spec } W_n & \xrightarrow{\sigma^n} & \text{Spec } W_n \end{array} \quad \begin{array}{l} \rho_{Z'} \curvearrowright \\ \rho_{Z,n} \curvearrowleft \end{array}$$

where we write σ^n instead of $W_n(F_{\text{Spec } k}^n)$.

Consider the following diagram, in which we set $\Lambda := W_n[-N_Z]$ and write $(-)^!$ instead of $(-)^{\Delta}$:

(1.11.3)

$$\begin{array}{ccccccc}
 i_{n*} \epsilon_{Z*} \omega_{Z'/W_n} & \xrightarrow{\cong} & i_{n*} \epsilon_{Z*} \rho_{Z'}^! \Lambda & \xrightarrow{\cong} & \epsilon_{X*} i'_* i'^! \rho_{X'}^! \Lambda & \xrightarrow{\text{Tr}_{i'}} & \epsilon_{X*} R\Gamma_Z \rho_{X'}^! \Lambda & \xrightarrow{\cong} & \epsilon_{X*} R\Gamma_Z \omega_{X'/W_n} [c] \\
 \downarrow & & \downarrow & & \downarrow & & \downarrow & & \downarrow \\
 i_{n*} \epsilon_{Z*} \mathcal{H}_{\text{DR}}^{N_Z}(Z') & \xrightarrow{\dots \exists} & i_{n*} \rho_{Z,n}^! \Lambda & \xrightarrow{\cong} & i_{n*} i_n^! \rho_{X,n}^! \Lambda & \xrightarrow{\text{Tr}_{i_n}} & R\Gamma_Z \rho_{X,n}^! \Lambda & \xleftarrow{\dots \exists} & \epsilon_{X*} R\Gamma_Z \mathcal{H}_{\text{DR}}^{N_X}(X') [c] \\
 \cong \uparrow & \nearrow \tau_{Z,n} & & & & & & \nwarrow \tau_{X,n} & \uparrow \cong \\
 i_{n*} W_n \omega_Z & & & & & & & & R\Gamma_Z(W_n \omega_X) [c] \\
 & & & & & & & & \uparrow \cong \\
 & & & & & & & & \mathcal{H}_Z^c(W_n \omega_X).
 \end{array}$$

We give some explanations: We have $\Lambda = \sigma^{n!} \Lambda$ (see Example 1.6.4, (1)) and the three vertical maps in the middle are the compositions

$$i_{n*} \epsilon_{Z*} \rho_{Z'}^! \Lambda \cong i_{n*} \epsilon_{Z*} \rho_{Z'}^! \sigma^{n!} \Lambda \cong i_{n*} \epsilon_{Z*} \epsilon_Z^! \rho_{Z,n}^! \Lambda \xrightarrow{\text{Tr}_{\epsilon_Z}} i_{n*} \rho_{Z,n}^! \Lambda,$$

$$\begin{aligned}
 \epsilon_{X*} i'_* i'^! \rho_{X'}^! \Lambda &\cong i_{n*} \epsilon_{Z*} i'^! \rho_{X'}^! \sigma^{n!} \Lambda \cong i_{n*} \epsilon_{Z*} i'^! \epsilon_X^! \rho_{X,n}^! \Lambda \\
 &\xrightarrow{\cong} i_{n*} \epsilon_{Z*} \epsilon_Z^! i_n^! \rho_{X,n}^! \Lambda \xrightarrow{\text{Tr}_{\epsilon_Z}} i_{n*} i_n^! \rho_{X,n}^! \Lambda,
 \end{aligned}$$

$$\epsilon_{X*} R\Gamma_Z \rho_{X'}^! \Lambda \cong R\Gamma_Z \epsilon_{X*} \rho_{X'}^! \sigma^{n!} \Lambda \cong R\Gamma_Z \epsilon_{X*} \epsilon_X^! \rho_{X,n}^! \Lambda \xrightarrow{\text{Tr}_{\epsilon_X}} R\Gamma_Z \rho_{X,n}^! \Lambda.$$

Now one easily checks that the two squares in the middle commute. In fact, to see that the first of the two middle squares commute, one only needs, that $c_{(f,g)} : (gf)^! \xrightarrow{\cong} f^! g^!$ is a natural transformation, which is compatible with triple compositions (see [Con00, p.139, (VAR1)]) and that $\text{Tr}_{\epsilon_Z} : \epsilon_{Z*} \epsilon_Z^! \rightarrow \text{id}$ is a natural transformation. For the commutativity of the second middle square we need the naturality of $\text{Tr}_{i'} : i'_* i'^! \rightarrow \text{id}$ and the formula (see [Con00, Lem 3.4.3, (TRA1)])

$$\begin{aligned}
 \text{Tr}_{i_n} \circ i_{n*} (\text{Tr}_{\epsilon_Z}) \circ (i_n \epsilon_Z)_* (c_{\epsilon_Z, i_n}) &= \text{Tr}_{i_n \epsilon_Z} \\
 &= \text{Tr}_{\epsilon_X i'} = \text{Tr}_{\epsilon_X} \circ \epsilon_{X*} (\text{Tr}_{i'}) \circ (\epsilon_X i')_* (c_{i', \epsilon_X}).
 \end{aligned}$$

That there exist two unique dotted arrows, which make the two outer diagrams commutative was proved by Ekedahl, see Remark 1.10.4. The two vertical maps in the two lower triangles are the isomorphisms from Lemma 1.10.2. Thus the two triangles commute by definition of $\tau_{Z,n}$ and $\tau_{X,n}$, see Remark 1.10.4. It follows that the whole diagram commutes.

Let σ_Z and σ_X be the following compositions

$$\sigma_Z : \epsilon_{Z*} \omega_{Z'/W_n} \rightarrow \epsilon_{Z*} \mathcal{H}_{\text{DR}}^{N_Z}(Z'/W_n) \xrightarrow{\cong} W_n \omega_Z,$$

$$\sigma_X : \epsilon_{X*} \mathcal{H}_Z^c(\omega_{X'/W_n}) \rightarrow \epsilon_{X*} \mathcal{H}_Z^c(\mathcal{H}_{\text{DR}}^{N_X}(X'/W_n)) \xrightarrow{\cong} \mathcal{H}_Z^c(W_n \omega_X).$$

We obtain maps

(1.11.4)

$$\begin{array}{ccc} & \text{Hom}(\epsilon_{X*}i_{n*}\omega_{Z'/W_n}, \epsilon_{X*}\mathcal{H}_Z^c(\omega_{X'/W_n})) & \\ & \downarrow \sigma_X & \\ \text{Hom}(i_{n*}W_n\omega_Z, \mathcal{H}_Z^c(W_n\omega_X)) & \xrightarrow{D(\sigma_Z)} & \text{Hom}(\epsilon_{X*}i_{n*}\omega_{Z'/W_n}, \mathcal{H}_Z^c(W_n\omega_X)), \end{array}$$

with the horizontal map being injective (since σ_Z is surjective). We denote by a the following composition

$$a := H^0(\tau_{X,n})^{-1} \circ H^0(\text{Tr}_{i_n} \circ c_{i_n, \rho_{X,n}} \circ \tau_{Z,n}) \in \text{Hom}(i_{n*}W_n\omega_Z, \mathcal{H}_Z^c(W_n\omega_X)).$$

The commutativity of the diagram in the statement of the proposition means, that a equals $\iota_{Z,n}$. Thus it is enough to show

$$(1.11.5) \quad D(\sigma_Z)(a) = D(\sigma_Z)(\iota_{Z,n}) \quad \text{in } \text{Hom}(\epsilon_{X*}i_{n*}\omega_{Z'/W_n}, \mathcal{H}_Z^c(W_n\omega_X)).$$

We define $a' \in \text{Hom}(i'_*\omega_{Z'/W_n}, \mathcal{H}_Z^c(\omega_{X'/W_n}))$ to be the following composition

$$a' : H^0 \left(i'_*\omega_{Z'/W_n} \simeq i'_*\rho_{Z'}^!\Lambda \simeq i'_*i'^!\rho_{X'}^!\Lambda \xrightarrow{\text{Tr}_{i'}} R\Gamma_Z(\rho_{X'}^!\Lambda) \simeq R\Gamma_Z(\omega_{X'/W_n})[c] \right).$$

Then diagram (1.11.3) says

$$(1.11.6) \quad D(\sigma_Z)(a) = \sigma_X(\epsilon_{X*}(a')).$$

We define $\iota_{Z'}$ by

$$\iota_{Z'} : i_{Z'*}\omega_{Z'/W_n} \rightarrow \mathcal{H}_Z^c(\omega_{X'/W_n}), \quad \beta \mapsto (-1)^c \begin{bmatrix} dt' \tilde{\beta} \\ t' \end{bmatrix},$$

with $\tilde{\beta} \in \Omega_{X'/W_n}^{N_Z}$ a lift of β . We have

$$(1.11.7) \quad D(\sigma_Z)(\iota_{Z,n}) = \sigma_X(\epsilon_{X*}(\iota_{Z'})).$$

Indeed, this follows from the concrete description of σ_X and σ_Z given by Lemma 1.10.2 and from the fact that by 1.11.1, 1) the following equality holds for all $\gamma \in W_{n+1}\Omega_{X'/W_n}^{N_Z}$

$$\begin{bmatrix} dt' F^n(\gamma) \\ t' \end{bmatrix} = \begin{bmatrix} t'^{p^n-1} dt' F^n(\gamma) \\ t'^{p^n} \end{bmatrix} = F^n \left(\begin{bmatrix} d[t']\gamma \\ [t'] \end{bmatrix} \right),$$

where we set $t'^{p^n-1} := t_1'^{p^n-1} \dots t_c'^{p^n-1}$. By (1.11.6) and (1.11.7) we are thus reduced to show

$$\iota_{Z'} = a' \quad \text{in } \text{Hom}(i'_*\omega_{Z'/W_n}, \mathcal{H}_Z^c(\omega_{X'/W_n})),$$

which is well-known (see e.g. [CR09, Lemma A.2.2]). □

2. PUSHFORWARD AND PULLBACK FOR HODGE-WITT COHOMOLOGY WITH SUPPORTS

In this section all schemes are assumed to be quasi-projective over k . We fix a k -scheme S .

2.1. RELATIVE HODGE-WITT COHOMOLOGY WITH SUPPORTS.

DEFINITION 2.1.1. We denote by (Sm_*/S) and (Sm^*/S) the following two categories: Both have as objects pairs (X, Φ) with X an S -scheme, which is smooth and *quasi-projective* over k (for short we will say X is a smooth k -scheme over S) and Φ a family of supports on X and the morphisms are given by

$$\mathrm{Hom}_{Sm_*}((X, \Phi), (Y, \Psi)) = \{f \in \mathrm{Hom}_S(X, Y) \mid f|_{\Phi} \text{ is proper and } f(\Phi) \subset \Psi\}$$

and

$$\mathrm{Hom}_{Sm^*}((X, \Phi), (Y, \Psi)) = \{f \in \mathrm{Hom}_S(X, Y) \mid f^{-1}(\Psi) \subset \Phi\}.$$

If X is a smooth k -scheme over S and $Z \subset X$ a closed subset, we write $(X, Z) = (X, \Phi_Z)$ and $X = (X, X) = (X, \Phi_X) \in \mathrm{Obj}(Sm_*) = \mathrm{Obj}(Sm^*)$ (see Definition 1.4.1 for the notation).

We will say that a morphism $f : (X, \Phi) \rightarrow (Y, \Psi)$ in (Sm_*/S) or (Sm^*/S) is étale, flat, smooth, etc. if the corresponding property holds for the underlying morphism of schemes $X \rightarrow Y$. We will say that a diagram

$$\begin{array}{ccc} (X', \Phi') & \longrightarrow & (Y', \Psi') \\ \downarrow & & \downarrow \\ (X, \Phi) & \longrightarrow & (Y, \Psi) \end{array}$$

is cartesian, if the underlying diagram of schemes is cartesian.

2.1.2. For $(X, \Phi) \in \mathrm{obj}(Sm_*/S) = \mathrm{obj}(Sm^*/S)$, with structure map $a : X \rightarrow S$ we denote by $\mathcal{H}((X, \Phi)/S)$ the de Rham-Witt system

$$\mathcal{H}((X, \Phi)/S) := \bigoplus_{i \geq 0} R^i a_{\Phi} W_{\bullet} \Omega_X \in \mathrm{dRW}_S$$

and its level n part by $\mathcal{H}_n((X, \Phi)/S)$. We denote by $\hat{\mathcal{H}}((X, \Phi)/S)$ the de Rham-Witt module

$$\hat{\mathcal{H}}((X, \Phi)/S) := \bigoplus_{i \geq 0} R^i \hat{a}_{\Phi} W_{\bullet} \Omega_X \in \widehat{\mathrm{dRW}}_S.$$

We write

$$\mathcal{H}^{i,q}((X, \Phi)/S) := R^i a_{\Phi} W_{\bullet} \Omega_X^q, \quad \hat{\mathcal{H}}^{i,q}((X, \Phi)/S) := R^i \hat{a}_{\Phi} W_{\bullet} \Omega_X^q.$$

By definition we have:

- (1) If $Z \subset X$ is closed then, $\hat{\mathcal{H}}((X, Z)/S) = \bigoplus R^i a_Z W \Omega_X$. In particular $\hat{\mathcal{H}}((X, Z)/X) = \bigoplus \mathcal{H}_Z^i(W \Omega_X)$, in particular $\hat{\mathcal{H}}(X/X) = W \Omega_X$.
- (2) Let $U \subset S$ be open and set $X_U := X \times_S U$ and $\Phi_U = \Phi \cap X_U$. Then

$$\mathcal{H}((X, \Phi)/S)|_U = \mathcal{H}((X_U, \Phi_U)/U), \quad \hat{\mathcal{H}}((X, \Phi)/S)|_U = \hat{\mathcal{H}}((X_U, \Phi_U)/U).$$

2.2. PULLBACK.

DEFINITION 2.2.1. Let $f : (X, \Phi) \rightarrow (Y, \Psi)$ be a morphism in (Sm^*/S) . Then applying $R\underline{\Gamma}_\Psi$ to the functoriality morphism $W_\bullet\Omega_Y \rightarrow Rf_*W_\bullet\Omega_X$ in $D^+(\mathrm{dRW}_Y)$ and composing it with the natural map $R\underline{\Gamma}_\Psi Rf_* = Rf_*R\underline{\Gamma}_{f^{-1}(\Psi)} \rightarrow Rf_*R\underline{\Gamma}_\Phi$ yields a morphism in $D^+(\mathrm{dRW}_Y)$

$$(2.2.1) \quad R\underline{\Gamma}_\Psi(W_\bullet\Omega_Y) \rightarrow R\underline{\Gamma}_\Psi Rf_*W_\bullet\Omega_X = Rf_*R\underline{\Gamma}_{f^{-1}(\Psi)}W_\bullet\Omega_X \rightarrow Rf_*R\underline{\Gamma}_\Phi W_\bullet\Omega_X.$$

Let $a : Y \rightarrow S$ be the structure morphism. Then we define

$$f^* := \bigoplus_i H^i(Ra_*(2.2.1)) : \mathcal{H}((Y, \Psi)/S) \rightarrow \mathcal{H}((X, \Phi)/S) \quad \text{in } \mathrm{dRW}_S$$

and

$$f^* := \bigoplus_i H^i(R\underline{\lim} Ra_*(2.2.1)) : \hat{\mathcal{H}}((Y, \Psi)/S) \rightarrow \hat{\mathcal{H}}((X, \Phi)/S) \quad \text{in } \widehat{\mathrm{dRW}}_S.$$

We call these morphisms the *pullback morphisms*. Notice that by definition f^* always factors as

$$f^* : \mathcal{H}((Y, \Psi)/S) \rightarrow \mathcal{H}((X, f^{-1}(\Psi)/S)) \xrightarrow{\mathrm{nat.}} \mathcal{H}((X, \Phi)/S),$$

same with $\hat{\mathcal{H}}$.

PROPOSITION 2.2.2. *The assignments*

$$\mathcal{H} : (Sm^*/S)^\circ \rightarrow \mathrm{dRW}_S, \quad (X, \Phi) \mapsto \mathcal{H}((X, \Phi)/S)$$

and

$$\hat{\mathcal{H}} : (Sm^*/S)^\circ \rightarrow \widehat{\mathrm{dRW}}_S, \quad (X, \Phi) \mapsto \hat{\mathcal{H}}((X, \Phi)/S)$$

define functors, where we set $\mathcal{H}(f) = f^*$ and $\hat{\mathcal{H}}(f) = f^*$, for a morphism $f : (X, \Phi) \rightarrow (Y, \Psi)$ in (Sm^*/S) .

Furthermore, if $U \subset S$ is open and $f_U : (X_U, \Phi_U) \rightarrow (Y_U, \Psi_U)$ is the pullback of $f : (X, \Phi) \rightarrow (Y, \Psi)$ over U , then $f_U^* = (f^*)|_U$ on $\mathcal{H}((Y_U, \Psi_U)/U) = \mathcal{H}((Y, \Psi)/S)|_U$ (resp. on $\hat{\mathcal{H}}$).

Proof. This is all straightforward. □

2.3. PUSHFORWARD. Let X be a smooth k -scheme of pure dimension N with structure map $\rho : X \rightarrow \mathrm{Spec} k$. Recall that we set $K_X = \rho_X^\Delta W_\bullet\omega$ and that this is isomorphic to $E(W_\bullet\omega_X)(N)[N]$, where E denotes the Cousin complex with respect to the codimension filtration (see Corollary 1.10.5). Furthermore we have the dualizing functor

$$D_X(-) = \mathcal{H}om(-, K_X)$$

(see 1.6.6), which we may view as a functor from $C(\mathrm{dRW}_{X,\mathrm{qc}})^\circ$ to $C(\mathrm{dRW}_X)$ or from $D(\mathrm{dRW}_{X,\mathrm{qc}})^\circ$ to $D(\mathrm{dRW}_X)$.

For any morphism $f : (X, \Phi) \rightarrow (Y, \Psi)$ in (Sm_*/S) the composition of the isomorphism $K_X \simeq f^\Delta K_Y$ (see Proposition 1.8.8) with ϑ_f from Proposition 1.9.4, defines a natural transformation of functors on $C(\mathrm{dRW}_{X,\mathrm{qc}})^\circ$

$$(2.3.1) \quad \theta_f : f_\Phi D_X(-) \rightarrow \underline{\Gamma}_\Psi D_Y(f_*(-)).$$

The map θ_f is compatible with compositions and étale base change. Furthermore, recall that the multiplication pairing $W_\bullet\Omega_X^i \otimes_W W_\bullet\Omega_X^j \rightarrow W_\bullet\Omega_X^{i+j}$ induces an isomorphism in $C^b(\mathrm{dRW}_{X,\mathrm{qc}})$ (see Corollary 1.10.6)

$$(2.3.2) \quad \mu_X : E(W_\bullet\Omega_X) \xrightarrow{\sim} D_X(W_\bullet\Omega_X)(-N)[-N].$$

The map μ_X is compatible with étale base change.

DEFINITION 2.3.1. Let $f : (X, \Phi) \rightarrow (Y, \Psi)$ be a morphism in (Sm_*/S) . We assume that X and Y are of pure dimension N_X and N_Y . Consider the following composition of morphisms of complexes of de Rham-Witt systems on Y

$$(2.3.3) \quad f_\Phi E(W_\bullet\Omega_X)(N_X)[N_X] \xrightarrow{\mu_X} f_\Phi D_X(W_\bullet\Omega_X) \xrightarrow{\theta_f} \underline{\Gamma}_\Psi D_Y(f_*W_\bullet\Omega_X) \\ \xrightarrow{D(f^*)} \underline{\Gamma}_\Psi D_Y(W_\bullet\Omega_Y) \xrightarrow{\mu_Y^{-1}} \underline{\Gamma}_\Psi E(W_\bullet\Omega_Y)(N_Y)[N_Y].$$

By Lemma 1.5.9 this induces a morphism in $D^b(\mathrm{dRW}_Y)$

$$(2.3.4) \quad Rf_\Phi W_\bullet\Omega_X \rightarrow R\underline{\Gamma}_\Psi W_\bullet\Omega_Y(-r)[-r],$$

where $r = N_X - N_Y$ is the relative dimension.

Let $a : Y \rightarrow S$ be the structure morphism. Then we define

$$f_* := \bigoplus_i H^i(Ra_*(2.3.4)) : \mathcal{H}((X, \Phi)/S) \rightarrow \mathcal{H}((Y, \Psi)/S)(-r) \quad \text{in } \mathrm{dRW}_S$$

and

$$f_* := \bigoplus_i H^i(R\underline{\lim} Ra_*(2.3.4)) : \hat{\mathcal{H}}((X, \Phi)/S) \rightarrow \hat{\mathcal{H}}((Y, \Psi)/S)(-r) \quad \text{in } \widehat{\mathrm{dRW}}_S.$$

We extend these definitions additively to all morphisms $f : (X, \Phi) \rightarrow (Y, \Psi)$ in (Sm_*/S) (where X and Y don't need to be pure dimensional). We call these morphisms the *push-forward morphisms*. Notice, that since $f|_\Phi$ is proper, $f(\Phi)$ is a family of supports on Y and f_* always factors as

$$f_* : \mathcal{H}((X, \Phi)/S) \rightarrow \mathcal{H}((Y, f(\Phi))/S)(-r) \xrightarrow{\mathrm{nat.}} \mathcal{H}((Y, \Psi)/S)(-r),$$

same with $\hat{\mathcal{H}}$.

Remark 2.3.2. Without supports and for a *proper* morphism between smooth k -schemes a similar push-forward was already defined in [Gro85, II, Def. 1.2.1]. (Although the verification, that it is compatible with F, V, d and π was omitted.)

PROPOSITION 2.3.3. *The assignments*

$$\mathcal{H} : (Sm_*/S) \rightarrow \mathrm{dRW}_S, \quad (X, \Phi) \mapsto \mathcal{H}((X, \Phi)/S)$$

and

$$\hat{\mathcal{H}} : (Sm_*/S) \rightarrow \widehat{\mathrm{dRW}}_S, \quad (X, \Phi) \mapsto \hat{\mathcal{H}}((X, \Phi)/S)$$

define functors, where we set $\mathcal{H}(f) = f_*$ and $\hat{\mathcal{H}}(f) = f_*$, for $f : (X, \Phi) \rightarrow (Y, \Psi)$ a morphism in (Sm_*/S) .

Furthermore: (1) If $U \subset S$ is open and $f_U : (X_U, \Phi_U) \rightarrow (Y_U, \Psi_U)$ is the pullback of $f : (X, \Phi) \rightarrow (Y, \Psi)$ over U , then $f_{U*} = (f_*)|_U$ on $\mathcal{H}((Y_U, \Psi_U)/U) = \mathcal{H}((Y, \Psi)/S)|_U$ (resp. on $\hat{\mathcal{H}}$). (2) If $j : (U, \Phi) \hookrightarrow (X, \Phi)$ is an open immersion in (Sm_*/S) , then j_* is the excision isomorphism.

Proof. This follows from the corresponding properties of (2.3.3), which follow from Proposition 1.9.4 and Corollary 1.10.6. \square

For later use we record:

PROPOSITION 2.3.4 ([Gro85, II, Prop. 4.2.9]). *Let $f : X \rightarrow Y$ be a finite morphism between two connected smooth k -schemes. We may view f as a morphism in (Sm_*/Y) or as a morphism in (Sm^*/Y) . Then the composition*

$$f_* \circ f^* : W_\bullet \Omega_Y \rightarrow W_\bullet \Omega_Y$$

equals multiplication with the degree of f .

2.4. COMPATIBILITY OF PUSHFORWARD AND PULLBACK.

PROPOSITION 2.4.1 (Gros). *Let $i : Z \hookrightarrow (X, Z)$ be a closed immersion of pure codimension c in (Sm_*/S) . Then*

$$R\Gamma_Z W_\bullet \Omega_X[c] \cong \mathcal{H}_Z^c(W_\bullet \Omega_X) \quad \text{in } D^+(\mathrm{dRW}_X).$$

Suppose further the ideal sheaf of Z in \mathcal{O}_X is generated by a regular sequence $t = t_1, \dots, t_c$ of global sections of X , then the projection of $i_ : \mathcal{H}(Z/X) \rightarrow \mathcal{H}((X, Z)/X)$ to the n -th level is given by*

$$(2.4.1) \quad W_n(i)_* W_n \Omega_Z \rightarrow \mathcal{H}_Z^c(W_n \Omega_X)(c), \quad \alpha \mapsto (-1)^c \begin{bmatrix} d[t] \tilde{\alpha} \\ [t] \end{bmatrix},$$

where we set $[t] := [t_1], \dots, [t_c]$ and $d[t] := d[t_1] \cdots d[t_c] \in W_n \Omega_X^c$, with $[t_i] \in W_n \mathcal{O}_X$ the Teichmüller lift of t_i , and $\tilde{\alpha}$ is any lift of $\alpha \in W_n(i)_ W_n \Omega_Z$ to $W_n \Omega_X$.*

The above proposition is proved in [Gro85, II, 3.4], but since the proof uses a result by Ekedahl, which is referred to as work in progress and which we could not find in the literature we reprove the above proposition (using Proposition 1.11.4, instead of a comparison of Ekedahl’s trace with Berthelot’s trace in crystalline cohomology as Gros does).

Proof. The first statement is proven as in (1.11.2). It remains to prove the explicit description (2.4.1). Let $\rho_{Z,n} : W_n Z \rightarrow \mathrm{Spec} W_n$ and $\rho_{X,n} : W_n X \rightarrow \mathrm{Spec} W_n$ be the structure maps, $i_n := W_n(i) : W_n Z \hookrightarrow W_n X$ the closed immersion (for this proof we don’t use the i_n defined in 1.1) and write $\dim X = N_X$, $\dim Z = N_Z$. By Definition 2.3.1 the projection to the n -th level of i_* is given

by the following composition in $D_{\text{qc}}^b(\mathcal{C}_{X,n})$

$$\begin{aligned} i_{n*}W_n\Omega_Z(N_Z)[N_Z] &\xrightarrow{\mu_{Z,n}, \simeq} i_{n*}R\mathcal{H}om(W_n\Omega_Z, K_{Z,n}) \\ &\xrightarrow{\theta_{i_n}} R\Gamma_Z R\mathcal{H}om(i_{n*}W_n\Omega_Z, K_{X,n}) \\ &\xrightarrow{D(i_n^*)} R\mathcal{H}om(W_n\Omega_X, \Gamma_Z K_{X,n}) \\ &\xrightarrow{\mu_{X,n}^{-1}, \simeq} R\Gamma_Z(W_n\Omega_X)(N_X)[N_X]. \end{aligned}$$

Here $\mu_{Z,n}$, $\mu_{X,n}$ and θ_{i_n} are images in the derived category of the projections to the n -th level of the corresponding maps from 2.3. We denote by $\iota_{Z,n}$ the following morphism

$$\iota_{Z,n} : (W_n i)_* W_n \omega_Z \longrightarrow \mathcal{H}_Z^c(W_n \omega_X)(c), \quad \alpha \mapsto (-1)^c \begin{bmatrix} d[t]\tilde{\alpha} \\ [t] \end{bmatrix},$$

with $\tilde{\alpha} \in W_n \Omega_X^{N_Z}$ any lift of α . Then it follows from Proposition 1.11.4, that in the derived category $(i_*)_n$ equals

$$\begin{aligned} i_{n*}W_n\Omega_Z(N_Z)[N_Z] &\xrightarrow{\text{multipl.}} \mathcal{H}om(i_{n*}W_n\Omega_Z, i_{n*}W_n\omega_Z(N_Z)[N_Z]) \\ &\xrightarrow{\iota_{Z,n}} \mathcal{H}om(i_{n*}W_n\Omega_Z, \mathcal{H}_Z^c(W_n\omega_X)(N_X)[N_Z]) \\ &\xrightarrow{D(i_n^*)} \mathcal{H}om(W_n\Omega_X, \mathcal{H}_Z^c(W_n\omega_X)(N_X)[N_Z]) \\ &\xrightarrow{\simeq, (*)} \mathcal{H}_Z^c(W_n\Omega_X)(N_X)[-c][N_X] \\ &\xrightarrow{\simeq} R\Gamma_Z(W_n\Omega_X)(N_X)[N_X]. \end{aligned}$$

Here the isomorphism $(*)$ is the inverse of

$$\mathcal{H}_Z^c(W_n\Omega_X)(N_X) \xrightarrow{\simeq} \mathcal{H}om(W_n\Omega_X, \mathcal{H}_Z^c(W_n\omega_X)), \quad \begin{bmatrix} \alpha \\ t^n \end{bmatrix} \mapsto \left(\beta \mapsto \begin{bmatrix} \alpha\beta \\ t^n \end{bmatrix} \right).$$

It is straightforward to check, that taking $H^{-N_Z}(-)$ of this composition gives (2.4.1) and hence the claim. \square

COROLLARY 2.4.2. *Let $i : Y \hookrightarrow X$ be a closed immersion between smooth affine k -schemes and assume that the ideal of Y in X is generated by a regular sequence t_1, \dots, t_c . Let $Z \subset Y$ be a closed subset which is equal to the vanishing set $V(f_1, \dots, f_i)$ of global sections $f_1, \dots, f_i \in \Gamma(Y, \mathcal{O}_Y)$. Denote by $i_Z : (Y, Z) \hookrightarrow (X, Z)$ the morphism in $(\text{Sm}_*/\text{Spec } k)$ induced by i . Then the projection to the n -th level of i_{Z*} is given by*

$$i_{Z*,n} : H_Z^i(Y, W_n\Omega_Y) \rightarrow H_Z^{i+c}(X, W_n\Omega_X)(c), \quad \begin{bmatrix} \alpha \\ [f] \end{bmatrix} \mapsto (-1)^c \begin{bmatrix} d[t]\tilde{\alpha} \\ [t], [f] \end{bmatrix},$$

where $\alpha \in \Gamma(Y, W_n\Omega_Y)$, $\tilde{\alpha} \in \Gamma(X, W_n\Omega_X)$ is some lift, $[f] = [f_1], \dots, [f_i]$ and $[t], d[t]$ as in Lemma 2.4.1.

Proof. Choose $\tilde{f}_1, \dots, \tilde{f}_i \in \Gamma(X, \mathcal{O}_X)$ lifts of f_1, \dots, f_i and define the set $\tilde{Z} \subset X$ as the vanishing locus of these \tilde{f}_i 's. Denote by $i_{Y/X}$ the closed immersion

$Y \hookrightarrow (X, Y)$ viewed as a morphism in (Sm_*/X) . Then it follows from the first part of Proposition 2.4.1 that $i_{Z^*,n}$ equals

$$H_{\mathbb{Z}}^i(X, i_{Y/X^*,n}) : H_{\mathbb{Z}}^i(Y, W_n\Omega_Y) \cong H_{\mathbb{Z}}^i(X, i_*W_n\Omega_Y) \longrightarrow H_{\mathbb{Z}}^i(X, \mathcal{H}_Y^c(W_n\Omega_X)) \cong H_{\mathbb{Z}}^{i+c}(X, W_n\Omega_X)[c].$$

Now denote by $[t] = [t_1, \dots, t_c]$ and $[\tilde{f}] = [\tilde{f}_1, \dots, \tilde{f}_i]$ the Teichmüller lifts of the sequences t and f to $\Gamma(X, W_n\mathcal{O}_X)$ and by $\mathcal{K}([t])$ and $\mathcal{K}([\tilde{f}])$ the corresponding sheafified Koszul complexes. We define a morphism of complexes

$$\varphi : W_n\Omega_X[0] \rightarrow \mathcal{H}om(\mathcal{K}^{-\bullet}([t]), W_n\Omega_X(c))[c], \quad \alpha \mapsto (e_1 \wedge \dots \wedge e_c \mapsto (-1)^c d[t]\alpha)$$

where e_1, \dots, e_c is the standard basis of $(W_n\mathcal{O}_X)^c$. Then the second part of Proposition 2.4.1 maybe rephrased by saying that the diagram

$$\begin{array}{ccc} W_n\Omega_X & \xrightarrow{\varphi} & \mathcal{H}om(\mathcal{K}^{-\bullet}([t]), W_n\Omega_X(c))[c] \\ i^* \downarrow & & \downarrow (1.11.1) \\ i_*W_n\Omega_Y & \xrightarrow{i_{Y/X^*}} & \mathcal{H}_Y^c(W_n\Omega_X)(c) \end{array}$$

commutes. This yields the following commutative diagram of complexes (in which the lower square is concentrated in degree 0):

$$\begin{array}{ccc} \mathcal{H}om(\mathcal{K}^{-\bullet}([\tilde{f}]), W_n\Omega_X)[i] & \xrightarrow{\varphi} & \mathcal{H}om(\mathcal{K}^{-\bullet}([\tilde{f}]), \mathcal{H}om(\mathcal{K}^{-\bullet}([t]), W_n\Omega_X(c))[c])[i] \\ (1.11.1) \downarrow & & \downarrow (1.11.1) \\ H_{\mathbb{Z}}^i(X, W_n\Omega_X) & \xrightarrow{\varphi} & H_{\mathbb{Z}}^i(X, \mathcal{H}om(\mathcal{K}^{-\bullet}([t]), W_n\Omega_X(c))[c]) \\ i^* \downarrow & & \downarrow (1.11.1) \\ H_{\mathbb{Z}}^i(X, i_*W_n\Omega_Y) & \xrightarrow{i_{Y/X^*}} & H_{\mathbb{Z}}^i(X, \mathcal{H}_Y^c(W_n\Omega_X)(c)). \end{array}$$

Now the claim follows from $\mathcal{K}([t]) \otimes_{W_n\mathcal{O}_X} \mathcal{K}([\tilde{f}]) = \mathcal{K}([t], [\tilde{f}])$ and the commutativity of the following diagram, in which we denote $W_n\Omega_X(c)$ simply by Λ

$$\begin{array}{ccc} \mathcal{H}om(\mathcal{K}^{-\bullet}([\tilde{f}]), \mathcal{H}om(\mathcal{K}^{-\bullet}([t]), \Lambda)[c])[i] & \xrightarrow{\cong} & \mathcal{H}om(\mathcal{K}^{-\bullet}([t], [\tilde{f}]), \Lambda)[c+i] \\ \downarrow & & \downarrow \\ H_{\mathbb{Z}}^i(X, \mathcal{H}_Y^c(\Lambda)) & \xrightarrow{\cong} & H_{\mathbb{Z}}^{i+c}(X, \Lambda). \end{array}$$

□

LEMMA 2.4.3. *Let Y be a smooth k -scheme, $f : X = \mathbb{P}_Y^r \rightarrow Y$ the projection and Ψ a family of supports on Y . We denote by $f_{\Psi/S} : (X, f^{-1}(\Psi)) \rightarrow (Y, \Psi)$ the morphism in (Sm_*/S) induced by f and by*

$$f_{\Psi/S^*} : \mathcal{H}((X, f^{-1}(\Psi))/S) \rightarrow \mathcal{H}((Y, \Psi)/S)(-r)$$

the corresponding push-forward (same for $\hat{\mathcal{H}}$). Let $a : Y \rightarrow S$ be the structure morphism. Then:

- (1) The higher direct images $R^i f_* W_\bullet \Omega_X$ vanish for $i \geq r + 1$. Therefore there is a natural morphism in $D^+(\mathrm{dRW}_Y)$

$$e : Rf_* W_\bullet \Omega_X \rightarrow R^r f_* W_\bullet \Omega_X[-r]$$

and the push-forward $f_{\Psi/S*}$ factors as

$$\bigoplus_i H^i(Ra_{\Psi} Rf_* W_\bullet \Omega_X) \xrightarrow{e} \bigoplus_i R^{i-r} a_{\Psi}(R^r f_* W_\bullet \Omega_X) \xrightarrow{f_{Y/Y*}} \bigoplus_i Ra_{\Psi}^{i-r} W_\bullet \Omega_Y(-r).$$

Similar for $\hat{\mathcal{H}}$ (replace a_{Ψ} by \hat{a}_{Ψ} in the above formula).

- (2) The push-forward $f_{Y/Y*} : \mathcal{H}(X/Y) \rightarrow \mathcal{H}(Y/Y)(-r)$ induces an isomorphism

$$f_{Y/Y*} : R^r f_* W_\bullet \Omega_X \xrightarrow{\simeq} W_\bullet \Omega_Y(-r),$$

with inverse induced by the push-forward $i_* : \mathcal{H}_\bullet(Y/Y)(-r) \rightarrow \mathcal{H}_\bullet(X/Y)$, where $i : Y \rightarrow X$ is any section of f .

Proof. First of all observe that it suffices to prove the statements on each finite level n separately.

- (1) By [Gro85, I, Cor 4.1.12] we have an isomorphism in $D(W_n \Omega_Y - \mathrm{dga})$ (with $W_n \Omega_Y - \mathrm{dga}$ the category of differential graded $W_n \Omega_Y$ -modules)

$$(2.4.2) \quad Rf_* W_n \Omega_X \simeq \bigoplus_{i=0}^r W_n \Omega_Y(-i)[-i].$$

This gives the first statement of (1) and in particular an isomorphism in $D(W_n \Omega_Y - \mathrm{dga})$

$$(2.4.3) \quad Rf_* W_n \Omega_X \simeq \bigoplus_{i=0}^r R^i f_* W_n \Omega_X[-i]$$

and the morphism e from (1) is the projection to the r -th summand. On the other hand $(f_{Y/Y*})_n$ is induced by a derived category morphism (2.3.4)

$$(2.4.4) \quad Rf_* W_n \Omega_X \rightarrow W_n \Omega_Y(-r)[-r].$$

Since the composition

$$R^i f_* W_n \Omega_X[-i] \xrightarrow{\text{via (2.4.3)}} Rf_* W_n \Omega_X \xrightarrow{(2.4.4)} W_n \Omega_Y(-r)[-r]$$

is an element in $\mathrm{Ext}^{i-r}(R^i f_* W_n \Omega_X, W_n \Omega_Y(-r))$, this composition is zero for all $i < r$. Thus the morphism (2.4.4) factors over e . This implies the second statement of (1).

- (2) It suffices to check that $f_{Y/Y*}$ induces an isomorphism; i_* will then automatically be the inverse for any section i of f . (Notice that by (2.4.2) we know $R^r f_* W_\bullet \Omega_X \simeq W_\bullet \Omega_Y(-r)$ but we don't know that this isomorphism is induced by $f_{Y/Y*}$.) To this end, let R be the Cartier-Dieudonné-Raynaud ring of k , see [IR83, I]. (R is a graded (non-commutative) W -algebra generated by symbols F and V in degree 0 and d in degree 1, satisfying the obvious relations). Set $R_n := R/V^n R + dV^n R$. We denote by $D^b(Y, R)$ the bounded derived category of R -modules. Any de Rham-Witt module is in particular an R -module. Thus

we may view $R^i \hat{f}_* W_\bullet \Omega_X = R^i f_* W \Omega_X$, $i \geq 1$, as an object in $D^b(Y, R)$. In $D^b(Y, R)$ we have the following isomorphisms

$$R_n \otimes_R^L R^i f_* W \Omega_X \simeq R_n \otimes_R^L W \Omega_Y(-i) \simeq W_n \Omega_Y(-i) \simeq R^i f_* W_n \Omega_X,$$

where the first and the last isomorphism follow from [Gro85, I, Thm. 4.1.11, Cor. 4.1.12] and the second isomorphism is [IR83, II, Thm (1.2)]. Since f_{Y/Y^*} is clearly a morphism of R -modules it follows that the projection of f_{Y/Y^*} to the n -th level, identifies with

$$R_n \otimes_R^L f_{Y/Y^*} : R_n \otimes_R^L R^r f_* W \Omega_X \rightarrow R_n \otimes_R^L W \Omega_Y(-r).$$

We thus have to show that this is an isomorphism for all n . Now Ekedahl's version of Nakayama's Lemma [Eke85, I, Prop. 1.1, Cor. 1.1.3] (also [Ill83, Prop. 2.3.7]) implies that it is in fact enough to prove that the projection of f_{Y/Y^*} to level 1 - in the following simply denoted by f_* - is an isomorphism, i.e. we have to show (for all $q \geq 0$)

$$f_* : R^r f_* \Omega_X^q \xrightarrow{\simeq} \Omega_Y^{q-r}.$$

In fact, f_* is given by H^r of the following morphism in $D_c^b(\mathcal{O}_Y)$

$$\begin{aligned} Rf_* \Omega_X^q &\simeq Rf_* R\mathcal{H}om(\Omega_X^{N_X-q}, \omega_X) \simeq R\mathcal{H}om(Rf_* \Omega_X^{N_X-q}, \omega_Y)[-r] \\ &\xrightarrow{D(f^*)[-r]} R\mathcal{H}om(\Omega_Y^{N_X-q}, \omega_Y)[-r] \simeq \Omega_Y^{q-r}[-r], \end{aligned}$$

with $N_X = \dim X$. Thus it is enough to show that $H^r(D(f^*)[-r]) = H^0(D(f^*))$ is an isomorphism. This follows from the well-known formula

$$Rf_* \Omega_X^{N_X-q} \simeq \bigoplus_{i=0}^r \Omega_Y^{N_X-q-i}[-i],$$

which might be proved using the Künneth decomposition $\Omega_X^q = \bigoplus_{i+j=q} f^* \Omega_Y^i \otimes_{\mathcal{O}_X} \Omega_{X/Y}^j$ and [SGA7(2), Exp. XI, Thm 1.1, (ii)]. \square

PROPOSITION 2.4.4. *Let*

$$\begin{array}{ccc} (X', \Phi') & \xrightarrow{f'} & (Y', \Psi') \\ g_X \downarrow & & \downarrow g_Y \\ (X, \Phi) & \xrightarrow{f} & (Y, \Psi) \end{array}$$

be a cartesian diagram, with f, f' morphisms in (Sm_*/S) and g_X, g_Y morphisms in (Sm^*/S) . Assume that one of the following conditions is satisfied

- (1) g_Y is flat, or
- (2) g_Y is a closed immersion and f is transversal to g_Y (i.e. $f'^* \mathcal{N}_{Y'/Y} = \mathcal{N}_{X'/X}$).

Then the following diagram commutes

$$(2.4.5) \quad \begin{array}{ccc} \mathcal{H}((X', \Phi')/S) & \xrightarrow{f'_*} & \mathcal{H}((Y', \Psi')/S) \\ g_X^* \uparrow & & \uparrow g_Y^* \\ \mathcal{H}((X, \Phi)/S) & \xrightarrow{f_*} & \mathcal{H}((Y, \Psi)/S) \end{array}$$

and also with \mathcal{H} replaced by $\hat{\mathcal{H}}$.

In [Gro85, II, Prop. 2.3.2 and p.49] a version of this Proposition (in the derived category) is proved in the case without supports with f a closed immersion and g_Y transversal to f . (This case is also covered here by factoring g_Y as a closed immersion followed by a smooth projection.)

Proof. We distinguish two cases.

1. case: f is a closed immersion and g_Y is either flat or transversal to f . Since Φ and Φ' are also families of supports on Y and Y' the push-forward f_* factors over $\mathcal{H}((Y, \Phi)/S)$ and f'_* factors over $\mathcal{H}((Y', \Phi')/S)$. Hence we may assume $\Psi = \Phi$ and $\Psi' = \Phi'$. Furthermore by Definition 2.2.1, the pull-back g_X^* factors over $\mathcal{H}((X', g_X^{-1}(\Phi))/S)$ and g_Y^* factors over $\mathcal{H}((Y', g_Y^{-1}(\Phi))/S)$. Hence we may assume that in the diagram (2.4.5) we have $\Psi = \Phi$, $\Phi' = g_X^{-1}(\Phi)$, $\Psi' = g_Y^{-1}(\Phi)$. We may further assume, that X, X', Y, Y' are equidimensional and set $c := \dim Y - \dim X = \dim Y' - \dim X'$ and $h := f \circ g_X = g_Y \circ f'$. We consider the following diagram in $D^+(\mathrm{dRW}_Y)$

$$(2.4.6) \quad \begin{array}{ccc} Rh_* W_\bullet \Omega_{X'} & \xrightarrow{Rg_{Y'}(f'_*)} & Rg_{Y'} R\Gamma_{X'} W_\bullet \Omega_{Y'}(c)[c] \\ Rf_*(g_X^*) \uparrow & & \uparrow R\Gamma_X(g_Y^*) \\ Rf_* W_\bullet \Omega_X & \xrightarrow{f_*} & R\Gamma_X W_\bullet \Omega_Y(c)[c], \end{array}$$

where the horizontal (resp. vertical) morphisms are induced by (2.3.4) (resp. (2.2.1)). By our assumption on the supports we obtain diagram (2.4.5) if we apply $\oplus H^i(Ra_\Phi(-))$ (resp. $\oplus H^i(R\varprojlim Ra_\Phi(-))$) to diagram (2.4.6), where $a : Y \rightarrow S$ is the structure morphism. Thus it suffices to show that (2.4.6) commutes. By Proposition 2.4.1 the diagram (2.4.6) is isomorphic to the outer square of the following diagram

$$(2.4.7) \quad \begin{array}{ccc} Rh_* W_\bullet \Omega_{X'} & \xrightarrow{Rg_{Y'}(f'_*)} & Rg_{Y'} \mathcal{H}_{X'}^c(W_\bullet \Omega_{Y'})(c) \\ \text{nat.} \uparrow & & \uparrow \text{nat.} \\ h_* W_\bullet \Omega_{X'} & \xrightarrow{g_{Y'}(f'_*)} & g_{Y'} \mathcal{H}_{X'}^c(W_\bullet \Omega_{Y'})(c) \\ g_X^* \uparrow & & \uparrow g_Y^* \\ f_* W_\bullet \Omega_X & \xrightarrow{f_*} & \mathcal{H}_X^c(W_\bullet \Omega_Y)(c). \end{array}$$

The upper square obviously commutes. Thus it suffices to check that the lower square of sheaves of de Rham-Witt systems commutes. This may be checked locally and we may hence assume that the ideal sheaf of X in \mathcal{O}_Y is generated by a regular sequence $t = t_1, \dots, t_c$. By our assumptions on g_Y , the sequence $g_Y^*t = g_Y^*t_1, \dots, g_Y^*t_c$ is again a regular sequence generating the ideal sheaf of X' in $\mathcal{O}_{Y'}$. Now the statement follows from the explicit formula (2.4.1).

2. case: $X = \mathbb{P}_Y^r$ and $f : X \rightarrow Y$ is the projection map. In this situation we also have $X' = \mathbb{P}_{Y'}^r$, and $f' : X' \rightarrow Y'$ is the projection map. The morphisms f and f' factor over $(X, f^{-1}(\Psi)) \rightarrow (Y, \Psi)$ and $(X', f'^{-1}(\Psi')) \rightarrow (Y', \Psi')$, which are morphisms in (Sm_*/S) . Similar to the first case we conclude that it is enough to consider the case $\Phi = f^{-1}(\Psi)$, $\Psi' = g_Y^{-1}(\Psi)$ and $\Phi' = f'^{-1}(\Psi')$. Now Lemma 2.4.3, (1) reduces us to show the commutativity of the following diagram (with $h = f \circ g_X = g_Y \circ f'$)

$$(2.4.8) \quad \begin{array}{ccc} R^r h_* W_\bullet \Omega_{X'} & \xrightarrow{H^0(Rg_{Y*}(f'_*))} & g_{Y*} W_\bullet \Omega_{Y'}(-r) \\ g_X^* \uparrow & & \uparrow g_Y^* \\ R^r f_* W_\bullet \Omega_X & \xrightarrow{H^0(f_*)} & W_\bullet \Omega_Y(-r). \end{array}$$

But now let $i : Y \rightarrow X$ be a section and $i' : Y' \rightarrow X'$ its pull-back, viewed as morphisms over Y . We obtain push-forwards

$$i_* : W_\bullet \Omega_Y(-r) \rightarrow R^r f_* W_\bullet \Omega_X, \quad i'_* : g_{Y*} W_\bullet \Omega_{Y'}(-r) \rightarrow R^r h_* W_\bullet \Omega_{X'}.$$

And since g_X is transversal to i we may apply case 1. to obtain

$$(g_Y^* f_* - f'_* g_X^*) \circ i_* = g_Y^* f_* i_* - f'_* i'_* g_Y^* = 0.$$

Thus the commutativity of (2.4.8) follows from Lemma 2.4.3, (2), which tells us that i_* is an isomorphism (with inverse f_*).

Proof in the general case. Since $f : (X, \Phi) \rightarrow (Y, \Psi)$ is quasi-projective we may factor it as follows

$$(X, \Phi) \xrightarrow{i} (U, \Phi) \xrightarrow{j} (P, \Phi) \xrightarrow{\pi} (Y, \Psi),$$

where $P = \mathbb{P}_Y^r$, $i : (X, \Phi) \rightarrow (U, \Phi)$ is a closed immersion, $j : (U, \Phi) \rightarrow (P, \Phi)$ is an open immersion and $\pi : (P, \Phi) \rightarrow (Y, \Psi)$ is the projection. Notice that we use the properness of $f|_\Phi$ to conclude that Φ is also a family of supports on U and on P and that all the maps above are in (Sm_*/S) . By base change we

obtain a diagram

$$\begin{array}{ccccccc}
 & & & & f' & & \\
 & & & & \curvearrowright & & \\
 (X', \Phi') & \xrightarrow{i'} & (U_{Y'}, \Phi') & \xrightarrow{j'} & (P_{Y'}, \Phi') & \xrightarrow{\pi'} & (Y', \Psi') \\
 g_X \downarrow & & g_U \downarrow & & g_P \downarrow & & g_Y \downarrow \\
 (X, \Phi) & \xrightarrow{i} & (U_Y, \Phi) & \xrightarrow{j} & (P, \Phi) & \xrightarrow{\pi} & (Y, \Psi) \\
 & & & & \curvearrowleft & & \\
 & & & & f & &
 \end{array}$$

with the horizontal maps in (Sm_*/S) and the vertical maps in (Sm^*/S) . Notice that if g_Y satisfies condition (1) or (2) from the statement of the Proposition, then so does g_U . By the functoriality of the push-forward the commutativity of the square (2.4.5) follows from the first two cases and the fact that j_* and j'_* are just the excision isomorphisms by Proposition 2.3.3, (2). This proves the proposition. \square

3. CORRESPONDENCES AND HODGE-WITT COHOMOLOGY

In this section all schemes are assumed to be quasi-projective over k . We fix a k -scheme S .

3.1. EXTERIOR PRODUCT.

3.1.1. Let X and Y be two k -schemes and M and N two complexes of de Rham-Witt systems on X and Y respectively. Then for all $n \geq 1$ we set

$$M_n \boxtimes N_n := p_1^{-1} M_n \otimes_{\mathbb{Z}} p_2^{-1} N_n \quad \text{as a complex of abelian sheaves on } X \times Y,$$

where $p_1 : X \times Y \rightarrow X$ and $p_2 : X \times Y \rightarrow Y$ are the two projection morphisms. It is obvious, that the maps $\pi_M \otimes \pi_N$ make the family $\{M_n \boxtimes N_n\}_{n \geq 1}$ into a pro-complex of abelian sheaves on $X \times Y$, which we denote by

$$(3.1.1) \quad M \boxtimes N.$$

3.1.2. *Godement resolution.* Let X be a k -scheme. For a sheaf of abelian groups \mathcal{A} on X we denote by $G(\mathcal{A})$ its Godement resolution. Then there is a natural way to equip the family $\{G(W_n \Omega_X)\}_{n \geq 1}$ with the structure of a complex of de Rham-Witt systems on X , which we denote by $G(W_\bullet \Omega_X)$. It follows from the exactness of $G(-)$ and the surjectivity of the transition maps $W_{n+1} \Omega_X \rightarrow W_n \Omega_X$, that each term $G^q(W_\bullet \Omega_X)$ is a flasque de Rham-Witt system (in the sense of 1.5.3). Thus the natural augmentation map $W_\bullet \Omega_X \rightarrow G(W_\bullet \Omega_X)$, makes $G(W_\bullet \Omega_X)$ a flasque resolution of de Rham-Witt systems of X .

3.1.3. *Exterior product for Godement resolutions.* Let X and Y be two k -schemes with families of supports. It follows from the general construction in [God73, II, §6. 1], that there is a morphism of *pro-complexes of abelian sheaves on $X \times Y$*

$$G(W_\bullet\Omega_X) \boxtimes G(W_\bullet\Omega_Y) \longrightarrow G(W_\bullet\Omega_X \boxtimes W_\bullet\Omega_Y).$$

Therefore, multiplication induces a morphism of complexes

$$(3.1.2) \quad G(W_\bullet\Omega_X) \boxtimes G(W_\bullet\Omega_Y) \longrightarrow G(W_\bullet\Omega_{X \times Y}).$$

DEFINITION 3.1.4. Let (X, Φ) and (Y, Ψ) be in (Sm_*/S) and denote the structure maps by $a : X \rightarrow S$ and $b : Y \rightarrow S$. Then we define the exterior products

$$\times : \mathcal{H}^i((X, \Phi)/S) \boxtimes \mathcal{H}^j((Y, \Psi)/S) \rightarrow \mathcal{H}^{i+j}((X \times Y, \Phi \times \Psi)/S \times S),$$

$$\times : \hat{\mathcal{H}}^i((X, \Phi)/S) \boxtimes \hat{\mathcal{H}}^j((Y, \Psi)/S) \rightarrow \hat{\mathcal{H}}^{i+j}((X \times Y, \Phi \times \Psi)/S \times S)$$

as the composition of $H^{i+j}((a \times b)_{\Phi \times \Psi}(3.1.2))$ (resp. $H^{i+j}(\widehat{(a \times b)}_{\Phi \times \Psi}(3.1.2))$) with the natural map

$$\mathcal{H}^i((X, \Phi)/S) \boxtimes \mathcal{H}^j((Y, \Psi)/S) \rightarrow H^{i+j}((a \times b)_{\Phi \times \Psi}(G(W_\bullet\Omega_X) \boxtimes G(W_\bullet\Omega_Y)))$$

(resp. with $(\hat{})$), where we use $\mathcal{H}^i((X, \Phi)/S) = H^i(a_\Phi G(W_\bullet\Omega_X))$, etc.

LEMMA 3.1.5. *Let $X = \text{Spec } A$ and $Y = \text{Spec } B$ be two smooth affine k -schemes and $Z_X \subset X$, $Z_Y \subset Y$ closed subsets. Assume there are sections $s_1, \dots, s_i \in A$ and $t_1, \dots, t_j \in B$ whose vanishing set equals Z_X and Z_Y respectively. Then for all $n \geq 1$ and $q, r \geq 0$ the morphism*

$$\times : H_{Z_X}^i(X, W_n\Omega_X^q) \otimes_{\mathbb{Z}} H_{Z_Y}^j(Y, W_n\Omega_Y^r) \rightarrow H_{Z_X \times Z_Y}^{i+j}(X \times Y, W_n\Omega_{X \times Y}^{q+r})$$

induced by Definition 3.1.4 is given by

$$\left[\begin{array}{c} \alpha \\ [s_1], \dots, [s_i] \end{array} \right] \times \left[\begin{array}{c} \beta \\ [t_1], \dots, [t_j] \end{array} \right] = \left[\begin{array}{c} p_1^* \alpha \cdot p_2^* \beta \\ p_1^*[s_1], \dots, p_1^*[s_i], p_2^*[t_1], \dots, p_2^*[t_j] \end{array} \right],$$

where we use the notation of 1.11.1, $\alpha \in W_n\Omega_A^q$, $\beta \in W_n\Omega_B^r$ and $p_1 : X \times Y \rightarrow X$ and $p_2 : X \times Y \rightarrow Y$ are the two projection maps.

Proof. We denote by C the tensor product $A \otimes_k B$ and by K_A , K_B and K_C the respective Koszul complexes (see 1.11.1) $K^\bullet([t], W_n\Omega_A^q)$, $K^\bullet([s], W_n\Omega_B^r)$ and $K^\bullet(p_1^*[t], p_2^*[s], W_n\Omega_C^{q+r})$. For any ordered tuple $L = (l_1 < \dots < l_\xi)$ of natural numbers $l_\nu \in [1, i+j]$ we denote $\sigma(L) := \max\{\nu \in [0, \xi] \mid l_\nu \leq i\}$, where we set $l_0 := 0$. Notice that $\sigma(L) = 0$ iff $l_1 > i$ or $L = \emptyset$. We define

$$L^{\leq i} = \begin{cases} (l_1 < \dots < l_{\sigma(L)}), & \text{if } \sigma(L) \in [1, \xi], \\ \emptyset, & \text{if } \sigma(L) = 0, \end{cases}$$

$$L^{> i} = \begin{cases} ((l_{\sigma(L)+1} - i) < \dots < (l_\xi - i)), & \text{if } \sigma(L) < \xi, \\ \emptyset, & \text{if } \sigma(L) = \xi \text{ or } L = \emptyset. \end{cases}$$

Thus the natural numbers appearing in the tuple $L^{\leq i}$ are contained in $[1, i]$, whereas those appearing in $L^{> i}$ are contained in $[1, j]$.

Let $\{e_1, \dots, e_i\}$, $\{f_1, \dots, f_j\}$ and $\{h_1, \dots, h_{i+j}\}$ be the standard bases of $W_n(A)^i$, $W_n(B)^j$ and $W_n(C)^{i+j}$, respectively. If $L = (l_1 < \dots < l_\xi)$ is an ordered tuple with $l_\nu \in [1, i]$, then we write $e_L = e_{l_1} \wedge \dots \wedge e_{l_\xi} \in \bigwedge^\xi W_n(A)^i$. In case L is the emptyset we define e_L to be $1 \in W_n(A)$. Similarly we define the notations f_L and h_L .

With this notation we can define the morphism

$$\epsilon : \bigoplus_{\mu+\nu=\xi} \left(\text{Hom}_{W_n(A)} \left(\bigwedge^\mu W_n(A)^i, W_n\Omega_A^q \right) \times \text{Hom}_{W_n(B)} \left(\bigwedge^\nu W_n(B)^j, W_n\Omega_B^r \right) \right) \longrightarrow \text{Hom}_{W_n(C)} \left(\bigwedge^\xi W_n(C)^{i+j}, W_n\Omega_C^{q+r} \right),$$

by sending a tuple of pairs of morphisms $\bigoplus_{\mu+\nu=\xi} (\varphi^\mu, \psi^\nu)$ to the morphism, which is uniquely determined by

$$h_L \mapsto p_1^* \varphi^{\sigma(L)}(e_{L \leq i}) \cdot p_2^* \psi^{\xi - \sigma(L)}(f_{L > i}).$$

It is straightforward to check, that ϵ induces a morphism of complexes $K_A \otimes_{\mathbb{Z}} K_B \rightarrow K_C$, which in degree zero equals

$$\epsilon^0 : W_n\Omega_A^q \otimes_{\mathbb{Z}} W_n\Omega_B^r \rightarrow W_n\Omega_C^{q+r}, \quad \alpha \otimes \beta \mapsto p_1^* \alpha \cdot p_2^* \beta$$

and in degree $i + j$ is given by

$$\epsilon^{i+j} : K_A^i \otimes_{\mathbb{Z}} K_B^j \rightarrow K_C^{i+j}, \quad \epsilon^{i+j}(\varphi \otimes \psi)(h_{[1, i+j]}) = p_1^* \varphi(e_{[1, i]}) \cdot p_2^* \psi(f_{[1, j]}).$$

By the very definition of the symbols $[\dots]_{[s_1], \dots, [s_i]}^\alpha$, etc. in 1.11.1 it remains to show, that the following diagram commutes:

$$(3.1.3) \quad \begin{array}{ccc} H^i(K_A) \otimes H^j(K_B) & \xrightarrow{\text{nat.}} & H^{i+j}(K_A \otimes K_B) \xrightarrow{\epsilon} H^{i+j}(K_C) \\ \downarrow (1.11.1) & & \downarrow (1.11.1) \\ H_{Z_X}^i \otimes H_{Z_Y}^j & \xrightarrow{\quad \times \quad} & H_{Z_X \times Z_Y}^{i+j} \end{array}$$

where we abbreviate $H_{Z_X}^i = H_{Z_X}^i(X, W_n\Omega_X^q)$, etc. This is a straightforward calculation. □

3.1.6. *Exterior product for Cousin complexes.* Let X and Y be smooth k -schemes and denote by Z_X^\bullet and Z_Y^\bullet their respective filtrations by codimension. Then we obtain from 3.1.2 a natural morphism

$$\frac{\Gamma_{Z_X^a} G(W_\bullet, \Omega_X) \boxtimes \Gamma_{Z_Y^b} G(W_\bullet, \Omega_Y)}{\Gamma_{Z_X^{a+1}} \boxtimes \Gamma_{Z_Y^b} + \Gamma_{Z_X^a} \boxtimes \Gamma_{Z_Y^{b+1}}} \rightarrow \frac{\Gamma_{Z_{X \times Y}^{a+b}} G(W_\bullet, \Omega_{X \times Y})}{\Gamma_{Z_{X \times Y}^{a+b+1}} G(W_\bullet, \Omega_{X \times Y})},$$

where we abbreviate the denominator on the left in the obvious way. With the notation from 1.5.8, there is an obvious morphism from

$$\mathcal{H}_{Z_X^a/Z_X^{a+1}}^i(W_\bullet, \Omega_X) \boxtimes \mathcal{H}_{Z_Y^b/Z_Y^{b+1}}^j(W_\bullet, \Omega_Y)$$

to the $(i + j)$ -th cohomology of the left hand side. In composition we obtain a morphism

$$\mathcal{H}_{Z_X^a/Z_X^{a+1}}^i(W_\bullet\Omega_X) \boxtimes \mathcal{H}_{Z_Y^b/Z_Y^{b+1}}^j(W_\bullet\Omega_Y) \rightarrow \mathcal{H}_{Z_{X \times Y}^{a+b}/Z_{X \times Y}^{a+b+1}}^{i+j}(W_\bullet\Omega_{X \times Y}),$$

which is compatible with Frobenius. It is straightforward to check that this pairing induces a morphism of *pro-complexes of sheaves of W -modules on $X \times Y$*

$$(3.1.4) \quad E(W_\bullet\Omega_X) \boxtimes E(W_\bullet\Omega_Y) \rightarrow E(W_\bullet\Omega_{X \times Y}),$$

where $E(-)$ denotes the Cousin complex with respect to the codimension filtration (see 1.5.8).

PROPOSITION 3.1.7. *The exterior products defined above satisfy the following properties:*

- (1) *The exterior products (for \mathcal{H} and $\hat{\mathcal{H}}$) can also be calculated by using the morphism (3.1.4) in Definition 3.1.4 instead of the morphism (3.1.2).*
- (2) *The exterior products are associative.*
- (3) *Let $f : (X, \Phi) \rightarrow (Y, \Psi)$ and $f' : (X', \Phi') \rightarrow (Y', \Psi')$ be two morphisms in (Sm^*/S) and $\alpha \in \mathcal{H}((Y, \Psi)/S)$, $\alpha' \in \mathcal{H}((Y', \Psi')/S)$, then*

$$(f^*\alpha) \times (f'^*\alpha') = (f \times f')^*(\alpha \times \alpha') \quad \text{in } \mathcal{H}((X \times X', \Phi \times \Phi')/S \times S).$$

Similar with $\hat{\mathcal{H}}$.

- (4) *Let $f : (X, \Phi) \rightarrow (Y, \Psi)$ be a morphism in (Sm_*/S) and $\alpha \in \mathcal{H}((X, \Phi)/S)$, $\alpha' \in \mathcal{H}((X', \Phi')/S)$, then*

$$(f_*\alpha) \times \alpha' = (f \times \text{id}_{X'})_*(\alpha \times \alpha') \quad \text{in } \mathcal{H}((Y \times X', \Psi \times \Phi')/S \times S).$$

Similar with $\hat{\mathcal{H}}$.

- (5) *Let (X, Φ) and (Y, Ψ) be smooth k -schemes over S with families of supports and $\alpha \in \mathcal{H}^{i,q}((X, \Phi)/S)$, $\beta \in \mathcal{H}^{j,r}((Y, \Psi)/S)$. Then (similar properties hold for $\hat{\mathcal{H}}$):*

- (a) *The switching isomorphism*

$$\mathcal{H}((X \times Y, \Phi \times \Psi)/S \times S) \cong \mathcal{H}((Y \times X, \Psi \times \Phi)/S \times S)$$

sends $\alpha \times \beta$ to $(-1)^{qr+ij}\beta \times \alpha$.

- (b)

$$F(\alpha) \times F(\beta) = F(\alpha \times \beta), \quad \pi(\alpha) \times \pi(\beta) = \pi(\alpha \times \beta),$$

$$d(\alpha \times \beta) = (d\alpha) \times \beta + (-1)^q\alpha \times d(\beta),$$

for $\alpha, \beta \in \mathcal{H}_n$.

- (c)

$$V(\alpha) \times \beta = V(\alpha \times F(\beta)), \quad \underline{p}(\alpha) \times \beta = \underline{p}(\alpha \times \pi(\beta))$$

for $\alpha \in \mathcal{H}_n, \beta \in \mathcal{H}_{n+1}$.

Proof. (1) follows from [God73, II, Thm. 6.2.1]. (2), (3) and (5) easily follow from the properties [God73, II, 6.5, (b), (d), (e)] and the corresponding properties of the multiplication map $W_\bullet\Omega_X \boxtimes W_\bullet\Omega_Y \rightarrow W_\bullet\Omega_{X \times Y}$. It remains to check (4). Since f is quasi-projective we may factor it as the composition of a regular closed immersion followed by an open embedding into a projective space over Y followed by the projection to Y . Using similar arguments as in the proof of Proposition 2.4.4 (together with Lemma 2.4.3), we can reduce to the case that $\Phi = \Psi$ and that $f : (X, \Phi) \hookrightarrow (Y, \Phi)$ is a regular closed immersion of pure codimension c . By (1) it thus suffices to show, that the following diagram commutes

$$\begin{CD} f_*(E(W_\bullet\Omega_X)) \boxtimes E(W_\bullet\Omega_{X'}) @>(3.1.4)>> E(W_\bullet\Omega_{X \times X'}) \\ @V f_* \boxtimes \text{id} VV @VV (f \times \text{id}_{X'})_* V \\ (\Gamma_X(E(W_\bullet\Omega_Y))(c)[c]) \boxtimes E(W_\bullet\Omega_{X'}) @>(3.1.4)>> \Gamma_{X \times X'} E(W_\bullet\Omega_{Y \times X'})(c)[c], \end{CD}$$

where we denote by f_* and $(f \times \text{id}_{X'})_*$ the morphism defined in (2.3.3). The question is local and we may therefore assume $X = \text{Spec } A$, $X' = \text{Spec } A'$ and $Y = \text{Spec } B$ are affine and that the ideal of Y in X is generated by a regular sequence τ_1, \dots, τ_c . Furthermore we may check the commutativity in each degree separately and thus by the description of the terms of the Cousin complex in 1.5.8, (2), we are reduced to show the commutativity of the following diagram:

$$\begin{CD} H_{Z_X}^i(X, W_n\Omega_X^q) \otimes H_{Z_{X'}}^i(X', W_n\Omega_{X'}^q) @>\times>> H_{Z_X \times Z_{X'}}^{i+j}(X \times X', W_n\Omega_{X \times X'}^{q+r}) \\ @V f_* \otimes \text{id} VV @VV (f \times \text{id})_* V \\ H_{Z_X}^{i+c}(Y, W_n\Omega_{Y \times X'}^{q+c}) \otimes H_{Z_{X'}}^i(X', W_n\Omega_{X'}^q) @>\times>> H_{Z_X \times Z_{X'}}^{i+j+c}(Y \times X', W_n\Omega_{Y \times X'}^{q+r+c}), \end{CD}$$

where $Z_X \subset X$ and $Z_{X'} \subset X'$ are integral closed subschemes of codimension i and j respectively. But this follows from the explicit formulas in Corollary 2.4.2 and Lemma 3.1.5. \square

Remark 3.1.8. Notice that in the situation of (4) above the two elements $\alpha' \times (f_*\alpha)$ and $(\text{id}_{X'} \times f)_*(\alpha' \times \alpha)$ are in general only equal up to a sign. This is the reason for the sign in (3.3.6).

3.2. THE CYCLE CLASS OF GROS. We review some results of [Gro85].

3.2.1. *Witt log forms.* Let X be a smooth k scheme. We denote by

$$W_n\Omega_{X,\log}^q$$

the abelian sheaf on the small étale site $X_{\text{ét}}$ of X defined in [Ill79, I, 5.7]; it is the abelian subsheaf of $W_n\Omega_{X_{\text{ét}}}^q$ which is locally generated by sections of the form

$$\frac{d[x_1]}{[x_1]} \dots \frac{d[x_q]}{[x_q]}, \quad \text{with } x_1, \dots, x_d \in \mathcal{O}_{X_{\text{ét}}}^\times.$$

By definition $W_n\Omega_{X,\log}^0 = \mathbb{Z}/p^n\mathbb{Z}$.

3.2.2. *The cycle class.* Let X be a smooth k -scheme and $Z \subset X$ a closed integral subscheme of codimension c . Then by [Gro85, II, (4.1.6)] the restriction map

$$(3.2.1) \quad H_Z^c(X_{\text{ét}}, W_n \Omega_{X, \log}^c) \xrightarrow{\cong} H_{Z \setminus Z_{\text{sing}}}^c((X \setminus Z_{\text{sing}})_{\text{ét}}, W_n \Omega_{X, \log}^c)$$

is an isomorphism (where Z_{sing} denotes the singular locus of Z). Therefore following Gros we can define the *log-cycle class of Z of level n*

$$cl_{\log, n}(Z) \in H_Z^c(X_{\text{ét}}, W_n \Omega_{X, \log}^c)$$

as the image of 1 under the Gysin isomorphism ([Gro85, II, Thm 3.5.8 and (3.5.19)])

$$(3.2.2) \quad H^0((Z \setminus Z_{\text{sing}})_{\text{ét}}, \mathbb{Z}/p^n \mathbb{Z}) \xrightarrow{\cong} H_{Z \setminus Z_{\text{sing}}}^c((X \setminus Z_{\text{sing}})_{\text{ét}}, W_n \Omega_{X, \log}^c)$$

composed with the inverse of (3.2.1) (see [Gro85, II, Def. 4.1.7.]). We define the *cycle class of Z of level n*

$$cl_n(Z) \in H_Z^c(X, W_n \Omega_X)$$

to be the image of $cl_{\log, n}(Z)$ under the natural morphism $H_Z^c(X_{\text{ét}}, W_n \Omega_{X, \log}^c) \rightarrow H_Z^c(X, W_n \Omega_X^c)$. We have

$$\pi(cl_{\log, n}(Z)) = cl_{\log, n-1}(Z), \quad \pi(cl_n(Z)) = cl_{n-1}(Z).$$

Therefore $(cl_{\log, n}(Z))_n$ (resp. $(cl_n(Z))_n$) give rise to an element in the projective system $H_Z^c(X_{\text{ét}}, W_\bullet \Omega_{X, \log}^c)$ (resp. $H_Z^c(X, W_\bullet \Omega_X^c)$), which we simply denote by $cl_{\log}(Z)$ (resp. $cl(Z)$).

Notice that we have an isomorphism

$$(3.2.3) \quad H_Z^c(X, W \Omega_X^c) \cong \varprojlim H_Z^c(X, W_n \Omega_X^c).$$

(Indeed by Proposition 1.5.6, (5) we have a spectral sequence

$$R^i \varprojlim H_Z^j(X, W_n \Omega_X^c) \implies H_Z^{i+j}(X, W \Omega_X^c),$$

which by Lemma 1.5.1, (1) gives a short exact sequence

$$0 \rightarrow R^1 \varprojlim H_Z^{c-1}(X, W_n \Omega_X^c) \rightarrow H_Z^c(X, W \Omega_X^c) \rightarrow \varprojlim H_Z^c(X, W_n \Omega_X^c) \rightarrow 0.$$

But by Lemma 1.5.9 $H_Z^{c-1}(X, W_n \Omega_X^c)$ equals 0 for all n . Hence (3.2.3).) We define

$$\hat{cl}(Z) \in H_Z^c(X, W \Omega_X^c)$$

as the image of $\varprojlim_n cl_n(Z)$ via (3.2.3).

3.2.3. *Properties of the cycle classes.* The above cycle classes have the following properties (we only list them for cl there are analogous properties for cl_{\log} and \hat{cl}):

- (1) Let $Z \subset X$ be a closed integral subscheme of codimension c and $U \subset X$ be open such that $Z \cap U$ is smooth. Denote by $i : Z \cap U \hookrightarrow (U, Z \cap U)$ the morphism in $(Sm_*/\text{Spec } k)$ which is induced by the inclusion of Z

in X and by $j : (U, Z \cap U) \hookrightarrow (X, Z)$ in $(Sm^*/\text{Spec } k)$ the morphism induced by the open immersion $U \subset X$. Then

$$j^*cl(Z) = cl(Z \cap U) = i_*(1) \quad \text{in } H_Z^c(X, W_\bullet \Omega_X^c),$$

where 1 is the multiplicative unit in the pro-ring $H^0(Z \cap U, W_\bullet \mathcal{O}_{Z \cap U})$.

- (2) Let Z and Z' be two closed integral subschemes of X of codimension c and c' respectively intersecting each other properly (i.e. each irreducible component of $Z \cap Z'$ has codimension $c + c'$ in X). Then

$$cl(Z.Z') = \Delta^*(cl(Z) \times cl(Z')) \quad \text{in } H_{Z \cap Z'}^{c+c'}(X, W_\bullet \Omega^{c+c'}).$$

Here $Z.Z'$ is the intersection cycle $\sum_T n_T [T]$, where the sum is over the irreducible components of $Z \cap Z'$ and n_T are the intersection multiplicities (computed via Serres tor-formula), $cl(Z.Z')$ is defined to be $\sum_T n_i cl(T)$ and $\Delta : (X, Z \cap Z') \rightarrow (X \times X, Z \times Z')$ in $(Sm^*/\text{Spec } k)$ is induced by the diagonal morphism.

- (3) For a line bundle \mathcal{L} on X we denote by $c_1(\mathcal{L})_{\log} \in H^1(X_{\text{ét}}, W_\bullet \Omega_{X, \log}^1)$ the sequence of elements $(dlog_n([\mathcal{L}]))_n$, where $[\mathcal{L}]$ denotes the class of \mathcal{L} in $H^1(X, \mathcal{O}_X^\times) = H^1(X_{\text{ét}}, \mathbb{G}_m)$ and $dlog_n$ is induced by taking H^1 of the map $\mathbb{G}_m \rightarrow W_n \Omega_{X, \log}^1, a \rightarrow dlog[a]$. We denote by $c_1(\mathcal{L})$ the image of $c_1(\mathcal{L})_{\log}$ in $H^1(X, W_\bullet \Omega_X^c)$.

Let $D \subset X$ be an integral subscheme of codimension 1, then

$$cl(D) = c_1(\mathcal{O}(D)) \quad \text{in } H^1(X, W_\bullet \Omega_X^1).$$

- (4) Let Z be as in (1), then:

$$F(cl(Z)) = \pi(cl(Z)), \quad V(cl(Z)) = \underline{p}(cl(Z)), \quad d(cl(Z)) = 0.$$

(1) follows from the fact, that the Gysin morphism (3.2.2) is induced by the pushforward i_{0*} , where $i_0 : Z \setminus Z_{\text{sing}} \rightarrow (X \setminus Z_{\text{sing}}, Z \setminus Z_{\text{sing}})$ in $(Sm_*/\text{Spec } k)$ is induced by the inclusion. This follows from [Gro85, II, 3.4.] (or alternatively from the description of the Gysin morphism in [Gro85, II, Prop. 3.5.6] together with Proposition 2.4.1.) (2) is [Gro85, II, Prop. 4.2.12.] and (3) is [Gro85, II, Prop. 4.2.1]. The first and the last equality in (4) follow from the definition, the second equality is implied by the first and $\pi(cl_n(Z)) = cl_{n-1}(Z)$.

3.3. HODGE-WITT COHOMOLOGY AS WEAK COHOMOLOGY THEORY WITH SUPPORTS.

3.3.1. *Weak cohomology theories with supports.* Weak cohomology theories with supports have been introduced in [CR09]. For the convenience of the reader we recall the definitions.

First, recall the definition of the categories $Sm_* = (Sm_*/k)$ and $Sm^* = (Sm^*/k)$ in Definition 2.1.1. For both categories Sm^* and Sm_* finite coproducts exist:

$$(X, \Phi) \coprod (Y, \Psi) = (X \coprod Y, \Phi \cup \Psi).$$

For (X, Φ) let $X = \coprod_i X_i$ be the decomposition into connected components, then

$$(X, \Phi) = \coprod_i (X_i, \Phi \cap \Phi_{X_i}).$$

In general products don't exist, and we define

$$(X, \Phi) \otimes (Y, \Psi) := (X \times Y, \Phi \times \Psi)$$

which together with the unit object $\mathbf{1} = \text{Spec}(k)$ and the obvious isomorphism $(X, \Phi) \otimes (Y, \Psi) \rightarrow (Y, \Psi) \otimes (X, \Phi)$ makes Sm_* and Sm^* to a symmetric monoidal category (see [ML98, VII.1]).

3.3.2. A weak cohomology theory with supports consists of the following data (F_*, F^*, T, e) :

- (1) Two functors $F_* : Sm_* \rightarrow \text{GRAB}$ and $F^* : (Sm^*)^{op} \rightarrow \text{GRAB}$ such that $F_*(X) = F^*(X)$ as abelian groups for every object $X \in \text{ob}(Sm_*) = \text{ob}(Sm^*)$. We will simply write $F(X) := F_*(X) = F^*(X)$. We use lower indexes for the grading on $F_*(X)$, i.e. $F_*(X) = \oplus_i F_i(X)$, and upper indexes for $F^*(X)$.
- (2) For every two objects $X, Y \in \text{ob}(Sm_*) = \text{ob}(Sm^*)$ a morphism of graded abelian groups (for both gradings):

$$T_{X,Y} : F(X) \otimes_{\mathbb{Z}} F(Y) \rightarrow F(X \otimes Y).$$

- (3) A morphism of abelian groups $e : \mathbb{Z} \rightarrow F(\text{Spec}(k))$. For all smooth schemes $\pi : X \rightarrow \text{Spec}(k)$ we denote by 1_X the image of $1 \in \mathbb{Z}$ via the map $\mathbb{Z} \xrightarrow{e} F^*(\text{Spec}(k)) \xrightarrow{F^*(\pi)} F^*(X)$.

3.3.3. These data are required to satisfy the following conditions:

- (1) The functor F_* preserves coproducts and F^* maps coproducts to products. Moreover, for $(X, \Phi_1), (X, \Phi_2) \in \text{ob}(Sm_*)$ with $\Phi_1 \cap \Phi_2 = \{\emptyset\}$ we require that the map $F^*(j_1) + F^*(j_2) : F^*(X, \Phi_1) \oplus F^*(X, \Phi_2) \rightarrow F^*(X, \Phi_1 \cup \Phi_2)$, with $j_1 : (X, \Phi_1 \cup \Phi_2) \rightarrow (X, \Phi_1)$ and $j_2 : (X, \Phi_1 \cup \Phi_2) \rightarrow (X, \Phi_2)$ in Sm^* , is an isomorphism.
- (2) The data (F_*, T, e) and (F^*, T, e) respectively define a (right-lax) symmetric monoidal functor.
- (3) (Grading) For (X, Φ) such that X is connected the equality

$$F_i(X, \Phi) = F^{2 \dim X - i}(X, \Phi)$$

holds for all i .

- (4) For all cartesian diagrams (cf. 2.1.1)

$$\begin{array}{ccc} (X', \Phi') & \xrightarrow{f'} & (Y', \Psi') \\ \downarrow g_X & & \downarrow g_Y \\ (X, \Phi) & \xrightarrow{f} & (Y, \Psi) \end{array}$$

with $g_X, g_Y \in Sm^*$ and $f, f' \in Sm_*$ such that either g_Y is smooth or g_Y is a closed immersion and f is transversal to g_Y the following equality holds:

$$F^*(g_Y) \circ F_*(f) = F_*(f') \circ F^*(g_X).$$

Recall that f is transversal to g_Y if $(f')^*N_{Y'/Y} = N_{X'/X}$ where N denotes the normal bundle. The case $X' = \emptyset$ is also admissible; in this case the equality 3.3.3(4) reads:

$$F^*(g_Y) \circ F_*(f) = 0.$$

In [CR09, p. 700] it is spelled out what it means for (F_*, T, e) (and (F^*, T, e)) to be a right-lax symmetric monoidal functor.

DEFINITION 3.3.4. If the data (F_*, F^*, T, e) as in 3.3.2 satisfy the conditions 3.3.3 then we call (F_*, F^*, T, e) a *weak cohomology theory with supports*.

DEFINITION 3.3.5. Let $(F_*, F^*, T, e), (G_*, G^*, U, \epsilon)$ be two weak cohomology theories with support. By a morphism

$$(3.3.1) \quad (F_*, F^*, T, e) \rightarrow (G_*, G^*, U, \epsilon)$$

we understand a morphism of graded abelian groups (for both gradings)

$$\phi : F(X) \rightarrow G(X) \quad \text{for every } X \in \text{ob}(V_*) = \text{ob}(V^*),$$

such that ϕ induces a natural transformation of (right-lax) symmetric monoidal functors

$$\phi : (F_*, T, e) \rightarrow (G_*, U, \epsilon) \text{ and } \phi : (F^*, T, e) \rightarrow (G^*, U, \epsilon),$$

i.e. ϕ induces natural transformations $F_* \rightarrow G_*, F^* \rightarrow G^*$, and

$$(3.3.2) \quad \phi \circ T = U \circ (\phi \otimes \phi), \quad \phi \circ e = \epsilon.$$

3.3.6. *Chow theory as a weak cohomology theory with supports.* An example of a weak cohomology theory with supports are the Chow groups (CH_*, CH^*, \times, e) [CR09, 1.1]. We will briefly recall the definitions for the convenience of the reader.

DEFINITION 3.3.7 (Chow groups with support). Let Φ be a family of supports on X . We define:

$$\text{CH}(X, \Phi) = \varinjlim_{W \in \Phi} \text{CH}(W).$$

The group $\text{CH}(X, \Phi)$ is graded by dimension. We set

$$\text{CH}_*(X, \Phi) = \bigoplus_{d \geq 0} \text{CH}_d(X, \Phi)[2d],$$

where the bracket $[2d]$ means that $\text{CH}_d(X, \Phi)$ is considered to be in degree $2d$. There is also a grading by codimension. Let $X = \coprod_i X_i$ be the decomposition into connected components then $\text{CH}^*(X, \Phi) = \bigoplus_i \text{CH}^*(X_i, \Phi \cap \Phi_{X_i})$ and

$$\text{CH}^*(X_i, \Phi \cap \Phi_{X_i}) = \bigoplus_{d \geq 0} \text{CH}^d(X_i, \Phi \cap \Phi_{X_i})[2d]$$

where $\text{CH}^d(X_i, \Phi \cap \Phi_{X_i})$ is generated by cycles $[Z]$ with $Z \in \Phi \cap \Phi_{X_i}$, Z irreducible, and $\text{codim}_{X_i}(Z) = d$.

Let $f : (X, \Phi) \rightarrow (Y, \Psi)$ be a morphism in Sm_* . There is a push-forward

$$(3.3.3) \quad \text{CH}_*(f) : \text{CH}(X, \Phi) \rightarrow \text{CH}(Y, \Psi),$$

defined by the usual push-forward of cycles $(f|_W)_* : \text{CH}(W) \rightarrow \text{CH}(f(W)) \rightarrow \text{CH}(Y, \Psi)$ (for this we need that $f|_W$ is proper and $f(\Phi) \subset \Psi$). We obtain a functor

$$(3.3.4) \quad \text{CH}_* : Sm_* \rightarrow \text{GRAB}, \quad \text{CH}_*(X, \Phi) := \text{CH}(X, \Phi), \quad f \mapsto \text{CH}_*(f).$$

In order to define a functor

$$\text{CH}^* : (Sm^*)^{op} \rightarrow \text{GRAB}$$

we use Fulton’s work on refined Gysin morphisms [Ful98, §6.6].

Let $f : X \rightarrow Y$ be a morphism between smooth schemes and let $V \subset Y$ be a closed subscheme, thus $f : (X, f^{-1}(V)) \rightarrow (Y, V)$ is a morphism in Sm^* . We define

$$\text{CH}^*(f) := f^! : \text{CH}(Y, V) = \text{CH}(V) \rightarrow \text{CH}(f^{-1}(V)) = \text{CH}(X, f^{-1}(V)).$$

For the general case let $f : (X, \Phi) \rightarrow (Y, \Psi)$ be any morphism in Sm^* . For every $V \in \Psi$ the map f induces $(X, f^{-1}(V)) \rightarrow (Y, V)$ in Sm^* . We may define

$$\text{CH}^*(f) : \text{CH}(Y, \Psi) = \varinjlim_{V \in \Psi} \text{CH}(Y, V) \rightarrow \varinjlim_{W \in \Phi} \text{CH}(X, W) = \text{CH}(X, \Phi).$$

The assignment

$$\text{CH}^* : (Sm^*)^{op} \rightarrow \text{GRAB}, \quad \text{CH}^*(X, \Phi) = \text{CH}(X, \Phi), \quad f \mapsto \text{CH}^*(f)$$

defines a functor. Together with the exterior product \times (see [Ful98, §1.10]) and the obvious unit $1 : \mathbb{Z} \rightarrow \text{CH}(\text{Spec}(k))$, we obtain a weak cohomology theory with supports $(\text{CH}_*, \text{CH}^*, \times, 1)$.

3.3.8. From Proposition 2.3.3 and Proposition 2.2.2 we obtain two functors

$$\begin{aligned} \hat{H}_* &: Sm_* \rightarrow \text{GRAB}, \\ \hat{H}^* &: (Sm^*)^{op} \rightarrow \text{GRAB}, \end{aligned}$$

where for an object $(X, \Phi) \in \text{ob}(Sm_*) = \text{ob}(Sm^*)$ with structure morphism $a : X \rightarrow \text{Spec}(k)$ we have

$$\hat{H}(X, \Phi) := \hat{H}_*(X, \Phi) := \hat{H}^*(X, \Phi) := \hat{\mathcal{H}}((X, \Phi)/\text{Spec } k) = \bigoplus_{p, q \geq 0} R^q \hat{a}_\Phi W_\bullet \Omega_X^p$$

as abelian groups (see (1.4.4) for the definition of \hat{a}_Φ). In view of Proposition 1.5.6 we get an equality

$$R^q \hat{a}_\Phi = H^q \circ R \varprojlim \circ Ra_* \circ R \Gamma_{\Phi}.$$

In particular, if $\Phi = \Phi_X$ is the set of all closed subsets of X then

$$\hat{H}(X) := \hat{H}(X, \Phi_X) = \bigoplus_{p, q \geq 0} H^q(R \varprojlim R \Gamma(W_\bullet \Omega_X^p)).$$

Note that $R\varprojlim \circ R\Gamma = R\Gamma \circ R\varprojlim$ and $R\varprojlim(W_\bullet \Omega_X^p) = \varprojlim(W_\bullet \Omega_X^p) = W\Omega_X^p$ by Lemma 1.5.1. Thus we obtain

$$\hat{H}(X) = \bigoplus_{p,q \geq 0} H^q(X, W\Omega_X^p).$$

In the following we will write $\hat{H}(X)$ for $\hat{H}(X, \Phi_X)$.

The grading for \hat{H}^* is defined by

$$\hat{H}^i(X, \Phi) = \bigoplus_{p+q=i} R^q \hat{a}_\Phi W_\bullet \Omega_X^p.$$

For $\hat{H}_*(X, \Phi)$ the grading is defined such that 3.3.3(4) holds.

For two objects $(X, \Phi), (Y, \Psi)$ with structure maps $a : X \rightarrow \text{Spec}(k), b : Y \rightarrow \text{Spec}(k)$, we define

$$(3.3.5) \quad T : \hat{H}(X, \Phi) \otimes_{\mathbb{Z}} \hat{H}(Y, \Psi) \rightarrow \hat{H}(X \times Y, \Phi \times \Psi)$$

by the formula

$$(3.3.6) \quad T(\alpha_{i,p} \otimes \beta_{j,q}) = (-1)^{(i+p) \cdot j} (\alpha_{i,p} \times \beta_{j,q}),$$

where $\alpha_{i,p} \in R^i \hat{a}_\Phi W_\bullet \Omega_X^p, \beta_{j,q} \in R^j \hat{b}_\Psi W_\bullet \Omega_Y^q$, and \times is the map in 3.1.4.

We define

$$e : \mathbb{Z} \rightarrow \hat{H}(\text{Spec}(k)) = W(k)$$

by $e(n) = n$ for all $n \in \mathbb{Z}$.

PROPOSITION 3.3.9. *The triples (\hat{H}_*, T, e) and (\hat{H}^*, T, e) define right-lax symmetric monoidal functors.*

Proof. The morphism T respects the grading \hat{H}^* and the grading \hat{H}_* . In the following we will work with the upper grading \hat{H}^* . All arguments will also work for the lower grading \hat{H}_* because the difference between lower and upper grading is an even integer.

The axioms which involve e do obviously hold. By using the associativity of \times (Proposition 3.1.7(2)) it is straightforward to prove the associativity of T . Let us prove the commutativity of T , i.e. that the diagram

$$(3.3.7) \quad \begin{array}{ccc} H(X, \Phi) \otimes H(Y, \Psi) & \xrightarrow{T} & H(X \times Y, \Phi \times \Psi) \\ \downarrow & & \downarrow \epsilon \\ H(Y, \Psi) \otimes H(X, \Phi) & \xrightarrow{T} & H(Y \times X, \Psi \times \Phi) \end{array}$$

is commutative. The left vertical map is defined by $\alpha \otimes \beta \mapsto (-1)^{\deg(\alpha) \deg(\beta)} \beta \otimes \alpha$. By using Proposition 3.1.7(5)(a) we obtain

$$\begin{aligned} \epsilon(T(\alpha_{i,p} \otimes \beta_{j,q})) &= \epsilon((-1)^{(i+p)j} \alpha_{i,p} \times \beta_{j,q}) = (-1)^{(i+p)j} (-1)^{pq+ij} \beta_{j,q} \times \alpha_{i,p} \\ &= (-1)^{pq+pj} \beta_{j,q} \times \alpha_{i,p}. \end{aligned}$$

On the other hand,

$$\begin{aligned} T((-1)^{(i+p)(j+q)} \beta_{j,q} \otimes \alpha_{i,p}) &= (-1)^{(i+p)(j+q)} (-1)^{(j+q)i} \beta_{j,q} \times \alpha_{i,p} \\ &= (-1)^{pq+pj} \beta_{j,q} \times \alpha_{i,p}. \end{aligned}$$

This proves the commutativity of diagram (3.3.7).

The functoriality of T with respect to \hat{H}^* follows immediately from Proposition 3.1.7(3). The functoriality of T with respect to \hat{H}_* follows from Proposition 3.1.7(4). \square

THEOREM 3.3.10. *The datum $(\hat{H}_*, \hat{H}^*, T, e)$ is a weak cohomology theory with supports (cf. Definition 3.3.4).*

Proof. We have to verify the properties in section 3.3.3. Property 3.3.3(1) is obvious. Proposition 3.3.9 implies 3.3.3(2). The compatibility of the gradings (3.3.3(3)) is satisfied by definition. Proposition 2.4.4 yields 3.3.3(4). \square

3.3.11. In the same way as in section 3.3.8 we define

$$\begin{aligned} P\hat{H}_* &: Sm_* \rightarrow \text{GRAB}, \\ P\hat{H}^* &: (Sm^*)^{op} \rightarrow \text{GRAB}, \end{aligned}$$

where for an object $(X, \Phi) \in \text{ob}(Sm_*) = \text{ob}(Sm^*)$ with structure morphism $a : X \rightarrow \text{Spec}(k)$ we have

$$P\hat{H}(X, \Phi) = P\hat{H}_*(X, \Phi) = P\hat{H}^*(X, \Phi) = \bigoplus_{p \geq 0} R^p \hat{a}_\Phi W_\bullet \Omega_X^p$$

as abelian groups. The grading is defined in the obvious way:

$$P\hat{H}^{2p}(X, \Phi) = R^p \hat{a}_\Phi W_\bullet \Omega_X^p,$$

and zero for all odd degrees. Of course, for $P\hat{H}_*(X, \Phi)$ the grading is defined such that 3.3.3(4) holds.

For two objects $(X, \Phi), (Y, \Psi)$, we get an induced map

$$T : P\hat{H}(X, \Phi) \otimes_{\mathbb{Z}} P\hat{H}(Y, \Psi) \rightarrow P\hat{H}(X \times Y, \Phi \times \Psi)$$

from (3.3.5).

Theorem 3.3.10 implies the following statement.

COROLLARY 3.3.12. *The datum $(P\hat{H}_*, P\hat{H}^*, T, e)$ is a weak cohomology theory with supports (cf. Definition 3.3.4). Induced by the inclusion $P\hat{H}(X, \Phi) \subset \hat{H}(X, \Phi)$, we get a morphism of weak cohomology theories with supports (cf. Definition 3.3.5)*

$$(P\hat{H}_*, P\hat{H}^*, T, e) \rightarrow (\hat{H}_*, \hat{H}^*, T, e).$$

3.4. CYCLE CLASSES. One of the main goals of [CR09] was to give a criterion when a weak cohomology theory (F_*, F^*, T, e) admits a morphism

$$(\mathrm{CH}_*, \mathrm{CH}^*, \times, 1) \rightarrow (F_*, F^*, T, e).$$

In order to apply [CR09, Theorem 1.2.3] in our situation, i.e. for $(P\hat{H}_*, P\hat{H}^*, T, e)$, we first need to prove semi-purity.

Recall that $(P\hat{H}_*, P\hat{H}^*, T, e)$ satisfies the semi-purity condition (cf. [CR09, Definition 1.2.1]) if the following holds:

- For all smooth schemes X and irreducible closed subsets $W \subset X$ the groups $P\hat{H}_i(X, W)$ vanish if $i > 2 \dim W$.
- For all smooth schemes X , closed subsets $W \subset X$, and open sets $U \subset X$ such that U contains the generic point of every irreducible component of W , we require the map

$$P\hat{H}^*(j) : P\hat{H}_{2 \dim W}(X, W) \rightarrow P\hat{H}_{2 \dim W}(U, W \cap U),$$

induced by $j : (U, W \cap U) \rightarrow (X, W)$ in Sm^* , to be *injective*.

PROPOSITION 3.4.1. *The weak cohomology theory with supports $(P\hat{H}_*, P\hat{H}^*, T, e)$ satisfies the semi-purity condition.*

Proof. To verify the two conditions we may assume that X is connected of dimension d . Since

$$P\hat{H}_{2i}(X, W) = H_W^{d-i}(X, W\Omega_X^{d-i}),$$

we need to show that $H_W^{d-i}(X, W\Omega_X^{d-i})$ vanishes if $2i > 2 \dim W$ (or equivalently, $d - i < \mathrm{codim}_X W$). Indeed, we have

$$H_W^{d-i}(X, W\Omega_X^{d-i}) = R^{d-i}a_* R\Gamma_W R\varprojlim(W_n\Omega_X^{d-i})$$

with $a : X \rightarrow \mathrm{Spec}(k)$ the structure map. Since $R\Gamma_W R\varprojlim = R\varprojlim R\Gamma_W$, we need to show that

$$(3.4.1) \quad R^j\Gamma_W W_n\Omega_X^{d-i} = 0,$$

for $j < \mathrm{codim}_X W$. This follows from Lemma 1.5.9 (which follows from the fact that the graded pieces of the standard filtration on $W_n\Omega_X^{d-i}$ are extensions of locally free \mathcal{O}_X -modules [Ill79, I, Cor. 3.9]).

For the second condition, let $W \subset X$ be a closed subset and $U \subset X$ open, such that U contains the generic point of every irreducible component of W . We need to prove that the restriction map

$$H_W^{d-\dim W}(X, W\Omega_X^{d-\dim W}) \rightarrow H_{U \cap W}^{d-\dim W}(U, W\Omega_U^{d-\dim W})$$

is injective. But the kernel is a quotient of $H_{W \setminus U}^{d-\dim W}(X, W\Omega_X^{d-\dim W})$, which vanishes because $d - \dim W < \mathrm{codim}_X(W \setminus U)$. \square

3.4.2. In view of [CR09, Theorem 1.2.3] there is at most one morphism of weak cohomology theories with support

$$(\mathrm{CH}_*, \mathrm{CH}^*, \times, 1) \rightarrow (P\hat{H}_*, P\hat{H}^*, T, e).$$

THEOREM 3.4.3. *There is exactly one morphism (in the sense of Definition 3.3.5)*

$$(\mathrm{CH}_*, \mathrm{CH}^*, \times, 1) \rightarrow (P\hat{H}_*, P\hat{H}^*, T, e).$$

Proof. We need to verify the criteria given in [CR09, Theorem 1.2.3]. The condition [CR09, Theorem 1.2.3](1) is satisfied by Proposition 2.3.4. For the second condition [CR09, Theorem 1.2.3](2) we need to show that for the 0-point $\iota_0 : \mathrm{Spec}(k) \rightarrow \mathbb{P}^1$ and the ∞ -point $\iota_\infty : \mathrm{Spec}(k) \rightarrow \mathbb{P}^1$ the following equality holds:

$$P\hat{H}_*(\iota_0) \circ e = P\hat{H}_*(\iota_\infty) \circ e.$$

By 3.2.3(1) the left hand side is $\hat{c}l(\{0\})$ and the right hand side equals $\hat{c}l(\{\infty\})$. In view of 3.2.3(3) we obtain

$$\hat{c}l(\{0\}) = c_1(\mathcal{O}_{\mathbb{P}^1}(1)) = \hat{c}l(\{\infty\}).$$

For $W \subset X$ an irreducible closed subset, we set $cl_{(X,W)} := \hat{c}l(W)$. Then 3.2.3(1) implies that condition [CR09, Theorem 1.2.3](4) holds.

Finally, we prove [CR09, Theorem 1.2.3](3). Let $\iota : X \rightarrow Y$ be a closed immersion between smooth schemes, and let $D \subset Y$ be an effective smooth divisor such that

- D meets X properly, thus $D \cap X := D \times_Y X$ is a divisor on X ,
- $D' := (D \cap X)_{red}$ is smooth and connected, and thus $D \cap X = n \cdot D'$ as divisors (for some $n \in \mathbb{Z}, n \geq 1$).

We denote by $\iota_X : X \rightarrow (Y, X), \iota_{D'} : D' \rightarrow (D, D')$ the morphisms in Sm_* induced by ι , and we define $g : (D, D') \rightarrow (Y, X)$ in Sm^* by the inclusion $D \subset Y$. Then the following equality is required to hold:

$$(3.4.2) \quad P\hat{H}^*(g)(P\hat{H}_*(\iota_X)(1_X)) = n \cdot P\hat{H}_*(\iota_{D'})(1_{D'}).$$

Obviously, we may assume that X and Y are connected; we set $c := \mathrm{codim}_Y X$. A priori, we need to prove the equality (3.4.2) in $H_{D'}^c(D, W\Omega_D^c)$. Since $\hat{H}^*(g), \hat{H}_*(\iota_X)$ and $\hat{H}_*(\iota_{D'})$ are morphisms in $\widehat{\mathrm{dRW}}_k$, they commute with Frobenius. Thus both sides of (3.4.2) are already contained in the part which is invariant under the Frobenius $H_{D'}^c(D, W\Omega_D^c)^F$. Denote by $f : (D, D') \rightarrow (Y, D')$ the morphism in Sm_* which is induced by the inclusion $D \subset Y$.

We claim that

$$P\hat{H}_*(f) : H_{D'}^c(D, W\Omega_D^c) \rightarrow H_{D'}^{c+1}(Y, W\Omega_Y^{c+1})$$

is injective on $H_{D'}^c(D, W\Omega_D^c)^F$.

Indeed, by [Ill79, I, Thm. 5.7.2] there is an exact sequence of pro-sheaves on $D_{\acute{e}t}$

$$0 \rightarrow W_\bullet \Omega_{D, \log}^c \rightarrow W_\bullet \Omega_D^c \xrightarrow{1-F} W_\bullet \Omega_D^c \rightarrow 0.$$

This yields an exact sequence of pro-groups

$$H_{D'}^{c-1}(D, W_{\bullet}\Omega_D^c) \rightarrow H_{D'}^c(D_{\acute{e}t}, W_{\bullet}\Omega_{D,\log}^c) \rightarrow H_{D'}^c(D, W_{\bullet}\Omega_D^c)^F \rightarrow 0.$$

(Notice that $H_{D'}^i(D, W_{\bullet}\Omega_D^q) = H_{D'}^i(D_{\acute{e}t}, W_{\bullet}\Omega_D^q)$, since the $W_n\Omega_D^q$ are quasi-coherent on W_nD and $D_{\acute{e}t} = (W_nD)_{\acute{e}t}$.) But $H_{D'}^{c-1}(D, W_{\bullet}\Omega_D^c) = 0$ (see e.g. (3.4.1)) and thus

$$\varprojlim_n H_{D'}^c(D_{\acute{e}t}, W_{\bullet}\Omega_{D,\log}^c) = \varprojlim_n H_{D'}^c(D, W_{\bullet}\Omega_D^c)^F \stackrel{(3.2.3)}{=} H_{D'}^c(D, W_{\bullet}\Omega_D^c)^F.$$

Combining this with Gros' Gysin isomorphism (3.2.2) we obtain an isomorphism

$$\mathbb{Z}_p \xrightarrow{\cong} H_{D'}^c(D, W_{\bullet}\Omega_D^c)^F, \quad 1 \mapsto \hat{c}l(D') \stackrel{3.2.3, (1)}{=} P\hat{H}_*(\iota_{D'})(1_{D'}).$$

In the same way we also obtain an isomorphism

$$\mathbb{Z}_p \xrightarrow{\cong} H_{D'}^{c+1}(Y, W_{\bullet}\Omega_Y^{c+1})^F, \quad 1 \mapsto \hat{c}l(D') \stackrel{3.2.3, (1)}{=} P\hat{H}_*(f) \circ P\hat{H}_*(\iota_{D'})(1_{D'}).$$

This yields the claim.

Thus it suffices to prove the following equality

$$P\hat{H}_*(f)P\hat{H}^*(g)(P\hat{H}_*(\iota_X)(1_X)) = n \cdot P\hat{H}_*(f)P\hat{H}_*(\iota_{D'})(1_{D'}).$$

For this denote by $g' : D \rightarrow (Y, D)$ in Sm_* and $\Delta_Y : (Y, D') \rightarrow (Y \times Y, D \times X)$ in Sm^* the morphisms induced by the inclusion and the diagonal respectively. Then

$$\begin{aligned} P\hat{H}_*(f)P\hat{H}^*(g)(P\hat{H}_*(\iota_X)(1_X)) &= P\hat{H}_*(g')(1_D) \cup P\hat{H}_*(\iota_X)(1_X) \\ &= P\hat{H}^*(\Delta_Y)(\hat{c}l(D) \times \hat{c}l(X)) && 3.2.3, (1) \\ &= \hat{c}l(D.X) && 3.2.3, (2) \\ &= n \cdot \hat{c}l(D') \\ &= n \cdot P\hat{H}_*(f \circ \iota_{D'})(1_{D'}) && 3.2.3, (1), \\ &= n \cdot P\hat{H}_*(f)P\hat{H}_*(\iota_{D'})(1_{D'}) \end{aligned}$$

where $a \cup b := P\hat{H}^*(\Delta_Y)(T(a \otimes b))$ and hence the first equality holds by the projection formula, see [CR09, Proposition 1.1.11]. This finishes the proof. \square

DEFINITION 3.4.4. We denote by

$$\hat{c}l : (\text{CH}_*, \text{CH}^*, \times, 1) \rightarrow (\hat{H}_*, \hat{H}^*, T, e)$$

the composition of the morphism in Theorem 3.4.3 and Corollary 3.3.12. Note that there is no conflict with the notation in section 3.2.

3.4.5. To a weak cohomology theory $F = (F_*, F^*, T, e)$ we can attach a graded additive category Cor_F ([CR09, Definition 1.3.5]). By definition we have $\text{ob}(Cor_F) = \text{ob}(Sm_*) = \text{ob}(Sm^*)$, and

$$\text{Hom}_{Cor_F}((X, \Phi), (Y, \Psi)) = F(X \times Y, P(\Phi, \Psi)),$$

where

$$P(\Phi, \Psi) := \{Z \subset X \times Y; Z \text{ is closed, } \text{pr}_2 \mid Z \text{ is proper, } \\ Z \cap \text{pr}_1^{-1}(W) \in \text{pr}_2^{-1}(\Psi) \text{ for every } W \in \Phi\}.$$

The composition for correspondences is as usual:

$$b \circ a = F_*(p_{13})(F^*(p_{12})(a) \cup F^*(p_{23})(b)),$$

with $p_{ij} : X_1 \times X_2 \times X_3 \rightarrow X_i \times X_j$ the projection; the product \cup is defined by $a \cup b := F^*(\Delta)(T(a \otimes b))$ (see [CR09, 1.3] for the details).

Moreover, there is a functor

$$\rho_F : Cor_F \rightarrow (\text{Ab})$$

to abelian groups defined by

$$\rho_F(X, \Phi) = F(X, \Phi) \\ \rho_F(\gamma) = (a \mapsto F_*(\text{pr}_2)(F^*(\text{pr}_1)(a) \cup \gamma)), \quad \text{for } \gamma \in F(X \times Y, P(\Phi, \Psi)),$$

(see [CR09, 1.3.9]).

The morphism

$$\hat{c}l : \text{CH} = (\text{CH}_*, \text{CH}^*, \times, 1) \rightarrow \hat{H} = (\hat{H}_*, \hat{H}^*, T, e)$$

(Definition 3.4.4) induces a functor (see [CR09, 1.3.6])

$$Cor(\hat{c}l) : Cor_{\text{CH}} \rightarrow Cor_{\hat{H}}.$$

For $(X, \Phi) \in Cor_{\hat{H}}$, the group $\rho_{\hat{H}}(X, \Phi) = \hat{H}(X, \Phi)$ is a de Rham-Witt module over k (i.e. an object in $\widehat{\text{dRW}}_k$). The next theorem states that the Chow correspondences act as morphism of de Rham-Witt modules.

THEOREM 3.4.6. *The composition*

$$Cor_{\text{CH}} \rightarrow Cor_{\hat{H}} \xrightarrow{\rho_{\hat{H}}} (\text{Ab})$$

induces a functor

$$(3.4.3) \quad Cor_{\hat{c}l} : Cor_{\text{CH}} \rightarrow \widehat{\text{dRW}}_k.$$

Proof. Let $(X, \Phi), (Y, \Psi) \in Sm$. Note that $Cor_{\text{CH}}((X, \Phi), (Y, \Psi))$ is graded and is only non-trivial in even degrees. By definition

$$Cor_{\text{CH}}^{2i}((X, \Phi), (Y, \Psi)) = \text{CH}^{\dim X+i}(X \times Y, P(\Phi, \Psi)),$$

(we may assume that X is equidimensional), where $\text{CH}^{\dim X+i}(X \times Y, P(\Phi, \Psi))$ is generated by cycles Z with $\text{codim}_{X \times Y} Z = \dim X + i$, as in 3.3.6.

We will show that a cycle $Z \in Cor_{\text{CH}}^{2i}((X, \Phi), (Y, \Psi))$ defines a morphism

$$\rho_{\hat{H}}(\hat{c}l(Z)) : \hat{H}(X, \Phi)(-i) \rightarrow \hat{H}(Y, \Psi)$$

in \widehat{dRW}_k (the shift by $-i$ concerns only the grading and the differential, which is multiplied with $(-1)^i$).

Because pullback is a morphism of de Rham-Witt modules (Definition 2.2.1) and pushforward is a morphism of de Rham-Witt modules up to the shift with the relative dimension (Definition 2.3.1), we can easily reduce the statement to the following claim. For all (X, Φ) and all irreducible closed subsets $Z \subset X$ of codimension c the map

$$\begin{aligned} \hat{H}(X, \Phi) &\rightarrow \hat{H}(X \times X, \Phi \times \Phi_Z)(c) \\ a &\mapsto T(a \otimes \hat{c}l(Z)) \end{aligned}$$

is a morphism in \widehat{dRW}_k . The compatibility with F and V follows immediately from Proposition 3.1.7(5) and the invariance $F(\hat{c}l(Z)) = \hat{c}l(Z)$ (results of Gros 3.2.3(4)). The compatibility with d also follows from Proposition 3.1.7(5) and $d(\hat{c}l(Z)) = 0$ (3.2.3(4)):

$$\begin{aligned} T(da \otimes \hat{c}l(Z)) &\stackrel{\text{dfn}}{=} (-1)^{\deg(da) \cdot c} da \times \hat{c}l(Z) && (3.3.6), \\ &= (-1)^{(\deg(a)+1) \cdot c} d(a \times \hat{c}l(Z)) && 3.1.7(5), 3.2.3(4), \\ &= (-1)^c dT(a \otimes \hat{c}l(Z)) && (3.3.6). \end{aligned}$$

□

PROPOSITION 3.4.7. (i) For every $f : (X, \Phi) \rightarrow (Y, \Psi)$ in Sm^* the transpose of the graph $\Gamma(f)^t \subset Y \times X$ defines an element in $Cor_{CH}^0((Y, \Psi), (X, \Phi))$ (i.e. has degree 0). The morphism

$$Cor_{\hat{c}l}(\Gamma(f)^t) : \hat{H}(Y, \Psi) \rightarrow \hat{H}(X, \Phi)$$

is the same as f^* in Definition 2.2.1.

(ii) For every $f : (X, \Phi) \rightarrow (Y, \Psi)$ in Sm_* the graph $\Gamma(f) \subset X \times Y$ defines an element in $Cor_{CH}^{-2r}((X, \Phi), (Y, \Psi))$ (i.e. has degree $-2r = 2(\dim(Y) - \dim(X))$). The morphism

$$Cor_{\hat{c}l}(\Gamma(f)) : \hat{H}(X, \Phi)(r) \rightarrow \hat{H}(Y, \Psi)$$

is the same as f_* in Definition 2.3.1.

Proof. For (i). Let $i_2 : (X, f^{-1}\Psi) \rightarrow (Y \times X, \text{pr}_Y^{-1}\Psi)$ be the morphism in Sm^* induced by the morphism of schemes (f, id_X) . Let $i_1 : X \rightarrow (Y \times X, \Gamma(f)^t)$ and $i_3 : (X, f^{-1}\Psi) \rightarrow (Y \times X, \Gamma(f)^t \cap \text{pr}_Y^{-1}\Psi)$ be the morphism in Sm_* induced by (f, id_X) . For all $a \in \hat{H}(Y, \Psi)$ we use the projection formula [CR09, Proposition 1.1.11] to obtain:

$$\begin{aligned} Cor_{\hat{c}l}(\Gamma(f)^t)(a) &= \hat{H}_*(\text{pr}_X)(\hat{H}^*(\text{pr}_Y)(a) \cup \hat{c}l(\Gamma(f)^t)) \quad \text{by definition,} \\ &= \hat{H}_*(\text{pr}_X)(\hat{H}^*(\text{pr}_Y)(a) \cup \hat{H}_*(i_1)(1)) \quad \text{functoriality of } \hat{c}l \text{ (cf. 3.4.4)} \\ &= \hat{H}_*(\text{pr}_X)(\hat{H}_*(i_3)(\hat{H}^*(i_2)\hat{H}^*(\text{pr}_Y)(a))) \quad \text{projection formula,} \\ &= \hat{H}_*(\text{pr}_X)(\hat{H}_*(i_3)(\hat{H}^*(f)(a))) \\ &= \hat{H}^*(f)(a). \end{aligned}$$

The proof of (ii) is similar. □

3.5. CORRESPONDENCE ACTION ON RELATIVE HODGE-WITT COHOMOLOGY. For a scheme S over k we need a relative version of the functor (3.4.3) with values in de Rham-Witt modules $\widehat{\mathrm{dRW}}_S$ over S .

Let $f : X \rightarrow S$ and $g : Y \rightarrow S$ be two schemes over S . Suppose that X, Y are smooth over k . Since $X \times_S Y \subset X \times Y$ defines a closed subset we can define the family of supports on $X \times Y$

$$(3.5.1) \quad P(X \times_S Y) := P(\Phi_X, \Phi_Y) \cap X \times_S Y,$$

this is the family of supports consisting of all closed subsets of $X \times_S Y$, which are proper over Y . (Recall that Φ_X denotes the family of all closed subsets of X .) If $h : Z \rightarrow S$ is another S -schemes which is smooth over k , then

$$\begin{cases} p_{13} \mid p_{12}^{-1}(P(X \times_S Y)) \cap p_{23}^{-1}(P(Y \times_S Z)) \text{ is proper,} \\ p_{13}(p_{12}^{-1}(P(X \times_S Y)) \cap p_{23}^{-1}(P(Y \times_S Z))) \subset P(X \times_S Z). \end{cases}$$

By using the fact that CH defines a weak cohomology theory with supports we obtain a composition (see [CR09, 1.3] for details)

$$(3.5.2) \quad \mathrm{CH}(X \times Y, P(X \times_S Y)) \times \mathrm{CH}(Y \times Z, P(Y \times_S Z)) \xrightarrow{\circ} \mathrm{CH}(X \times Z, P(X \times_S Z)),$$

$$(a, b) \mapsto b \circ a := \mathrm{CH}_*(p_{13})(\mathrm{CH}^*(p_{12})(a) \cup \mathrm{CH}^*(p_{23})(b)),$$

with morphisms p_{ij} induced by the projections:

$$\begin{aligned} p_{12} &: (X \times Y \times Z, P(X \times_S Y) \times Z) \rightarrow (X \times Y, P(X \times_S Y)) \in \mathrm{Sm}^*, \\ p_{23} &: (X \times Y \times Z, X \times P(Y \times_S Z)) \rightarrow (Y \times Z, P(Y \times_S Z)) \in \mathrm{Sm}^*, \\ p_{13} &: (X \times Y \times Z, P(X \times_S Y) \cap P(Y \times_S Z)) \rightarrow (X \times Z, P(X \times_S Z)) \in \mathrm{Sm}_*. \end{aligned}$$

DEFINITION 3.5.1. We define \mathcal{C}_S to be the (graded) additive category whose objects are given by k -morphisms $f : X \rightarrow S$, where X is a smooth scheme over k . Sometimes we will by abuse of notation write X instead of $f : X \rightarrow S$. The morphisms are defined by

$$\mathrm{Hom}_{\mathcal{C}_S}(X, Y) := \mathrm{CH}(X \times Y, P(X \times_S Y)),$$

and the composition is as in (3.5.2). For $f : X \rightarrow S$ the identity element in $\mathrm{Hom}_{\mathcal{C}_S}(X, X)$ is the diagonal Δ_X .

The verification that the composition defines a category is a straightforward calculation. Notice that if $X \rightarrow S$ and $Y \rightarrow S$ are proper, then $\mathrm{Hom}_{\mathcal{C}_S}(X, Y) = \mathrm{CH}(X \times_S Y)$. The full subcategory of \mathcal{C}_S whose objects are proper S -schemes, which are smooth over k , has been defined and studied in [CH00, Definition 2.8].

3.5.2. Let $a \in \text{Hom}_{\mathcal{C}_S}(f : X \rightarrow S, g : Y \rightarrow S) = \text{CH}(X \times Y, P(X \times_S Y))$, with X and Y integral. Suppose that a is of degree i , i.e. is contained in $\text{CH}^{\dim X+i}$. For every open subset $U \subset S$ we get by restriction an induced cycle

$$a_U \in \text{CH}(f^{-1}(U) \times g^{-1}(U), P(f^{-1}(U) \times_U g^{-1}(U))).$$

By using the functor Cor_{cl} (3.4.3) we obtain a morphism in $\widehat{\text{dRW}}_k$:

$$\text{Cor}_{\text{cl}}(a_U) : \hat{H}(f^{-1}(U))(-i) \rightarrow \hat{H}(g^{-1}(U)).$$

Recall that $\hat{H}(f^{-1}(U)) = \bigoplus_{p,q \geq 0} H^q(f^{-1}(U), W\Omega_{f^{-1}(U)}^p)$; we denote by

$$U \mapsto \mathcal{F}(U) = \hat{H}(f^{-1}(U))(-i), \quad U \mapsto \mathcal{G}(U) := \hat{H}(g^{-1}(U)),$$

the obvious presheaves on S .

LEMMA 3.5.3. *Let $a \in \text{Hom}_{\mathcal{C}_S}(f : X \rightarrow S, g : Y \rightarrow S) = \text{CH}(X \times Y, P(X \times_S Y))$. The collection $(\text{Cor}_{\text{cl}}(a_U))_{U \subset S}$ defines a morphism of presheaves*

$$(\text{Cor}_{\text{cl}}(a_U))_{U \subset S} : \mathcal{F} \rightarrow \mathcal{G}.$$

By sheafification we obtain a morphism of sheaves

$$(3.5.3) \quad (\text{Cor}_{\text{cl}}(a_U))_{U \subset S} : \bigoplus_{p,q \geq 0} R^q f_* W\Omega_X^p(-i) \rightarrow \bigoplus_{p,q \geq 0} R^q g_* W\Omega_Y^p.$$

Proof. Let U, V be two open subsets of S such that $V \subset U$. We denote by $\Gamma(f^{-1}(V) \rightarrow f^{-1}(U))^t \subset f^{-1}(U) \times f^{-1}(V)$ the transpose of the graph of the inclusion $f^{-1}(V) \rightarrow f^{-1}(U)$, and similarly for $\Gamma(g^{-1}(V) \rightarrow g^{-1}(U))^t$. Since the restrictions $\mathcal{F}(U) \rightarrow \mathcal{F}(V)$ and $\mathcal{G}(U) \rightarrow \mathcal{G}(V)$ are induced by the correspondences $\text{Cor}_{\text{cl}}(\Gamma(f^{-1}(V) \rightarrow f^{-1}(U))^t)$ and $\text{Cor}_{\text{cl}}(\Gamma(g^{-1}(V) \rightarrow g^{-1}(U))^t)$, respectively, we only need to prove that

$$\Gamma(g^{-1}(V) \rightarrow g^{-1}(U))^t \circ a_U = a_V \circ \Gamma(f^{-1}(V) \rightarrow f^{-1}(U))^t$$

as morphisms in Cor_{CH} , i.e. as cycles in

$$\text{CH}(f^{-1}(U) \times g^{-1}(V), P(f^{-1}(U) \times_S g^{-1}(V))).$$

Via the identification $f^{-1}(U) \times_S g^{-1}(V) = f^{-1}(V) \times_S g^{-1}(V)$ both sides equal a_V .

Since the sheafification of \mathcal{F} is $\bigoplus_{p,q \geq 0} R^q f_* W\Omega_X^p(-i)$, and similarly for \mathcal{G} , we obtain (3.5.3). □

PROPOSITION 3.5.4. *The assignment*

$$\begin{aligned} \hat{\mathcal{H}}(?/S) &: \mathcal{C}_S \rightarrow \widehat{\text{dRW}}_S, \\ \hat{\mathcal{H}}(X/S) &:= \bigoplus_{p,q \geq 0} R^q f_* W\Omega_X^p, \\ \hat{\mathcal{H}}(a/S) &:= (\text{Cor}_{\text{cl}}(a_U))_{U \subset S}, \end{aligned}$$

(cf. Lemma 3.5.3) defines a functor to de Rham-Witt modules over S .

For the proof of the Proposition we will need the following lemma.

LEMMA 3.5.5. *Let $S = \text{Spec}(R)$. Let X, Y be S -schemes which are smooth over k . We denote by $f^* : W(R) \rightarrow H^0(X, W\mathcal{O}_X)$ and $g^* : W(R) \rightarrow H^0(Y, W\mathcal{O}_Y)$ the maps induced by $f : X \rightarrow S$ and $g : Y \rightarrow S$, respectively. We denote by $\text{pr}_1 : X \times Y \rightarrow X$ and $\text{pr}_2 : X \times Y \rightarrow Y$ the projections.*

Let $Z \subset X \times_S Y$ be an irreducible closed subset which is proper over Y ; we set $c = \text{codim}_{X \times Y} Z$. Then

$$\hat{H}^*(\text{pr}_2)(g^*(r)) \cup \hat{c}l([Z]) = \hat{H}^*(\text{pr}_1)(f^*(r)) \cup \hat{c}l([Z])$$

in $H_Z^c(X \times Y, W\Omega_{X \times Y}^c)$.

Proof. Choose an open set $U \subset X \times Y$ such that $Z \cap U$ is nonempty and smooth. Since the natural map

$$H_Z^c(X \times Y, W\Omega_{X \times Y}^c) \rightarrow H_{Z \cap U}^c(U, W\Omega_U^c)$$

is injective (Proposition 3.4.1), it suffices to check the equality on $H_{Z \cap U}^c(U, W\Omega_U^c)$. We write $\iota_1 : Z \cap U \rightarrow (U, Z \cap U)$ in Sm_* and $\iota_2 : Z \cap U \rightarrow U$ in Sm^* for the obvious morphisms. By using the projection formula and $\hat{c}l([Z \cap U]) = \hat{H}_*(\iota_1)(1)$ (3.2.3(1)) we reduce to the statement

$$\hat{H}^*(\iota_2)\hat{H}^*(\text{pr}_2)(g^*(r)) = \hat{H}^*(\iota_2)\hat{H}^*(\text{pr}_1)(f^*(r)).$$

This follows from $g \circ \text{pr}_2 \circ \iota_2 = f \circ \text{pr}_1 \circ \iota_1$. □

Proof of Proposition 3.5.4. For two composable morphisms a, b in \mathcal{C}_S , we clearly have

$$(b \circ a)_U = b_U \circ a_U$$

for every open $U \subset S$. This implies $\hat{H}(b \circ a/S) = \hat{H}(b/S) \circ \hat{H}(a/S)$.

Moreover, $\text{Cor}_{\hat{c}l}(a_U)$ is a morphism in $\widehat{\text{dRW}}_k$ for all U , thus $\hat{H}(a/S)$ commutes with F, V, d . Finally, we need to show that $\hat{H}(a/S)$ induces a morphism of $W(\mathcal{O}_S)$ -modules. For this, we may assume that $S = \text{Spec}(R)$ is affine, and it suffices to show that

$$\text{Cor}_{\hat{c}l}(a) : \hat{H}(X)(-i) \rightarrow \hat{H}(Y)$$

is $W(R)$ -linear. We proceed as in the proof of [CR09, Proposition 3.2.4]. The ring homomorphism $f^* : W(R) \rightarrow H^0(X, W\mathcal{O}_X)$ and $g^* : W(R) \rightarrow H^0(Y, W\mathcal{O}_Y)$ induce the $W(R)$ -module structures on $\hat{H}(X)$ and $\hat{H}(Y)$ via the \cup -product:

$$r \cdot a = f^*(r) \cup a,$$

for all $r \in W(R)$ and $a \in \hat{H}(X)$; similarly for $\hat{H}(Y)$. We have to prove the following equality for all $r \in R, a \in \hat{H}(X)$, and irreducible closed subsets $Z \subset X \times_S Y$:

$$g^*(r) \cup \hat{H}_*(\text{pr}_2)(\hat{H}^*(\text{pr}_1)(a) \cup \hat{c}l([Z])) = \hat{H}_*(\text{pr}_2)(\hat{H}^*(\text{pr}_1)(f^*(r) \cup a) \cup \hat{c}l([Z])).$$

For this, it is enough to show that

$$\hat{H}^*(\text{pr}_2)(g^*(r)) \cup \hat{c}l([Z]) = \hat{H}^*(\text{pr}_1)(f^*(r)) \cup \hat{c}l([Z])$$

in $H_Z^c(X \times Y, W\Omega_{X \times Y}^c)$, with c the codimension of Z . Lemma 3.5.5 implies the claim. \square

PROPOSITION 3.5.6. *Let X and Y be two S -schemes which are smooth over k .*

- (i) *If Z is the transpose of the graph of a morphism $h : X \rightarrow Y$ over S then $\hat{\mathcal{H}}([Z]/S)$ is the pullback morphism defined in 2.2.1.*
- (ii) *If Z is the graph of a proper morphism $h : X \rightarrow Y$ over S then $\hat{\mathcal{H}}([Z]/S)$ is the pushforward morphism defined in 2.3.1.*

Proof. The statement follows from Proposition 3.4.7. \square

3.5.7. *Local cup product.* Let X be a smooth equidimensional k -scheme and $Z \subset X$ an integral closed subscheme of codimension c . We have (see e.g. (3.4.1))

$$\mathcal{H}_Z^i(X, W\Omega_X) = 0, \quad \text{for all } i < c.$$

Hence there is a natural morphism in the derived category of de Rham-Witt modules on X

$$(3.5.4) \quad \mathcal{H}_Z^c(W\Omega_X) \rightarrow R\underline{\Gamma}_Z W\Omega_X[c]$$

inducing an isomorphism

$$(3.5.5) \quad H_Z^c(X, W\Omega_X) \cong H^0(X, \mathcal{H}_Z^c(W\Omega_X)).$$

We may thus define a local version of the cup product with the cycle class of Z , $\hat{c}l(Z) \in H^0(X, \mathcal{H}_Z^c(W\Omega_X))$

$$(3.5.6) \quad W\Omega_X \rightarrow \mathcal{H}_Z^c(W\Omega_X)(c), \quad \alpha \mapsto \alpha \cup \hat{c}l([Z]) := \Delta^*((-1)^{c \cdot \deg \alpha}(\alpha \times \hat{c}l([Z])))$$

as the composition

$$W\Omega_X \xrightarrow{\text{nat.}} \text{pr}_{1*} \text{pr}_1^{-1} W\Omega_X \xrightarrow{(-1)^{c \cdot \deg \alpha}(-\times \hat{c}l([Z]))} \text{pr}_{1*} \mathcal{H}_{X \times Z}^c(W\Omega_{X \times X})(c) \xrightarrow{\Delta^*} \text{pr}_{1*} \Delta_* \mathcal{H}_Z^c(W\Omega_X)(c) = \mathcal{H}_Z^c(W\Omega_X)(c)$$

where $\Delta : (X, Z) \rightarrow (X \times X, X \times Z)$ in $(Sm^*/X \times X)$ is induced by the diagonal, Δ^* is the pullback constructed in 2.2.1 and \times is the exterior product from 3.1.4. Notice that by Lemma 3.1.5 $\alpha \cup \hat{c}l(Z)$ equals $(-1)^{c \cdot \deg \alpha} \alpha \cdot \hat{c}l([Z])$.

LEMMA 3.5.8. *In the above situation the cup product with $\hat{c}l([Z])$,*

$$H^i(X, W\Omega_X) \rightarrow H_Z^{i+c}(X, W\Omega_X)(c), \quad a \mapsto a \cup \hat{c}l([Z]),$$

factors via the local cup product, i.e. equals the composition

$$H^i(X, W\Omega_X) \xrightarrow{(3.5.6)} H^i(X, \mathcal{H}_Z^c(W\Omega_X)(c)) \xrightarrow{(3.5.4)} H_Z^{i+c}(X, W\Omega_X)(c).$$

Proof. Recall that for $a \in H^i(X, W\Omega_X^q)$ the cup product $a \cup \hat{c}l([Z])$ equals

$$H^*(\Delta)((-1)^{(i+q)c}(a \times \hat{c}l([Z])),$$

where $\Delta : (X, Z) \rightarrow (X \times X, X \times Z)$ in $(Sm^*/\text{Spec } k)$ is the diagonal. Therefore it suffices to show that the exterior product

$$(3.5.7) \quad \times \hat{c}l([Z]) : H^i(X, W\Omega_X^q) \rightarrow H_{X \times Z}^{i+c}(X \times X, W\Omega_{X \times X}^{q+c})$$

factors via $H^i(X, -)$ applied to

$$\text{pr}_1^{-1}W\Omega_X^q \rightarrow \mathcal{H}_{X \times Z}^c(W\Omega_{X \times X}), \quad \alpha \mapsto (-1)^{ic}(\alpha \times \hat{c}l([Z])),$$

composed with the natural map $\mathcal{H}_{X \times Z}^c(W\Omega_{X \times X}) \rightarrow R\Gamma_{X \times Z}W\Omega_{X \times X}[c]$. Let $E(W_\bullet\Omega_X)$ be the Cousin complex of $W_\bullet\Omega_X$ (see 1.5.8, Lemma 1.5.9), then the complex $\Gamma_Z E(W_\bullet\Omega_X)$ equals zero in all degrees $< c$, hence there is a morphism of complexes

$$(3.5.8) \quad \mathcal{H}_Z^c(W_\bullet\Omega_X) \rightarrow \Gamma_Z E(W_\bullet\Omega_X^c)[c],$$

which, after applying \varprojlim to it, represents (3.5.4). We obtain the following commutative diagram

$$\begin{array}{ccccc} \text{pr}_1^{-1}E(W_\bullet\Omega_X^q) & \xrightarrow{\boxtimes \hat{c}l([Z])} & E(W_\bullet\Omega_X^q) \boxtimes \mathcal{H}_Z^c(W_\bullet\Omega_X^c) & \longrightarrow & \Gamma_{X \times Z}E(W_\bullet\Omega_{X \times X}^{q+c})[c] \\ \uparrow & & \uparrow & & \uparrow (3.5.8) \\ \text{pr}_1^{-1}W_\bullet\Omega_X^q & \xrightarrow{\boxtimes \hat{c}l([Z])} & W_\bullet\Omega_X^q \boxtimes \mathcal{H}_Z^c(W_\bullet\Omega_X^c) & \xrightarrow{\times} & \mathcal{H}_{X \times Z}^c(W_\bullet\Omega_{X \times X}^{q+c}), \end{array}$$

where the top right arrow is the composition

$$\begin{aligned} E(W_\bullet\Omega_X^q) \boxtimes \mathcal{H}_Z^c(W_\bullet\Omega_X^c) &\xrightarrow{\text{id} \boxtimes (3.5.8)} E(W_\bullet\Omega_X^q) \boxtimes \Gamma_Z E(W_\bullet\Omega_X^c)[c] \\ &\xrightarrow{\cong} (E(W_\bullet\Omega_X^q) \boxtimes \Gamma_Z E(W_\bullet\Omega_X^c))[c] \xrightarrow{(3.1.4)} \Gamma_{X \times Z}E(W_\bullet\Omega_{X \times X}^{q+c})[c]. \end{aligned}$$

Notice that the isomorphism

$$(3.5.9) \quad (E \boxtimes \Gamma_Z E[c]) \cong (E \boxtimes \Gamma_Z E)[c]$$

is given by multiplication with $(-1)^{ic}$ on $E^i \boxtimes \Gamma_Z E^j$, for all i, j (see the sign convention in [Con00, (1.3.6)]). Now we apply $H^i(X, -) \circ \varprojlim$ to the above diagram, use Proposition 3.1.7, (1), take care about the sign from (3.5.9) and use (3.2.3) and we see that (3.5.7) factors as desired. \square

3.5.9. Let $h : S \rightarrow T$ be a morphism of k -schemes. Then any two objects $X, Y \in \mathcal{C}_S$ naturally define objects in \mathcal{C}_T (via h) and $X \times_S Y \subset X \times_T Y$ is a closed subscheme. This gives a natural map $\text{CH}(X \times Y, P(X \times_S Y)) \rightarrow \text{CH}(X \times Y, P(X \times_T Y))$. In this way h induces a functor

$$\mathcal{C}_S \rightarrow \mathcal{C}_T.$$

If h is fixed, we denote the image of $a \in \text{Hom}_{\mathcal{C}_S}(X, Y)$ in $\text{Hom}_{\mathcal{C}_T}(X, Y)$ via this functor again by a . But notice that this functor is in general not faithful.

PROPOSITION 3.5.10. *Let $h : S \rightarrow T$ be a morphism of k -schemes. Let $f : X \rightarrow S$ and $g : Y \rightarrow S$ be two objects in \mathcal{C}_S and assume that X and Y are integral and f and g are affine. Let $Z \subset X \times_S Y$ be a closed integral subscheme which*

is finite and surjective over Y , therefore giving rise to a morphism in $\widehat{\mathrm{dRW}}_S$ (by Proposition 3.5.4)

$$\hat{\mathcal{H}}([Z]/S) : f_* W\Omega_X \rightarrow g_* W\Omega_Y.$$

Then we have the following equality of morphisms in $\widehat{\mathrm{dRW}}_T$

$$\hat{\mathcal{H}}([Z]/T) = \bigoplus_i R^i h_* \hat{\mathcal{H}}([Z]/S) : \bigoplus_i R^i (hf)_* W\Omega_X \rightarrow \bigoplus_i R^i (hg)_* W\Omega_Y.$$

Proof. We consider the following composition in the derived category of abelian sheaves on S :

$$\begin{aligned} (3.5.10) \quad f_* W\Omega_X &\xrightarrow{\mathrm{pr}_1^*} R(f\mathrm{pr}_1)_* W\Omega_{X \times Y} \\ &\xrightarrow{(3.5.6)} R(f\mathrm{pr}_1)_* \mathcal{H}_Z^c(W\Omega_{X \times Y})(c) \\ &\xrightarrow{\simeq, Z \subset X \times_S Y} R(g\mathrm{pr}_2)_* \mathcal{H}_Z^c(W\Omega_{X \times Y})(c) \\ &\xrightarrow{(3.5.4)} g_* R\mathrm{pr}_{2*} R\underline{\Gamma}_Z(W\Omega_{X \times Y})c \\ &\xrightarrow{(2.3.4)} g_* W\Omega_Y. \end{aligned}$$

Notice that the third arrow only exists in the category of abelian sheaves, it is not respecting the $W\mathcal{O}_S$ -module structure. We claim that the composition (3.5.10) equals $\hat{\mathcal{H}}([Z]/S)$ and that $\bigoplus_i R^i h_*$ (3.5.10) equals $\hat{\mathcal{H}}([Z]/T)$. This will prove the statement. Clearly it suffices to prove the last claimed equality, the first then follows with $h = \mathrm{id}$.

To this end, let $U \subset T$ be an open subset. We denote by X_U , Y_U and Z_U the pullbacks of X , Y and Z over U . Then $\mathrm{Cor}_{\mathrm{el}}([Z_U]) : H^i(X_U, W\Omega_{X_U}) \rightarrow H^i(Y_U, W\Omega_{Y_U})(c)$ is given by the following composition

$$\begin{aligned} H^i(X_U, W\Omega_{X_U}) &\xrightarrow{\mathrm{pr}_1^*} H^i(X_U, R\mathrm{pr}_{1*} W\Omega_{X_U \times Y_U}) \\ &\xrightarrow{\simeq} H^i(Y_U, R\mathrm{pr}_{2*} W\Omega_{X_U \times Y_U}) \\ &\xrightarrow{\cup_{\mathrm{el}}([Z_U])} H^i(Y_U, R\mathrm{pr}_{2*} R\underline{\Gamma}_{Z_U} W\Omega_{X_U \times Y_U}c) \\ &\xrightarrow{(2.3.4)} H^i(Y_U, W\Omega_{Y_U}). \end{aligned}$$

Since f and g are affine this composition equals by Lemma 3.5.8 the composition $H^i(h^{-1}(U), (3.5.10))$. By definition $\hat{\mathcal{H}}([Z]/T)$ is the sheafification of $U \mapsto \mathrm{Cor}_{\mathrm{el}}([Z_U])$ and the sheafification of $U \mapsto H^i(h^{-1}(U), (3.5.10))$ is $R^i h_*$ ((3.5.10)). This proves the claim. \square

3.6. VANISHING RESULTS. Recall from Proposition 3.5.4 that we have a functor

$$\begin{aligned} \hat{\mathcal{H}}(?/S) : \mathcal{C}_S &\rightarrow \widehat{\mathrm{dRW}}_S, \\ \hat{\mathcal{H}}(X/S) &:= \bigoplus_{p,q \geq 0} R^q f_* W\Omega_X^p \end{aligned}$$

(cf. Definition 3.5.1 for \mathcal{C}_S). Let $f : X \rightarrow S, g : Y \rightarrow S$ be objects in \mathcal{C}_S , i.e. S -schemes which are smooth over k . For simplicity let us assume that X and Y are connected. For an equidimensional closed subset $Z \subset X \times_S Y$ which is proper over Y and has codimension $\dim X + i$ in $X \times Y$ we obtain a morphism in $\widehat{\text{dRW}}_S$:

$$\hat{\mathcal{H}}([Z]/S) : \left(\bigoplus_{p,q \geq 0} R^q f_* W\Omega_X^p \right) (-i) \rightarrow \bigoplus_{p,q \geq 0} R^q g_* W\Omega_Y^p.$$

LEMMA 3.6.1. *Let $Z \subset X \times_S Y$ be closed subset which is proper over Y . Let $r \geq 0$ be an integer. Suppose that for every point $z \in Z$ the image $\text{pr}_2(z)$ is a point of codimension $\geq r$ in Y . Then there is a natural number $N \geq 1$ such that*

$$N \cdot \text{image}(\hat{\mathcal{H}}([Z]/S)) \subset \bigoplus_{p \geq r, q \geq r} R^q g_* W\Omega_Y^p.$$

In other words, the projection of $\text{image}(\hat{\mathcal{H}}([Z]/S))$ to

$$\bigoplus_{p < r \text{ or } q < r} R^q g_* W\Omega_Y^p$$

is killed by N .

Proof. We may assume that Z is irreducible and X, Y are connected. We set $i = \dim Y - \dim Z = \text{codim}_{X \times Y} Z - \dim X$.

By using an alteration we can find a smooth equidimensional scheme D of dimension $\dim Y - r$ together with a proper morphism $\pi : D \rightarrow Y$ such that $\pi(D) \supset \text{pr}_2(Z)$. In particular, Z is contained in the image of the map $\text{id}_X \times_S \pi : X \times_S D \rightarrow X \times_S Y$.

Let $Z_D \subset X \times_S D$ be an irreducible closed subset with $\dim(Z_D) = \dim(Z)$ and $(\text{id}_X \times \pi)(Z_D) = Z$. (Z_D is automatically proper over D .) We define N to be the degree of the field extension $k(Z) \subset k(Z_D)$.

We obtain three maps:

$$\begin{aligned} \hat{\mathcal{H}}(Z/S) &: \hat{\mathcal{H}}(X/S)(-i) \rightarrow \hat{\mathcal{H}}(Y/S), \\ \hat{\mathcal{H}}(Z_D/S) &: \hat{\mathcal{H}}(X/S)(-i) \rightarrow \hat{\mathcal{H}}(D/S)(-r), \\ \hat{\mathcal{H}}(\Gamma(\pi)/S) &: \hat{\mathcal{H}}(D/S)(-r) \rightarrow \hat{\mathcal{H}}(Y/S), \end{aligned}$$

($\Gamma(\pi)$ denotes the graph of π). We claim that

$$N \cdot \hat{\mathcal{H}}(Z/S) = \hat{\mathcal{H}}(\Gamma(\pi)/S) \circ \hat{\mathcal{H}}(Z_D/S).$$

Indeed, by functoriality it is sufficient to prove

$$N \cdot [Z] = [\Gamma(\pi)] \circ [Z_D]$$

where \circ is the composition in \mathcal{C}_S (see (3.5.2)). This is an easy computation in intersection theory.

Thus it is sufficient to show that

$$\text{image}(\hat{\mathcal{H}}([\Gamma(\pi)]/S)) \subset \bigoplus_{p \geq r, q \geq r} R^q g_* W\Omega_Y^p.$$

Proposition 3.5.6 implies that $\hat{\mathcal{H}}([\Gamma(\pi)]/S)$ is the push-forward π_* defined in 2.3.1. Thus

$$\hat{\mathcal{H}}([\Gamma(\pi)]/S)(R^q(g \circ \pi)_* W\Omega_D^p) \subset R^{q+r} g_* W\Omega_Y^{p+r},$$

for all (p, q) , which completes the proof. □

LEMMA 3.6.2. *Let $Z \subset X \times_S Y$ be closed subset, which is proper over Y . Suppose that X is equidimensional of dimension d . Let $r \geq 0$ be an integer. Suppose that for every point $z \in Z$ the image $\text{pr}_1(z)$ is a point of codimension $\geq r$ in X . Then there is a natural number $N \geq 1$ such that*

$$N \cdot \left(\bigoplus_{p > d-r \text{ or } q > d-r} R^q f_* W\Omega_X^p \right) \subset \ker(\hat{\mathcal{H}}([Z]/S)).$$

Proof. The proof is analogous to the proof of Lemma 3.6.1.

We may assume that Z is irreducible and Y connected. We set $i = \dim Y - \dim Z = \text{codim}_{X \times Y} Z - \dim X$.

By using an alteration we can find a smooth equidimensional scheme D of dimension $d - r$ together with a proper morphism $\pi : D \rightarrow X$ such that $\pi(D) \supset \text{pr}_1(Z)$. In particular, Z is contained in the image of the map $\pi \times_S \text{id}_Y : D \times_S Y \rightarrow X \times_S Y$.

Let $Z_D \subset D \times_S Y$ be an irreducible closed subset with $\dim(Z_D) = \dim(Z)$ and $(\pi \times \text{id}_Y)(Z_D) = Z$. (Z_D is automatically proper over Y .) We define N to be the degree of the field extension $k(Z) \subset k(Z_D)$.

We obtain three maps:

$$\begin{aligned} \hat{\mathcal{H}}(Z/S) &: \hat{\mathcal{H}}(X/S)(-i) \rightarrow \hat{\mathcal{H}}(Y/S), \\ \hat{\mathcal{H}}(Z_D/S) &: \hat{\mathcal{H}}(D/S)(-i) \rightarrow \hat{\mathcal{H}}(Y/S), \\ \hat{\mathcal{H}}(\Gamma(\pi)^t/S) &: \hat{\mathcal{H}}(X/S) \rightarrow \hat{\mathcal{H}}(D/S), \end{aligned}$$

$(\Gamma(\pi)^t$ denotes the transpose of the graph of π). We claim that

$$N \cdot \hat{\mathcal{H}}(Z/S) = \hat{\mathcal{H}}(Z_D/S) \circ \hat{\mathcal{H}}(\Gamma(\pi)^t/S).$$

Indeed, by functoriality it is sufficient to prove

$$N \cdot [Z] = [Z_D] \circ [\Gamma(\pi)^t]$$

where \circ is the composition in \mathcal{C}_S (see (3.5.2)). This is a straightforward calculation in intersection theory.

Because $\dim D = d - r$, the map $\hat{\mathcal{H}}(\Gamma(\pi)^t/S)$ vanishes on

$$\bigoplus_{p > d-r \text{ or } q > d-r} R^q f_* W\Omega_X^p,$$

which proves the statement. □

Remark 3.6.3. Resolution of singularities implies that Lemma 3.6.1 and 3.6.2 hold for $N = 1$.

3.7. DE RHAM-WITT SYSTEMS AND MODULES MODULO TORSION.

3.7.1. Let \mathcal{A} be an abelian category (only $\mathcal{A} = \text{dRW}_X$ and $\mathcal{A} = \widehat{\text{dRW}}_X$ will be important for us). There are two natural Serre subcategories attached to torsion objects of \mathcal{A} . We define

$$\mathcal{A}_{b\text{-tor}} := \{X \in \text{ob}(\mathcal{A}) \mid \exists n \in \mathbb{Z} \setminus \{0\} : n \cdot \text{id}_X = 0\}$$

as full subcategory of \mathcal{A} . We define

$$\mathcal{A}_{\text{tor}} := \{X \in \text{ob}(\mathcal{A}) \mid \exists (X_i)_{i \in I} \in \mathcal{A}_{b\text{-tor}}^I \exists \phi : \bigoplus_{i \in I} X_i \rightarrow X \text{ epimorphism}\}$$

as full subcategory of \mathcal{A} . Note that the index set I is not finite in general. Obviously,

$$\mathcal{A}_{b\text{-tor}} \subset \mathcal{A}_{\text{tor}}.$$

3.7.2. If \mathcal{A} is well-powered, i.e. the family of subobjects of any object is a set (e.g. $\mathcal{A} = \widehat{\text{dRW}}_X$), then the quotient categories $\mathcal{A}/\mathcal{A}_{b\text{-tor}}$ and $\mathcal{A}/\mathcal{A}_{\text{tor}}$ exist and are abelian categories. We refer to [Gab62, Chapitre III] for quotient categories. The functors

$$q : \mathcal{A} \rightarrow \mathcal{A}/\mathcal{A}_{b\text{-tor}}, \quad q' : \mathcal{A} \rightarrow \mathcal{A}/\mathcal{A}_{\text{tor}},$$

are exact. Moreover, $q(X) = 0$ if and only if $X \in \mathcal{A}_{b\text{-tor}}$; the same statement holds for q' and \mathcal{A}_{tor} . There is an obvious factorization

$$\mathcal{A} \rightarrow \mathcal{A}/\mathcal{A}_{b\text{-tor}} \rightarrow \mathcal{A}/\mathcal{A}_{\text{tor}}.$$

If \mathcal{A} is the category of (left) modules over a ring R then

$$\mathcal{A}/\mathcal{A}_{\text{tor}} \cong (R \otimes_{\mathbb{Z}} \mathbb{Q}\text{-modules}).$$

We define

$$\mathcal{A}_{\mathbb{Q}} := \mathcal{A}/\mathcal{A}_{b\text{-tor}}.$$

For future reference, we record the following special case.

DEFINITION 3.7.3. Let X be a scheme over k . We define

$$\widehat{\text{dRW}}_{X,\mathbb{Q}} := \widehat{\text{dRW}}_X / \widehat{\text{dRW}}_{X,b\text{-tor}}$$

as quotient category. We denote by q the projection functor

$$q : \widehat{\text{dRW}}_X \rightarrow \widehat{\text{dRW}}_{X,\mathbb{Q}}.$$

We use the same definitions for dRW_X .

The main reason for working with the quotient $\mathcal{A}/\mathcal{A}_{b\text{-tor}}$ instead of $\mathcal{A}/\mathcal{A}_{\text{tor}}$ is that the Homs are well-behaved.

PROPOSITION 3.7.4. *Let \mathcal{A} be a well-powered abelian category. Let $X, Y \in \text{ob}(\mathcal{A})$. Then*

$$\text{Hom}_{\mathcal{A}_{\mathbb{Q}}}(q(X), q(Y))$$

is naturally a \mathbb{Q} -module and the map

$$\text{Hom}_{\mathcal{A}}(X, Y) \otimes_{\mathbb{Z}} \mathbb{Q} \rightarrow \text{Hom}_{\mathcal{A}_{\mathbb{Q}}}(q(X), q(Y))$$

is an isomorphism.

Proof. For any $n \in \mathbb{Z} \setminus \{0\}$ the morphism $q(Y) \xrightarrow{\cdot n} q(Y)$ in $\mathcal{A}_{\mathbb{Q}}$ is invertible; therefore

$$\mathbb{Q} \subset \text{Hom}_{\mathcal{A}_{\mathbb{Q}}}(q(Y), q(Y)).$$

Via the composition

$$\text{Hom}(q(X), q(Y)) \times \text{Hom}(q(Y), q(Y)) \rightarrow \text{Hom}(q(X), q(Y))$$

we see that $\text{Hom}(q(X), q(Y))$ is a \mathbb{Q} -module. The \mathbb{Q} -module structure induced by $\mathbb{Q} \subset \text{Hom}(q(X), q(X))$ and

$$\text{Hom}(q(X), q(X)) \times \text{Hom}(q(X), q(Y)) \rightarrow \text{Hom}(q(X), q(Y))$$

is the same, because

$$f \circ (q(X) \xrightarrow{\cdot n} q(X)) = (q(Y) \xrightarrow{\cdot n} q(Y)) \circ f = nf.$$

We need to show that the canonical map

$$\text{Hom}_{\mathcal{A}}(X, Y) \otimes_{\mathbb{Z}} \mathbb{Q} \rightarrow \text{Hom}_{\mathcal{A}_{\mathbb{Q}}}(q(X), q(Y))$$

is an isomorphism.

For the injectivity it is sufficient to prove that for all $f \in \text{Hom}_{\mathcal{A}}(X, Y)$ with $q(f) = 0$ it follows that $n \cdot f = 0$ for some $n \in \mathbb{Z} \setminus \{0\}$. Indeed, $q(f) = 0$ implies $\text{image}(f) \in \mathcal{A}_{b\text{-tor}}$; thus there is an integer $n \neq 0$ with $n \cdot \text{id}_{\text{image}(f)} = 0$. It follows that $nf = 0$.

For the surjectivity, let $f : X \rightarrow Y$ be a morphism in \mathcal{A} such that $\ker(f), \text{coker}(f) \in \mathcal{A}_{b\text{-tor}}$; equivalently $q(f)$ is an isomorphism. We need to show that the inverse map $q(f)^{-1}$ is contained in the image of $\text{Hom}_{\mathcal{A}}(Y, X) \otimes_{\mathbb{Z}} \mathbb{Q}$. Choose an integer $n_1 \neq 0$ such that $n_1 \cdot \text{id}_{\ker(f)} = 0$. Then there exists $g_1 : \text{im}(f) \rightarrow X$ such that $g_1 \circ f = n_1 \cdot \text{id}_X$. Let $n_2 \neq 0$ be an integer such that $n_2 \cdot \text{id}_{\text{coker}(f)} = 0$. Then $Y \xrightarrow{\cdot n_2} Y$ factors through $\text{im}(f)$ and we obtain $g_2 : Y \xrightarrow{\cdot n_2} \text{im}(f) \xrightarrow{g_1} X$. Thus the image of $g_2 \otimes (n_1 \cdot n_2)^{-1}$ is the inverse of $q(f)$. \square

COROLLARY 3.7.5. *Let $F : \mathcal{A} \rightarrow \mathcal{B}$ be a functor between well-powered abelian categories. There is a natural functor $F_{\mathbb{Q}} : \mathcal{A}_{\mathbb{Q}} \rightarrow \mathcal{B}_{\mathbb{Q}}$ defined by*

$$F_{\mathbb{Q}}(q(X)) = F(X)$$

for every object X in \mathcal{A} , and

$$F_{\mathbb{Q}} : \text{Hom}_{\mathcal{A}_{\mathbb{Q}}}(q(X), q(Y)) \rightarrow \text{Hom}_{\mathcal{B}_{\mathbb{Q}}}(F_{\mathbb{Q}}q(X), F_{\mathbb{Q}}q(Y))$$

$$F_{\mathbb{Q}} := F \otimes_{\mathbb{Z}} \text{id}_{\mathbb{Q}}$$

via the isomorphism of Proposition 3.7.4.

Remark 3.7.6. The statement of Corollary 3.7.5 also follows from

$$F(\mathcal{A}_{b\text{-tor}}) \subset \mathcal{B}_{b\text{-tor}}$$

by using the universal property of the quotient category.

PROPOSITION 3.7.7. *Let $F : \mathcal{A} \rightarrow \mathcal{B}$ be a left-exact functor between well-powered abelian categories. Suppose that the left derived functor*

$$RF : D^+(\mathcal{A}) \rightarrow D^+(\mathcal{B})$$

exists. Suppose that there exist sufficiently many F -acyclic objects in \mathcal{A} . Then

$$F_{\mathbb{Q}} : \mathcal{A}_{\mathbb{Q}} \rightarrow \mathcal{B}_{\mathbb{Q}}$$

is left-exact and the left derived functor

$$RF_{\mathbb{Q}} : D^+(\mathcal{A}_{\mathbb{Q}}) \rightarrow D^+(\mathcal{B}_{\mathbb{Q}})$$

exists. Moreover, the diagram

$$\begin{array}{ccc} D^+(\mathcal{A}) & \xrightarrow{RF} & D^+(\mathcal{B}) \\ D^+(q) \downarrow & & \downarrow D^+(q) \\ D^+(\mathcal{A}_{\mathbb{Q}}) & \xrightarrow{RF_{\mathbb{Q}}} & D^+(\mathcal{B}_{\mathbb{Q}}) \end{array}$$

is commutative.

Proof. Let

$$(3.7.1) \quad 0 \rightarrow X' \xrightarrow{f'} Y' \xrightarrow{g'} Z' \rightarrow 0$$

be an exact sequence in $\mathcal{A}_{\mathbb{Q}}$. By [Gab62, Chapitre III, §1, Corollaire 1] we can find an exact sequence

$$(3.7.2) \quad 0 \rightarrow X \xrightarrow{f} Y \xrightarrow{g} Z \rightarrow 0$$

in \mathcal{A} and isomorphisms $X' \xrightarrow{\cong} q(X), Y' \xrightarrow{\cong} q(Y), Z' \xrightarrow{\cong} q(Z)$, such that the diagram

$$\begin{array}{ccccccccc} 0 & \longrightarrow & X' & \xrightarrow{f'} & Y' & \xrightarrow{g'} & Z' & \longrightarrow & 0 \\ & & \downarrow \cong & & \downarrow \cong & & \downarrow \cong & & \\ 0 & \longrightarrow & q(X) & \xrightarrow{f} & q(Y) & \xrightarrow{g} & q(Z) & \longrightarrow & 0 \end{array}$$

is commutative. Since q is exact and $qF = F_{\mathbb{Q}}q$, it follows that $F_{\mathbb{Q}}$ is left exact. We define

$$P_{\mathbb{Q}} := \{X' \in \text{ob}(\mathcal{A}_{\mathbb{Q}}) \mid \exists X \in \text{ob}(\mathcal{A}) : q(X) \cong X', R^i F(X) \in \mathcal{B}_{b\text{-tor}} \text{ for all } i > 0\}$$

as a full subcategory of $\mathcal{A}_{\mathbb{Q}}$. If $X \in \text{ob}(\mathcal{A})$ is an F -acyclic object then $q(X) \in P_{\mathbb{Q}}$. Therefore every object $Y \in \text{ob}(\mathcal{A}_{\mathbb{Q}})$ admits a monomorphism $\iota : Y \rightarrow X$ with $X \in \text{ob}(P_{\mathbb{Q}})$.

Suppose that in the exact sequence (3.7.1) we know that $X' \in \text{ob}(P_{\mathbb{Q}})$. We claim that the following holds

$$Y' \in \text{ob}(P_{\mathbb{Q}}) \Leftrightarrow Z' \in \text{ob}(P_{\mathbb{Q}}).$$

As above, we may prove the claim for (3.7.2) instead of (3.7.1). It is easy to see that $R^i F(X) \in \mathcal{B}_{b\text{-tor}}$ for all $i > 0$, and we conclude that

$$qR^i F(Y) \xrightarrow{\cong} qR^i F(Z) \quad \text{for all } i > 0.$$

This implies the claim.

In the same way we can see that if (3.7.1) is an exact sequence with objects in $P_{\mathbb{Q}}$ then

$$0 \rightarrow F_{\mathbb{Q}}(X') \rightarrow F_{\mathbb{Q}}(Y') \rightarrow F_{\mathbb{Q}}(Z') \rightarrow 0$$

is exact. From [Har66, I, Cor. 5.3, β] it follows that the left derived functor

$$RF_{\mathbb{Q}} : D^+(\mathcal{A}_{\mathbb{Q}}) \rightarrow D^+(\mathcal{B}_{\mathbb{Q}})$$

exists. Moreover, if X is F -acyclic then $q(X)$ is $F_{\mathbb{Q}}$ acyclic. Therefore the diagram

$$\begin{array}{ccc} D^+(\mathcal{A}) & \xrightarrow{RF} & D^+(\mathcal{B}) \\ D^+(q) \downarrow & & \downarrow D^+(q) \\ D^+(\mathcal{A}_{\mathbb{Q}}) & \xrightarrow{RF_{\mathbb{Q}}} & D^+(\mathcal{B}_{\mathbb{Q}}) \end{array}$$

is commutative. □

Remark 3.7.8. Let $f : X \rightarrow Y$ be a morphism between k -schemes and Φ a family of supports on X . The assumptions of the Proposition are satisfied for the functors

$$\begin{aligned} \underline{\Gamma}_{\Phi} &: \widehat{\text{dRW}}_X \rightarrow \widehat{\text{dRW}}_X, \\ f_* &: \widehat{\text{dRW}}_X \rightarrow \widehat{\text{dRW}}_Y, \\ \underline{\lim} &: \widehat{\text{dRW}}_X \rightarrow \widehat{\text{dRW}}_X, \\ f_{\Phi} &: \widehat{\text{dRW}}_X \rightarrow \widehat{\text{dRW}}_Y, \\ \hat{f}_{\Phi} &: \widehat{\text{dRW}}_X \rightarrow \widehat{\text{dRW}}_Y, \end{aligned}$$

of Proposition 1.5.6.

NOTATION 3.7.9. In general, we will denote by a subscript \mathbb{Q} the image of an object of $\widehat{\text{dRW}}_X$ (resp. of $D^+(\widehat{\text{dRW}}_X)$) under the localization functor q (resp. $D^+(q)$). If $F : \widehat{\text{dRW}}_X \rightarrow \widehat{\text{dRW}}_Y$ is a functor we will by abuse of notation denote $F_{\mathbb{Q}}$ again by F and $RF_{\mathbb{Q}}$ again by RF . Thus $q(Rf_* W\Omega_X)$ will be denoted by $Rf_* W\Omega_{X, \mathbb{Q}}$, etc. (Warning: $W\Omega_{X, \mathbb{Q}}$ is not the same as $W\Omega_X \otimes_{\mathbb{Z}} \mathbb{Q}$.)

4. WITT-RATIONAL SINGULARITIES

All schemes in this section are quasi-projective over k .

4.1. THE WITT CANONICAL SYSTEM.

4.1.1. Recall from Definition 1.8.3 the notion of a Witt residual complex. Let X be a k -scheme with structure map $\rho_X : X \rightarrow \text{Spec } k$, then there is a canonical Witt residual complex $K_X = \rho_X^\Delta W_\bullet \omega$ (see Notation 1.9.2). This complex has the following properties:

- (1) If X is smooth of pure dimension d , then there is a quasi-isomorphism of graded complexes $\tau_X : W_\bullet \omega_X \rightarrow K_X(-d)[-d]$, which is compatible with localization. (Ekedahl, see Theorem 1.10.3.)
- (2) If $j : U \hookrightarrow X$ is an open subscheme, then $j^* K_X \cong K_U$ (see Proposition 1.8.8).
- (3) If $f : X \rightarrow Y$ is a morphism of k -schemes, then there is a canonical isomorphism $f^\Delta K_Y \cong K_X$ induced by c_{f, ρ_Y} , where ρ_Y is the structure map of Y . This isomorphism is compatible with composition and localization and in case f is an open embedding also with the isomorphism in (2) (via $f^* \cong f^\Delta$). (See Proposition 1.8.8.)
- (4) For a proper k -morphism $f : X \rightarrow Y$, there is a trace map $\text{Tr}_f : f_* K_X \rightarrow K_Y$, which is a morphism of complexes of Witt quasi-dualizing systems (see Definition 1.6.3); it is compatible with composition and localization (see Lemma 1.8.9).
- (5) We have a functor

$$D_X := \mathcal{H}om(-, K_X) : C(\text{dRW}_{X, \text{qc}})^o \rightarrow C(\text{dRW}_X).$$

It preserves quasi-isomorphisms and hence also induces a functor from $D(\text{dRW}_{X, \text{qc}})^o$ to $D(\text{dRW}_X)$ (see 1.9.1).

- (6) Let $f : X \rightarrow Y$ be a finite morphism. We denote by

$$\vartheta_f : f_* D_X(-) \rightarrow D_Y(f_*(-))$$

the composition

$$f_* \mathcal{H}om(-, K_X) \xrightarrow{\text{nat.}} \mathcal{H}om(f_*(-), f_* K_X) \xrightarrow{\text{Tr}_f} \mathcal{H}om(-, K_Y).$$

Then ϑ_f is an isomorphism of functors on $C^-(\text{dRW}_{X, \text{qc}})$; it is compatible with composition and localization. (It is an isomorphism on each level by duality theory, see 1.7.3, (7); for the other assertions see Proposition 1.9.4.)

DEFINITION 4.1.2. Let X be a k -scheme of pure dimension d . The *Witt canonical system on X* is defined to be the $(-d)$ -th cohomology of K_X sitting in degree d and is denoted by $W_\bullet \omega_X$, i.e.

$$W_\bullet \omega_X := H^{-d}(K_X)(-d).$$

Since $W_\bullet \mathcal{O}_X$ is a Witt system (i.e. a de Rham-Witt system with zero differential), $W_\bullet \omega_X$ inherits the structure of a Witt system from the canonical isomorphism $K_X \cong \mathcal{H}om(W_\bullet \mathcal{O}_X, K_X)$ and 4.1.1, (5). We denote the limit with respect to π by

$$W\omega_X := \varprojlim_{\pi} W_\bullet \omega_X.$$

Remark 4.1.3. Notice that in case X is smooth the above definition of $W_\bullet\omega_X$ coincides (up to canonical isomorphism) with our earlier definition of $W_\bullet\omega_X = W_\bullet\Omega_X^d$, by 4.1.1, (1). But observe, in Example 1.6.4, (2) we viewed it as a Witt dualizing system, whereas now as a Witt system. This will not cause any confusion.

PROPOSITION 4.1.4. *Let X be a k -scheme of pure dimension d . Then $W_\bullet\omega_X$ has the following properties:*

- (1) *The sheaf $W_1\omega_X$ equals the usual canonical sheaf on X if X is normal.*
- (2) *The complex K_X is concentrated in degree $[-d, 0]$, hence there is a natural morphism of complexes*

$$W_\bullet\omega_X \rightarrow K_X(-d)[-d].$$

This morphism is a quasi-isomorphism if X is Cohen-Macaulay (CM).

- (3) *For each n the sheaf $W_n\omega_X$ is a coherent sheaf on W_nX , which satisfies Serre's condition S_2 .*
- (4) *Let $j : U \hookrightarrow X$ be an open subscheme which contains all 1-codimensional points of X . Then we have an isomorphism of Witt systems*

$$W_\bullet\omega_X \xrightarrow{\simeq, \text{nat.}} j_*j^*W_\bullet\omega_X = j_*W_\bullet\omega_U.$$

If X is normal, U can be chosen to be smooth, in which case we have an isomorphism

$$j_*W_\bullet\omega_U \simeq j_*W_\bullet\Omega_U^d,$$

which is induced by 4.1.1, (1) and (2). In particular the transition maps $W_n\omega_X \rightarrow W_{n-1}\omega_X$ are surjective if X is normal.

- (5) *Assume X is normal. Then there is a natural isomorphism for all n*

$$\mathcal{H}\text{om}(W_n\omega_X, W_n\omega_X) \cong H^{-d}(D_{X,n}(W_n\omega_X)(-d)),$$

where $D_{X,n}(-) = \mathcal{H}\text{om}(-, K_{X,n})$. Therefore $\mathcal{H}\text{om}(W_\bullet\omega_X, W_\bullet\omega_X)$ is naturally equipped with the structure of a Witt system and multiplication induces an isomorphism of Witt systems

$$W_\bullet\mathcal{O}_X \xrightarrow{\simeq} \mathcal{H}\text{om}(W_\bullet\omega_X, W_\bullet\omega_X).$$

- (6) *Let $f : X \rightarrow Y$ be a proper morphism between k -schemes, which are both of pure dimension d . Then we define the pushforward $f_* : f_*W_\bullet\omega_X \rightarrow W_\bullet\omega_Y$ as the composition in dRW_Y*

$$f_* : f_*W_\bullet\omega_X = f_*H^{-d}(K_X)(-d) = H^{-d}(f_*K_X)(-d) \xrightarrow{\text{Tr}_f} H^{-d}(K_Y)(-d) = W_\bullet\omega_Y$$

This morphism is compatible with composition and localization and in case X and Y are smooth coincides with the pushforward defined in Definition 2.3.1 (for $S = Y$.)

- (7) *The sequence of $W_n\mathcal{O}_X$ -modules*

$$0 \rightarrow i_{n*}W_{n-1}\omega_X \xrightarrow{p} W_n\omega_X \xrightarrow{F^{n-1}} i_*F_{X*}^{n-1}\omega_X$$

is exact for any $n \geq 1$. Furthermore, if X is CM, then the map on the right is also surjective. (Here we write $\omega_X := W_1\omega_X$.)

Proof. (1) is clear by construction. (In the derived category of coherent sheaves $K_{X,1}$ is isomorphic to $\rho_X^!k$, where $\rho_X : X \rightarrow \text{Spec } k$ is the structure map.) It suffices to prove the other statements on a fixed finite level n .

(2). The codimension function associated to $K_{X,n}$ (see 1.7.1) is given by

$$d_{K_{X,n}}(x) = -\text{trdeg}(k(x)/k) = \dim \mathcal{O}_{X,x} - d$$

(see [Con00, (3.2.4)]). This already gives the first statement of (2). For the second statement the same argument as in [Eke84, I, 2.] works (there for smooth schemes). For the convenience of the reader we recall the argument. Let X be CM. We have to show that $H^i(K_{X,n}) = 0$ for all $i \neq -d$. For $n = 1$ this follows from (1). We have an exact sequence of coherent $W_n\mathcal{O}_X$ -modules

$$0 \rightarrow i_*F_{X^*}^{n-1}\mathcal{O}_X \xrightarrow{V^{n-1}} W_n\mathcal{O}_X \xrightarrow{\pi} i_{n*}W_{n-1}\mathcal{O}_X,$$

where F_X denotes the absolute Frobenius on X and i, i_n denote the closed embeddings $X \hookrightarrow W_nX, W_{n-1}X \hookrightarrow W_nX$. Applying $R\mathcal{H}om_{W_n\mathcal{O}_X}(-, K_{X,n})$ to it and using duality for the finite morphisms iF_X^{n-1} and i_n (cf. 4.1.1, (6)) we obtain a triangle in $D(W_n\mathcal{O}_X)$

$$(4.1.1) \quad i_{n*}K_{X,n-1} \rightarrow K_{X,n} \rightarrow (iF_X^{n-1})_*K_{X,1} \rightarrow i_{n*}K_{X,n-1}[1].$$

Therefore the statement follows by induction.

(3). Since $K_{X,n}$ is a residual complex, it has coherent cohomology by definition. To prove the S_2 property of $W_n\omega_X$ it suffices to show

$$(4.1.2) \quad \begin{cases} \text{Ext}_{W_n\mathcal{O}_{X,x}}^0(k(x), W_n\omega_{X,x}) = 0, & \text{for all } x \in X^{(1)} \\ \text{Ext}_{W_n\mathcal{O}_{X,x}}^i(k(x), W_n\omega_{X,x}) = 0, & \text{for } i = 0, 1 \text{ and all } x \in X^{(2)}, \end{cases}$$

where $X^{(c)}$ denotes the points of codimension c . By the formula for the codimension function associated to $K_{X,n}$ above and [Har66, V, §7], we have

$$(4.1.3) \quad \text{Ext}_{W_n\mathcal{O}_{X,x}}^i(k(x), K_{X,n}) = \begin{cases} 0, & \text{if } x \notin X^{(i+d)}, \\ k(x), & \text{if } x \in X^{(i+d)}. \end{cases}$$

Thus the vanishing (4.1.2) can easily be deduced from the spectral sequence

$$E_2^{i,j} = \text{Ext}^i(k(x), H^j(K_{X,n})) \Rightarrow \text{Ext}^{i+j}(k(x), K_{X,n})$$

and the vanishing $E_2^{i,j} = 0$ if $i < 0$ or $j \notin [-d, 0]$ (by (2)).

(4). The first morphism is bijective by (3) and [SGA2, Exp. III, Cor. 3.5].

(5). The first isomorphism is obtained by considering the spectral sequence

$$\text{Ext}^i(W_n\omega_X, H^j(K_{X,n})) \Rightarrow \text{Ext}^{i+j}(W_n\omega_X, K_{X,n}) = H^{i+j}(D_{X,n}(W_n\omega_X))$$

and using that $H^j(K_{X,n}) \neq 0$ only for $j \in [-d, 0]$, by (2). The second isomorphism can easily be deduced from (4), the isomorphism $W_n\mathcal{O}_X \xrightarrow{\sim} j_*W_n\mathcal{O}_U$ ($W_n\mathcal{O}_X$ is S_2) and the corresponding statement for smooth schemes, see 1.10.1. (6) The equality $f_*H^{-d}(K_X) = H^{-d}(f_*K_X)$ follows from the spectral sequence $R^i f_* H^j(K_X) \Rightarrow R^{i+j} f_* K_X$, (2) above and from $Rf_* K_X = f_* K_X$. The other statements follow from 4.1.1, (4).

(7) Recall that F and \underline{p} on $W_\bullet\omega_X$ are defined by the isomorphism $K_X \cong \mathcal{H}om(W_\bullet\mathcal{O}_X, K_X)$ and the formulas in 1.6.6. Thus the exact sequence in (7) is the result of applying H^{-d} to the triangle (4.1.1) above and using the isomorphisms $\underline{p} \circ \epsilon_i$ and $V \circ \epsilon_{\sigma_i}$ from 1.6.5. \square

DEFINITION 4.1.5. Let $f : X \rightarrow Y$ be a finite and surjective k -morphism between integral normal schemes both of dimension d .

(1) We define a pullback morphism in dRW_Y

$$f^* : W_\bullet\omega_Y \rightarrow f_*W_\bullet\omega_X$$

as follows: Choose open and smooth subschemes $j_X : U \hookrightarrow X$ and $j_Y : V \hookrightarrow Y$, which contain all 1-codimensional points of X and Y respectively and such that f restricts to a morphism $f' : U \rightarrow V$. We define f^* as the composition

$$W_\bullet\omega_Y \xrightarrow{\simeq, 4.1.4, (5)} j_{Y*}W_\bullet\Omega_V^d \xrightarrow{f'^*} j_{Y*}f'_*W_\bullet\Omega_U^d \xrightarrow{\simeq, 4.1.3} f_*j_{X*}W_\bullet\omega_U \xrightarrow{\simeq, 4.1.4, (5)} f_*W_\bullet\omega_X.$$

It is straightforward to check, that this morphism is independent of the choice of U . We also write f^* for the sum of the natural pullback on $W_\bullet\mathcal{O}$ with the just defined pullback, i.e.

$$f^* : W_\bullet\mathcal{O}_Y \oplus W_\bullet\omega_Y \rightarrow f_*(W_\bullet\mathcal{O}_X \oplus W_\bullet\omega_X).$$

Taking the limit we obtain a pullback in $\widehat{\text{dRW}}_Y$

$$f^* : W\mathcal{O}_Y \oplus W\omega_Y \rightarrow f_*(W\mathcal{O}_X \oplus W\omega_X).$$

(2) We define a pushforward in dRW_Y

$$f_* : f_*W_\bullet\mathcal{O}_X \rightarrow W_\bullet\mathcal{O}_Y$$

as the composition

$$\begin{aligned} f_*W_\bullet\mathcal{O}_X &\xrightarrow{\simeq, 4.1.4, (5)} f_*\mathcal{H}om(W_\bullet\omega_X, W_\bullet\omega_X) \\ &\xrightarrow{\simeq, 4.1.4, (5)} H^{-d}(f_*D_X(W_\bullet\omega_X)(-d)) \\ &\xrightarrow{\simeq, \vartheta_f} H^{-d}(D_Y(f_*W_\bullet\omega_X)(-d)) \\ &\xrightarrow{D_Y(f^*), (1)} H^{-d}(D_Y(W_\bullet\omega_Y)(-d)) \\ &\xrightarrow{\simeq, 4.1.4, (5)} \mathcal{H}om(W_\bullet\omega_Y, W_\bullet\omega_Y) \\ &\xrightarrow{\simeq, 4.1.4, (5)} W_\bullet\mathcal{O}_Y. \end{aligned}$$

We also write f_* for the sum of the pushforward on $f_*W_\bullet\omega_X$ defined in Proposition 4.1.4, (6) with the just defined pushforward. Taking the limit we obtain a pushforward in $\widehat{\text{dRW}}_Y$

$$f_* : f_*(W\mathcal{O}_X \oplus W\omega_X) \rightarrow W\mathcal{O}_Y \oplus W\omega_Y.$$

LEMMA 4.1.6. *Let $f : X \rightarrow Y$ be a finite and surjective k -morphism between normal integral schemes of dimension d .*

- (1) *Let $j_Y : V \hookrightarrow Y$ be an open smooth subscheme, which contains all 1-codimensional points of Y and such that $U := f^{-1}(V) \subset X$ is smooth and contains all 1-codimensional points of X (e.g. $V = Y \setminus f(X_{\text{sing}})$). Denote by $j_X : U \hookrightarrow X$ and $f' : U = f^{-1}(V) \rightarrow V$ the pullbacks. Then*

$$j_X^* \circ f^* = f'^* \circ j_Y^*, \quad j_Y^* \circ f_* = f'_* \circ j_X^*,$$

where f'^* is the pullback defined in Definition 2.2.1 and f'_* is the push-forward defined in Definition 2.3.1 (with $S = V$).

- (2) *The composition $f_* \circ f^*$ equals the multiplication with the degree of f .*

Proof. The first part of (1) follows from the fact that all the maps in the definitions of f_* and f^* are compatible with localization; the second part follows immediately from the definitions. For (2) we have to check that $f_* \circ f^* - \deg f = 0$ in $\mathcal{H}om(W_\bullet \mathcal{O}_Y, W_\bullet \mathcal{O}_Y) = W_\bullet \mathcal{O}_Y$ and in $\mathcal{H}om(W_\bullet \omega_Y, W_\bullet \omega_Y) = W_\bullet \mathcal{O}_Y$. It suffices to check this on some open $V \subset Y$ and hence the assertion follows from (1) and Proposition 2.3.4 (Gros). \square

4.2. TOPOLOGICAL FINITE QUOTIENTS.

4.2.1. *Universal homeomorphisms.* Recall that a morphism of k -schemes $u : X \rightarrow Y$ is a *universal homeomorphism* if for any $Y' \rightarrow Y$ the base change morphism $u' : X \times_Y Y' \rightarrow Y'$ is a homeomorphism. By [EGAIV(4), Cor. (18.12.11)] this is equivalent to say that u is finite, surjective and radical. In case X and Y are integral and Y is normal it follows from [EGAI, Prop. (3.5.8)], that u is a universal homeomorphism if and only if u is finite, surjective and purely inseparable (i.e. $k(Y) \subset k(X)$ is purely inseparable).

4.2.2. *Relative Frobenius.* We denote by $\sigma : \text{Spec } k \rightarrow \text{Spec } k$ the Frobenius (we will not use the notation from section 1.2 any longer); for a k -scheme X we denote by $X^{(n)}$ the pullback of X along $\sigma^n : \text{Spec } k \rightarrow \text{Spec } k$ and by $\sigma_X^n : X^{(n)} \rightarrow X$ the projection. Notice that since k is perfect, σ_X^n is an isomorphism of \mathbb{F}_p -schemes. The n -th relative Frobenius of X over k is by definition the unique k -morphism $F_{X/k}^n : X \rightarrow X^{(n)}$ which satisfies $\sigma_X^n \circ F_{X/k}^n = F_X^n$, where $F_X : X \rightarrow X$ is the absolute Frobenius morphism of X . Clearly $F_{X/k}^n$ is a universal homeomorphism.

LEMMA 4.2.3. *Let $u : X \rightarrow Y$ be a morphism between integral k -schemes, which is a universal homeomorphism and assume that Y is normal. Then $\deg u = p^n$ for some natural number n and there exists a universal homeomorphism $v : Y \rightarrow$*

$X^{(n)}$ such that the following diagram commutes

$$\begin{array}{ccc} X & \xrightarrow{F_{X/k}^n} & X^{(n)}, \\ u \downarrow & \nearrow v & \downarrow u \times \text{id}_{\text{Spec } k} \\ Y & \xrightarrow{F_{Y/k}^n} & Y^{(n)}. \end{array}$$

Proof. We may assume $X = \text{Spec } B$ and $Y = \text{Spec } A$. Then by 4.2.1 $u^* : A \hookrightarrow B$ is an inclusion of k -algebras which makes $k(A) \subset k(B)$ a purely inseparable field extension of degree p^n . $F_{X/k}^{n*}$ is given by $B \otimes_{k, \sigma^n} k \rightarrow B$, $b \otimes \lambda \mapsto b^{p^n} \lambda$. But $b^{p^n} \in k(A) \cap B = A$ by the normality of Y . Therefore $F_{X/k}^{n*}$ factors via $u^* : A \hookrightarrow B$. We obtain a homomorphism of k -algebras $B \otimes_{k, \sigma^n} k \rightarrow A$, which gives rise to a k -morphism $v : Y \rightarrow X^{(n)}$. It follows that v is a universal homeomorphism, which makes the diagram in the statement commutative. \square

LEMMA 4.2.4. *Let $u : X \rightarrow Y$ be a universal homeomorphism between two integral and normal schemes. Let u_* and u^* be the pushforward and the pullback from Definition 4.1.5. Then*

$$u_* u^* = \deg u \cdot \text{id}_{(W\mathcal{O}_Y \oplus W\omega_Y)}, \quad u^* u_* = \deg u \cdot \text{id}_{u_*(W\mathcal{O}_X \oplus W\omega_X)}.$$

Proof. The equality on the left is a particular case of Lemma 4.1.6, (2). To prove the equality on the right we may assume that X and Y are smooth (by Proposition 4.1.4 and the corresponding statement for $W\mathcal{O}$). Then by Lemma 4.1.6, (1) and Proposition 3.5.6, we have $u_* = \hat{\mathcal{H}}([\Gamma]/Y)$ and $u^* = \hat{\mathcal{H}}([\Gamma^t]/Y)$, where $\Gamma \subset X \times Y$ is the graph of u and Γ^t its transpose. Therefore we are reduced to show

$$(4.2.1) \quad [\Gamma^t] \circ [\Gamma] = \deg u \cdot [\Delta_X] \quad \text{in } \text{CH}(X \times X, P(X \times_Y X)),$$

where $\Delta_X \subset X \times X$ is the diagonal. But since u is flat (being a finite and surjective morphism between integral and smooth schemes) and a homeomorphism, we have

$$[\Gamma^t] \circ [\Gamma] = [X \times_Y X] = \dim_{k(X)}(k(X) \otimes_{k(Y)} k(X)) \cdot [\Delta_X] = \deg u \cdot [\Delta_X].$$

\square

DEFINITION 4.2.5. Let X be a normal and equidimensional scheme. We say that X is a *finite quotient* if there exists a finite and surjective morphism from a smooth scheme $Y \rightarrow X$. We say that X is a *tame finite quotient* if this morphism can be chosen to have its degree (as a locally constant function on Y) not divisible by p . We say that X is a *topological finite quotient* if there exists a universal homeomorphism $u : X \rightarrow X'$ to a finite quotient X' .

Remark 4.2.6. If X' is a finite quotient, so is $X'^{(n)}$. It follows from Lemma 4.2.3 that if X is normal and equidimensional and if there exists a universal homeomorphism $u : X' \rightarrow X$ with a source a finite quotient X' , then X is a topological finite quotient.

Topological finite quotients are good to handle because of the following Proposition.

PROPOSITION 4.2.7. *Consider morphisms*

$$\begin{array}{ccc} & & Z \\ & & \downarrow u \\ X & \xrightarrow{f} & Y, \end{array}$$

where X is smooth and integral, Y, Z are normal and integral, f is finite and surjective and u is a universal homeomorphism. Let $a : Y \rightarrow S$ be a morphism to some k -scheme S . Set

$\beta := \text{pr}_{1,3,*}[X \times_Y Z \times_Y X]$ in $\text{Hom}_{\mathcal{C}_S}^0(X, X) = \text{CH}^{\dim X}(X \times X, P(X \times_S X))$, where $\text{pr}_{1,3} : X \times Z \times X \rightarrow X \times X$ is the projection (see 3.5 for the notation). Then for all i the composition

$$f^* \circ u_* : R^i(au)_*(W\mathcal{O}_Z \oplus W\omega_Z)_{\mathbb{Q}} \rightarrow R^i(af)_*(W\mathcal{O}_X \oplus W\omega_X)_{\mathbb{Q}}$$

induces an isomorphism in $\widehat{\text{dRW}}_S$

$$R^i(au)_*(W\mathcal{O}_Z \oplus W\omega_Z)_{\mathbb{Q}} \cong \hat{\mathcal{H}}(\beta/S)(R^i(af)_*(W\mathcal{O}_X \oplus W\omega_X)_{\mathbb{Q}}),$$

where $\hat{\mathcal{H}}(\beta/S)$ is the morphism from Proposition 3.5.4 (see Notation 3.7.9 for the meaning of the subscript \mathbb{Q}).

Proof. First of all notice that both f and u are finite and universally equidimensional. It follows that β defines an element in $\text{CH}^{\dim X}(X \times X, P(X \times_Y X))$, a fortiori in $\text{CH}^{\dim X}(X \times X, P(X \times_S X))$. In particular, Proposition 3.5.4 yields a morphism

$$\hat{\mathcal{H}}(\beta/Y) : f_*(W\mathcal{O}_X \oplus W\omega_X) \rightarrow f_*(W\mathcal{O}_X \oplus W\omega_X).$$

We claim

$$(4.2.2) \quad \hat{\mathcal{H}}(\beta/Y) = f^*u_*u^*f_*$$

with f_*, u_*, u^*, f^* as in Definition 4.1.5. By Proposition 4.1.4, (4) and Lemma 4.1.6, (1) and since $\hat{\mathcal{H}}(\beta/Y)$ is compatible with localization in Y (just by construction), we may assume, that X, Y and Z are smooth. Then

$$\beta = [\Gamma_f^t] \circ [\Gamma_u] \circ [\Gamma_u^t] \circ [\Gamma_f],$$

where we denote by $\Gamma_f^t \subset Y \times X$ the transpose of the graph of f , etc. Therefore claim (4.2.2) follows from Proposition 3.5.6. Thus Proposition 3.5.10 implies that

$$\begin{aligned} & \hat{\mathcal{H}}(\beta/S)(R^i(af)_*(W\mathcal{O}_X \oplus W\omega_X)) \\ &= \text{Image}(f^*u_*u^*f_* : R^i(af)_*(W\mathcal{O}_X \oplus W\omega_X) \rightarrow R^i(af)_*(W\mathcal{O}_X \oplus W\omega_X)). \end{aligned}$$

Thus the assertion follows from $(u^*f_*)(f^*u_*) = \text{deg } f \text{ deg } u \cdot \text{id}$ (by Lemma 4.1.6, (a) and Lemma 4.2.4). \square

4.3. QUASI-RESOLUTIONS AND RELATIVE HODGE-WITT COHOMOLOGY.

DEFINITION 4.3.1. We say that a morphism between two integral k -schemes $f : X \rightarrow Y$ is a *quasi-resolution* if the following conditions are satisfied:

- (1) X is a topological finite quotient,
- (2) f is projective, surjective, and generically finite,
- (3) the extension of the function fields $k(Y) \subset k(X)$ is purely inseparable.

Condition (2) and (3) are for example satisfied if f is projective and birational. Let X, Y be integral and normal. In general, every projective, surjective and generically finite morphism $f : X \rightarrow Y$ can be factored through the normalization Y' of Y in the function field $k(X)$ of X :

$$f : X \xrightarrow{f'} Y' \xrightarrow{u} Y.$$

The morphism f' is birational and u is finite. If $k(Y) \subset k(X)$ is purely inseparable then u is a universal homeomorphism.

Remark 4.3.2. Let X be an integral k -scheme. It follows from the proof of [dJ97, Cor. 5.15] (cf. also [dJ96, Cor. 7.4]), that there exists a finite and surjective morphism from a normal integral scheme $u : X' \rightarrow X$, such that $k(X) \subset k(X')$ is purely inseparable and a smooth, integral and quasi-projective scheme X'' , with a finite group G acting on it such that there is a projective and birational morphism $f : X''/G \rightarrow X'$. In particular X has a quasi-resolution

$$X''/G \xrightarrow{f} X' \xrightarrow{u} X.$$

THEOREM 4.3.3. *Let Y be a topological finite quotient and $f : X \rightarrow Y$ a quasi-resolution. Then the pullback f^* and the pushforward (see Proposition 4.1.4, (6)) induce isomorphisms in $D^b(\widehat{\mathrm{dRW}}_{Y, \mathbb{Q}})$*

$$f^* : W\mathcal{O}_{Y, \mathbb{Q}} \xrightarrow{\cong} Rf_* W\mathcal{O}_{X, \mathbb{Q}}, \quad Rf_* W\omega_{X, \mathbb{Q}} \cong f_* W\omega_{X, \mathbb{Q}}[0] \xrightarrow{f^*, \cong} W\omega_{Y, \mathbb{Q}}.$$

Proof. We can assume that X and Y are integral schemes of dimension d and (by Lemma 4.2.4) also that they are finite quotients. Thus there exist smooth integral schemes X' and Y' together with finite and surjective morphisms $a : X' \rightarrow X$ and $b : Y' \rightarrow Y$. Let

$$X \xrightarrow{\pi} X_1 \xrightarrow{u} Y,$$

be a factorization of f , with π projective and birational, X_1 normal and u a universal homeomorphism. We can find a non-empty smooth open subscheme $U_0 \subset Y$, such that $U_1 := u^{-1}(U_0)$, $U_2 := \pi^{-1}(U_1)$, $U'_0 := b^{-1}(U_0)$ and $U'_2 := a^{-1}(U_2)$ are smooth and $\pi|_{U_2}$ is an isomorphism. Notice that $a|_{U'_2}$, $b|_{U'_0}$ and $u|_{U_1}$ are then automatically flat. Set $Z'_0 := Y' \setminus U'_0$ and $Z'_2 := X' \setminus U'_2$. We define

$$\begin{aligned} \alpha &:= [X' \times_X X'] \in \mathrm{CH}^d(X' \times X', P(X' \times_X X')), \\ \beta &:= [Y' \times_Y Y'] \in \mathrm{CH}^d(Y' \times Y', P(Y' \times_Y Y')), \\ \gamma &:= [X' \times_Y Y'] \in \mathrm{CH}^d(X' \times Y', P(X' \times_Y Y')). \end{aligned}$$

(See 3.5 for the notation.) These cycles are well-defined since $a : X' \rightarrow X$ and $b : Y' \rightarrow Y$ are universally equidimensional. We claim

$$(4.3.1) \quad \deg b \cdot (\gamma \circ \alpha) - \deg a \cdot (\beta \circ \gamma) \in \text{image}(\text{CH}(Z'_2 \times_Y Z'_0)),$$

$$(4.3.2) \quad \deg a \cdot (\gamma^t \circ \beta) - \deg b \cdot (\alpha \circ \gamma^t) \in \text{image}(\text{CH}(Z'_0 \times_Y Z'_2)),$$

$$(4.3.3) \quad (\gamma^t \circ \gamma \circ \alpha) - (\deg a \deg b \deg u) \cdot \alpha \in \text{image}(\text{CH}(Z'_2 \times_Y Z'_2)),$$

$$(4.3.4) \quad (\gamma \circ \gamma^t \circ \beta) - (\deg a \deg b \deg u) \cdot \beta \in \text{image}(\text{CH}(Z'_0 \times_Y Z'_0)).$$

Observe that $(U'_2 \times_Y Y') \cup (X' \times_Y U'_0) = U'_2 \times_{U_0} U'_0$ etc. Thus using the localization sequence we see that it suffices to prove

$$\begin{aligned} \deg b \cdot (\gamma|_{U'_2 \times U'_0} \circ \alpha|_{U'_2 \times U'_0}) &= \deg a \cdot (\beta|_{U'_0 \times U'_0} \circ \gamma|_{U'_2 \times U'_0}) \in \text{CH}(U'_2 \times_{U_0} U'_0), \\ \deg a \cdot (\gamma^t|_{U'_0 \times U'_2} \circ \beta|_{U'_0 \times U'_0}) &= \deg b \cdot (\alpha|_{U'_2 \times U'_2} \circ \gamma^t|_{U'_0 \times U'_2}) \in \text{CH}(U'_0 \times_{U_0} U'_2), \\ (\gamma^t|_{U'_0 \times U'_2} \circ \gamma|_{U'_2 \times U'_0} \circ \alpha|_{U'_2 \times U'_2}) &= (\deg a \deg b \deg u) \cdot \alpha|_{U'_2 \times U'_2} \in \text{CH}(U'_2 \times_{U_0} U'_2), \\ (\gamma|_{U'_2 \times U'_0} \circ \gamma^t|_{U'_0 \times U'_2} \circ \beta|_{U'_0 \times U'_0}) &= (\deg a \deg b \deg u) \cdot \beta|_{U'_0 \times U'_0} \in \text{CH}(U'_0 \times_{U_0} U'_0). \end{aligned}$$

Obviously

$$\begin{aligned} \alpha|_{U'_2 \times U'_2} &= [\Gamma_{a|U'_2}^t] \circ [\Gamma_{a|U'_2}], \\ \beta|_{U'_0 \times U'_0} &= [\Gamma_{b|U'_0}^t] \circ [\Gamma_{b|U'_0}], \\ \gamma|_{U'_2 \times U'_0} &= [\Gamma_{b|U'_0}^t] \circ [\Gamma_{u|U_1}] \circ [\Gamma_{\pi|U_2}] \circ [\Gamma_{a|U'_2}]. \end{aligned}$$

Thus the claim follows from

$$[\Gamma_{u|U_1}] \circ [\Gamma_{u|U_1}^t] = \deg u \cdot [\Delta_{U_0}], \quad [\Gamma_{u|U_1}^t] \circ [\Gamma_{u|U_1}] = \deg u \cdot [\Delta_{U_1}];$$

see (4.2.1) for the equality on the right.

In view of the vanishing Lemmas 3.6.1 and 3.6.2, we see that (4.3.1) implies, that $\hat{\mathcal{H}}(\gamma/Y)$ induces a morphism

$$\hat{\mathcal{H}}(\alpha/Y) \left(\bigoplus_i R^i(fa)_*(W\mathcal{O}_{X'} \oplus W\omega_{X'})_{\mathbb{Q}} \right) \rightarrow \hat{\mathcal{H}}(\beta/Y) (b_*(W\mathcal{O}_{Y'} \oplus W\omega_{Y'})_{\mathbb{Q}})$$

and (4.3.2) implies that $\hat{\mathcal{H}}(\gamma^t/Y)$ induces a morphism

$$\hat{\mathcal{H}}(\beta/Y) (b_*(W\mathcal{O}_{Y'} \oplus W\omega_{Y'})_{\mathbb{Q}}) \rightarrow \hat{\mathcal{H}}(\alpha/Y) \left(\bigoplus_i R^i(fa)_*(W\mathcal{O}_{X'} \oplus W\omega_{X'})_{\mathbb{Q}} \right).$$

By (4.3.3) and (4.3.4) these morphisms are inverse to each other, up to multiplication with $(\deg a \deg b \deg u)$. By Proposition 4.2.7 a^* induces for all i an isomorphism $R^i f_*(W\mathcal{O}_X \oplus W\omega_X)_{\mathbb{Q}} \rightarrow \hat{\mathcal{H}}(\alpha/Y)(R^i(fa)_*(W\mathcal{O}_{X'} \oplus W\omega_{X'})_{\mathbb{Q}})$ and b^* induces an isomorphism $(W\mathcal{O}_Y \oplus W\omega_Y)_{\mathbb{Q}} \rightarrow \hat{\mathcal{H}}(\beta/Y)(b_*(W\mathcal{O}_{Y'} \oplus W\omega_{Y'})_{\mathbb{Q}})$. This gives $R^i f_* W\mathcal{O}_{X, \mathbb{Q}} = 0 = R^i f_* W\omega_{X, \mathbb{Q}}$, for all $i \geq 1$. It also gives isomorphisms in cohomological degree 0, but it is not immediately clear that these coincide with pullback and pushforward. But since X_1 is normal and $\pi : X \rightarrow X_1$ is birational, the pullback $\pi^* : W\mathcal{O}_{X_1} \rightarrow \pi_* W\mathcal{O}_X$ clearly is an

isomorphism and hence so is $f^* : W\mathcal{O}_{Y,\mathbb{Q}} \rightarrow f_*W\mathcal{O}_{X,\mathbb{Q}}$, by Lemma 4.2.4. Now the statement follows from the next lemma. \square

LEMMA 4.3.4. *Let $f : X \rightarrow Y$ be a proper morphism between k -schemes of the same pure dimension d and assume that the pullback morphism $f^* : W\mathcal{O}_{Y,\mathbb{Q}} \xrightarrow{\sim} Rf_*W\mathcal{O}_{X,\mathbb{Q}}$ is an isomorphism in $D(\widehat{dRW}_{Y,\mathbb{Q}})$. Then the pushforward (see Proposition 4.1.4, (6))*

$$f_* : f_*W\omega_{X,\mathbb{Q}} \xrightarrow{\sim} W\omega_{Y,\mathbb{Q}}$$

is an isomorphism. More precisely, there exists a natural number $N \geq 1$ such that kernel and cokernel of $f_*W_n\omega_X \rightarrow W_n\omega_Y$ are N -torsion for all $n \geq 1$.

Proof. By assumption we find a natural number $N \geq 1$ such that $R^i f_*W\mathcal{O}_X$, $i \geq 1$, as well as kernel and cokernel of $f^* : W\mathcal{O}_Y \rightarrow f_*W\mathcal{O}_X$ are all N -torsion. It follows from the short exact sequence

$$0 \rightarrow W\mathcal{O} \xrightarrow{V^n} W\mathcal{O} \rightarrow W_n\mathcal{O} \rightarrow 0$$

that there exists a natural number $M \geq 1$ (e.g. $M = N^2$) such that $R^i f_*W_\bullet\mathcal{O}_X$, $i \geq 1$, as well as kernel and cokernel of $f^* : W_\bullet\mathcal{O}_Y \rightarrow f_*W_\bullet\mathcal{O}_X$ are M -torsion. Let C_n be the cone of $f^* : W_n\mathcal{O}_Y \rightarrow Rf_*W_n\mathcal{O}_X$ in $D_{\text{qc}}^b(W_n\mathcal{O}_Y)$. (Here we write f_* instead of $W_n(f)_*$ etc.) Then the above can be rephrased by saying, that $H^i(C_n)$ is M -torsion for all $i \in \mathbb{Z}$ and all $n \geq 1$. Now applying the dualizing functor $D_{W_n Y}$ to the triangle in $D_{\text{qc}}^b(W_n\mathcal{O}_Y)$

$$W_n\mathcal{O}_Y \rightarrow Rf_*W_n\mathcal{O}_X \rightarrow C_n \rightarrow W_n\mathcal{O}_Y[1]$$

and using the duality isomorphism $D_{W_n Y}Rf_* \cong Rf_*D_{W_n X}$ yields a triangle in $D_{\text{qc}}^b(W_n\mathcal{O}_Y)$

$$D_{W_n Y}(C_n) \rightarrow Rf_*K_{X,n} \rightarrow K_{Y,n} \rightarrow D_{W_n Y}(C_n[-1]).$$

Taking H^{-d} , we obtain an exact sequence

$$\mathcal{E}\text{xt}^{-d}(C_n, K_{Y,n}) \rightarrow R^{-d}f_*K_{X,n} \rightarrow W_n\omega_Y \rightarrow \mathcal{E}\text{xt}^{-d}(C_n[-1], K_{Y,n}).$$

As in Proposition 4.1.4, (6) the morphism in the middle is just the pushforward $f_* : f_*W_n\omega_X \rightarrow W_n\omega_Y$. Consider the spectral sequence

$$E_2^{i,j} = \mathcal{E}\text{xt}^i(H^{-j}(C_n), K_{Y,n}) \Rightarrow \mathcal{E}\text{xt}^{i+j}(C_n, K_{Y,n}).$$

The filtration on $\mathcal{E}\text{xt}^{i+j}(C_n, K_{Y,n})$ induced by this spectral sequence is finite and the E_2 -terms are M -torsion. Thus the groups $\mathcal{E}\text{xt}^{-d}(C_n[-1], K_{Y,n})$ and $\mathcal{E}\text{xt}^{-d}(C_n, K_{Y,n})$ are M^r -torsion, for some $r \gg 0$. In fact r only depends on the length of the filtration of the above spectral sequence and since this length is bounded for all $n \geq 1$, we may choose r to work for all n . It follows that kernel and cokernel of $f_* : f_*W_n\omega_X \rightarrow W_n\omega_Y$ are M^r -torsion; hence kernel and cokernel of the limit $f_* : f_*W\omega_X \rightarrow W\omega_Y$ as well. This yields the statement. \square

4.4. RATIONAL AND WITT-RATIONAL SINGULARITIES. A special class of singularities which appear naturally in higher dimensional geometry are the *rational* singularities. Essentially, rational singularities do not affect the cohomological properties of the structure sheaf.

DEFINITION 4.4.1 ([KKMS73, p. 50]). Let S be a normal variety and $f : X \rightarrow S$ a resolution of singularities (i.e. f is projective and birational and X is smooth). We say that f is a *rational resolution* if

- (1) $R^i f_* \mathcal{O}_X = 0$, for $i > 0$,
- (2) $R^i f_* \omega_X = 0$ for $i > 0$ (this always holds by Grauert-Riemenschneider if the characteristic of the ground field is zero (see e.g. [KM98, Cor.2.68]), but is needed in positive characteristic).

We say that Y has *rational singularities* if a rational resolution exists.

An immediate problem with the definition of rational singularities in positive characteristic is that the existence of a resolution of singularities is assumed. For example, tame quotient singularities are rational singularities provided that a resolution of singularities exists [CR09, Theorem 2].

If an integral normal scheme Y over a field has one rational resolution, then all resolutions are rational, i.e. rational singularities are an intrinsic property of Y . (In characteristic zero, this was proved by Hironaka, see [CR09, Theorem 1] for the characteristic p case.)

In characteristic zero, Kovács [Kov00] observed that one can replace condition (1) in Definition 4.4.1 by the following condition: there is an alteration $f : X \rightarrow S$ such that the natural morphism

$$\mathcal{O}_S \rightarrow Rf_* \mathcal{O}_X$$

admits a splitting in the derived category of coherent sheaf on S . The main tool in the proof is Grauert-Riemenschneider vanishing, and this characterization does not hold in positive characteristic.

In order to study congruence formulas for the number of points of a variety over a finite field, Blickle and Esnault [BE08] introduced the notion of Witt-rational singularities.

DEFINITION 4.4.2 ([BE08, Def. 2.3]). Let S be an integral k -scheme and $f : X \rightarrow S$ a generically étale alteration with X a smooth k -scheme. We say that S has *BE-Witt-rational singularities* if the natural morphism

$$W\mathcal{O}_S \otimes_{\mathbb{Z}} \mathbb{Q} \rightarrow Rf_* W\mathcal{O}_X \otimes_{\mathbb{Z}} \mathbb{Q}$$

admits a splitting in the derived category of sheaves of abelian groups on X .

We call the singularities defined in [BE08] BE-Witt-rational singularities, rather than Witt-rational singularities, because we will redefine Witt-rational singularities in 4.4.4. We remark:

PROPOSITION 4.4.3. *The notion of BE-Witt-rational singularities is independent of the chosen generically étale alteration. More precisely, if an integral scheme S has BE-Witt-rational singularities, then for any alteration*

(not necessarily generically étale) $g : Y \rightarrow S$, with Y smooth, the pullback $g^* : W\mathcal{O}_S \otimes_{\mathbb{Z}} \mathbb{Q} \rightarrow Rg_*W\mathcal{O}_Y \otimes_{\mathbb{Z}} \mathbb{Q}$ admits a splitting in the derived category of sheaves of abelian groups on X .

Proof. Obviously it suffices to prove that if $f : X \rightarrow Y$ is an alteration between smooth schemes, then the composition

$$W\mathcal{O}_Y \xrightarrow{f^*} Rf_*W\mathcal{O}_X \xrightarrow{f_*} W\mathcal{O}_Y$$

is multiplication with $[k(X) : k(Y)]$, where f_* is the pushforward from Definition 2.3.1. It suffices to check this on some non-empty open subscheme U of Y such that $f|f^{-1}(U)$ is finite. Thus the statement follows from Proposition 2.3.4 (Gros). \square

DEFINITION 4.4.4. We say that an integral k -scheme S has *Witt-rational singularities* if for any quasi-resolution (see Definition 4.3.1) $f : X \rightarrow S$ the following conditions are satisfied:

- (1) $f^* : W\mathcal{O}_{S,\mathbb{Q}} \xrightarrow{\cong} f_*W\mathcal{O}_{X,\mathbb{Q}}$ is an isomorphism.
- (2) $R^i f_*W\mathcal{O}_{X,\mathbb{Q}} = 0$, for all $i \geq 1$.
- (3) $R^i f_*W\omega_{X,\mathbb{Q}} = 0$, for all $i \geq 1$.

In case only the first two properties are satisfied we say S has *Witt-rational singularities*. (See Notation 3.7.9 for the meaning of the subscript \mathbb{Q} .)

Remark 4.4.5. Notice that if S is normal, then condition (1) above is automatically satisfied. Indeed, each quasi-resolution $f : X \rightarrow S$ can be factored as $X \xrightarrow{\pi} X_1 \xrightarrow{u} S$ with X_1 normal, π projective and birational and u an universal homeomorphism; thus condition (1) is satisfied by Lemma 4.2.4.

PROPOSITION 4.4.6. *Let S be an integral k -scheme. Then the following statements are equivalent:*

- (1) S has Witt-rational singularities.
- (2) There exists a quasi-resolution $f : X \rightarrow S$ satisfying (1), (2), (3) of Definition 4.4.4.
- (3) There exists a quasi-resolution $f : X \rightarrow S$, such that there are isomorphisms in $D^b(\widehat{\mathrm{dRW}}_{S,\mathbb{Q}})$

(4.4.1)

$$f^* : W\mathcal{O}_{S,\mathbb{Q}} \xrightarrow{\cong} Rf_*W\mathcal{O}_{X,\mathbb{Q}}, \quad Rf_*W\omega_{X,\mathbb{Q}} \cong f_*W\omega_{X,\mathbb{Q}}[0] \xrightarrow{f_*} W\omega_{S,\mathbb{Q}}.$$

- (4) For all quasi-resolutions $f : X \rightarrow S$ the morphisms (4.4.1) are isomorphisms.

Proof. Clearly (1) \Rightarrow (2) and (4) \Rightarrow (1). (2) \Rightarrow (3) follows from Lemma 4.3.4. For (3) \Rightarrow (4) notice that by de Jong (see Remark 4.3.2) any two quasi-resolutions of S can be dominated by a third one. Thus the statement follows from Theorem 4.3.3. \square

There is an obvious analog of this proposition for Witt-rational singularities.

COROLLARY 4.4.7. *Topological finite quotients over k have Witt-rational singularities.*

Remark 4.4.8. In characteristic 0 finite quotients always have rational singularities (see e.g. [KM98, Prop. 5.13]). In characteristic $p > 0$ this is not the case. Indeed let $G = \mathbb{Z}/p^n\mathbb{Z}$ act linearly on a finite dimensional k -vector space, where k is assumed to be algebraically closed. Then it is shown in [ES80], that $\mathbb{A}(V)/G$ is not CM, provided that $\dim_k V > \dim_k V^G + 2$. In particular $\mathbb{A}(V)/G$ cannot have rational singularities in the sense of Definition 4.4.1, which are always CM. This also shows that Witt-rational singularities do not need to be CM.

PROPOSITION 4.4.9. *Let $u : Y \rightarrow Y'$ be a universal homeomorphism between normal schemes. Then Y has Witt-rational singularities if and only if Y' has Witt-rational singularities.*

Proof. If $f : X \rightarrow Y$ is a quasi-resolution then $u \circ f$ is a quasi-resolution. For all $i > 0$ we get


$$R^i(u \circ f)_*W\mathcal{O}_{X,\mathbb{Q}} = u_*R^if_*W\mathcal{O}_{X,\mathbb{Q}}, \quad R^i(u \circ f)_*W\omega_{X,\mathbb{Q}} = u_*R^if_*W\omega_{X,\mathbb{Q}},$$

and thus

$$\begin{aligned} R^i(u \circ f)_*W\mathcal{O}_{X,\mathbb{Q}} = 0 &\Leftrightarrow R^if_*W\mathcal{O}_{X,\mathbb{Q}} = 0, \\ R^i(u \circ f)_*W\omega_{X,\mathbb{Q}} = 0 &\Leftrightarrow R^if_*W\omega_{X,\mathbb{Q}} = 0. \end{aligned}$$

□

DEFINITION 4.4.10. Let S be a k -scheme and X and Y two integral S -schemes. We say that X and Y are *quasi-birational over S* if there exists a commutative diagram



with π_X and π_Y quasi-resolutions (see Definition 4.3.1). We say that the triple (Z, π_X, π_Y) (or just Z if we do not need to specify π_X and π_Y) is a *quasi-birational correspondence between X and Y* .

Since quasi-resolutions always exist (see Remark 4.3.2), two integral *projective* S -schemes X and Y are quasi-birational over S if and only if the generic points of X and Y map to the same point η in S and there exists a field L with a homomorphism $\mathcal{O}_{S,\eta} \rightarrow L$ and inclusions of $\mathcal{O}_{S,\eta}$ -algebras $k(X) \hookrightarrow L$, $k(Y) \hookrightarrow L$, which make L a finite and purely inseparable field extension of $k(X)$ and $k(Y)$. In particular this is the case if $k(X)$ and $k(Y)$ are isomorphic as $\mathcal{O}_{S,\eta}$ -algebras.

COROLLARY 4.4.11. *Let S be a k -scheme and $f : X \rightarrow S$ and $g : Y \rightarrow S$ two S -schemes, which are integral, have Witt-rational singularities and are quasi-birational over S to each other. Then the choice of a quasi-birational correspondence between X and Y induces isomorphisms in $D^b(\widehat{\text{dRW}}_S)$*

$$(4.4.2) \quad Rf_*W\mathcal{O}_{X,\mathbb{Q}} \cong Rg_*W\mathcal{O}_{Y,\mathbb{Q}}, \quad Rf_*W\omega_{X,\mathbb{Q}} \cong Rg_*W\omega_{Y,\mathbb{Q}}.$$

Moreover, two quasi-birational correspondences Z and Z' induce the same isomorphisms if there exists a field L with a homomorphism $\mathcal{O}_{S,\eta} \rightarrow L$ ($\eta \in S$ being the common image of the generic points of X and Y) and inclusions of $\mathcal{O}_{S,\eta}$ -algebras $k(Z) \hookrightarrow L$, $k(Z') \hookrightarrow L$, which make L a finite and purely inseparable field extension of $k(Z)$ and $k(Z')$, such that the composite inclusions $k(X), k(Y) \hookrightarrow k(Z) \hookrightarrow L$ and $k(X), k(Y) \hookrightarrow k(Z') \hookrightarrow L$ are equal.

Proof. A quasi-birational correspondence (Z, π_X, π_Y) between X and Y induces an isomorphism

$$Rf_*W\mathcal{O}_{X,\mathbb{Q}} \xrightarrow{\pi_X^* \simeq} Rf_*R\pi_{X*}W\mathcal{O}_{Z,\mathbb{Q}} \cong Rg_*R\pi_{Y*}W\mathcal{O}_{Z,\mathbb{Q}} \xleftarrow{\pi_Y^* \simeq} Rg_*W\mathcal{O}_{Y,\mathbb{Q}},$$

and similar for $W\omega$. For the second statement first notice that if (Z, π_X, π_Y) and (Z', π'_X, π'_Y) are two quasi-birational correspondences between X and Y and if there is a quasi-resolution $a : Z' \rightarrow Z$ such that $\pi'_X = \pi_X \circ a$ and $\pi'_Y = \pi_Y \circ a$, then they induce the same isomorphisms (4.4.2). If we are given two arbitrary quasi-birational correspondences Z and Z' between X and Y and a field L as is the statement of the corollary, then we can take a quasi-resolution of the closure of the image of $\text{Spec } L \rightarrow Z \times_S Z'$ to obtain a quasi-birational correspondence Z'' between X and Y mapping via a quasi-resolution to Z and Z' and is compatible with $\pi_X, \pi'_X, \pi_Y, \pi'_Y$ in the obvious sense. This proves the statement. \square

COROLLARY 4.4.12. *In the situation of Corollary 4.4.11 assume that S is integral and f and g are generically finite and purely inseparable. Then any quasi-birational correspondence between X and Y induces the same isomorphism in $D^b(\widehat{\text{dRW}}_S)$*

$$Rf_*W\mathcal{O}_{X,\mathbb{Q}} \cong Rg_*W\mathcal{O}_{Y,\mathbb{Q}}, \quad Rf_*W\omega_{X,\mathbb{Q}} \cong Rg_*W\omega_{Y,\mathbb{Q}}.$$

COROLLARY 4.4.13. *Let S be a k -scheme and $f : X \rightarrow S$ be an integral and projective S -scheme, which has Witt-rational singularities. Let $k(X)^{\text{perf}}$ be the perfect closure of $k(X)$ and $\eta \in S$ the image of the generic point of X . Then $\text{End}_{\mathcal{O}_{S,\eta}\text{-alg}}(k(X)^{\text{perf}})$ is acting on $Rf_*W\mathcal{O}_{X,\mathbb{Q}}$ and $Rf_*W\omega_{X,\mathbb{Q}}$ as objects in $D^b(\widehat{\text{dRW}}_S)$.*

Proof. An element σ in $\text{End}_{\mathcal{O}_{S,\eta}\text{-alg}}(k(X)^{\text{perf}})$ will when composed with $k(X) \hookrightarrow k(X)^{\text{perf}}$ factor over a finite and purely inseparable extension L of $k(X)$. By the remark after Definition 4.4.10 it hence gives rise to a quasi-birational correspondence of X with itself and thus yields the promised well-defined action by Corollary 4.4.11. \square

Remark 4.4.14. The above corollaries have obvious analogs for $W\mathcal{O}$ -rational singularities.

COROLLARY 4.4.15. *Let X and Y be two integral k -schemes, which have $W\mathcal{O}$ -rational singularities and are quasi-birational over k . Then in $\widehat{\text{dRW}}_{k,\mathbb{Q}}$*

$$H^i(X, W\mathcal{O}_{X,\mathbb{Q}}) \cong H^i(Y, W\mathcal{O}_{Y,\mathbb{Q}}), \quad \text{for all } i \geq 0.$$

In particular, if X and Y are projective then [BBE07, Thm 1.1] yields a Frobenius equivariant isomorphism

$$H_{\text{rig}}^i(X/K)^{<1} \cong H_{\text{rig}}^i(Y/K)^{<1} \quad \text{for all } i \geq 0,$$

where $K = \text{Frac}(W(k))$ and $H_{\text{rig}}^i(X/K)^{<1}$ denotes the part of rigid cohomology on which the Frobenius acts with slope < 1 .

Proof. Apply Corollary 4.4.11 in the case $S = k$. □

We will also give some results on the torsion, see Theorem 5.1.10 and Theorem 5.1.13.

COROLLARY 4.4.16. *Let k be a finite field. Let X and Y be two quasi-birational integral and projective k -schemes, which have $W\mathcal{O}$ -rational singularities. Then for any finite field extension k' of k we have*

$$|X(k')| \equiv |Y(k')| \pmod{|k'|}.$$

Proof. This follows from Corollary 4.4.15 and [BBE07, Cor. 1.3]. □

In the case where X and Y are smooth, integral and proper, the above corollary was proved in [Eke83, Cor. 3, (i)]. In case there is a morphism $f : X \rightarrow Y$, which is birational and X is smooth and projective and $Y = Z/G$ is the quotient under a finite group G of a smooth projective scheme Z , this was proved in [Cha09, Thm 4.5].

We investigate the properties of Witt-rational singularities a little bit further.

PROPOSITION 4.4.17. *Consider the following properties on an integral k -scheme S :*

- (1) S has rational singularities.
- (2) S has Witt-rational singularities.
- (3) S has $W\mathcal{O}$ -rational singularities.
- (4) S has BE-Witt-rational singularities.

Then

$$(1) \Rightarrow (2) \Rightarrow (3) \Rightarrow (4).$$

Furthermore the first implication is strict, by Remark 4.4.8 above.

Proof. (1) \Rightarrow (2): By assumption there exists a resolution $f : X \rightarrow S$. The exact sequences

$$0 \rightarrow W_{n-1}\mathcal{O}_X \xrightarrow{V} W_n\mathcal{O}_X \rightarrow \mathcal{O}_X \rightarrow 0$$

and (see Proposition 4.1.4, (7))

$$0 \rightarrow W_{n-1}\omega_X \xrightarrow{p} W_n\omega_X \xrightarrow{F^{n-1}} \omega_X \rightarrow 0$$

give us $R^i f_* W_n \mathcal{O}_X = 0 = R^i f_* W_n \omega_X$ for all $n, i \geq 1$ and also $W_n \mathcal{O}_S \cong f_* W_n \mathcal{O}_X$. Since S is CM, the isomorphism $W_n \mathcal{O}_S \cong Rf_* W_n \mathcal{O}_X$ also gives the isomorphism $f_* W_n \omega_X \cong W_n \omega_S$ via duality (see Proposition 4.1.4, (2)). Now the statement follows from the exact sequence (see Lemma 1.5.1)

$$0 \rightarrow R^1 \varprojlim_n R^{i-1} f_* E_n \rightarrow R^i f_* (\varprojlim_n E_n) \rightarrow \varprojlim_n R^i f_* E_n \rightarrow 0,$$

where $E_n \in \{W_n \mathcal{O}_X, W_n \omega_X\}$. (For the vanishing of $R^1 f_* W_n \omega_X$ use that $f_* W_n \omega_X \cong W_n \omega_S$ and hence the projection maps $f_* W_n \omega_X \rightarrow f_* W_{n-1} \omega_X$ are surjective, by Proposition 4.1.4, (4).)

(2) \Rightarrow (3): trivial.

(3) \Rightarrow (4): By Remark 4.3.2 we find a quasi-resolution of S of the form $f : X \rightarrow S$ with X a finite quotient. We thus find a smooth scheme X' with a finite and surjective morphism $g : X' \rightarrow X$. Then $h := fg : X' \rightarrow S$ is an alteration and $h^* : W\mathcal{O}_{S, \mathbb{Q}} \rightarrow Rg_* W\mathcal{O}_{X', \mathbb{Q}}$ splits by Proposition 4.4.6, (1) \Rightarrow (2), and Lemma 4.1.6, (2); a fortiori $h^* : W\mathcal{O}_S \otimes \mathbb{Q} \rightarrow Rg_* W\mathcal{O}_{X'} \otimes \mathbb{Q}$ splits. \square

4.5. COMPLEXES AND SHEAVES ATTACHED TO SINGULARITIES OF SCHEMES.

COROLLARY 4.5.1. *Let S be an integral scheme and $f : X \rightarrow S$ and $g : Y \rightarrow S$ be quasi-resolutions. Then X and Y are quasi-birational over S and any quasi-birational correspondence between X and Y induces the same isomorphism in the derived category $D^b(\widehat{\text{dRW}}_S)$*

$$Rf_* W\mathcal{O}_{X, \mathbb{Q}} \cong Rg_* W\mathcal{O}_{Y, \mathbb{Q}}, \quad Rf_* W\omega_{X, \mathbb{Q}} \cong Rg_* W\omega_{Y, \mathbb{Q}}.$$

Furthermore, if Z is another integral scheme and $h : Z \rightarrow S$ a quasi-resolution, then the isomorphisms

$$Rf_* W\mathcal{O}_{X, \mathbb{Q}} \cong Rh_* W\mathcal{O}_{Z, \mathbb{Q}}, \quad Rf_* W\omega_{X, \mathbb{Q}} \cong Rh_* W\omega_{Z, \mathbb{Q}}$$

induced by any quasi-birational correspondence between X and Z equals the isomorphism obtained by composing the isomorphisms induced by quasi-birational correspondences between X and Y and between Y and Z .

Proof. First we show that X and Y are quasi-birational over S . For this notice that $k(X) \otimes_{k(S)} k(Y)$ is a local Artin algebra. Denote by L its residue field. Then we can take a quasi-resolution of the closure of the image of $\text{Spec } L$ in $X \times_S Y$ to obtain a quasi-birational correspondence between X and Y . Let V and V' be two quasi-birational correspondences between X and Y . Denote by L'' the residue field of the local Artin algebra $k(V) \otimes_{k(S)} k(V')$. This is a purely inseparable field extension of $k(S)$. Hence there is only one embedding over $k(S)$ of $k(X)$ and $k(Y)$ into L'' . Thus by Corollary 4.4.11 the two isomorphisms induced by V and V' are equal. Finally if we have the three quasi-resolutions

f, g, h , we find a diagram of S -morphisms

$$\begin{array}{ccccc}
 V'' & \longrightarrow & V' & \longrightarrow & Z \\
 \downarrow & & \downarrow & & \\
 V & \longrightarrow & Y & & \\
 \downarrow & & & & \\
 X & & & &
 \end{array}$$

in which V is a quasi-birational correspondence between X and Y , V' is a quasi-birational correspondence between Y and Z and V'' is a quasi-birational correspondence between V and V' (and also between X and Z). The last statement of the Corollary follows. \square

DEFINITION 4.5.2. Let S be an integral k -scheme of dimension d . We define in $D^b(\widehat{\text{dRW}}_{S,\mathbb{Q}})$

$$\mathcal{WS}_{0,S} := Rf_*W\mathcal{O}_{X,\mathbb{Q}}, \quad \mathcal{WS}_{d,S} := Rf_*W\omega_{X,\mathbb{Q}},$$

where $f : X \rightarrow S$ is any quasi-resolution. This definition is independent (up to a canonical isomorphism) of the choice of the quasi-resolution f by Corollary 4.5.1.

It follows, that S has $W\mathcal{O}$ -rational singularities if and only if $\mathcal{WS}_{0,S} \cong W\mathcal{O}_S$ and it has Witt-rational singularities if and only if in addition we have $H^i(\mathcal{WS}_{d,S}) = 0$ for all $i \geq 1$ (which is equivalent to $\mathcal{WS}_{d,S} \cong W\omega_{S,\mathbb{Q}}$). Next we want to give a characterization of Witt-rational singularities using alterations.

PROPOSITION 4.5.3. Let S be an integral k -scheme of dimension d and $f : X \rightarrow S$ an alteration with X smooth. Set

$$\epsilon_f := \frac{1}{\text{deg } f} [X \times_S X]^0 \in \text{CH}^d(X \times X, P(X \times_S X))_{\mathbb{Q}},$$

where $[X \times_S X]^0$ is the cycle associated to the closure of $X_{\eta} \times_{\eta} X_{\eta}$ in $X \times_S X$ with η the generic point of S . Further set

$$\hat{\mathcal{H}}^{*,0}(X/S) := \bigoplus_i R^i f_* W\mathcal{O}_X, \quad \hat{\mathcal{H}}^{*,d}(X/S) := \bigoplus_i R^i f_* W\Omega_X^d,$$

$$\hat{\mathcal{H}}^{*,(0,d)}(X/S) := \hat{\mathcal{H}}^{*,0}(X/S) \oplus \hat{\mathcal{H}}^{*,d}(X/S).$$

Then:

- (1) The restriction of $\hat{\mathcal{H}}(\epsilon_f/S)$ to $\hat{\mathcal{H}}^{*,(0,d)}(X/S)_{\mathbb{Q}}$ is a projector, which we denote by e_f . (See Proposition 3.5.4 for the notation.)
- (2) The pullback f^* induces a natural morphism of Witt modules over S

$$f^* : W\mathcal{O}_{S,\mathbb{Q}} \rightarrow e_f \hat{\mathcal{H}}^{*,(0,d)}(X/S)_{\mathbb{Q}}.$$

(3) If $g : Y \rightarrow S$ is another alteration with Y smooth. Then set

$$\gamma_{f,g} := \frac{1}{\deg f} [X \times_S Y]^0 \in \text{CH}^d(X \times Y, P(X \times_S Y)),$$

where $[X \times_S Y]^0$ is the cycle associated to the closure of $X_\eta \times_\eta Y_\eta$ in $X \times_S Y$. Then $\hat{\mathcal{H}}(\gamma_{f,g}/S) : \hat{\mathcal{H}}(X/S) \rightarrow \hat{\mathcal{H}}(Y/S)$ induces an isomorphism

$$(4.5.1) \quad \hat{\mathcal{H}}(\gamma_{f,g}/S) : e_f(\hat{\mathcal{H}}^{*,(0,d)}(X/S))_{\mathbb{Q}} \xrightarrow{\cong} e_g(\hat{\mathcal{H}}^{*,(0,d)}(Y/S))_{\mathbb{Q}},$$

which is compatible with the pullback morphism from (2).

Proof. Take a non-empty open and smooth subset U of S such that the pullback of f - and in case (3) also of g - over U is finite and surjective and hence also flat; the pullbacks are denoted by $f_U : X_U \rightarrow U$ and $g_U : Y_U \rightarrow U$. Notice that the restriction of $[X \times_S X]^0$ and $[X \times_S Y]^0$ over U equal $[X_U \times_U X_U]$ and $[X_U \times_U Y_U]$.

(1) We have to show $e_f \circ e_f - e_f = 0$ on $\hat{\mathcal{H}}^{*,(0,d)}(X/S)_{\mathbb{Q}}$. By the same argument as in the proof of Theorem 4.3.3 (using the vanishing Lemmas 3.6.1 and 3.6.2) it suffices to prove

$$[X_U \times_U X_U] \circ [X_U \times_U X_U] - \deg f \cdot [X_U \times_U X_U] = 0.$$

This follows immediately from $[X_U \times_U X_U] = [\Gamma_{f_U}^t] \circ [\Gamma_{f_U}]$.

(2) The morphism e_f , in particular gives a projector $e_f : f_* W\mathcal{O}_{X,\mathbb{Q}} \rightarrow f_* W\mathcal{O}_{S,\mathbb{Q}}$. Thus we need to show, that $e_f \circ f^* = f^*$ on $W\mathcal{O}_{S,\mathbb{Q}}$. It suffices to prove this over U ; thus the statement follows from $\deg f \cdot e_{f_U} = f_U^* \circ f_{U*}$.

(3) To prove that $\gamma_{f,g}$ is an isomorphism on $e_f(\hat{\mathcal{H}}^{*,(0,d)}(X/S)_{\mathbb{Q}})$ with inverse $\frac{1}{\deg g} \hat{\mathcal{H}}([Y \times_S X]^0/S)$ it suffices to show that the following cycles in $\text{CH}^d(X \times Y, P(X \times_S Y))$, etc.,

$$\begin{aligned} & \deg g \cdot [X \times_S Y]^0 \circ [X \times_S X]^0 - \deg f \cdot [Y \times_S Y]^0 \circ [X \times_S Y]^0 \\ & \deg f \cdot [Y \times_S X]^0 \circ [Y \times_S Y]^0 - \deg g \cdot [X \times_S X]^0 \circ [Y \times_S X]^0 \\ & [Y \times_S X]^0 \circ [X \times_S Y]^0 \circ [X \times_S X]^0 - (\deg f \deg g) \cdot [X \times_S X]^0 \\ & [X \times_S Y]^0 \circ [Y \times_S X]^0 \circ [Y \times_S Y]^0 - (\deg f \deg g) \cdot [Y \times_S Y]^0 \end{aligned}$$

act as zero on $\hat{\mathcal{H}}^{*,(0,d)}(X/S)_{\mathbb{Q}}$ and $\hat{\mathcal{H}}^{*,(0,d)}(Y/S)_{\mathbb{Q}}$ respectively. By the same argument as in the proof of Theorem 4.3.3 (using the vanishing Lemmas 3.6.1 and 3.6.2) it suffices to prove that the pullback along U of the above cycles vanish. This follows easily from $[X_U \times_U X_U] = [\Gamma_{f_U}^t] \circ [\Gamma_{f_U}]$, $[X_U \times_U Y_U] = [\Gamma_{g_U}^t] \circ [\Gamma_{f_U}]$ and $[\Gamma_{f_U}] \circ [\Gamma_{f_U}^t] = \deg f \cdot [\Delta_U]$, etc. This yields the isomorphism (4.5.1). It is compatible with the pullback from (2), since on $W\mathcal{O}_{S,\mathbb{Q}}$ we have $\hat{\mathcal{H}}(\gamma_{f,g}/S)f^* = g^*$. Indeed over U we have $\hat{\mathcal{H}}(\gamma_{f_U,g_U}/U) = \frac{1}{\deg f} \cdot g_U^* \circ f_{U*}$; thus it holds on U and hence on all of S . \square

DEFINITION 4.5.4. Let S be an integral k -scheme of dimension d and $f : X \rightarrow S$ an alteration with X smooth. Then using the notations from Proposition 4.5.3

we define

$$\mathcal{A}^0(X/S) := \operatorname{coker}(W\mathcal{O}_{S,\mathbb{Q}} \xrightarrow{f^*} e_f \hat{\mathcal{H}}^{*,0}(X/S)_{\mathbb{Q}}) = \frac{f_* W\mathcal{O}_{X,\mathbb{Q}}}{W\mathcal{O}_{S,\mathbb{Q}}} \bigoplus_{i \geq 1} e_f R^i f_* W\mathcal{O}_{X,\mathbb{Q}}$$

and

$$\mathcal{A}^d(X/S) := e_f \hat{\mathcal{H}}^{>0,d}(X/S) = \bigoplus_{i \geq 1} e_f R^i f_* W\Omega_{X,\mathbb{Q}}^d.$$

Then by Proposition 4.5.3 $\mathcal{A}^0(X/S)$ and $\mathcal{A}^d(X/S)$ are independent of $f : X \rightarrow S$ up to a canonical isomorphism and are therefore denoted

$$\mathcal{A}^0(S) := \mathcal{A}^0(X/S), \quad \mathcal{A}^d(S) := \mathcal{A}^d(X/S), \quad \mathcal{A}(S) := \mathcal{A}^0(S) \oplus \mathcal{A}^d(S).$$

Remark 4.5.5. If S is normal, then $\mathcal{A}(S)$ is a direct summand of $\mathcal{H}^{*,(0,d)}(X/S)_{\mathbb{Q}}$ for any alteration $f : X \rightarrow S$ with X smooth.

THEOREM 4.5.6. *Let S be an integral k -scheme. Then there are isomorphisms in $\widehat{\operatorname{dRW}}_{S,\mathbb{Q}}$*

$$\mathcal{A}^0(S) \cong \frac{H^0(\mathcal{WS}_{0,S})}{W\mathcal{O}_{S,\mathbb{Q}}} \bigoplus_{i \geq 1} H^i(\mathcal{WS}_{0,S}), \quad \mathcal{A}^d(S) \cong \bigoplus_{i \geq 1} H^i(\mathcal{WS}_{d,S}).$$

In particular:

- (1) S has $W\mathcal{O}$ -rational singularities $\iff \mathcal{A}^0(S) = 0$.
- (2) S has Witt-rational singularities $\iff \mathcal{A}(S) = 0$.

Proof. By de Jong (see Remark 4.3.2) there exists an alteration $f : X \rightarrow S$ which factors as

$$f : X \xrightarrow{h} Y \xrightarrow{g} S,$$

where X is smooth, $Y = X/G$, for G a finite group, h is the quotient map and g is a quasi-resolution. Thus the following equalities hold by definition for all i

$$H^i(\mathcal{WS}_{0,S}) = R^i g_* W\mathcal{O}_{Y,\mathbb{Q}}, \quad H^i(\mathcal{WS}_{d,S}) = R^i g_* W\omega_{Y,\mathbb{Q}}.$$

Since Y is normal and h is finite and surjective it is also universally equidimensional. Thus $[X \times_Y X] \in \operatorname{CH}^d(X \times X, P(X \times_Y X))$, where $d = \dim S$. We denote by $[X \times_S X]^0$ the cycle associated to the closure of $X_\eta \times_\eta X_\eta$ in $X \times_S X$, where η is the generic point of S . We claim

$$(4.5.2) \quad [X \times_S X]^0 = \operatorname{deg} g \cdot [X \times_Y X] \quad \text{in } \operatorname{CH}^d(X \times X, P(X \times_S X)).$$

Indeed it suffices to check this over a smooth and dense open subscheme U of S , over which g is a universal homeomorphism and Y is smooth. But then

$$\begin{aligned} [X_U \times_U X_U]^0 &= [X_U \times_U X_U] \\ &= [\Gamma_{f_U}^t] \circ [\Gamma_{f_U}] \\ &= [\Gamma_{h_U}^t] \circ [\Gamma_{g_U}^t] \circ [\Gamma_{g_U}] \circ [\Gamma_{h_U}] \\ &= \operatorname{deg} g \cdot [\Gamma_{h_U}^t] \circ [\Gamma_{h_U}] && \text{by (4.2.1)} \\ &= \operatorname{deg} g \cdot [X_U \times_{Y_U} X_U]. \end{aligned}$$

Hence the claim. Now $\hat{\mathcal{H}}([X \times_Y X]/Y)$ induces an endomorphism of $h_*(W\mathcal{O}_X \oplus W\Omega_X^d)$ which actually equals $h^* \circ h_*$, for h_* and h^* as in Definition 4.1.5 (see (4.2.2).) It thus follows from Proposition 3.5.10 and (4.5.2) that $e_f = \frac{1}{\deg f} \hat{\mathcal{H}}([X \times_S X]^0/S)$ factors for each $i \geq 0$ as

$$\begin{array}{ccc}
 R^i f_*(W\mathcal{O}_X \oplus W\Omega_X^d)_{\mathbb{Q}} & \xrightarrow{e_f} & R^i f_*(W\mathcal{O}_X \oplus W\Omega_X^d)_{\mathbb{Q}} \\
 \searrow \frac{1}{\deg h} \cdot h_* & & \nearrow h^* \\
 & R^i g_*(W\mathcal{O}_Y \oplus W\omega_Y)_{\mathbb{Q}} &
 \end{array}$$

Taking into account that $h_* \circ h^*$ is multiplication with the degree of h , this yields the statement of the theorem. \square

4.6. ISOLATED SINGULARITIES. In this section we will relate the sheaf $\mathcal{A}^0(S)$ (Definition 4.5.4) for a normal variety S with an isolated singularity to the Witt vector cohomology of the exceptional set in a suitably good resolution of singularities \tilde{S} . For this, we need to compute the higher direct images of $W(\mathcal{O}_{\tilde{S}})_{\mathbb{Q}}$.

PROPOSITION 4.6.1. *Let $f : \tilde{X} \rightarrow \tilde{Y}$ be a proper morphism of schemes. Let $Y_0 \subset \tilde{Y}$ be a closed subset, we denote by $Y = \tilde{Y} \setminus Y_0$ the complement. We consider the cartesian diagrams*

$$\begin{array}{ccccc}
 X & \longrightarrow & \tilde{X} & \longleftarrow & X_0 \\
 \downarrow & & \downarrow & & \downarrow \\
 & \square & & \square & \\
 Y & \longrightarrow & \tilde{Y} & \longleftarrow & Y_0.
 \end{array}$$

Let $\mathcal{I} \subset \mathcal{O}_{\tilde{X}}$ be a sheaf of ideals for X_0 . Suppose that $R^i f_* \mathcal{O}_X = 0$ for all $i > 0$. Then

$$R^i f_* W(\mathcal{I})_{\mathbb{Q}} = 0 \quad \text{for all } i > 0.$$

In order to prove Proposition 4.6.1 we need several Lemmas.

LEMMA 4.6.2. *Let X be a scheme and $\mathcal{I} \subset \mathcal{O}_X$ a sheaf of ideals. For all integers $a \geq 1$ the natural map*

$$W(\mathcal{I}^a)_{\mathbb{Q}} \rightarrow W(\mathcal{I})_{\mathbb{Q}}$$

is an isomorphism.

Proof. The proof is the same as in [BBE07, Proposition 2.1(ii)]. \square

LEMMA 4.6.3. *With the assumptions of Proposition 4.6.1. There are $N, a \geq 0$ such that for all $r \geq N$ and all $n \geq 1$ the morphism*

$$R^i f_* W_n(\mathcal{I}^{r+a}) \rightarrow R^i f_* W_n(\mathcal{I}^r),$$

induced by $\mathcal{I}^{r+a} \subset \mathcal{I}^r$, vanishes for all integers $i > 0$.

Proof. [BBE07, Lemma 2.7(i)]. \square

Proof of Proposition 4.6.1. Choose N as in Lemma 4.6.3. By using Lemma 4.6.2 we may replace \mathcal{I} by \mathcal{I}^N . Thus we may assume that $N = 1$. The image of the Frobenius acting on $W_n(\mathcal{I})$ is contained in $W_n(\mathcal{I}^p)$:

$$\begin{array}{ccc} W_n(\mathcal{I}) & \xrightarrow{F} & W_n(\mathcal{I}) \\ & \searrow \exists! & \nearrow \\ & & W_n(\mathcal{I}^p) \end{array}$$

and therefore F^a (with a as in Lemma 4.6.3) is zero on $R^i f_* W_n(\mathcal{I})$ for all $i > 0$. We continue as in the proof of [BBE07, Theorem 2.4]. Since F^a acts as zero on $\varprojlim_n R^i f_* W_n(\mathcal{I})$ and $R^1 \varprojlim_n R^i f_* W_n(\mathcal{I})$ for all $i \geq 1$, we obtain via the exact sequence

$$0 \rightarrow R^1 \varprojlim_n R^{i-1} f_* W_n(\mathcal{I}) \rightarrow R^i f_*(W(\mathcal{I})) \rightarrow \varprojlim_n R^i f_* W_n(\mathcal{I}) \rightarrow 0$$

that F^{2a} acts as zero on $R^i f_* W(\mathcal{I})$ for all $i > 0$ (we use that $R^1 \varprojlim_n f_* W_n(\mathcal{I}) = 0$). Thus the relation $FV = p$ implies that p^{2a} kills $R^i f_* W(\mathcal{I})$ for all $i > 0$. \square

COROLLARY 4.6.4. *Let S be an integral scheme with an isolated singularity at the closed point $s \in S$. Let $f : X \rightarrow S$ be projective and birational. Suppose that f is an isomorphism over $S \setminus \{s\}$; we denote by E the closed set $f^{-1}(s)$ equipped with some scheme structure. Then*

$$R^i f_* W\mathcal{O}_{X, \mathbb{Q}} = H^i(E, W\mathcal{O}_{E, \mathbb{Q}}) \quad \text{for all } i > 0,$$

where $H^i(E, W\mathcal{O}_{E, \mathbb{Q}})$ is considered as skyscraper sheaf supported in s . Moreover, the morphism is compatible with the Frobenius and Verschiebung action.

Proof. Let \mathcal{I} be the sheaf of ideals for E . We obtain an exact sequence

$$0 \rightarrow W\mathcal{I}_{\mathbb{Q}} \rightarrow W\mathcal{O}_{X, \mathbb{Q}} \rightarrow W\mathcal{O}_{E, \mathbb{Q}} \rightarrow 0.$$

Proposition 4.6.1 implies that the higher direct images of $W\mathcal{I}_{\mathbb{Q}}$ vanish, which proves the statement. \square

LEMMA 4.6.5. *Let E be a scheme. Write $E = \cup_{i=1}^r E_i$ with E_i irreducible for all i . There is a spectral sequence*

$$E_1^{s,t} = \bigoplus_{1 \leq \iota_0 < \iota_1 < \dots < \iota_s \leq r} H^t(\cap_{j=0}^s E_{\iota_j}, W\mathcal{O}_{\cap_{j=0}^s E_{\iota_j}, \mathbb{Q}}) \Rightarrow H^{s+t}(E, W\mathcal{O}_{E, \mathbb{Q}}).$$

The spectral sequence is compatible with the Frobenius and Verschiebung operation.

Proof. By [BBE07, Proposition 2.1(i)] the sheaf $W\mathcal{O}_{\cap_{j=0}^s E_{\iota_j}, \mathbb{Q}}$ doesn't depend on the choice of the scheme structure for $\cap_{j=0}^s E_{\iota_j}$.

From [BBE07, Corollary 2.3] we get an exact sequence

$$W\mathcal{O}_{E, \mathbb{Q}} \rightarrow \bigoplus_{\iota_0} W\mathcal{O}_{E_{\iota_0}, \mathbb{Q}} \rightarrow \bigoplus_{\iota_0 < \iota_1} W\mathcal{O}_{E_{\iota_0} \cap E_{\iota_1}, \mathbb{Q}} \rightarrow \dots,$$

where the maps are sums of restriction maps and thus compatible with F and V .

For every affine open set $U \subset E$ we have

$$H^t(U \cap_{j=0}^s E_{i_j}, W\mathcal{O}) = 0, \quad \text{for all } t > 0,$$

because $U \cap_{j=0}^s E_{i_j}$ is affine. Therefore we obtain the spectral sequence in the statement. \square

PROPOSITION 4.6.6. *Let E be a projective scheme over a finite field $k = \mathbb{F}_{p^a}$. Write $E = \cup_{i=1}^r E_i$ with E_i irreducible for all i . Suppose that for all $s \geq 1$ and all $1 \leq \iota_0 < \dots < \iota_s \leq r$ the set theoretic intersection $\cap_{j=0}^s E_{i_j}$, equipped with the reduced scheme structure, is smooth.*

For all $n \geq 2$ the differential of $E_n^{s,t}$, induced by the spectral sequence of Lemma 4.6.5, vanishes. In other words, $E_2^{s,t} \Rightarrow H^{s+t}(E, W\mathcal{O}_{E,\mathbb{Q}})$ is degenerated and

$$E_\infty^{s,t} = E_2^{s,t} \quad \text{for all } s, t.$$

Proof. By definition $E_n^{s,t}$ is a subquotient of $E_1^{s,t}$. By assumption, $\cap_{j=0}^s E_{i_j}$ is smooth and projective. We know that

$$H^t(\cap_{j=0}^s E_{i_j}, W\mathcal{O}_{\cap_{j=0}^s E_{i_j}, \mathbb{Q}}) \cong H_{\text{crys}}^t(\cap_{j=0}^s E_{i_j}/K)_{[0,1]},$$

where the right hand side is the slope < 1 -part of crystalline cohomology ($K = W(k)$). This isomorphism is compatible with the F -operation. It follows from [KM74] that the characteristic polynomial of F^a acting on $H_{\text{crys}}^t(\cap_{j=0}^s E_{i_j}/K)$ is the characteristic polynomial of the geometric Frobenius acting on $H_{\text{ét}}^t((\cap_{j=0}^s E_{i_j}) \times_k \bar{k}, \mathbb{Q}_\ell)$ for any prime $\ell \neq p$. Thus F^a acting on $H_{\text{crys}}^t(\cap_{j=0}^s E_{i_j}/K)_{[0,1]}$ has algebraic eigenvalues with absolute value $p^{\frac{ta}{2}}$ with respect to any embedding of K into \mathbb{C} . In particular, this holds for the eigenvalues of $E_n^{s,t}$ for $n \geq 1$ and all s . Thus the differential

$$E_n^{s-n, t+n-1} \rightarrow E_n^{s,t} \rightarrow E_n^{s+n, t-n+1}$$

vanishes if $n \geq 2$. \square

THEOREM 4.6.7. *Let k be a finite field. Let S be a normal integral scheme with an isolated singularity at the closed point $s \in S$. Let $f : X \rightarrow S$ be projective and birational. Suppose that f is an isomorphism over $S \setminus \{s\}$; we denote by E the closed set $f^{-1}(s)$. Write $E = \cup_{i=1}^r E_i$ with E_i irreducible for all i . Suppose that for all $s \geq 1$ and all $1 \leq \iota_0 < \dots < \iota_s \leq r$ the set theoretic intersection $\cap_{j=0}^s E_{i_j}$, equipped with the reduced scheme structure, is smooth.*

Then S has $W\mathcal{O}$ -rational singularities (Definition 4.4.4) if and only if the spectral sequence of Lemma 4.6.5 satisfies

$$E_2^{s,t} = 0 \quad \text{for all } (s, t) \neq (0, 0).$$

Proof. It follows from Corollary 4.6.4 that S has $W\mathcal{O}$ -rational singularities if and only if

$$H^i(E, W\mathcal{O}_{E,\mathbb{Q}}) = 0 \quad \text{for all } i > 0.$$

The assertion follows from Proposition 4.6.6. \square

The case $t = 0$ is a combinatorial condition on the exceptional set E . For any smooth and proper scheme X over k we have

$$H^0(X, W\mathcal{O}_X) \otimes_{W(k)} W(\bar{k}) \cong H^0_{\text{ét}}(X \times_k \bar{k}, \mathbb{Q}_p) \otimes_{\mathbb{Q}_p} W(\bar{k}).$$

Therefore the condition $E_2^{s,0} = 0$, for all $s \geq 1$, is equivalent to the vanishing of the cohomology of the complex

$$\begin{aligned} \bigoplus_{\iota_0} H^0(E_{\iota_0} \times_k \bar{k}, \mathbb{Q}_p) &\rightarrow \bigoplus_{\iota_0 < \iota_1} H^0((E_{\iota_0} \cap E_{\iota_1}) \times_k \bar{k}, \mathbb{Q}_p) \rightarrow \\ &\bigoplus_{\iota_0 < \iota_1 < \iota_2} H^0((E_{\iota_0} \cap E_{\iota_1} \cap E_{\iota_2}) \times_k \bar{k}, \mathbb{Q}_p) \rightarrow \dots \end{aligned}$$

in degree ≥ 1 . Of course, the cohomology in degree $= 0$ equals $H^0(E \times_k \bar{k}, \mathbb{Q}_p) = \mathbb{Q}_p$, because S is normal.

For a surface S the conditions for $t \geq 1$ are equivalent to $E_{i,\text{red}} \times_k \bar{k} \cong \coprod_j \mathbb{P}^1$ for all i . Indeed, for a smooth curve E_i we have

$$\dim H^1(E_i, W\mathcal{O}_{E_i, \mathbb{Q}}) \geq \dim H^1(E_i, \mathcal{O}_{E_i}),$$

and $H^1(E_i, \mathcal{O}_{E_i}) = 0$ if and only if $E_i \times_k \bar{k}$ is a disjoint union of \mathbb{P}^1 s.

Therefore S has $W\mathcal{O}$ -rational singularities if and only if the exceptional divisor of a minimal resolution over \bar{k} is a tree of \mathbb{P}^1 s.

4.7. CONES AND WITT-RATIONAL SINGULARITIES. In [BE08, Ex. 2.3] Blickle and Esnault give an example of an \mathbb{F}_p -scheme which has BE-rational singularities but not rational singularities. Their proof in fact shows, that their example also has $W\mathcal{O}$ -rational singularities. In this section we slightly generalize their example and show that it also has Witt-rational singularities.

4.7.1. Let X be a proper k -scheme, \mathcal{L} an invertible sheaf on X and $V(\mathcal{L}) = \text{Spec}(\bigoplus_{n \geq 0} \mathcal{L}^{\otimes n})$. We denote by $s_0 : X \hookrightarrow V(\mathcal{L})$ the zero-section. By Grauert's criterion (see [EGAII, Cor. (8.9.2)]) \mathcal{L} is ample on X if and only if there exists a k -scheme C together with a k -rational point $v \in C$ and a proper morphism $q : V(\mathcal{L}) \rightarrow C$, such that q induces an isomorphism $V(\mathcal{L}) \setminus s_0(X) \xrightarrow{\cong} C \setminus \{v\}$ and $q^{-1}(v)_{\text{red}} = s_0(X)_{\text{red}}$. If \mathcal{L} is ample, we call any triple (C, q, v) as above a *contractor of the zero-section of $V(\mathcal{L})$* . One can choose C for example to be the cone $\text{Spec } S$, where S is the graded ring $k \oplus_{n \geq 1} H^0(X, \mathcal{L}^{\otimes n})$ and v is the vertex, i.e. the point corresponding to the ideal S_+ .

The following proposition is well-known; we prove it for lack of reference.

PROPOSITION 4.7.2. *Let X be a proper and smooth k -scheme and \mathcal{L} an ample sheaf on X . Then the following statements are equivalent:*

- (1) *For any $n \geq 1$ and all contractors (C, q, v) of the zero-section of $V(\mathcal{L}^{\otimes n})$ we have $R^i q_* \omega_{V(\mathcal{L}^{\otimes n})} = 0$ for all $i \geq 1$.*
- (2) *$H^i(X, \omega_X \otimes \mathcal{L}^{\otimes n}) = 0$ for all $i, n \geq 1$.*

Proof. Let (C, q, v) be a contractor of the zero-section of $V(\mathcal{L})$. Denote by \mathcal{I} the ideal sheaf of the zero section of X in $V(\mathcal{L})$. We have

$$s_0^*\mathcal{I} = \mathcal{L}, \quad s_{0*}\mathcal{L}^{\otimes n} = \mathcal{I}^n/\mathcal{I}^{n+1}.$$

We set $Y_n := \text{Spec}(\mathcal{O}_{V(\mathcal{L})}/\mathcal{I}^n)$; in particular $Y_1 = s_0(X)$. The sheaves $R^i q_* \omega_{V(\mathcal{L})}$ have support in $\{v\}$ and since $\sqrt{\mathfrak{m}_v \mathcal{O}_{V(\mathcal{L})}} = \mathcal{I}$, where $\mathfrak{m}_v \subset \mathcal{O}_{X,v}$ is the ideal of v , the theorem on formal functions yields

$$R^i q_* \omega_{V(\mathcal{L})} = 0 \iff \varprojlim_n H^i(Y_n, \omega_{V(\mathcal{L})|Y_n}) = 0.$$

Further $\omega_{V(\mathcal{L})|Y_1} \cong \omega_X \otimes \mathcal{L}$, hence tensoring the exact sequence $0 \rightarrow \mathcal{I}^n/\mathcal{I}^{n+1} \rightarrow \mathcal{O}_{Y_{n+1}} \rightarrow \mathcal{O}_{Y_n} \rightarrow 0$ with $\omega_{V(\mathcal{L})|Y_{n+1}}$ yields the exact sequence

$$(4.7.1) \quad 0 \rightarrow \omega_X \otimes \mathcal{L}^{\otimes n+1} \rightarrow \omega_{V(\mathcal{L})|Y_{n+1}} \rightarrow \omega_{V(\mathcal{L})|Y_n} \rightarrow 0.$$

(2) \Rightarrow (1): We have $H^i(Y_1, \omega_{V(\mathcal{L})|Y_1}) = H^i(X, \omega_X \otimes \mathcal{L}) = 0$ for all $i \geq 1$, by assumption. Now the statement for (C, q, v) follows from (4.7.1) and induction. Since \mathcal{L} is any ample sheaf, we can replace \mathcal{L} by $\mathcal{L}^{\otimes n}$ in the above argument and obtain the statement also for a contractor of the zero-section of $V(\mathcal{L}^{\otimes n})$.

(1) \Rightarrow (2): Let d be the dimension of X . Then the maps $H^d(Y_{n+1}, \omega_{V(\mathcal{L})|Y_{n+1}}) \rightarrow H^d(Y_n, \omega_{V(\mathcal{L})|Y_n})$ are surjective for all $n \geq 1$. Since $\varprojlim_n H^d(Y_n, \omega_{V(\mathcal{L})|Y_n}) = 0$ by assumption, we get in particular

$$H^d(Y_1, \omega_{V(\mathcal{L})|Y_1}) = H^d(X, \omega_X \otimes \mathcal{L}) = 0.$$

Replacing \mathcal{L} by $\mathcal{L}^{\otimes n}$ in the above argument thus gives us

$$H^d(X, \omega_X \otimes \mathcal{L}^{\otimes n}) = 0, \quad \text{for all } n \geq 1.$$

Assume we proved $H^{i+1}(X, \omega_X \otimes \mathcal{L}^{\otimes n}) = 0$, for all $n \geq 1$. Then (4.7.1) yields that $H^i(Y_{n+1}, \omega_{V(\mathcal{L})|Y_{n+1}}) \rightarrow H^i(Y_n, \omega_{V(\mathcal{L})|Y_n})$ is surjective and as above we conclude that in particular $H^i(Y_1, \omega_{V(\mathcal{L})|Y_1}) = H^i(X, \omega_X \otimes \mathcal{L})$ equals zero (if $i \geq 1$). Again we can do the argument with \mathcal{L} replaced by $\mathcal{L}^{\otimes n}$ and obtain $H^i(X, \omega_X \otimes \mathcal{L}^{\otimes n}) = 0$ for all $n \geq 1$. This finishes the proof. \square

Remark 4.7.3. Notice that in characteristic zero condition (1) is always satisfied because of the Grauert-Riemenschneider vanishing theorem; condition (2) is always satisfied because of the Kodaira vanishing theorem.

THEOREM 4.7.4. *Let X_0 be a smooth, projective and geometrically connected k -scheme and \mathcal{L} an ample sheaf on X_0 . Assume (X_0, \mathcal{L}) satisfies the following condition:*

$$(4.7.2) \quad H^i(X_0, \omega_{X_0} \otimes \mathcal{L}^{\otimes n}) = 0, \quad \text{for all } n, i \geq 1.$$

Let X be the projective cone of (X_0, \mathcal{L}) , i.e. $X = \text{Proj}(S[z])$, where S denotes the graded k -algebra $k \oplus_{n \geq 1} H^0(X_0, \mathcal{L}^{\otimes n})$.

Then X is integral and normal; it has Witt-rational singularities if and only if

$$(4.7.3) \quad H^i(X_0, W\mathcal{O}_{X_0})_{\mathbb{Q}} = 0, \quad \text{for all } i \geq 1.$$

Furthermore if \mathcal{L} is very ample, then X is CM if and only if the following condition is satisfied:

$$(4.7.4) \quad H^i(X_0, \mathcal{L}^{\otimes n}) = 0, \quad \text{for all } n \in \mathbb{Z} \text{ and all } 1 \leq i \leq \dim X_0 - 1.$$

Proof. Let v be the k -rational point of X , corresponding to the homogenous ideal in $S[z]$ generated by S_+ . Then the pointed projective cone $X \setminus \{v\}$ is an \mathbb{A}^1 -bundle over $X_0 = \text{Proj } S$. Thus X has an isolated singularity at v and it suffices to consider the affine cone $C = \text{Spec } S$ of (X_0, \mathcal{L}) . Since X_0 is integral, projective and geometrically connected and k is perfect we have $H^0(X_0, \mathcal{O}_{X_0}) = k$, by Zariski's connectedness theorem. Thus C is integral and normal by [EGAII, Prop. (8.8.6), (ii)]. Set $V := V(\mathcal{L}) = \text{Spec}(\oplus_{n \geq 0} \mathcal{L}^{\otimes n})$; it is an \mathbb{A}^1 -bundle over X_0 , hence is smooth. By [EGAII, Cor. (8.8.4)] there exists a projective morphism $q : V \rightarrow C$, such that the triple (C, q, v) becomes a contractor of the zero-section of V . In particular, $q : V \rightarrow C$ is a projective resolution of singularities of C . (In case \mathcal{L} is very ample, $q : V \rightarrow C$ is the blow-up of C in the closed point v , see [EGAII, Rem. (8.8.3)].) By Proposition 4.4.6 and Remark 4.4.5 it suffices to prove

$$R^i q_* W \mathcal{O}_{V, \mathbb{Q}} = 0, \quad R^i q_* W \omega_{V, \mathbb{Q}} = 0, \quad \text{for all } i \geq 1.$$

By Corollary 4.6.4 the vanishing $R^i q_* W \mathcal{O}_{V, \mathbb{Q}} = 0$ is equivalent to the vanishing $H^i(X_0, W \mathcal{O}_{X_0})_{\mathbb{Q}} = 0$. Thus it suffices to prove the vanishing $R^i q_* W \omega_{V, \mathbb{Q}} = 0$ under the assumptions (4.7.2) and $Rq_* W \mathcal{O}_{V, \mathbb{Q}} \cong W \mathcal{O}_{C, \mathbb{Q}}$. Proposition 4.7.2 yields $R^i q_* \omega_V = 0$ for all $i \geq 1$. Hence also $R^i q_* W_n \omega_V = 0$ for all $i, n \geq 1$, by the exact sequence in Proposition 4.1.4, (7) and induction. Therefore the exact sequence

$$0 \rightarrow R^1 \varprojlim_n R^{i-1} q_* W_n \omega_V \rightarrow R^i q_* W \omega_V \rightarrow \varprojlim_n R^i q_* W_n \omega_V \rightarrow 0$$

immediately yields

$$(4.7.5) \quad R^i q_* W \omega_{V, \mathbb{Q}} = 0, \quad \text{for all } i \geq 2.$$

To conclude also the vanishing of $R^1 q_* W \omega_{V, \mathbb{Q}}$ we have to prove the vanishing $(R^1 \varprojlim_n q_* W_n \omega_V)_{\mathbb{Q}} = 0$. To this end denote by I_{\bullet} , K_{\bullet} and C_{\bullet} the image, the kernel and the cokernel of $q_* W_{\bullet} \omega_V \rightarrow W_{\bullet} \omega_C$, respectively. Notice that K_n and C_n are coherent $W_n \mathcal{O}_C$ -modules, whose support is concentrated in the closed point v ; hence these modules have finite length. Therefore K_{\bullet} and C_{\bullet} satisfy the Mittag-Leffler condition, in particular $R^1 \varprojlim C_n = 0 = R^1 \varprojlim K_n$. Furthermore, $R^2 \varprojlim K_n = 0$ by Lemma 1.5.1, (1). Thus the exact sequence $0 \rightarrow K_{\bullet} \rightarrow q_* W_{\bullet} \omega_V \rightarrow I_{\bullet} \rightarrow 0$ gives

$$R^1 \varprojlim q_* W_n \omega_V \cong R^1 \varprojlim I_n.$$

We also have the vanishing $R^1 \varprojlim W_n \omega_C = 0$, since the transition maps are surjective (by Proposition 4.1.4, (4)) and therefore the exact sequence $0 \rightarrow I_{\bullet} \rightarrow W_{\bullet} \omega_C \rightarrow C_{\bullet} \rightarrow 0$ gives a surjection

$$\varprojlim C_n \rightarrow R^1 \varprojlim I_n \rightarrow 0.$$

By Lemma 4.3.4 $(\varprojlim C_n)_{\mathbb{Q}} = 0$, hence also $(R^1 \varprojlim q_* W_n \omega_V)_{\mathbb{Q}} = 0$.

For the last statement notice that X is CM if and only if $H_v^i(C, \mathcal{O}_C) = 0$ for all $i \leq d - 1 = \dim X_0$. Since C is affine the long exact localization sequence gives us an exact sequence

$$0 \rightarrow H_v^0(C, \mathcal{O}_C) \rightarrow H^0(C, \mathcal{O}_C) \rightarrow H^0(C \setminus \{v, \}, \mathcal{O}_C) \rightarrow H_v^1(C, \mathcal{O}_C) \rightarrow 0$$

and isomorphisms

$$H^{i-1}(C \setminus \{v, \}, \mathcal{O}_C) \cong H_v^i(C, \mathcal{O}_C), \quad \text{for all } i \geq 2.$$

Let $\pi : C \setminus \{v\} \rightarrow X_0 = \text{Proj } S$ be the pointed affine cone over $X_0 = \text{Proj } S$. Since \mathcal{L} is very ample the graded ring S is generated by S_1 and we obtain

$$\pi_* \mathcal{O}_{C \setminus \{v\}} = \bigoplus_{n \in \mathbb{Z}} \mathcal{L}^{\otimes n}.$$

Therefore $H^0(C \setminus \{v, \}, \mathcal{O}_C) = \bigoplus_{n \geq 0} H^0(X_0, \mathcal{L}^{\otimes n}) = S = H^0(C, \mathcal{O}_C)$. Thus $H_v^0(C, \mathcal{O}_C)$ and $H_v^1(C, \mathcal{O}_C)$ always vanish and $H_v^i(C, \mathcal{O}_C)$ vanishes for $2 \leq i \leq \dim X_0$ if and only if $\bigoplus_{n \in \mathbb{Z}} H^{i-1}(X_0, \mathcal{L}^{\otimes n})$ vanishes, which is exactly condition (4.7.4). This proves the Theorem. \square

Remark 4.7.5. (1) Condition (4.7.2) in the above theorem holds in characteristic zero by Kodaira vanishing, which in general is wrong in positive characteristic; for a counter example see [Ray78]. By [DI87, Cor. 2.8.] this condition holds if X_0 lifts to a smooth $W_2(k)$ -scheme and has dimension $\leq p$. Notice also that condition (4.7.2) and (4.7.4) are always satisfied if X_0 is a smooth hypersurface in some \mathbb{P}_k^n and $\mathcal{L} = \mathcal{O}_{\mathbb{P}_k^n}(1)|_{X_0}$.

(2) The vanishing (4.7.3) is satisfied if the degree map induces an isomorphism $\text{CH}_0(X_0 \times_k \bar{k}(X_0))_{\mathbb{Q}} \cong \mathbb{Q}$, which is for example the case if $X_{0, \bar{k}}$ is rationally chain connected, where \bar{k} is an algebraic closure of k . See [BE08, Ex. 2.3], alternatively one can also use Bloch's decomposition of the diagonal and the vanishing Lemmas 3.6.1 and 3.6.2. Another case where (4.7.3) holds is when X_0 has a smooth projective model \mathcal{X}_0 over a complete discrete valuation ring R of mixed characteristic and with residue field k , such that the generic fiber $\mathcal{X}_{0, \eta}$ satisfies $H^i(\mathcal{X}_{0, \eta}, \mathcal{O}_{\mathcal{X}_{0, \eta}}) = 0$ for all $i \geq 1$. This follows from p -adic Hodge theory ("the Newton polygon of $H_{\text{crys}}^i(X_0/W) \otimes \text{Frac}(R) \cong H_{dR}^i(\mathcal{X}_{0, \eta})$ lies above the Hodge polygon").

(3) The example in [BE08] alluded to at the beginning of this section is the following: By [Shi74, Prop. 3] the Fermat hypersurfaces $X_0 \subset \mathbb{P}_k^{2r+1}$ given by $x_0^n + \dots + x_{2r+1}^n$, where n is such that $p^\nu \equiv -1 \pmod{n}$, for some $\nu \geq 1$, are unirational over \bar{k} . Hence X_0 satisfies the conditions (4.7.2), (4.7.3) and (4.7.4) of the theorem and we obtain that the projective cone X of X_0 is normal, CM and has Witt-rational singularities. (In [BE08, Ex. 2.3.] it was shown that X is $W\mathcal{O}$ -rational.) Choosing X_0 of degree larger than $2r + 2$ we see, that $H^{\dim X_0}(X_0, \mathcal{O}_{X_0}) \neq 0$ and it follows that X cannot have rational singularities.

- (4) Let X be as in the theorem and let $\pi : Y \rightarrow X$ be any resolution of singularities. Then $R^i\pi_*W\omega_{Y,\mathbb{Q}} = 0$ for all $i \geq 1$. As it follows from the proof and [CR09, Thm. 1], we even have the stronger vanishing $R^i\pi_*\omega_Y = 0$ for all $i \geq 1$, which is implied by condition (4.7.2) solely. It is an obvious question whether the vanishing $R^i\pi_*W\omega_{Y,\mathbb{Q}} = 0$ also holds without condition (4.7.2), i.e. does some version of Grauert-Riemenschneider vanishing hold. In view of Proposition 4.7.2, this question should be linked to some kind of Kodaira vanishing, of which at the moment even the formulation is not clear. (But see [BBE07, Cor. 1.2] for a first result in this direction.)

4.8. MORPHISMS WITH RATIONALLY CONNECTED GENERIC FIBRE. The goal of this section is to prove the following theorem.

THEOREM 4.8.1. *Let X be an integral scheme with Witt-rational singularities. Let $f : X \rightarrow Y$ be a projective morphism to an integral and normal scheme Y . We denote by η the generic point of Y , and X_η denotes the generic fibre of f . Suppose that X_η is smooth and for every field extension $L \supset k(\eta)$ the degree map*

$$\mathrm{CH}_0(X_\eta \times_{k(\eta)} L) \otimes_{\mathbb{Z}} \mathbb{Q} \rightarrow \mathbb{Q}$$

is an isomorphism. Then

$$\bigoplus_{i>0} R^i f_* W\mathcal{O}_{X,\mathbb{Q}} \cong \bigoplus_{i>0} H^i(\mathrm{WS}_{0,Y}), \quad \bigoplus_{i>0} R^i f_* W\omega_{X,\mathbb{Q}} \cong \bigoplus_{i>0} H^i(\mathrm{WS}_{\dim Y,Y}),$$

where WS is defined in Definition 4.5.2.

4.8.2. Let $f : X \rightarrow Y$ be as in the assumptions of Theorem 4.8.1. We can choose a factorization (cf. Remark 4.3.2)

$$X' \xrightarrow{\pi} X'/G \xrightarrow{h} X'' \xrightarrow{u} X,$$

such that

- X' is smooth, integral and quasi-projective, G is a finite group acting on X' ,
- h is birational and projective,
- X'' is normal and u is a universal homeomorphism.

We obtain a commutative diagram

$$(4.8.1) \quad \begin{array}{ccc} X' & & \\ \downarrow \pi & & \\ X'/G & & \\ \downarrow h & & \\ X'' & \longrightarrow & Y'' \\ \downarrow u & & \downarrow u' \\ X & \xrightarrow{f} & Y \end{array}$$

where Y'' is the Stein factorization of $X'' \rightarrow Y$. Therefore u' is a universal homeomorphism. Note that Y'' is also the Stein factorization of $X'/G \rightarrow Y$.

LEMMA 4.8.3. *With the assumptions of Theorem 4.8.1. Set $f' = f \circ u \circ h \circ \pi$ with the notation as in diagram (4.8.1). We consider X' via f' as a scheme over Y . Let $P = \sum_{g \in G} [\Gamma(g)] \in \text{CH}(X' \times_Y X')$, with $\Gamma(g)$ the graph of g . Then there is a natural isomorphism*

$$\hat{\mathcal{H}}(P/Y) \left(\bigoplus_{i>0} R^i f'_*(W\mathcal{O}_{X',\mathbb{Q}} \oplus W\omega_{X',\mathbb{Q}}) \right) \rightarrow \bigoplus_{i>0} R^i f_*(W\mathcal{O}_{X,\mathbb{Q}} \oplus W\omega_{X,\mathbb{Q}})$$

Proof. The cycle $P = \sum_{g \in G} [\Gamma(g)]$ is already defined in $\text{CH}(X' \times_{X'/G} X')$. Theorem 4.5.6 and Corollary 4.4.7 imply

$$(4.8.2) \quad \hat{\mathcal{H}}(P/(X'/G)) (\pi_* W\mathcal{O}_{X',\mathbb{Q}} \oplus \pi_* W\omega_{X',\mathbb{Q}}) \cong W\mathcal{O}_{X'/G,\mathbb{Q}} \oplus W\omega_{X'/G,\mathbb{Q}}.$$

It follows from Proposition 3.5.10 that

$$\hat{\mathcal{H}}(P/Y) = \bigoplus_i R^i (f \circ u \circ h)_* \hat{\mathcal{H}}(P/(X'/G)).$$

Since $P^2 = \#G \cdot P$, we obtain from (4.8.2) that

$$\hat{\mathcal{H}}(P/Y) \left(\bigoplus_{i>0} R^i f'_*(W\mathcal{O}_{X',\mathbb{Q}} \oplus W\omega_{X',\mathbb{Q}}) \right) = \bigoplus_{i>0} R^i (f \circ u \circ h)_*(W\mathcal{O}_{X'/G,\mathbb{Q}} \oplus W\omega_{X'/G,\mathbb{Q}}).$$

Because X has Witt-rational singularities and $X'/G \xrightarrow{u \circ h} X$ is a quasi-resolution we get

$$R^j (u \circ h)_*(W\mathcal{O}_{X'/G,\mathbb{Q}} \oplus W\omega_{X'/G,\mathbb{Q}}) = \begin{cases} 0 & \text{for all } j > 0 \\ W\mathcal{O}_{X,\mathbb{Q}} \oplus W\omega_{X,\mathbb{Q}} & \text{for } j = 0, \end{cases}$$

which implies the statement. \square

PROPOSITION 4.8.4. *Let $f' : X' \rightarrow Y$ be a projective and surjective morphism. Suppose that Y is normal, and X' is smooth and connected. We denote by η the generic point of Y . Let $Y' \subset X'$ be a closed irreducible subset such that $Y' \rightarrow Y$ is generically finite. We denote by A_1 and A_2 the closure of $Y'_\eta \times_\eta X'_\eta$ and $X'_\eta \times_\eta Y'_\eta$ in $X' \times_Y X'$, respectively. Then there are natural isomorphisms*

$$(4.8.3) \quad \hat{\mathcal{H}}([A_1]/Y) \left(\bigoplus_{i>0} R^i f'_* W\mathcal{O}_{X',\mathbb{Q}} \right) \cong \mathcal{A}^0(Y)$$

$$(4.8.4) \quad \hat{\mathcal{H}}([A_2]/Y) \left(\bigoplus_{i>0} R^i f'_* W\omega_{X',\mathbb{Q}} \right) \cong \mathcal{A}^{\dim Y}(Y).$$

(See Definition 4.5.4 for \mathcal{A} .)

Proof. We will prove (4.8.4), the identity (4.8.3) can be proved in the same way.

Let $\tilde{Y} \rightarrow Y'$ be an alteration of Y' , such that \tilde{Y} is smooth. We denote by $\iota : \tilde{Y} \rightarrow X'$ the composition. The graph of ι defines a morphism

$$\hat{\mathcal{H}}(\Gamma(\iota)/Y) : \hat{\mathcal{H}}(\tilde{Y}/Y)(-r) \rightarrow \hat{\mathcal{H}}(X'/Y),$$

with $r := \dim X' - \dim Y$. The closure Z of $X'_\eta \times_\eta \tilde{Y}_\eta$ in $X' \times_Y \tilde{Y}$ defines a morphism

$$\hat{\mathcal{H}}(Z/Y) : \hat{\mathcal{H}}(X'/Y) \rightarrow \hat{\mathcal{H}}(\tilde{Y}/Y)(-r).$$

We have $(\text{id}_{X'} \times \iota)_*([Z]) = d \cdot [A_2]$ for some $d \neq 0$. Finally, we define Q to be the closure of $\tilde{Y}_\eta \times_\eta \tilde{Y}_\eta$ in $\tilde{Y} \times_Y \tilde{Y}$. We have the following relations

$$\begin{aligned} [\Gamma(\iota)] \circ [Z] - d[A_2] &\in \ker(\text{CH}(X' \times_Y X') \rightarrow \text{CH}(X'_\eta \times_\eta X'_\eta)), \\ [Z] \circ [\Gamma(\iota)] - [Q] &\in \ker(\text{CH}(\tilde{Y} \times_Y \tilde{Y}) \rightarrow \text{CH}(\tilde{Y}_\eta \times_\eta \tilde{Y}_\eta)), \\ [A_2] \circ [A_2] - e[A_2] &\in \ker(\text{CH}(X' \times_Y X') \rightarrow \text{CH}(X'_\eta \times_\eta X'_\eta)), \\ [Q] \circ [Q] - g[Q] &\in \ker(\text{CH}(\tilde{Y} \times_Y \tilde{Y}) \rightarrow \text{CH}(\tilde{Y}_\eta \times_\eta \tilde{Y}_\eta)), \end{aligned}$$

for some $e, g \neq 0$. Because of Lemma 3.6.2 we conclude that

$$\hat{\mathcal{H}}([A_2]/Y) \left(\bigoplus_{i>0} R^i f_* W\omega_{X',\mathbb{Q}} \right) \cong \hat{\mathcal{H}}([Q]/Y) \left(\bigoplus_{i>0} R^i (f' \circ \iota)_* W\omega_{\tilde{Y},\mathbb{Q}} \right).$$

By Definition 4.5.4 the right hand side is $\mathcal{A}^{\dim Y}(Y)$. □

Proof of Theorem 4.8.1. We use diagram 4.8.1 and the constructions of 4.8.2. The first step is to prove

$$(4.8.5) \quad \text{CH}_0(X'_\eta \times_{k(\eta)} L)^G \otimes \mathbb{Q} \cong \mathbb{Q}$$

for every field extension $L \supset k(\eta)$. We have a push-forward map

$$\alpha : \text{CH}_0(X'_\eta \times_{k(\eta)} L) \otimes \mathbb{Q} \rightarrow \text{CH}_0(X_\eta \times_{k(\eta)} L) \otimes \mathbb{Q},$$

and because X_η is smooth over $k(\eta)$ we have a pull-back map

$$\beta : \text{CH}_0(X_\eta \times_{k(\eta)} L) \otimes \mathbb{Q} \rightarrow \text{CH}_0(X'_\eta \times_{k(\eta)} L) \otimes \mathbb{Q}.$$

We have the following formula for the composition

$$\beta \circ \alpha = \text{deg}(u) \sum_{g \in G} g_*,$$

provided that G acts faithfully on X' (we may assume this). Thus (4.8.5) follows from the assumptions.

Let

$$P = \sum_{g \in G} [\Gamma(g)] \in \text{CH}(X' \times_Y X'),$$

with $\Gamma(g)$ the graph of g . We denote by P_η the image of P in $\text{CH}(X'_\eta \times_\eta X'_\eta)$. The cycle P_η is invariant under the action of G on the left and the right factor of $X'_\eta \times_\eta X'_\eta$.

By the same arguments as in [BS83, Proposition 1] we see that there are non-zero integers N_1, N_2, M_1, M_2 , effective divisors D_1, D_2 on X'_η , and a closed point α of X'_η , such that

$$(4.8.6) \quad N_1 P_\eta + N_2 [\alpha \times_\eta X'_\eta] \in \text{image}(\text{CH}(X'_\eta \times_\eta D_2)),$$

$$(4.8.7) \quad M_1 P_\eta + M_2 [X'_\eta \times_\eta \alpha] \in \text{image}(\text{CH}(D_1 \times_\eta X'_\eta)).$$

Let \bar{D}_1, \bar{D}_2 , and Y_α be the closures of D_1, D_2 , and α in X' . The map $Y_\alpha \rightarrow Y$ is generically finite and $Y_{\alpha, \eta} = \alpha$. Let A_1 and A_2 be the closures of $\alpha \times_\eta X'_\eta$ and $X'_\eta \times_\eta \alpha$ in $X' \times_Y X'$. Because of the formulas (4.8.6), (4.8.7) over the generic point η , there is a Weil divisor $S \subset Y$ such that

$$N_1 P + N_2 [A_1] \in \text{image}(\text{CH}(X' \times_Y \bar{D}_2)) + \text{image}(\text{CH}(X'_S \times_S X'_S)),$$

$$M_1 P + M_2 [A_2] \in \text{image}(\text{CH}(\bar{D}_1 \times_Y X')) + \text{image}(\text{CH}(X'_S \times_S X'_S)).$$

Lemma 3.6.1 implies that $\hat{\mathcal{H}}(P/Y)$ acts as $\hat{\mathcal{H}}([A_1]/Y)$ on

$$\bigoplus_{i>0} R^i f'_* W \mathcal{O}_{X', \mathbb{Q}}.$$

From Lemma 3.6.2 we conclude that $\hat{\mathcal{H}}(P/Y)$ acts as $\hat{\mathcal{H}}([A_2]/Y)$ on

$$\bigoplus_{i>0} R^i f'_* W \omega_{X', \mathbb{Q}}.$$

In view of Proposition 4.8.4, Theorem 4.5.6, and Lemma 4.8.3 this implies the statement of the theorem. \square

5. FURTHER APPLICATIONS

5.1. RESULTS ON TORSION.

5.1.1. *The Cartier-Dieudonné-Raynaud algebra.* Recall from [IR83, I, (1.1)] that the Cartier-Dieudonné-Raynaud algebra is the graded (non-commutative) W -algebra, generated by formal symbols F and V in degree 0 and by d in degree 1 which satisfy the following relations

$$F \cdot a = F(a) \cdot F, \quad a \cdot V = V \cdot F(a), \quad (a \in W), \quad F \cdot V = p = V \cdot F$$

$$a \cdot d = d \cdot a \quad (a \in W), \quad F \cdot d \cdot V = d, \quad d \cdot d = 0.$$

(Here F is a formal symbol, whereas $F(a)$ denotes the Frobenius on the Witt vectors of k applied to the Witt vector a .) Thus R is concentrated in degree 0 and 1. Notice that any de-Rham-Witt module and any Witt-module on a scheme X naturally becomes an R -module (the latter with d acting as 0). Therefore we have an exact functor

$$\widehat{\text{dRW}}_X \rightarrow \text{Sh}(X, R) := (\text{sheaves-of } R\text{-modules on } X)$$

which trivially derives to a functor

$$(5.1.1) \quad \phi : D^b(\widehat{\text{dRW}}_X) \rightarrow D^b(\text{Sh}(X, R)) := D^b(X, R).$$

We set

$$R_n := R/(V^n \cdot R + d \cdot V^n \cdot R).$$

Notice that this is a left R -module. We obtain a functor

$$R_n \otimes_R : \text{Sh}(X, R) \rightarrow \text{Sh}(X, W_n[d]) := (\text{sheaves of } W_n[d]\text{-modules on } X),$$

where $W_n[d]$ is the graded W_n -algebra $W_n \oplus W_n \cdot d$, with $d^2 = 0$. By [IR83, Prop. (3.2)] the following sequence is an exact sequence of right R -modules

$$(5.1.2) \quad 0 \rightarrow R(-1) \xrightarrow{(F^n, -F^n d)} R(-1) \oplus R \xrightarrow{dV^n + V^n} R \rightarrow R_n \rightarrow 0.$$

This allows us to calculate the left-derived functor

$$R_n \otimes_R^L - : D^-(\text{Sh}(X, R)) \rightarrow D^-(\text{Sh}(X, W_n[d])) := D^-(X, W_n[d]).$$

One obtains the following.

- (1) Assume X is smooth. Then $R_n \otimes_R^L W\Omega_X \cong W_n\Omega_X$.
- (2) Let $f : X \rightarrow Y$ be a morphism of k -schemes, then

$$R_n \otimes_R^L Rf_*(-) \cong Rf_*(R_n \otimes_R^L (-)) : D^b(X, R) \rightarrow D^b(Y, W_n[d]).$$

- (3) Let X be a scheme and $Z \subset X$ be a closed subscheme, then

$$R_n \otimes_R^L R\Gamma_Z(-) \cong R\Gamma_Z(R_n \otimes_R^L (-)) : D^b(X, R) \rightarrow D^b(X, W_n[d]).$$

The first statement is [IR83, II, Thm (1.2)], the last two statements follow directly from (5.1.2).

PROPOSITION 5.1.2 ([Eke85, I, Prop. 1.1.]). *Let S be a k -scheme and $M \in D^b(S, R)$ a complex of R -modules, which is bounded in both directions, i.e. there exists a natural number m , such that $H^i(M)^j$ is non-zero only for $(i, j) \in [-m, m] \times [-m, m]$.*

- (1) *Assume there exist integers $r, s \in \mathbb{Z}$, such that $H^i(R_1 \otimes_R^L M)^j = 0$ for all pairs (i, j) satisfying one of the following conditions*

$$(5.1.3) \quad (i+j = r, i \geq s) \text{ or } (i+j = r-1, i \geq s+1) \text{ or } (i+j = r+1, i \geq s+1).$$

Then

$$H^i(R_n \otimes_R^L M)^j = 0, \quad \text{for all } n \text{ and all } (i, j) \text{ with } i+j = r, i \geq s.$$

- (2) *Assume there exist integers $r, s \in \mathbb{Z}$, such that $H^i(R_1 \otimes_R^L M)^j = 0$ for all pairs (i, j) satisfying one of the following conditions*

$$(5.1.4) \quad (i+j = r, i \leq s) \text{ or } (i+j = r+1, i \leq s) \text{ or } (i+j = r-1, i \leq s-1).$$

Then

$$H^i(R_n \otimes_R^L M)^j = 0 \quad \text{for all } n \text{ and all } (i, j) \text{ with } i+j = r, i \leq s.$$

This proposition is called Ekedahl's Nakayama Lemma because applying it to the cone of a morphism $f : M \rightarrow N$ in $D^b(X, R)$ between two complexes, which are bounded in both directions, we obtain that $R_n \otimes_R f$ is an isomorphism for all n , if $R_1 \otimes_R^L f$ is.

COROLLARY 5.1.3. *Let S be a k -scheme and $f : X \rightarrow S, g : Y \rightarrow S$ two S -schemes which are smooth over k . Let $\varphi : Rf_*W\Omega_X \rightarrow Rg_*W\Omega_Y$ be a morphism in $D^b(S, R)$. Then $R_1 \otimes_R \varphi$ is a morphism*

$$\varphi_1 := R_1 \otimes_R^L \varphi : Rf_*\Omega_X \rightarrow Rg_*\Omega_Y \quad \text{in } D^b(S, k[d])$$

and furthermore:

- (1) *If there exists a natural number $a \geq 0$ such that $\varphi_1 : \bigoplus_{j \leq a} Rf_*\Omega_X^j \xrightarrow{\cong} \bigoplus_{j \leq a} Rg_*\Omega_Y^j$ is an isomorphism, then*

$$\hat{\varphi} := R\varprojlim_n (R_n \otimes_R \varphi) : \bigoplus_{j \leq a} Rf_*W\Omega_X^j \xrightarrow{\cong} \bigoplus_{j \leq a} Rg_*W\Omega_Y^j$$

is an isomorphism.

- (2) *If there exists a natural number $a \geq 0$ such that $\varphi_1 : \bigoplus_{j \geq a} Rf_*\Omega_X^j \xrightarrow{\cong} \bigoplus_{j \geq a} Rg_*\Omega_Y^j$ is an isomorphism, then*

$$\hat{\varphi} = R\varprojlim_n (R_n \otimes_R \varphi) : \bigoplus_{j \geq a} Rf_*W\Omega_X^j \xrightarrow{\cong} \bigoplus_{j \geq a} Rg_*W\Omega_Y^j$$

is an isomorphism.

Proof. First of all notice, that by 5.1.1, (1) and (2) $R_n \otimes_R^L \varphi$ indeed is a morphism $Rf_*W_n\Omega_X \rightarrow Rg_*W_n\Omega_Y$ and that $R\varprojlim_n (R_n \otimes_R^L \varphi)$ is a morphism $Rf_*W\Omega_X \rightarrow Rg_*W\Omega_Y$, since $R\varprojlim_n Rf_*W_n\Omega_X = Rf_*R\varprojlim_n W_n\Omega_X = Rf_*W\Omega_X$.

Denote by C the cone of φ in $D^b(S, R)$. It is clearly bounded in both directions (in the sense of Proposition 5.1.2). Assume we are in the situation of (1). Then $H^i(R_1 \otimes_R^L C)^j = 0$ for all $i \in \mathbb{Z}$ and $j \leq a$. Choose $i_0 \in \mathbb{Z}$ and $j_0 \leq a$ and set $s := i_0$ and $r := j_0 + i_0$. Then $H^i(R_1 \otimes_R^L C)^j = 0$ for all (i, j) as in (5.1.3). Thus $H^{i_0}(R_n \otimes_R^L C)^{j_0} = 0$ for all n . Therefore $R_n \otimes_R^L \varphi$ is an isomorphism for all n in degree $\leq a$, which gives (1). Now assume we are in the situation of (2). Then $H^i(R_1 \otimes_R^L C)^j = 0$ for all $i \in \mathbb{Z}$ and $j \geq a$. Choose $i_0 \in \mathbb{Z}$ and $j_0 \geq a$ and set $s := i_0$ and $r := i_0 + j_0$. Then $H^i(R_1 \otimes_R^L C)^j = 0$ for (i, j) as in (5.1.4). Thus $H^{i_0}(R_n \otimes_R^L C)^{j_0} = 0$, which implies (2) as above. \square

LEMMA 5.1.4. *Let X be a smooth scheme and $E(W_\bullet\Omega_X)$ the Cousin complex of $W_\bullet\Omega$ with respect to the codimension filtration on X (see 1.5.8). Then there is a natural commutative diagram of isomorphisms in $D^b(X, W_n[d])$*

$$\begin{array}{ccc} R_n \otimes_R^L W\Omega_X & \xrightarrow{\cong} & R_n \otimes_R^L \varprojlim E(W_\bullet\Omega_X) \\ \downarrow \cong & & \downarrow \cong \\ W_n\Omega_X & \xrightarrow{\cong} & E(W_n\Omega_X). \end{array}$$

Proof. There is an obvious map

$$R_n \otimes_R \varprojlim E(W_\bullet\Omega_X) = \varprojlim E(W_\bullet\Omega_X)/(V^n + dV(\varprojlim E(W_\bullet\Omega_X))) \rightarrow E(W_n\Omega),$$

yielding a morphism $R_n \otimes_R^L \varprojlim E(W_\bullet \Omega_X) \rightarrow E(W_n \Omega)$, which clearly fits into a diagram as in the statement. By Lemma 1.5.9, $E(W_\bullet \Omega_X)$ is a flasque resolution of $W_\bullet \Omega_X$ and thus, by Lemma 1.5.4, (4), $\varprojlim E(W_\bullet \Omega_X)$ is a resolution of $W \Omega_X$. Hence the two horizontal arrows in the diagram are isomorphisms, and by 5.1.1(1) the vertical arrow on the left is an isomorphism. This proves the claim. \square

5.1.5. Let $f : X \rightarrow Y$ be a morphism between k -schemes. Then we have the pullback in $D^b(\mathrm{dRW}_Y)$ (see Definition 2.2.1)

$$f^* : W_\bullet \Omega_Y \rightarrow Rf_* W_\bullet \Omega_X.$$

Using (5.1.2) and 5.1.1, (1), (2) one checks

$$f_n^* = R_n \otimes_R^L (\phi(R \varprojlim f^*)) \quad \text{in } D^b(Y, W_n[d]),$$

where ϕ is the forgetful functor from (5.1.1), f_n^* denotes the image of f^* under the projection from $D^b(\mathrm{dRW}_Y)$ to $D^b(Y, W_n[d])$ and $R \varprojlim : D^b(\mathrm{dRW}_Y) \rightarrow D^b(\widehat{\mathrm{dRW}}_Y)$ (see Proposition 1.5.6.)

5.1.6. Let $f : X \rightarrow Y$ be a morphism of pure relative dimension r between smooth schemes and let $Z \subset X$ be a closed subscheme, such that $f|_Z$ is proper. Then we have the pushforward in $D^b(\mathrm{dRW}_Y)$ (see (2.3.4))

$$f_* : Rf_* R\Gamma_Z W_\bullet \Omega_X \rightarrow W_\bullet \Omega_Y(-r)[-r].$$

Using (2.3.3) and Lemma 5.1.4 one checks

$$f_{*,n} = R_n \otimes_R^L (\phi(R \varprojlim f_*)) \quad \text{in } D^b(Y, W_n[d]).$$

5.1.7. Let X be smooth equidimensional scheme and $Z \subset X$ an integral closed subscheme of codimension c . Then we have the cup-product with $cl([Z])$ (see 3.2.2) in $D^b(\mathrm{dRW}_X)$

$$(-) \cup cl([Z]) : W_\bullet \Omega_X \xrightarrow{(3.5.6)} \mathcal{H}_Z^c(W_\bullet \Omega)(c) \xrightarrow{(3.5.4)} R\Gamma_Z(W_\bullet \Omega)(c)[c].$$

Notice that in 3.5.7 we defined this maps only in the limit, but they can clearly be constructed on each level and the limit of the above gives 3.5.7. Also notice that the first map is given by $\alpha \mapsto (-1)^{c \cdot \deg \alpha} \alpha \cdot cl([Z])$, which is a morphism of de Rham-Witt systems (since $F(cl([Z])) = \pi(cl([Z]))$ and $d(cl([Z])) = 0$.) Using (5.1.2) and 5.1.1, (1), (3) one checks

$$(- \cup cl_n([Z])) = R_n \otimes_R^L \phi(R \varprojlim ((-) \cup cl([Z]))) \quad \text{in } D^b(X, W_n[d]).$$

5.1.8. Let S be a k -scheme, $f : X \rightarrow S$ and $g : Y \rightarrow S$ be two S -schemes which are smooth over k and of the same pure dimension N . Let $Z \subset X \times_S Y$ be a closed integral subscheme of dimension N , which is proper over Y . Then we define the morphism

$$\mathcal{R}([Z]/S) : Rf_* W \Omega_X \rightarrow Rg_* W \Omega_Y \quad \text{in } D^b(S, R)$$

as the composition

$$\begin{aligned}
 Rf_*W\Omega_X &\xrightarrow{\text{pr}_1^*} Rf_*R\text{pr}_{1*}W\Omega_{X\times Y} \\
 &\xrightarrow{\cup \text{cl}([Z])} Rf_*R\text{pr}_{1*}R\Gamma_ZW\Omega_{X\times Y}(N)[N] \\
 &\xrightarrow{\simeq, Z\subset X\times_S Y} Rg_*R\text{pr}_{2*}R\Gamma_ZW\Omega_{X\times Y}(N)[N] \\
 &\xrightarrow{\text{pr}_{2*}} Rg_*W\Omega_Y.
 \end{aligned}$$

Here we simply write pr_1^* instead of $\phi(R\varprojlim(\text{pr}_1^*))$ etc. Notice that the third arrow does not exist in $D^b(\widehat{\text{dRW}}_S)$, since it is not compatible with the $W\mathcal{O}_S$ -action. By 5.1.1 and 5.1.5, 5.1.6, 5.1.7, $R_n \otimes_R^L \mathcal{R}([Z]/S)$ equals the following composition in $D^b(S, W_n[d])$, for all $n \geq 1$

$$\begin{aligned}
 (5.1.5) \quad Rf_*W_n\Omega_X &\xrightarrow{\text{pr}_{1,n}^*} Rf_*R\text{pr}_{1*}W_n\Omega_{X\times Y} \\
 &\xrightarrow{\cup \text{cl}_n([Z])} Rf_*R\text{pr}_{1*}R\Gamma_ZW_n\Omega_{X\times Y}(N)[N] \\
 &\xrightarrow{\simeq, Z\subset X\times_S Y} Rg_*R\text{pr}_{2*}R\Gamma_ZW_n\Omega_{X\times Y}(N)[N] \\
 &\xrightarrow{\text{pr}_{2,n*}} Rg_*W_n\Omega_Y.
 \end{aligned}$$

This also shows that

$$(5.1.6) \quad \mathcal{R}([Z]/S) = R\varprojlim(R_n \otimes_R^L \mathcal{R}([Z]/S)).$$

LEMMA 5.1.9. *In the situation of 5.1.8 we have*

$$\oplus_i H^i(\mathcal{R}([Z]/S)) = \hat{\mathcal{H}}([Z]/S) : \hat{\mathcal{H}}(X/S) \rightarrow \hat{\mathcal{H}}(Y/S),$$

in particular $H^i(\mathcal{R}([Z]/S))$ is $W\mathcal{O}_S$ -linear. (See Proposition 3.5.4 for the notation.) Further,

$$\oplus_i H^i(R_1 \otimes_R^L \mathcal{R}([Z]/S)) : \oplus_i R^i f_*\Omega_X \rightarrow \oplus_i R^i g_*\Omega_Y$$

equals the the morphism $\rho_H(Z/S)$ from [CR09, Prop. 3.2.4]; in particular it is \mathcal{O}_S -linear.

Proof. By Lemma 3.5.8 (and its proof for the level $n = 1$ case) the global cup product $(-) \cup \hat{\text{cl}}([Z]) : H^i(X \times Y, W\Omega_{X \times Y}) \rightarrow H_Z^{i+N}(X \times Y, W\Omega_{X \times Y}(N))$ (resp. $(-) \cup \text{cl}_1([Z]) : H^i(X \times Y, \Omega_{X \times Y}) \rightarrow H_Z^{i+N}(X \times Y, \Omega_{X \times Y}(N))$) is given by applying $H^i(X \times Y, R\varprojlim(-))$ to the cup product of 5.1.7 (resp. applying $H^i(X \times Y, R_1 \otimes_R^L (-))$ to 5.1.7). Now the lemma follows from going through the definitions (cf. also the proof of Proposition 3.5.10 and of [CR09, Lem. 4.1.3].) \square

THEOREM 5.1.10. *Let S be a k -scheme and $f : X \rightarrow S$ and $g : Y \rightarrow S$ be two S -schemes which are integral and smooth over k and have dimension N . Assume X and Y are properly birational over S , i.e. there exists a closed integral subscheme $Z \subset X \times_S Y$, such that the projections $Z \rightarrow X$ and $Z \rightarrow Y$ are*

proper and birational. Then $R_n \otimes_R^L \mathcal{R}([Z]/S)$ (see 5.1.8) induces isomorphisms in $D^b(S, W_n)$, for all $n \geq 1$

$$Rf_* W_n \mathcal{O}_X \cong Rg_* W_n \mathcal{O}_Y, \quad Rf_* W_n \Omega_X^N \cong Rg_* W_n \Omega_Y^N.$$

Therefore $\mathcal{R}([Z]/S) = R\varprojlim(R_n \otimes_R^L \mathcal{R}([Z]/S))$ induces isomorphisms in $D^b(S, R)$

$$Rf_* W \mathcal{O}_X \cong Rg_* W \mathcal{O}_Y, \quad Rf_* W \Omega_X^N \cong Rg_* W \Omega_Y^N.$$

Taking cohomology we obtain isomorphisms of $W\mathcal{O}_S$ -modules which are compatible with Frobenius and Verschiebung

$$R^i f_* W \mathcal{O}_X \cong R^i g_* W \mathcal{O}_Y, \quad R^i f_* W \Omega_X^N \cong R^i g_* W \Omega_Y^N, \quad \text{for all } i \geq 0.$$

Proof. By Ekedahl’s Nakayama Lemma (Corollary 5.1.3) it suffices, to show that

$$H^i(R_1 \otimes_R^L \mathcal{R}([Z]/S)) : R^i f_* \Omega_X \rightarrow R^i g_* \Omega_Y$$

is an isomorphism for all $i \geq 0$. This follows from Lemma 5.1.9 and [CR09, Thm. 3.2.6]. The second statement follows again from Lemma 5.1.9. \square

Remark 5.1.11. Notice that we do not need to assume here that any of the schemes X, Y or S are quasi-projective. We had to assume it in section 2 and 3 since, we can only prove the compatibility of $\mathcal{H}(-/S)$ with composition of correspondences, in the quasi-projective case. But in the argument above we only need that the maps exist in $D^b(S, R)$ and can then reduce to the result of [CR09], where no quasi-projectiveness assumption is needed.

Modulo torsion the $W\mathcal{O}$ -part of the following corollary was proved by Ekedahl in [Eke83].

COROLLARY 5.1.12. *Let X and Y be two smooth and proper k -schemes, which are birational and of pure dimension N . Then there are isomorphisms of $W[F, V]$ -modules*

$$H^i(X, W\mathcal{O}_X) \cong H^i(Y, W\mathcal{O}_Y), \quad H^i(X, W\Omega_X^N) \cong H^i(Y, W\Omega_Y^N), \quad \text{for all } i \geq 0$$

and also for all n , isomorphisms of W_n -modules

$$H^i(X, W_n \mathcal{O}_X) \cong H^i(Y, W_n \mathcal{O}_Y), \quad H^i(X, W_n \Omega_X^N) \cong H^i(Y, W_n \Omega_Y^N), \quad \text{for all } i \geq 0.$$

In the case where X and Y are tame finite quotients (see Definition 4.2.5) we have no map like $\mathcal{R}([Z]/S)$. This is why we have to assume that there exists a morphism in this case:

THEOREM 5.1.13. *Let $f : X \rightarrow Y$ be a proper and birational k -morphism between two tame finite quotients. Then we have isomorphisms*

$$f^* : W\mathcal{O}_Y \xrightarrow{\cong} Rf_* W\mathcal{O}_X, \quad Rf_* W\omega_X \cong f_* W\omega_X[0] \xrightarrow{\cong, f_*} W\omega_Y,$$

where $W\omega$ is defined in Definition 4.1.2 and f_* is the pushforward from Proposition 4.1.4, (6). There are also corresponding isomorphisms on each finite level.

Proof. By [CR09, Cor. 4.3.3] we have isomorphisms

$$f^* : \mathcal{O}_Y \xrightarrow{\cong} Rf_*\mathcal{O}_X, \quad Rf_*\omega_X \cong f_*\omega_X[0] \xrightarrow{\cong, f_*} \omega_Y.$$

By [KM98, Prop. 5.7] tame finite quotients are CM. Now the statement follows by induction from the two exact sequences (where X is any pure dimensional CM scheme)

$$0 \rightarrow W_{n-1}\mathcal{O}_X \xrightarrow{V} W_n\mathcal{O}_X \rightarrow \mathcal{O}_X \rightarrow 0$$

and (see Proposition 4.1.4, (7))

$$0 \rightarrow W_{n-1}\omega_X \xrightarrow{p} W_n\omega_X \xrightarrow{F^{n-1}} \omega_X \rightarrow 0.$$

□

Notice that the $W\mathcal{O}$ -part of the theorem is a direct consequence of [CR09, Cor. 4.3.3] and does not need any of the techniques developed in this paper.

COROLLARY 5.1.14. *In the situation of Theorem 5.1.13 we have isomorphisms of $W[V, F]$ -modules*

$$H^i(X, W\mathcal{O}_X) \cong H^i(Y, W\mathcal{O}_Y), \quad H^i(X, W\omega_X) \cong H^i(Y, W\omega_Y), \quad \text{for all } i \geq 0.$$

There are also corresponding isomorphisms on each finite level.

REFERENCES

- [BBE07] Pierre Berthelot, Spencer Bloch, and Hélène Esnault. On Witt vector cohomology for singular varieties. *Compositio Mathematica*, 143(2):363–392, 2007.
- [BER10] Pierre Berthelot, Hélène Esnault, and Kay Rülling. Rational points over finite fields for regular models of algebraic varieties of Hodge type ≥ 1 . *Annals of Mathematics*, 176(1):413–508, 2012.
- [BE08] Manuel Blickle and Hélène Esnault. Rational singularities and rational points. *Pure Appl. Math. Q.*, 4(3, part 2):729–741, 2008.
- [BS83] Spencer Bloch and Vasudevan Srinivas. Remarks on correspondences and algebraic cycles. *Amer. J. Math.*, 105(5):1235–1253, 1983.
- [Cha09] Andre Chatzistamatiou. First coniveau notch of the Dwork family and its mirror. *Mathematical Research Letters*, 16(4):563–575, 2009.
- [CR09] Andre Chatzistamatiou and Kay Rülling. Higher direct images of the structure sheaf in positive characteristic. *Algebra & Number Theory*, 5(6):693–775, 2011.
- [Con00] Brian Conrad. *Grothendieck duality and base change*, volume 1750 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 2000.
- [CH00] Alessio Corti and Masaki Hanamura. Motivic decomposition and intersection Chow groups. I. *Duke Math. J.*, 103(3):459–522, 2000.
- [dJ96] A. Johan de Jong. Smoothness, semi-stability and alterations. *Institut des Hautes Études Scientifiques. Publications Mathématiques*, (83):51–93, 1996.

- [dJ97] A. Johan de Jong. Families of curves and alterations. *Université de Grenoble. Annales de l'Institut Fourier*, 47(2):599–621, 1997.
- [DI87] Pierre Deligne and Luc Illusie. Relèvements modulo p^2 et décomposition du complexe de de Rham. *Inventiones Mathematicae*, 89(2):247–270, 1987.
- [Eke83] Torsten Ekedahl. Sur le groupe fondamental d'une variété unirrationnelle. *Comptes Rendus des Séances de l'Académie des Sciences. Série I. Mathématique*, 297(12):627–629, 1983.
- [Eke84] Torsten Ekedahl. On the multiplicative properties of the de Rham-Witt complex. I. *Arkiv för Matematik*, 22(2):185–239, 1984.
- [Eke85] Torsten Ekedahl. On the multiplicative properties of the de Rham-Witt complex. II. *Arkiv för Matematik*, 23(1):53–102, 1985.
- [ES80] Geir Ellingsrud and Tor Skjelbred. Profondeur d'anneaux d'invariants en caractéristique p . *Compositio Mathematica*, 41(2):233–244, 1980.
- [Ful98] William Fulton. *Intersection theory*, volume 2 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics]*. Springer-Verlag, Berlin, second edition, 1998.
- [Gab62] Pierre Gabriel. Des catégories abéliennes. *Bull. Soc. Math. France*, 90:323–448, 1962.
- [God73] Roger Godement. *Topologie algébrique et théorie des faisceaux*. Hermann, Paris, 1973. Troisième édition revue et corrigée, Publications de l'Institut de Mathématique de l'Université de Strasbourg, XIII, Actualités Scientifiques et Industrielles, No. 1252.
- [Gro85] Michel Gros. Classes de Chern et classes de cycles en cohomologie de Hodge-Witt logarithmique. *Mémoires de la Société Mathématique de France. Nouvelle Série*, (21):87, 1985.
- [EGAI] Alexander Grothendieck. Éléments de géométrie algébrique. I. Le langage des schémas. *Institut des Hautes Études Scientifiques. Publications Mathématiques*, (4):228, 1960.
- [EGAI] Alexander Grothendieck. Éléments de géométrie algébrique. II. Étude globale élémentaire de quelques classes de morphismes. *Institut des Hautes Études Scientifiques. Publications Mathématiques*, (8):222, 1961.
- [EGAIV(4)] Alexander Grothendieck. Éléments de géométrie algébrique. IV. Étude locale des schémas et des morphismes de schémas IV. *Institut des Hautes Études Scientifiques. Publications Mathématiques*, (32):361, 1967.
- [SGA2] Alexander Grothendieck. *Cohomologie locale des faisceaux cohérents et théorèmes de Lefschetz locaux et globaux (SGA 2)*. Documents Mathématiques (Paris) 4. Société Mathématique de France, Paris,

2005. Séminaire de Géométrie Algébrique du Bois Marie, 1962, Augmenté d'un exposé de Michèle Raynaud. With a preface and edited by Yves Laszlo, Revised reprint of the 1968 French original.
- [Har66] Robin Hartshorne. *Residues and duality*. Lecture notes of a seminar on the work of A. Grothendieck, given at Harvard 1963/64. With an appendix by Pierre Deligne. Lecture Notes in Mathematics, No. 20. Springer-Verlag, Berlin, 1966.
- [Ill79] Luc Illusie. Complexe de de Rham-Witt et cohomologie cristalline. *Annales Scientifiques de l'École Normale Supérieure. Quatrième Série*, 12(4):501–661, 1979.
- [Ill83] Luc Illusie. Finiteness, duality, and Künneth theorems in the cohomology of the de Rham-Witt complex. In *Algebraic geometry (Tokyo/Kyoto, 1982)*, volume 1016 of *Lecture Notes in Math.*, pages 20–72. Springer, Berlin, 1983.
- [IR83] Luc Illusie and Michel Raynaud. Les suites spectrales associées au complexe de de Rham-Witt. *Institut des Hautes Études Scientifiques. Publications Mathématiques*, (57):73–212, 1983.
- [KM74] Nicholas M. Katz and William Messing. Some consequences of the Riemann hypothesis for varieties over finite fields. *Invent. Math.*, 23:73–77, 1974.
- [KKMS73] George Kempf, Finn Faye Knudsen, David Mumford, and Bernard Saint-Donat. *Toroidal embeddings. I*. Lecture Notes in Mathematics, Vol. 339. Springer-Verlag, Berlin, 1973.
- [KM98] János Kollár and Shigefumi Mori. *Birational geometry of algebraic varieties*, volume 134 of *Cambridge Tracts in Mathematics*. Cambridge University Press, Cambridge, 1998. With the collaboration of C. H. Clemens and A. Corti, Translated from the 1998 Japanese original.
- [Kov00] Sándor J. Kovács. A characterization of rational singularities. *Duke Math. J.*, 102(2):187–191, 2000.
- [LZ04] Andreas Langer and Thomas Zink. De Rham-Witt cohomology for a proper and smooth morphism. *Journal of the Institute of Mathematics of Jussieu. JIMJ. Journal de l'Institut de Mathématiques de Jussieu*, 3(2):231–314, 2004.
- [ML98] Saunders Mac Lane. *Categories for the working mathematician*, volume 5 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, second edition, 1998.
- [Ray78] Michel Raynaud. Contre-exemple au “vanishing theorem” en caractéristique $p > 0$. In *C. P. Ramanujam – a tribute*, volume 8 of *Tata Inst. Fund. Res. Studies in Math.*, pages 273–278. Springer, Berlin, 1978.
- [Sai10] Shuji Saito. Cohomological Hasse principle and motivic cohomology of arithmetic schemes. *Proceedings of the International Congress of Mathematicians*, 2010.
- [Shi74] Tetsuji Shioda. An example of unirational surfaces in characteristic p . *Mathematische Annalen*, 211:233–236, 1974.

- [SGA7(2)] *Groupes de monodromie en géométrie algébrique. II*. Lecture Notes in Mathematics, Vol. 340. Springer-Verlag, Berlin, 1973. Séminaire de Géométrie Algébrique du Bois-Marie 1967–1969 (SGA 7 II), Dirigé par Pierre Deligne et Nicholas Katz.
- [Wei] Charles A. Weibel. *An introduction to homological algebra*, volume 38 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 1994.

Andre Chatzistamatiou
Fachbereich Mathematik
Universität Duisburg-Essen
45117 Essen
Germany
a.chatzistamatiou@uni-
due.de

Kay Rülling
Fachbereich Mathematik
Universität Duisburg-Essen
45117 Essen
Germany
kay.ruelling@uni-due.de

