

Mixed semicontinuous perturbation of a second order nonconvex sweeping process

Dalila Azzam-Laouir

Laboratoire de Mathématiques Pures et Appliquées

Université de Jijel, Algérie

Email: azzam_d@yahoo.com

Abstract

We prove a theorem on the existence of solutions of a second order differential inclusion governed by a class of nonconvex sweeping process with a mixed semicontinuous perturbation.

Keywords and phrases: Differential inclusions; Fixed-point theorem; Mixed semicontinuity; Sweeping process.

2000 Mathematics Subject Classification: 34A60, 34B15, 47H10.

1 Introduction

The existence of solutions for the second order differential inclusions governed by the sweeping process

$$(\mathcal{P}_F) \begin{cases} -\ddot{u}(t) \in N_{K(u(t))}(\dot{u}(t)) + F(t, u(t), \dot{u}(t)), & \text{a.e. } t \in [0, T], \\ \dot{u}(t) \in K(u(t)), \\ u(0) = u_0; \quad \dot{u}(0) = v_0, \end{cases}$$

where $N_{K(u(t))}(\cdot)$ denotes the normal cone to $K(u(t))$, has been thoroughly studied (when the sets $K(x)$ are convex or nonconvex) by Castaing for the first time when $F \equiv \{0\}$ see [5], and later by many other authors see for example [2], [3], [6], [10] and [12]. Note that in this literature some existence results are established for the problem (\mathcal{P}_F) with lower and upper semicontinuous perturbations.

Our aim in this paper is to prove existence results for (\mathcal{P}_F) when F is a mixed semicontinuous set-valued map and $K(x)$ are nonconvex sets. For the first order sweeping process with a mixed semicontinuous perturbation we refer the reader to [9], and to [13] for the sweeping process with non regular sets and to [1] for second order differential inclusions with mixed semicontinuous perturbations.

After some preliminaries, we present our main result in the finite dimensional space H whenever the sets $K(x)$ are uniformly ρ -prox-regular ($\rho > 0$) and the set-valued mapping F is mixed semicontinuous, that is, $F(\cdot, \cdot, \cdot)$ is measurable and for every $t \in [0, T]$, at each $(x, y) \in H \times H$ where $F(t, x, y)$ is convex the set-valued map $F(t, \cdot, \cdot)$ is upper semicontinuous, and whenever $F(t, x, y)$ is not convex $F(t, \cdot, \cdot)$ is lower semicontinuous on some neighborhood of (x, y) .

2 Definition and preliminary results

Let H be a real Hilbert space and let S be a nonempty closed subset of H . We denote by $d(\cdot, S)$ the usual distance function associated with S , i.e., $d(u, S) := \inf_{y \in S} \|u - y\|$. For any $x \in H$ and $r \geq 0$ the closed ball centered at x with radius r will be denoted by $\overline{\mathbf{B}}_H(x, r)$. For $x = 0$ and $r = 1$ we will put $\overline{\mathbf{B}}_H$ in place of $\overline{\mathbf{B}}_H(0, 1)$. $\mathcal{L}([0, T])$ is the σ -algebra of Lebesgue-measurable sets of $[0, T]$ and $\mathcal{B}(H)$ is the σ -algebra of Borel subsets of H . By $\mathbf{L}_H^1([0, T])$ we denote the space of all Lebesgue-Bochner integrable H -valued mappings defined on $[0, T]$ and by $\mathbf{C}_H([0, T])$ the Banach space of all continuous mappings $u : [0, T] \rightarrow H$, endowed with the sup norm

We need first to recall some notation and definitions that will be used in all the paper. Let x be a point in S . We recall (see [8]) that the proximal normal cone to S at x is defined by $N_S^P(x) := \partial^P \psi_S(x)$, where ψ_S denotes the indicator function of S , i.e., $\psi_S(x) = 0$ if $x \in S$ and $+\infty$ otherwise. Note that the proximal normal cone is also given by

$$N_S^P(x) = \{\xi \in H : \exists \alpha > 0 \text{ s.t. } x \in \text{Proj}_S(x + \alpha\xi)\},$$

where

$$\text{Proj}_S(u) := \{y \in S : d(u, S) := \|u - y\|\}.$$

Recall now that for a given $\rho \in]0, +\infty]$ the subset S is uniformly ρ -prox-regular (see [11]) or equivalently ρ -proximally smooth (see [8]) if and only if every nonzero proximal normal to S can be realized by ρ -ball, this means that for all $\bar{x} \in S$ and all $0 \neq \xi \in N_S^P(\bar{x})$ one has

$$\left\langle \frac{\xi}{\|\xi\|}, x - \bar{x} \right\rangle \leq \frac{1}{2\rho} \|x - \bar{x}\|^2,$$

for all $x \in S$. We make the convention $\frac{1}{\rho} = 0$ for $\rho = +\infty$. Recall that for $\rho = +\infty$ the uniform ρ -prox-regularity of S is equivalent to the convexity of S . The following proposition summarizes some important consequences of the uniform prox-regularity needed in the sequel. For the proof of these results we refer the reader to [11].

Proposition 2.1 *Let S be a nonempty closed subset in H and $x \in S$. The following assertions hold:*

- (1) $\partial^P d(x, S) = N_S^P(x) \cap \overline{\mathbf{B}}_H$;
 (2) let $\rho \in]0, +\infty]$. If S is uniformly ρ -prox-regular, then
 (2.1) for all $x \in H$ with $d(x, S) < \rho$; one has $\text{Proj}_S(x) \neq \emptyset$;
 (2.2) the proximal subdifferential of $d(\cdot, S)$ coincides with its Clarke subdifferential at all points $x \in H$ satisfying $d(x, S) < \rho$. So, in such a case, the subdifferential $\partial d(x, S) := \partial^P d(x, S) = \partial^C d(x, S)$ is a closed convex set in H ;
 (2.3) for all $x_i \in S$ and all $v_i \in N_S^P(x_i)$ with $\|v_i\| \leq \rho$ ($i = 1, 2$) one has

$$\langle v_1 - v_2, x_1 - x_2 \rangle \geq -\|x_1 - x_2\|^2.$$

As a consequence of (2.3) we get that for uniformly ρ -prox-regular sets, the proximal normal cone to S coincides with all the normal cones contained in the Clarke normal cone at all points $x \in S$, i.e., $N_S^P(x) = N_S^C(x)$. In such a case, we put $N_S(x) := N_S^P(x) = N_S^C(x)$. Here and above $\partial^C d(x, S)$ and $N_S^C(x)$ denote respectively the Clarke subdifferential of $d(\cdot, S)$ and the Clarke normal cone to S (see [8]).

Now, we recall some preliminaries concerning set-valued mappings. Let $T > 0$. Let $C : [0, T] \rightrightarrows H$ and $K : H \rightrightarrows H$ be two set-valued mappings. We say that C is absolutely continuous provided that there exists an absolutely continuous nonnegative function $a : [0, T] \rightarrow \mathbb{R}_+$ with $a(0) = 0$ such that

$$|d(x, C(t)) - d(y, C(s))| \leq \|x - y\| + |a(t) - a(s)|$$

for all $x, y \in H$ and all $s, t \in [0, T]$.

We will say that K is Hausdorff-continuous (resp. Lipschitz with ratio $\lambda > 0$) if for any $x \in H$ one has

$$\lim_{x' \rightarrow x} \mathcal{H}(K(x), K(x')) = 0$$

(resp. if for any $x, x' \in H$ one has

$$\mathcal{H}(K(x), K(x')) \leq \lambda \|x - x'\|.)$$

We close this section with the following theorem in [4], which is an important closedness property of the subdifferential of the distance function associated with a set-valued mapping.

Theorem 2.1 *Let $\rho \in]0, +\infty]$, Ω be an open subset in H , and $K : \Omega \rightrightarrows H$ be a Hausdorff-continuous set-valued mapping. Assume that $K(z)$ is uniformly ρ -prox-regular for all $z \in \Omega$. Then for a given $0 < \delta < \rho$, the following holds:*

"for any $\bar{z} \in \Omega$, $\bar{x} \in K(\bar{z}) + (\rho - \delta)\overline{\mathbf{B}}_H$, $x_n \rightarrow \bar{x}$, $z_n \rightarrow \bar{z}$ with $z_n \in \Omega$ (x_n not necessarily in $K(z_n)$) and $\xi_n \in \partial d(x_n, K(z_n))$ with $\xi_n \rightarrow^w \bar{\xi}$ one has $\bar{\xi} \in \partial d(\bar{x}, K(\bar{z}))$."

Here \rightarrow^w means the weak convergence in H .

Remark 2.1 *As a direct consequence of this theorem, we have for every $\rho \in]0, +\infty]$, for a given $0 < \delta < \rho$, and for every set-valued mapping $K : \Omega \rightrightarrows H$ with uniformly ρ -prox regular values, the set-valued mapping $(z, x) \mapsto \partial d(x, K(z))$ is upper semicontinuous from $\{(z, x) \in \Omega \times H : x \in K(z) + (\rho - \delta)\}$ to H endowed with the weak topology, which is equivalent to the upper semicontinuity of the function $(z, x) \mapsto \sigma(\partial d(x, K(z)), p)$ on $\{(z, x) \in \Omega \times H : x \in K(z) + (\rho - \delta)\}$, for any $p \in H$. Here $\sigma(S, p)$ denotes the support function to S defined by $\sigma(S, p) = \sup_{s \in S} \langle s, p \rangle$.*

3 Existence results under mixed semicontinuous perturbation.

Our existence result is stated in a finite dimensional space H under the following assumptions.

(H_1) For each $x \in H$, $K(x)$ is a nonempty closed subset in H and uniformly ρ -prox-regular for some fixed $\rho \in]0, +\infty[$;

(H_2) K is Lipschitz with ratio $\lambda > 0$;

(H_3) $l = \sup_{x \in H} |K(x)| < +\infty$.

The proof of our main theorem uses existence results for the first order sweeping process, the selection theorem proved in Tolstonogov [14] and the Kakutani fixed point theorem for set-valued mappings. We begin by recalling them.

Proposition 3.1 (Proposition 1.1 in [7]) *Let H be a finite dimensional space, $T > 0$ and let $C : I := [0, T] \rightrightarrows H$ be a nonempty closed valued set-valued mapping satisfying the following assumptions.*

(A_1) *For each $t \in I$, $C(t)$ is ρ -prox-regular for some fixed $\rho \in]0, +\infty[$;*

(A_2) *$C(t)$ varies in an absolutely continuous way, that is, there exists a nonnegative absolutely continuous function $v : I \rightarrow \mathbb{R}$ such that*

$$|d(x, C(t)) - d(y, C(s))| \leq \|x - y\| + |v(t) - v(s)|$$

for all $x, y \in H$ and $s, t \in I$.

Then for any mapping $h \in \mathbf{L}_H^1([0, T])$, the differential inclusion

$$\begin{cases} -\dot{u}(t) \in N_{C(t)}(u(t)) + h(t), & \text{a.e. } t \in [0, T], \\ u(0) = u_0 \in C(0) \end{cases}$$

admits one and only one absolutely continuous solution $u(\cdot)$ and

$$\|\dot{u}(t) + h(t)\| \leq |\dot{v}(t)| + \|h(t)\|.$$

Further, let m be a nonnegative Lebesgue-integrable function defined on $[0, T]$ and let

$$\mathcal{K} = \{h \in \mathbf{L}_H^1([0, T]) : \|h(t)\| \leq m(t) \text{ a.e.}\}.$$

Then the solutions set $\{u_h : h \in \mathcal{K}\}$, where u_h is the unique absolutely continuous solution of the above inclusion, is compact in $\mathbf{C}_H([0, T])$, and the mapping $h \mapsto u_h$ is continuous on \mathcal{K} when \mathcal{K} is endowed with the weak topology $w(\mathbf{L}_H^1([0, T]), \mathbf{L}_H^\infty([0, T]))$.

For the proof of our theorem we will also need the following theorem which is a direct consequence of Theorem 2.1 in [14].

Theorem 3.1 *Let H be a finite dimensional space and let $M : [0, T] \times H \times H \rightrightarrows H$ be a closed valued set-valued mapping satisfying the following hypotheses.*

(i) *M is $\mathcal{L}([0, T]) \otimes \mathcal{B}(H) \otimes \mathcal{B}(H)$ -measurable;*

(ii) for every $t \in [0, 1]$, at each $(x, y) \in H \times H$ such that $M(t, x, y)$ is convex, $M(t, \cdot, \cdot)$ is upper semicontinuous, and whenever $M(t, x, y)$ is not convex, $M(t, \cdot, \cdot)$ is lower semicontinuous on some neighborhood of (x, y) ;

(iii) there exists a Caratheodory function $\zeta : [0, 1] \times H \times H \rightarrow \mathbb{R}_+$ which is integrably bounded and such that $M(t, x, y) \cap \overline{\mathbf{B}}_H(0, \zeta(t, x, y)) \neq \emptyset$ for all $(t, x, y) \in [0, 1] \times H \times H$. Then for any $\varepsilon > 0$ and any compact set $K \subset \mathbf{C}_H([0, T])$ there is a nonempty closed convex valued multifunction $\Phi : K \rightrightarrows \mathbf{L}_H^1([0, T])$ which has a strongly-weakly sequentially closed graph such that for any $u \in K$ and $\phi \in \Phi(u)$ one has

$$\phi(t) \in M(t, u(t), \dot{u}(t));$$

$$\|\phi(t)\| \leq \zeta(t, u(t), \dot{u}(t)) + \varepsilon,$$

for almost every $t \in [0, T]$.

Now we are able to prove our main result.

Theorem 3.2 Let H be a finite dimensional space, $K : H \rightrightarrows H$ be a set-valued mapping satisfying assumptions (H_1) , (H_2) and (H_3) . Let $T > 0$ and let $F : [0, T] \times H \times H \rightrightarrows H$ be a set-valued mapping satisfying hypotheses (i) and (ii) of Theorem 3.1 and the following one

(iv) there exist nonnegative Lebesgue-integrable functions m, p and q defined on $[0, T]$ such that

$$F(t, x, y) \subset (m(t) + p(t)\|x\| + q(t)\|y\|)\overline{\mathbf{B}}_H$$

for all $(t, x, y) \in [0, T] \times H \times H$.

Then for all $u_0 \in H$ and $v_0 \in K(u_0)$, there exist two Lipschitz mappings $u, v : [0, T] \rightarrow H$ such that

$$\begin{cases} u(t) = u_0 + \int_0^t v(s)ds, & \forall t \in [0, T]; \\ -\dot{v}(t) \in N_{K(u(t))}(v(t)) + F(t, u(t), v(t)), & \text{a.e. on } [0, T]; \\ v(t) \in K(u(t)), & \forall t \in [0, T]; \\ u(0) = u_0; & v(0) = v_0 \end{cases}$$

with $\|\dot{v}(t)\| \leq \lambda + 2(m(t) + p(t)(\|u_0\| + lT) + q(t)l)$ a.e. $t \in [0, T]$.

In other words, there is a Lipschitz solution $u : [0, T] \rightarrow H$ to the Cauchy problem (\mathcal{P}_F) .

Proof. Step 1. Put $I := [0, T]$, $M(t) = m(t) + p(t)(\|u_0\| + lT) + q(t)l$, and let us consider the sets

$$\mathcal{X} = \{u \in \mathbf{C}_H(I) : u(t) = u_0 + \int_0^t \dot{u}(s)ds, \forall t \in I \text{ and } \|\dot{u}(t)\| \leq l, \text{ a.e. on } I\},$$

$$\mathcal{U} = \{v \in \mathbf{C}_H(I) : v(t) = v_0 + \int_0^t \dot{v}(s)ds, \forall t \in I \text{ and } \|\dot{v}(t)\| \leq \lambda + 2M(t), \text{ a.e. on } I\},$$

$$\mathcal{K} = \{h \in \mathbf{L}_H^1(I) : \|h(t)\| \leq M(t), \text{ a.e. on } I\}.$$

It is clear that \mathcal{K} is a convex $w(\mathbf{L}_H^1(I), \mathbf{L}_H^\infty(I))$ -compact subset of $\mathbf{L}_H^1(I)$, and by Ascoli-Arzelà theorem \mathcal{X} and \mathcal{U} are convex compact sets in $\mathbf{C}_H(I)$. Observe now, that for all $f \in \mathcal{X}$ the set valued mapping $K \circ f$ is Lipschitz with ratio λl . Indeed, for all $t, t' \in I$

$$\begin{aligned} \mathcal{H}((K \circ f)(t), (K \circ f)(t')) &= \mathcal{H}(K(f(t)), K(f(t'))) \\ &\leq \lambda \|f(t) - f(t')\| \\ &= \lambda \|x_0 + \int_0^t \dot{f}(s) ds - x_0 - \int_0^{t'} \dot{f}(s) ds\| \\ &\leq \lambda \int_{t'}^t |\dot{f}(s)| ds \leq \lambda l |t - t'|. \end{aligned}$$

By Proposition 3.1, for all $(f, h) \in \mathcal{X} \times \mathcal{K}$, there exists a unique solution $u_{f,h}$ to the problem

$$(P) \begin{cases} -\dot{u}_{f,h}(t) \in N_{K(f(t))}(u_{f,h}(t)) + h(t), & \text{a.e. on } I; \\ u_{f,h}(t) \in K(f(t)), \forall t \in I; \\ u_{f,h}(0) = v_0, \end{cases}$$

and for almost all $t \in I$, $\|\dot{u}_{f,h}(t)\| \leq \lambda l + 2M(t)$, i.e., $u_{f,h} \in \mathcal{U}$.

Let us consider the mapping $\mathcal{A} : \mathcal{X} \times \mathcal{K} \rightarrow \mathcal{U}$ defined by $\mathcal{A}(f, h) = u_{f,h}$, where $u_{f,h}$ is the unique solution of (P). We wish to show that \mathcal{A} is continuous. Let $(f_n, h_n)_n$ be a sequence in $\mathcal{X} \times \mathcal{K}$ such that $(f_n)_n$ converges uniformly to $f \in \mathcal{X}$ and $(h_n)_n$ converges $w(\mathbf{L}_H^1(I), \mathbf{L}_H^\infty(I))$ to $h \in \mathcal{K}$, and since $(u_{f_n, h_n})_n \subset \mathcal{U}$ we may suppose that it converges uniformly to some mapping $v \in \mathcal{U}$. For each $n \in \mathbb{N}$ we have

$$(P_n) \begin{cases} -\dot{u}_{f_n, h_n}(t) \in N_{K(f_n(t))}(u_{f_n, h_n}(t)) + h_n(t), & \text{a.e. on } I; \\ u_{f_n, h_n}(t) \in K(f_n(t)), \forall t \in I; \\ u_{f_n, h_n}(0) = v_0. \end{cases}$$

Since $u_{f_n, h_n}(t) \in K(f_n(t))$ for all $t \in I$, it follows from the Lipschitz property of K

$$d(u_{f_n, h_n}(t), K(f(t))) \leq \lambda \|f_n(t) - f(t)\| \rightarrow 0$$

and hence, one obtains $v(t) \in K(f(t))$, because the set $K(f(t))$ is closed. According to (P_n) one has

$$\dot{u}_{f_n, h_n}(t) + h_n(t) \in -N_{K(f_n(t))}(u_{f_n, h_n}(t)),$$

and

$$\|\dot{u}_{f_n, h_n}(t) + h_n(t)\| \leq \lambda l + M(t) := c(t), \text{ i.e., } \dot{u}_{f_n, h_n}(t) + h_n(t) \in c(t)\overline{\mathbf{B}}_H.$$

Therefore we get by Proposition 2.1 (1)

$$\dot{u}_{f_n, h_n}(t) + h_n(t) \in -c(t)\partial d(u_{f_n, h_n}(t), K(f_n(t))), \text{ a.e. on } I.$$

Now, as $(\dot{u}_{f_n, h_n} + h_n)_n$ converges weakly to $\dot{v} + h \in \mathbf{L}_H^1(I)$, Mazur's lemma ensures that for a.e. $t \in I$

$$\dot{v}(t) + h(t) \in \bigcap_n \overline{\text{co}}\{\dot{u}_{f_k, h_k}(t) + h_k(t) : k \geq n\}.$$

Fix such t in I and any μ in H , then the last relation gives

$$\begin{aligned} \langle \dot{v}(t) + h(t), \mu \rangle &\leq \limsup_n \sigma(-c(t)\partial d(u_{f_n, h_n}(t), K(f_n(t))), \mu) \\ &\leq \sigma(-c(t)\partial d(v(t), K(f(t))), \mu), \end{aligned}$$

where the second inequality follows from Remark 2.1 and Theorem 2.1. As the set $\partial d(v(t), K(f(t)))$ is closed and convex (see Proposition 2.1), we obtain

$$\dot{v}(t) + h(t) \in -c(t)\partial d(v(t), K(f(t))) \subset -N_{K(f(t))}(v(t)),$$

because $v(t) \in K(f(t))$. This can be rephrased as

$$\begin{cases} -\dot{v}(t) \in N_{K(f(t))}(v(t)) + h(t), & \text{a.e. on } I; \\ v(t) \in K(f(t)), \forall t \in I; \\ v(0) = v_0. \end{cases}$$

In other words, v is of the form $u_{f, h}$ with

$$\begin{cases} -\dot{u}_{f, h}(t) \in N_{K(f(t))}(u_{f, h}(t)) + h(t), & \text{a.e. on } I; \\ u_{f, h}(t) \in K(f(t)), \forall t \in I; \\ u_{f, h}(0) = v_0. \end{cases}$$

We conclude that \mathcal{A} is continuous. Hence, the mapping $P : \mathcal{X} \times \mathcal{K} \rightarrow \mathbf{C}_H(I)$ defined by $P(f, h)$, where for all $t \in I$

$$P(f, h)(t) = u_0 + \int_0^t \mathcal{A}(f, h)(s)ds = u_0 + \int_0^t u_{f, h}(s)ds$$

is also continuous when \mathcal{X} is endowed with the topology of uniform convergence and \mathcal{K} is endowed with the weak topology. Observe that for all $t \in I$, $u_{f, h}(t) \in K(f(t))$ and then by (H_3) , we have $\|u_{f, h}(t)\| \leq l$, we conclude that $P(f, h) \in \mathcal{X}$.

Step 2. By Theorem 3.1, there is a nonempty closed convex valued set-valued mapping $\Phi : \mathcal{X} \rightrightarrows \mathbf{L}_H^1(I)$ such that for any $u \in \mathcal{X}$ and $\phi \in \Phi(u)$

$$\phi(t) \in F(t, u(t), \dot{u}(t)) \quad \text{and} \quad \|\phi(t)\| \leq m(t) + p(t)\|u(t)\| + q(t)\|\dot{u}(t)\|$$

for almost all $t \in I$. Since $u \in \mathcal{X}$, we have $\|\dot{u}(t)\| \leq l$ and

$$\|u(t)\| = \|u_0 + \int_0^t \dot{u}(s)ds\| \leq \|u_0\| + lT,$$

hence

$$\|\phi(t)\| \leq m(t) + p(t)(\|u_0\| + lT) + q(t)l = M(t). \quad (3.1)$$

The relation (3.1) shows that Φ has $w(\mathbf{L}_H^1(I), \mathbf{L}_H^\infty(I))$ -compact values in $\mathbf{L}_H^1(I)$. Now, let us consider the set-valued mapping $\Psi : \mathcal{X} \rightrightarrows \mathcal{X}$ defined by

$$\Psi(f) = \{P(f, h) : h \in \Phi(f)\}.$$

It is clear that Ψ has nonempty convex values since Φ has nonempty convex values. Furthermore, for all $f \in \mathcal{X}$, $\Psi(f)$ is compact in \mathcal{X} . Indeed, Let $(v_n)_n$ be a sequence in $\Psi(f)$, then, for each n , there is $h_n \in \Phi(f)$ such that $v_n = P(f, h_n)$. Since $(h_n)_n \subset \Phi(f)$, by extracting a subsequence (that we do not relabel) we may suppose that $(h_n)_n$ $w(\mathbf{L}_H^1(I), \mathbf{L}_H^\infty(I))$ -converges to some mapping $h \in \Phi(f)$, and by the continuity of P we get $v_n = P(f, h_n) \rightarrow v = P(f, h) \in \mathcal{X}$. This shows the compactness of $\Psi(f)$.

We will prove that Ψ is upper semicontinuous, or equivalently the graph of Ψ $\text{gph}(\Psi) = \{(x, y) \in \mathcal{X} \times \mathcal{X} : y \in \Psi(x)\}$ is closed. Let $(x_n, y_n)_n$ be a sequence in $\text{gph}(\Psi)$ converging to $(x, y) \in \mathcal{X} \times \mathcal{X}$. For all $n \in \mathbb{N}$, $y_n \in \Psi(x_n)$, so, there is $h_n \in \Phi(x_n)$ such that $y_n = P(x_n, h_n)$. since $(h_n)_n \subset \mathcal{K}$, by extracting a subsequence (that we do not relabel) we may suppose that $(h_n)_n$ $w(\mathbf{L}_H^1(I), \mathbf{L}_H^\infty(I))$ -converges to some mapping $h \in \mathcal{K}$. As the sequence $(x_n)_n$ converges uniformly to $x \in \mathcal{X}$ and since $\text{gph}(\Phi)$ is strongly-weakly sequentially closed we conclude that $h \in \Phi(x)$. On the other hand, by the continuity of the mapping P we get

$$y = \lim_{n \rightarrow +\infty} y_n = \lim_{n \rightarrow +\infty} P(x_n, h_n) = P(x, h).$$

Hence $(x, y) \in \text{gph}(\Psi)$. This says that Ψ is upper semicontinuous. An application of Kakutani Theorem gives a fixed point of Ψ , that is, there is $f \in \mathcal{X}$ such that $f \in \Psi(f)$, which means that there is $h \in \Phi(f)$ such that $f = P(f, h)$. Consequently

$$f(t) = u_0 + \int_0^t u_{f,h}(s)ds \text{ and } h(t) \in F(t, f(t), u_{f,h}(t))$$

with

$$\begin{cases} -\dot{u}_{f,h}(t) \in N_{K(f(t))}(u_{f,h}(t)) + h(t), & \text{a.e. on } I; \\ u_{f,h}(t) \in K(f(t)), \forall t \in I; \\ u_{f,h}(0) = v_0, \end{cases}$$

or, by putting $u = f$

$$\begin{cases} -\ddot{u}(t) \in N_{K(u(t))}(\dot{u}(t)) + F(t, u(t), \dot{u}(t)), & \text{a.e. on } I; \\ \dot{u}(t) \in K(u(t)), \forall t \in I; \\ u(0) = u_0; \dot{u}(0) = v_0, \end{cases}$$

with, for almost all $t \in I$, $\|\ddot{u}(t)\| \leq \lambda + 2M(t)$. This finish the proof of our theorem. \blacksquare

References

- [1] D. Azzam-Laouir, S. Lounis and L. Thibault, Existence solutions for second-order differential inclusions with nonconvex perturbations. *Appl. Anal.* Vol. 86. No. 10. October 2007, 1199-1210.

- [2] M. Bounkhel, General existence results for second order nonconvex sweeping process with unbounded perturbations. *Portugaliae Mathematica* 60 (3) (2003), 269-304.
- [3] M. Bounkhel and D. Laouir-Azzam, Existence results on the second-order nonconvex sweeping process with perturbations. *Set-Valued Anal.* **12**: 291-318, 2004.
- [4] M. Bounkhel and L. Thibault, Nonconvex sweeping process and prox-regularity in Hilbert space, *J. Nonlinear Convex Anal.* Volume 6, Number 2, 2005.
- [5] C. Castaing, Quelques problèmes d'évolution du second ordre, *Sem. Anal. Convexe*, Montpellier. 1998, exposé No. 5.
- [6] C. Castaing, T. X. Duc Ha and M. Valadier, Evolution equations governed by the sweeping process, *Set-Valued Anal.* **1** (1993), 109-139.
- [7] C. Castaing, A. Salvadori and L. Thibault, Functional evolution equations governed by nonconvex sweeping process. *J. Nonlinear Convex Anal.* Vol 2, number 1, 2001.
- [8] F.H. Clarke, R.L. Stern and P.R. Wolenski, Proximal smoothness and the lower C^2 property. *J. Convex Anal.* Vol 2 (1995), No 1/2, 117-144.
- [9] T. Haddad and L. Thibault, Mixed semicontinuous perturbation of nonconvex sweeping process. *to appear*.
- [10] M. D. P. Monteiro-Marques, Differential inclusions in nonsmooth mechanical problem, shocks and dry friction. *Birkhauser*, 1995.
- [11] R.A. Poliquin, R.T. Rockafellar and L. Thibault, Local differentiability of distance functions. *Trans. Math. Soc.*, Vol. 352 (2000), No 11, 5231-5249.
- [12] A. Syam, Contributions aux inclusions différentielles. *Thèse, Université Montpellier II*. Montpellier 1993.
- [13] L. Thibault, Sweeping process with regular and nonregular sets. *J. Diff. Equa.* 193, 1-26 (2003).
- [14] A. A. Tolstonogov, Solutions of a differential inclusion with unbounded right-hand side. (Russian) *Sibirsk. Mat. Zh.* **29** no. 5, 212-225, 241 translation in *Siberian. Math. J.* **29**, no. 5, (1988). 857-868.

(Received June 29, 2008)