

**ON THE PERIODIC BOUNDARY-VALUE PROBLEM FOR  
SYSTEMS OF SECOND-ORDER NONLINEAR ORDINARY  
DIFFERENTIAL EQUATIONS**

G. GAPRINDASHVILI

ABSTRACT. The periodic boundary-value problem for systems of second-order ordinary nonlinear differential equations is considered. Sufficient conditions for the existence and uniqueness of a solution are established.

§ 1. STATEMENT OF THE MAIN RESULTS

Consider the periodic boundary-value problem

$$x'' = f(t, x, x'), \tag{1.1}$$

$$x(a) = x(b), \quad x'(a) = x'(b), \tag{1.2}$$

where the vector-function  $f : [a, b] \times \mathbb{R}^{2n} \rightarrow \mathbb{R}^n$  ( $\mathbb{R}^n$  denotes the  $n$ -dimensional Euclidean space with the norm  $\|\cdot\|$ ) satisfies the local Caratheodory conditions, i.e.,  $f(\cdot, x, y) : [a, b] \rightarrow \mathbb{R}^n$  is measurable for each  $(x, y) \in \mathbb{R}^{2n}$ ,  $f(t, \cdot) : \mathbb{R}^{2n} \rightarrow \mathbb{R}^n$  is continuous for almost all  $t \in [a, b]$ , and the function

$$f_r(\cdot) = \sup\{\|f(\cdot, x, y)\| : \|x\| + \|y\| \leq r\}$$

is Lebesgue integrable on  $[a, b]$  for each positive  $r$ .

By a solution of the problem (1.1), (1.2) we mean a vector-function  $x : [a, b] \rightarrow \mathbb{R}^n$  which has the absolutely continuous first derivative on  $[a, b]$  and satisfies the differential system (1.1) almost everywhere in  $[a, b]$ , as well as the boundary conditions (1.2).

For the literature on (1.1),(1.2) we refer to [1,2] and the references cited therein. Note that [1] deals with the scalar variant of the boundary-value problem (1.1),(1.2) (i.e., when  $n = 1$ ).

---

1991 *Mathematics Subject Classification.* 34B15, 34C25.

*Key words and phrases.* Second-order differential equation, periodic boundary-value problem, Nagumo pair.

Below, the sufficient conditions for solvability and unique solvability of the boundary value problem (1.1), (1.2) are given. They supplement some of those mentioned above.

We use the following notation:

$x \cdot y$  is the inner product of vectors  $x, y \in \mathbb{R}^n$ ;

$\mathbb{R} = \mathbb{R}^1$ ,  $\mathbb{R}_+ = [0, +\infty[$ ;

for each positive number  $r$  and vector  $y \in \mathbb{R}^n$

$$\eta_r(y) = \begin{cases} 0 & \text{for } \|y\| \leq r, \\ \frac{y}{\|y\|} & \text{for } \|y\| > r; \end{cases}$$

$U_\delta(t_0)$  is the  $\delta$ -neighborhood of  $t_0 \in \mathbb{R}$ ;

$\tilde{C}^1([a, b]; S)$  ( $S \subset \mathbb{R}^n$ ) is the set of vector-functions  $x : [a, b] \rightarrow S$  which have an absolutely continuous first derivative on  $[a, b]$ ;

$C(S_1; S_2)$  ( $S_1 \subset \mathbb{R}, S_2 \subset \mathbb{R}^n$ ) is the set of continuous vector-functions  $x : S_1 \rightarrow S_2$ ;

$L([a, b]; S)$  ( $S \subset \mathbb{R}$ ) is the set of functions  $x : [a, b] \rightarrow S$  which are Lebesgue integrable on  $[a, b]$ .

**Definition 1.1** (see [3, Definition 1.1] or [4, Definition 1.2]). Suppose that the functions  $\varphi : ]a, b[ \rightarrow ]0, \infty[$  and  $z : ]a, b[ \rightarrow \mathbb{R}^n$  have a first derivative which is absolutely continuous on every segment contained in  $]a, b[$ . A pair of functions  $(\varphi, z)$  is said to be a Nagumo pair of the differential system (1.1) if the condition

$$(x - z(t)) \cdot (f(t, x, y) - z''(t)) + \|y - z'(t)\|^2 - (\varphi'(t))^2 \geq \varphi(t)\varphi''(t)$$

$$\text{for } a < t < b, \|x - z(t)\| = \varphi(t) \text{ and } (x - z(t)) \cdot (y - z'(t)) = \varphi(t)\varphi'(t)$$

is satisfied, the function  $\|z''(t)\| + \varphi(t)$  being essentially bounded from above on every segment contained in  $]a, b[$ .

*Remark 1.1.* The Nagumo pair of differential system (1.1) serves as a vector analog for the upper and lower functions of the scalar equation (1.1), which were introduced by Nagumo [5] and which since then have been widely adopted in the theory of boundary-value problems (see [1] and the references cited therein; also [4, Remark 1.2], [3, Remark 1.1]). Namely, if  $n = 1$  and  $\sigma_1$  and  $\sigma_2$  are, respectively, the upper and lower solutions of the differential equation (1.1), then the pair  $(\varphi, z)$  defined by

$$\varphi(t) = \frac{\sigma_2(t) - \sigma_1(t)}{2} \quad \text{and} \quad z(t) = \frac{\sigma_2(t) + \sigma_1(t)}{2} \quad (1.3)$$

is the Nagumo pair of (1.1) (and vice versa).

Note also that the condition

$$x \cdot f(t, x, y) + \|y\|^2 \geq 0 \quad \text{for } \|x\| = r_0 \quad \text{and } x \cdot y = 0 \quad (1.4)$$

(see [2, Theorem 3.1]) is necessary and sufficient for  $(\varphi, z)$  to be a Nagumo pair of (1.1), where  $z(t) \equiv 0$  and  $\varphi(t) \equiv r_0 > 0$ .

**Definition 1.2.** A Nagumo pair  $(\varphi, z)$  of the differential system (1.1) is said to be a Nagumo pair of the problem (1.1),(1.2) if  $\varphi \in \tilde{C}^1([a, b]; \mathbb{R}_+)$ ,  $z \in \tilde{C}^1([a, b]; \mathbb{R}^n)$  and the conditions

$$\varphi(a) = \varphi(b), \quad z(a) = z(b) \quad (1.5_1)$$

and

$$\|z'(a) - z'(b)\| \leq \varphi'(b) - \varphi'(a) \quad (1.5_2)$$

are satisfied.

*Remark 1.2.* In the scalar case, (1.5<sub>1</sub>) – (1.5<sub>2</sub>) are equivalent to the conditions

$$\sigma_i(a) = \sigma_i(b), \quad (-1)^i(\sigma'_i(a) - \sigma'_i(b)) \leq 0 \quad (i = 1, 2),$$

assuming that (1.3) is satisfied. See these conditions in [1, § 16].

**Definition 1.3** (see [3, Definition 2.1] or [4, Definition 1.1]). Suppose that  $\varphi \in C([a, b]; \mathbb{R}_+)$  and  $z \in C([a, b]; \mathbb{R}^n)$ . A vector-function  $f$  is said to have the property  $V([a, b], \varphi, z)$  if there exist positive constants  $r$  and  $r_1$  such that if  $a \leq t_1 < t_2 \leq b$ ,  $\chi \in C(\mathbb{R}_+; [0, 1])$  and  $x \in \tilde{C}^1([t_1, t_2]; \mathbb{R}^n)$  is an arbitrary solution of the differential system

$$x'' = \chi(\|x'\|)f(t, x, x') \quad (1.6)$$

satisfying the inequalities

$$\|x(t) - z(t)\| \leq \varphi(t) \quad \text{for } t_1 \leq t \leq t_2 \quad (1.7)$$

and

$$\|x'(t)\| \geq r \quad \text{for } t_1 \leq t \leq t_2, \quad (1.8)$$

then  $x$  admits the estimate

$$\int_{t_1}^{t_2} \|x'(t)\| dt \leq r_1. \quad (1.9)$$

*Remark 1.3.* It is clear that each scalar function has the property  $V([a, b], \varphi, z)$  taking an arbitrary positive number for  $r$  and  $2 \max\{\varphi(t) + \|z(t)\| : a \leq t \leq b\}$  for  $r_1$ . The class of vector-functions with the property  $V([a, b], \varphi, z)$  is introduced just to unify the approach to the problem (1.1), (1.2) in both the scalar and the vector cases. Some other boundary-value problems were also studied using this approach (see [3,4] and the references cited therein).

Effective sufficient conditions for a vector-function  $f$  to have the property  $V([a, b], \varphi, z)$  are contained in [4, Propositions 1.1, 1.2] and [3, Proposition 2.1]. For example, if

$$(f(t, x, y) \cdot y)(x \cdot y) - (x \cdot f(t, x, y))\|y\|^2 \leq l(t)\|y\|^3 + k\|y\|^4 \quad (1.10)$$

for  $a \leq t \leq b$ ,  $\|x - z(t)\| \leq \varphi(t)$  and  $\|y\| > \rho$ ,

where  $l \in L([a, b]; \mathbb{R}_+)$ ,  $k < 1$  and  $\rho > 0$ , then  $f$  has the property  $V([a, b], \varphi, z)$ .

**Theorem 1.1<sub>1</sub>.** *Suppose that  $(\varphi, z)$  is a Nagumo pair of (1.1), (1.2), the vector-function  $f$  has the property  $V([a, b], \varphi, z)$ , and the inequality*

$$f(t, x, y) \cdot \eta_\rho(y) \leq w(\|y\|)(l(t) + \|y\|) \quad (1.11)$$

*is satisfied on the set*

$$\{(t, x, y) : a < t < b, \|x - z(t)\| \leq \varphi(t)\}, \quad (1.12)$$

*where  $\rho > 0$ ,  $l \in L([a, b]; \mathbb{R}_+)$ ,  $\omega \in C(\mathbb{R}_+; ]0, +\infty[)$ , and*

$$\int_0^{+\infty} \frac{ds}{\omega(s)} = +\infty. \quad (1.13)$$

*Then the boundary-value problem (1.1), (1.2) has at least one solution  $x \in \tilde{C}^1([a, b]; \mathbb{R}^n)$  satisfying the estimate*

$$\|x(t) - z(t)\| \leq \varphi(t) \quad \text{for } a \leq t \leq b. \quad (1.14)$$

**Theorem 1.1<sub>2</sub>.** *The conclusion of Theorem 1.1<sub>1</sub> remains valid if (1.11) is replaced by*

$$f(t, x, y) \cdot \eta_\rho(y) \geq -\omega(\|y\|)(l(t) + \|y\|). \quad (1.15)$$

**Theorem 1.2.** *Suppose that  $(\varphi, z)$  is a Nagumo pair of (1.1), (1.2), the vector-function  $f$  has the property  $V([a, b], \varphi, z)$ , the inequality (1.11) is satisfied on the set*

$$\{(t, x, y) : a_0 \leq t \leq b, \quad \|x - z(t)\| \leq \varphi(t)\},$$

and the inequality (1.15) on the set

$$\{(t, x, y) : a < t < b_0, \quad \|x - z(t)\| \leq \varphi(t)\},$$

where  $\rho > 0$ ,  $a \leq a_0 < b_0 \leq b$ ,  $l \in L([a, b]; \mathbb{R}_+)$ ,  $\omega \in C(\mathbb{R}_+; ]0, +\infty[)$ , and (1.13) holds. Then the boundary-value problem (1.1), (1.2) has at least one solution  $x \in \tilde{C}^1([a, b]; \mathbb{R}^n)$  satisfying the estimate (1.14).

**Theorem 1.3.** *Suppose that  $(\varphi, z)$  is a Nagumo pair of (1.1), (1.2), the vector-function  $f$  has the property  $V([a, b], \varphi, z)$ , the inequality (1.11) is satisfied on the set*

$$\{(t, x, y) : t \in ]a_1, t_0[ \cup ]b_2, b[, \quad \|x - z(t)\| \leq \varphi(t)\},$$

and the inequality (1.15) on the set

$$\{(t, x, y) : t \in ]a, a_2[ \cup ]t_0, b_1[, \quad \|x - z(t)\| \leq \varphi(t)\},$$

where  $\rho > 0$ ,  $a < a_1 < a_2 < t_0 < b_2 < b_1 < b$ ,  $l \in L([a, b]; \mathbb{R}_+)$ ,  $\omega \in C(\mathbb{R}_+; ]0, +\infty[)$ , and (1.13) holds. Then the boundary-value problem (1.1), (1.2) has at least one solution  $x \in \tilde{C}^1([a, b]; \mathbb{R}^n)$  satisfying the estimate (1.14).

*Remark 1.4.* Theorems 1.1–1.3 extend Theorem 3.1 [2] in the case of periodic boundary-value problem. As an example, define  $f_i(t, x, y) = -y_i \|y\|^m + 1 - \|x\|$  and  $f = (f_i)_{i=1}^n$ , where  $m$  is an arbitrary natural number. Let us verify the conditions of, e.g., Theorem 1.1<sub>1</sub> assuming that  $z(t) \equiv 0$ ,  $\varphi \equiv 1$ ,  $\rho = 1$ ,  $l(t) \equiv 1$ , and  $\omega \equiv 1$ . First, according to (1.4) where  $r_0 = 1$ ,  $(\varphi, z)$  is the Nagumo pair of (1.1), (1.2). Further, according to (1.10) where  $k = 0$ , the vector-function  $f$  has the property  $V([a, b], \varphi, z)$ . Finally, the correctness of (1.11), as well as of (1.13), is evident. On the other hand, Theorem 3.1 [2] fails for this example when  $m > 2$ .

Theorem 1.2 can also be considered as a vector analog of Theorem 16.2 from [1].

**Theorem 1.4.** *Suppose that for each positive  $r$  there exist  $l_i(t, r) \in L([a, b]; \mathbb{R}_+)$  ( $i = 1, 2$ ) such that  $l_1(t, r)$  differs from zero on a subset of positive measure of the interval  $]a, b[$  and*

$$\begin{aligned} & [f(t, x_1, y_1) - f(t, x_2, y_2)](x_1 - x_2) \geq \\ & \geq l_1(t, r) \|x_1 - x_2\|^2 - l_2(t, r) |(x_1 - x_2) \cdot (y_1 - y_2)| \\ & \text{for } \|x_k\| \leq r, \quad \|y_k\| \leq r \quad (k = 1, 2). \end{aligned} \tag{1.16}$$

Then the boundary-value problem (1.1), (1.2) has at most one solution in the class  $\tilde{C}^1([a, b]; \mathbb{R}^n)$ .

Theorem 1.4 can be considered as a vector analog of Theorem 16.4 from [1].

## § 2. SOME AUXILIARY RESULTS

**Lemma 2.1.** *Suppose that a vector-function  $q : [a, b] \times \mathbb{R}^{2n} \rightarrow \mathbb{R}^n$  satisfies the local Caratheodory conditions and the inequality*

$$\|q(t, x, y)\| \leq l(t)$$

holds on  $[a, b] \times \mathbb{R}^{2n}$  where  $l \in L([a, b]; \mathbb{R}_+)$ . Then the differential system

$$x'' = x + q(t, x, y)$$

has at least one solution  $x \in \tilde{C}^1([a, b]; \mathbb{R}^n)$  satisfying the boundary conditions (1.2).

*Proof.* It is easy to verify that the differential system

$$x'' = x$$

has no nontrivial solution satisfying the boundary conditions (1.2). Thus Lemma 2.1 immediately follows from Proposition 2.3 [1].  $\square$

The next result deals with the solvability of an auxiliary differential system

$$x'' = g(t, x, x'). \quad (2.1)$$

**Lemma 2.2.** *Suppose that  $(\varphi, z)$  is a Nagumo pair of the boundary-value problem (2.1), (1.2) and on  $[a, b] \times \mathbb{R}^{2n}$*

$$\|g(t, x, y)\| \leq h(t, x), \quad (2.2)$$

where the vector-functions  $g : [a, b] \times \mathbb{R}^{2n} \rightarrow \mathbb{R}^n$  and  $h : [a, b] \times \mathbb{R}^n \rightarrow \mathbb{R}^n$  satisfy the local Caratheodory conditions. Then the boundary-value problem (2.1), (1.2) has at least one solution  $x \in \tilde{C}^1([a, b]; \mathbb{R}^n)$  satisfying the estimate (1.14).

*Proof.* Put

$$\sigma(s, t) = \begin{cases} 0 & \text{for } s \leq 0 \text{ and } \tau \in \mathbb{R}, \\ \tau & \text{for } |\tau| < s, \\ s \operatorname{sign} \tau & \text{for } |\tau| \geq s > 0, \end{cases}$$

$$\gamma(t, x) = \begin{cases} 1 & \text{for } \|x - z(t)\| \leq \varphi(t), \\ \frac{\varphi(t)}{\|x - z(t)\|} & \text{for } \|x - z(t)\| > \varphi(t), \end{cases}$$

$$\begin{aligned}
\sigma_1(t, x, y) &= \sigma(\|x - z(t)\| - \varphi(t), \varphi'(t)\|x - z(t)\| - (x - z(t)) \cdot (y - z'(t))), \\
\tilde{y}(t, x, y) &= \begin{cases} y & \text{for } \|x - z(t)\| \leq \varphi(t), \\ y + \frac{\sigma_1(t, x, y)}{\|x - z(t)\|^2}(x - z(t)) & \text{for } \|x - z(t)\| > \varphi(t), \end{cases} \\
g_1(t, x, y) &= x - z(t) + \gamma(t, x)[f(t, z(t) + \gamma(t, x)(x - z(t)), \\
&\quad \tilde{y}(t, x, y)) - x + z(t)] - (\gamma(t, x) - 1)z''(t), \\
\sigma_2(t, x, y) &= \sigma\left[\|x - z(t)\| - \varphi(t), \|\tilde{y}(t, x, y) - z'(t)\|^2 - \right. \\
&\quad \left. - \|y - z'(t)\|^2 - (\varphi'(t))^2 + \left(\frac{(x - z(t)) \cdot (y - z'(t))}{\|x - z(t)\|}\right)^2\right], \\
g_2(t, x, y) &= \begin{cases} 0 & \text{for } \|x - z(t)\| \leq \varphi(t), \\ \frac{\sigma_2(t, x, y) + (\|x - z(t)\| - \varphi(t))(\varphi''(t) + 1)}{\|x - z(t)\|^2}(x - z(t)) & \text{for } \|x - z(t)\| > \varphi(t) \end{cases}
\end{aligned}$$

and

$$\tilde{g}(t, x, y) = g_1(t, x, y) + g_2(t, x, y).$$

By the condition (2.2) on  $[a, b] \times \mathbb{R}^{2n}$  we have

$$\|\tilde{g}(t, x, y) - x\| \leq h^*(t) + |\varphi''(t)| + \|z''(t)\| + 2 + \|z(t)\| + \varphi(t)$$

where

$$h^*(t) = \sup\{h(t, x) : \|x - z(t)\| \leq \varphi(t)\}.$$

Therefore according to Lemma 2.1 the differential system

$$x'' = \tilde{g}(t, x, x')$$

has at least one solution  $x \in \tilde{C}^1([a, b]; \mathbb{R}^n)$  satisfying the boundary conditions (1.2). Due to the definition of  $\tilde{g}$  it remains to show that  $x$  admits the estimate (1.14). Assume, on the contrary, that (1.14) is violated. Then there exists  $t_0 \in [a, b]$  where the function

$$u(t) = \|x(t) - z(t)\| - \varphi(t)$$

reaches its positive maximum on  $[a, b]$ . Assume first that  $t_0 \in ]a, b[$ . Then by the Fermat theorem we have

$$u'(t_0) = \frac{(x(t_0) - z(t_0)) \cdot (x'(t_0) - z'(t_0))}{\|x(t_0) - z(t_0)\|} - \varphi'(t_0) = 0.$$

Hence

$$\lim_{t \rightarrow t_0} \tilde{y}(t, x(t), x'(t)) = x'(t_0)$$

and for a certain positive  $\delta$  the inequalities

$$\begin{aligned}
|\varphi'(t)\|x(t) - z(t)\| - (x(t) - z(t)) \cdot (x'(t) - z'(t))| &< \\
&< \|x(t) - z(t)\| - \varphi(t)
\end{aligned} \tag{2.3}$$

and

$$\begin{aligned} & \|\tilde{y}(t, x(t), x'(t)) - z'(t)\|^2 - \|x'(t) - z'(t)\|^2 - (\varphi'(t))^2 + \\ & + \left( \frac{(x(t) - z(t)) \cdot (x'(t) - z'(t))}{\|x(t) - z(t)\|} \right)^2 < \|x(t) - z(t)\| \end{aligned}$$

hold in  $U_\delta(t_0)$ . Therefore due to the definition of  $\tilde{g}$  we obtain

$$\begin{aligned} u''(t) &= \frac{(x(t) - z(t)) \cdot (x''(t) - z''(t)) + \|x'(t) - z'(t)\|^2}{\|x(t) - z(t)\|} - \\ & - \frac{\left[ \frac{(x(t) - z(t)) \cdot (x'(t) - z'(t))}{\|x(t) - z(t)\|} \right]^2}{\|x(t) - z(t)\|} - \varphi''(t) = \\ & = \frac{(x(t) - z(t)) \cdot (g_1(t, x(t), x'(t)) - z''(t))}{\|x(t) - z(t)\|} + \\ & + \frac{\|\tilde{y}(t, x(t), x'(t)) - z'(t)\|^2 - \varphi(t)\varphi''(t) - (\varphi'(t))^2}{\|x(t) - z(t)\|} + \\ & + \frac{\|x(t) - z(t)\| - \varphi(t)}{\|x(t) - z(t)\|} \quad \text{for } t \in U_\delta(t_0). \end{aligned}$$

Furthermore, by (2.3) for each  $t \in U_\delta(t_0)$  we have

$$\|\gamma(t, x(t))(x(t) - z(t))\| = \varphi(t)$$

and

$$\gamma(t, x(t))(x(t) - z(t)) \cdot (\tilde{y}(t, x(t), x'(t)) - z'(t)) = \varphi(t)\varphi'(t).$$

Taking into account the last three equalities and Definition 1.1, it can be shown that  $u''(t)$  is positive for each  $t \in U_\delta(t_0)$ . But this is impossible, since  $t_0 \in ]a, b[$  and  $t_0$  is a point of maximum for  $u$ . Thus  $t_0 \notin ]a, b[$ . In view of (1.2) and (1.5<sub>1</sub>) both  $a$  and  $b$  are the points of maxima for the function  $u$ . Therefore  $u'(a) \leq 0$  and  $u'(b) \geq 0$ . Assuming  $u'(a) = 0$  or  $u'(b) = 0$ , an argument similar to the one carried out above leads us to a contradiction. Thus

$$u'(a) = \frac{(x(a) - z(a)) \cdot (x'(a) - z'(a))}{\|x(a) - z(a)\|} - \varphi'(a) < 0$$

and

$$u'(b) = \frac{(x(b) - z(b)) \cdot (x'(b) - z'(b))}{\|x(b) - z(b)\|} - \varphi'(b) > 0.$$

But since  $x'(a) = x'(b)$  and

$$\frac{x(a) - z(a)}{\|x(a) - z(a)\|} = \frac{x(b) - z(b)}{\|x(b) - z(b)\|},$$

the last two inequalities yield

$$\frac{x(a) - z(a)}{\|x(a) - z(a)\|} \cdot (z'(a) - z'(b)) > \varphi'(b) - \varphi'(a),$$

which contradicts (1.5<sub>2</sub>). Therefore the estimate (1.14) is proved.  $\square$

**Lemma 2.3.** *Suppose that  $\rho, r_1$ , and  $\rho'$  are positive constants,  $\omega \in C(\mathbb{R}_+; ]0, +\infty[)$ ,  $l \in L([t_1, t_2]; \mathbb{R}_+)$  and*

$$\int_{\rho}^{\rho'} \frac{ds}{\omega(s)} > r_1 + \int_{t_1}^{t_2} l(t) dt. \quad (2.4)$$

Then an arbitrary  $x \in \tilde{C}^1([t_1, t_2]; \mathbb{R}^n)$ , satisfying (1.9) and the inequalities

$$\|x'(t_i)\| \leq \rho \quad (2.5_i)$$

and

$$\begin{aligned} (-1)^{i-1} x''(t) \cdot \eta_{\rho}(x'(t)) &\leq \omega(\|x'(t)\|)(l(t) + \|x'(t)\|) \\ \text{for } t_1 \leq t \leq t_2 \end{aligned} \quad (2.6_i)$$

with  $i \in \{1, 2\}$ , admits the estimate

$$\|x'(t)\| \leq \rho' \quad \text{for } t_1 \leq t \leq t_2. \quad (2.7)$$

*Proof.* Assume for definiteness that  $i = 1$ . Admit to the contrary that (2.7) is violated, i.e., there exists  $t^* \in ]t_1, t_2]$  such that

$$\|x'(t^*)\| > \rho'. \quad (2.8)$$

By (2.5<sub>1</sub>) there exists  $t_* \in [t_1, t^*[$  such that

$$\|x'(t)\| = \rho \quad \text{and} \quad \|x'(t)\| > \rho' \quad \text{for } t_* \leq t \leq t^*.$$

Hence, taking into account the definition of the function  $\eta_r$ , from (2.6<sub>1</sub>) we get

$$\|x'(t)\|' \leq \omega(\|x'(t)\|)(l(t) + \|x'(t)\|) \quad \text{for } t_* \leq t \leq t^*.$$

Dividing this inequality by  $\omega(\|x'(t)\|)$ , integrating from  $t_*$  to  $t^*$ , and using (1.9) and (2.5<sub>1</sub>), we obtain

$$\int_{\rho}^{\|x'(t^*)\|} \frac{ds}{\omega(s)} \leq r_1 + \int_{t_1}^{t_2} l(t) dt,$$

which, on account of (2.8), contradicts (2.4).  $\square$

**Definition 2.1.** Suppose that  $r$  and  $r_1$  are positive constants. A vector-function  $x : [\alpha, \beta] \rightarrow \mathbb{R}^n$  is said to belong to the set  $W_n([\alpha, \beta], r, r_1)$  if (1.8) implies the estimate (1.9) for arbitrary  $t_1 \in [\alpha, \beta]$  and  $t_2 \in ]t_1, \beta]$ .

**Lemma 2.4<sub>1</sub>.** Suppose that  $r$  and  $r_1$  are positive constants,  $l \in L([t_1, t_2]; \mathbb{R}_+)$ ,  $\omega \in C(\mathbb{R}_+; ]0, +\infty[)$ , and (1.13) holds. Then there exists a positive constant  $r'$  such that if  $\delta \in ]0, \frac{b-a}{4}[$  and an arbitrary  $x \in \tilde{C}^1([a, b]; \mathbb{R}^n) \cap W_n([a+\delta, b-\delta], r, r_1)$  satisfies the boundary conditions (1.2), the inequality

$$\|x''(t)\| \leq l(t) \quad \text{for } t \in ]a, a+\delta[ \cup ]b-\delta, b[, \quad (2.9)$$

and furthermore the inequality

$$x''(t) \cdot \eta_r(x'(t)) \leq \omega(\|x'(t)\|)(l(t) + \|x'(t)\|) \quad (2.10)$$

on the set  $[a+\delta, b-\delta]$ , then  $x$  admits an estimate

$$\|x'(t)\| \leq r' \quad \text{for } a \leq t \leq b. \quad (2.11)$$

*Proof.* Due to Definition 2.1, without loss of generality we may assume that  $r(b-a) > 2r_1$ . In view of (1.13) there exist positive numbers  $r^*$  and  $r'$  such that

$$\int_r^{r^*} \frac{ds}{\omega(s)} > r_1 + \int_a^b l(t) dt \quad (2.12)$$

and

$$\int_\mu^{r'} \frac{ds}{\omega(s)} > r_1 + \int_a^b l(t) dt \quad (2.13)$$

where

$$\mu = r^* + \int_a^b l(t) dt.$$

Let us show that  $r'$  is a suitable constant.

First note that for a certain  $t_0 \in [\frac{3a+b}{4}, \frac{a+3b}{4}]$  we have

$$\|x'(t_0)\| \leq r. \quad (2.14)$$

Indeed, assuming the contrary implies that

$$\|x'(t)\| > r \quad \text{for } \frac{3a+b}{4} \leq t \leq \frac{a+3b}{4}.$$

Therefore, taking into account the inequality  $r(b-a) \geq 2r_1$ , we obtain

$$\int_{\frac{3a+b}{4}}^{\frac{a+3b}{4}} \|x'(t)\| dt > r_1.$$

But this is impossible, since  $x \in W_n([a+\delta, b-\delta], r, r_1)$ . Thus (2.14) is proved.

Now let us show that

$$\|x'(t)\| \leq r^* \quad \text{for } t_0 \leq t \leq b-\delta. \quad (2.15)$$

Suppose to the contrary that for arbitrary  $t_2 \in ]t_0, b-\delta]$  we have

$$\|x'(t_2)\| > r^*. \quad (2.16)$$

Then by (2.14) there exists  $t_1 \in [t_0, t_2[$  such that

$$\|x'(t_1)\| = r \quad \text{and} \quad \|x'(t)\| > r \quad \text{for } t_1 < t \leq t_2.$$

Hence, taking into account  $x \in W_n([a+\delta, b-\delta], r, r_1)$ , we get the estimate (1.9). Assuming  $i = 1$ ,  $\rho = r$ , and  $\rho' = r^*$ , it is easy to verify that  $x$  satisfies the other conditions of Lemma 2.4<sub>1</sub> too. Therefore  $x$  admits the estimate (2.7), which contradicts (2.16). Thus (2.15) is proved. In view of (1.2) and (2.9) it implies that

$$\|x'(t)\| \leq \mu \quad \text{for } t \in [a, a+\delta] \cup [b-\delta, b]. \quad (2.17)$$

In particular,  $\|x'(a+\delta)\| \leq \mu$ . Applying Lemma 2.3 where  $\rho = \mu$ ,  $\rho' = r'$  and  $i = 1$ , an argument similar to the one carried out above yields the estimate

$$\|x'(t)\| \leq r' \quad \text{for } a+\delta \leq t \leq t_0. \quad (2.18)$$

Finally, from (2.15), (2.17), and (2.18) we obtain the estimate (2.11).  $\square$

In a similar manner we can prove

**Lemma 2.4<sub>2</sub>.** *Suppose that the conditions of Lemma 2.4<sub>1</sub> are satisfied, except that the inequality (2.10) is replaced by*

$$x''(t) \cdot \eta_r(x'(t)) \geq -\omega(\|x'(t)\|)(l(t) + \|x'(t)\|). \quad (2.19)$$

*Then  $x$  admits the estimate (2.11).*

**Lemma 2.5.** *Suppose that  $r$  and  $r_1$  are positive constants,  $a \leq a_0 < b_0 \leq b$ ,  $l \in L([t_1, t_2]; \mathbb{R}_+)$ ,  $\omega \in C(\mathbb{R}_+; ]0, +\infty[)$ , and (1.13) holds. Then there exists a positive constant  $r'$  such that if  $0 < \delta < \min\{a_0 - a, b - b_0\}$  and an arbitrary  $x \in \tilde{C}^1([a, b]; \mathbb{R}^n) \cap W_n([a + \delta, b - \delta], r, r_1)$  satisfies the conditions (2.9), the inequality (2.10) on the set  $]a_0, b - \delta[$ , and, furthermore, the inequality (2.19) on the set  $]a + \delta, b_0[$ , then  $x$  admits the estimate (2.11).*

The proof of Lemma 2.5 is similar to the one carried out above for Lemma 2.4<sub>1</sub>, so we shall note only the main points.

Due to Definition 2.1, without loss of generality we may assume that  $r(b_0 - a_0) \geq r_1$ . In view of (1.13) there exists a positive number  $r^*$  such that (2.12) holds. The constant

$$r' = r^* + \int_a^b l(t) dt$$

is just the suitable one.

First, taking into account the conditions  $x \in W_n([a + \delta, b - \delta], r, r_1)$ , we conclude that for a certain  $t_0 \in [a_0, b_0]$  (2.14) is satisfied. Further, applying Lemma 2.3 ( $i = 1$ ) and the inequality (2.10), we get the estimate (2.15). Finally, from (2.9) and (2.15) it follows that we have  $\|x'(t)\| \leq r'$  on the set  $[b - \delta, b]$ . Thus the last estimate holds on  $[t_0, b]$ . Analogously, applying Lemma 2.3 ( $i = 2$ ) and the inequality (2.19), it can be proved on  $[a, t_0]$ .

In a similar manner we can prove

**Lemma 2.6.** *Suppose that  $r$  and  $r_1$  are positive constants,  $a < a_1 < a_2 < t_0 < b_2 < b_1 < b$ ,  $l \in L([t_1, t_2]; \mathbb{R}_+)$ ,  $\omega \in C(\mathbb{R}_+; ]0, +\infty[)$ , and (1.13) holds. Then there exists a positive constant  $r'$  such that if  $0 < \delta < \min\{a_1 - a, b - b_1\}$  and an arbitrary  $x \in \tilde{C}^1([a, b]; \mathbb{R}^n) \cap W_n([a + \delta, b - \delta], r, r_1)$  satisfies the conditions (2.9), the inequality (2.10) on the set  $]a_1, t_0[ \cup ]b_2, b[$ , and furthermore the inequality (2.19) on the set  $]a, a_2[ \cup ]t_0, b_1[$ , then  $x$  admits the estimate (2.11).*

### § 3. PROOF OF THE MAIN RESULTS

*Proof of Theorem 1.1<sub>1</sub>.* Without loss of generality we can assume that

$$l(t) \geq \|z''(t)\| + |\varphi''(t)| \quad \text{for } a < t < b.$$

Put

$$a_k = a + \frac{b - a}{4k}, \quad b_k = b - \frac{b - a}{4k}$$

and

$$\chi_\rho(s) = \begin{cases} 0 & \text{for } s \leq \rho, \\ \frac{2\rho-s}{\rho} & \text{for } \rho < s < 2\rho, \\ 0 & \text{for } 2\rho \leq s. \end{cases}$$

By Definition 1.1 there exists a sequence  $(\rho_k)_{k=1}^{+\infty}$  such that  $\lim_{k \rightarrow +\infty} \rho_k = +\infty$  and for each  $k \in \{1, 2, \dots\}$

$$(x - z(t)) \cdot (\chi_{\rho_k}(\|y\|)f(t, x, y) - z''(t)) + \|y - z'(t)\|^2 \geq \varphi(t)\varphi''(t)$$

for  $a_k < t < b_k$ ,  $\|x - z(t)\| = \varphi(t)$  and  $(x - z(t)) \cdot (y - z'(t)) = \varphi(t)\varphi'(t)$ .

Put

$$h(t, x, y) = \begin{cases} z''(t) + \frac{|\varphi''(t)|}{\varphi(t)}(x - z(t)) & \text{for } \|x - z(t)\| \leq \varphi(t), \\ z''(t) + \frac{|\varphi''(t)|}{\varphi(t)}(x - z(t)) & \text{for } \|x - z(t)\| > \varphi(t) \end{cases}$$

and

$$f_k(t, x, y) = \begin{cases} h(t, x, y) & \text{for } t \notin [a_k, b_k], \\ \chi_{\rho_k}(\|y\|)f(t, x, y) & \text{for } t \in [a_k, b_k] \end{cases}$$

( $k = 1, 2, \dots$ ). It is easy to verify that for each  $k \in \{1, 2, \dots\}$  the vector-function  $g(t, x, y) = f_k(t, x, y)$  satisfies all the conditions of Lemma 2.2. Therefore the differential system

$$x'' = f_k(t, x, x') \quad (3.1_k)$$

has at least one solution  $x_k \in \tilde{C}^1([a, b]; \mathbb{R}^n)$  satisfying the boundary conditions (1.2) and the estimate

$$\|x_k(t) - z(t)\| \leq \varphi(t) \quad \text{for } a \leq t \leq b \quad (k = 1, 2, \dots). \quad (3.2_k)$$

Choose the positive constants  $r$  and  $r_1$  according to Definition 1.3 and the constant  $r'$  according to Lemma 2.4<sub>1</sub>, assuming without loss of generality that  $r \geq \rho$ . Then by Lemma 2.4<sub>1</sub> we obtain

$$\|x'_k(t)\| \leq r' \quad \text{for } a \leq t \leq b \quad (k = 1, 2, \dots). \quad (3.3_k)$$

In view of (3.2<sub>k</sub>) and (3.3<sub>k</sub>) the sequences  $(x_k)_{k=1}^{+\infty}$  and  $(x'_k)_{k=1}^{+\infty}$  are uniformly bounded and equicontinuous on  $[a, b]$ . So due to the well-known Arzela–Ascoli theorem there exists a sequence  $(k_j)_{j=1}^{+\infty}$  such that  $(x_{k_j})_{j=1}^{+\infty}$  and  $(x'_{k_j})_{j=1}^{+\infty}$  uniformly converge on  $[a, b]$ . Put

$$x(t) = \lim_{j \rightarrow \infty} x_{k_j}(t) \quad \text{for } a \leq t \leq b.$$

Due to the definition of the functions  $f_k$  ( $k = 1, 2, \dots$ )  $x$  belongs to the set  $\tilde{C}^1([a, b]; \mathbb{R}^n)$  and is a solution of (1.1), (1.2).  $\square$

The *proof of Theorems 1.1<sub>2</sub>, 1.2, and 1.3* is similar to the one carried out above for Theorem 1.1<sub>1</sub>. The only difference is that one has to apply Lemmas 2.4<sub>2</sub>, 2.5, and 2.6 respectively instead of Lemma 2.4<sub>1</sub>.

*Remark 3.1.* Theorem 1.1<sub>1</sub> (Theorem 1.1<sub>2</sub>) can be strengthened by a slight complication of Lemma 2.4<sub>1</sub> (Lemma 2.4<sub>2</sub>). Namely, we can assume that the vector-function  $f$  has the property  $V([\alpha, \beta], \varphi, z)$  for each segment  $[\alpha, \beta]$  contained in the interval  $]a, b[$  (in the interval  $[a, b]$ ). In that case the vector-function  $f$  may fail to have the property  $V([a, b], \varphi, z)$ .

*Proof of Theorem 1.4.* Assume to the contrary that  $x_i \in \tilde{C}^1([a, b]; \mathbb{R}^n)$  ( $i = 1, 2$ ) are solutions of the boundary-value problem (1.1), (1.2) and  $x_1(t) \not\equiv x_2(t)$ . Put

$$\begin{aligned} x(t) &= x_1(t) - x_2(t) \quad \text{for } a \leq t \leq b, \\ u(t) &= \|x(t)\| \end{aligned}$$

and

$$r = \max\left\{\sum_{i=1}^2 \|x_i(t)\| + \|x'_i(t)\| : a \leq t \leq b\right\}.$$

Choose the functions  $l_i(t, r)$  ( $i = 1, 2$ ) according to the condition of Theorem 1.4. First prove that  $u'(t) \not\equiv 0$ . Indeed, assuming the contrary we have

$$\begin{aligned} 0 = u''(t) &= \frac{x''(t) \cdot x(t) + \|x'(t)\|^2}{\|x(t)\|} - \frac{(x(t) \cdot x'(t))^2}{\|x(t)\|^3} \geq \\ &\geq \frac{x''(t) \cdot x(t)}{\|x(t)\|} \quad \text{for } a < t < b. \end{aligned}$$

Hence by (1.16) we obtain

$$l_1(t, r) \leq 0 \quad \text{for } a < t < b.$$

But this is impossible, since  $l_1(t, r)$  is nonnegative and differs from zero on a subset of positive measure of the interval  $]a, b[$ . Thus  $u'(t) \not\equiv 0$ . Therefore, there exists  $t_0 \in ]a, b[$  such that either

$$u(t_0) > 0, \quad u'(t_0) > 0 \tag{3.4}$$

or

$$u(t_0) > 0, \quad u'(t_0) < 0. \tag{3.5}$$

Without loss of generality assume that (3.4) holds. Then on  $[t_0, b]$

$$u(t) > 0, \quad u'(t) > 0. \tag{3.6}$$

Indeed, if this is not so, then there exists  $t_1 \in ]t_0, b]$  such that  $u'(t_1) = 0$  and (3.6) holds on  $[t_0, t_1[$ . Applying (1.16) once more we obtain

$$\begin{aligned} u''(t) &\geq \frac{x''(t) \cdot x(t)}{\|x(t)\|} \geq l_1(t, r)u(t) - l_2(t, r)|u'(t)| \geq \\ &\geq -l_2(t, r)|u'(t)| \quad \text{for } t_0 < t < t_1. \end{aligned}$$

According to the Gronwall–Bellman lemma (see e.g. [6]) the last inequality yields

$$u'(t_1) \geq u'(t_0) \exp \left[ - \int_{t_0}^{t_1} l_2(t, r) dt \right] > 0.$$

The obtained contradiction shows that (3.6) holds on  $[t_0, b]$ . Hence, taking into account the equalities

$$u(a) = u(b), \quad u'(a) = u'(b) \tag{3.7}$$

which follow from the boundary conditions (1.2), we get

$$u(a) > 0, \quad u'(a) > 0.$$

Repeating the argument that was carried out above we can show the validity of (3.6) on  $[a, b]$ , but this contradicts (3.7).  $\square$

#### REFERENCES

1. I. T. Kiguradze, Some singular boundary-value problems for ordinary differential equations. (Russian) *Tbilisi University Press, Tbilisi*, 1975.
2. L. H. Erbe and P. K. Palamides, Boundary-value problems for second-order differential equations. *J. Math. Anal. Appl.* **127**(1987), 80-92.
3. G. D. Gavrindashvili, On Solvability of the Dirichlet boundary-value problem for systems of ordinary nonlinear differential equations with singularities. (Russian) *Differentsial'nye Uravneniya* **27**(1991), No. 9, 1521-1525.
4. G. D. Gavrindashvili, On certain boundary-value problems for systems of second-order nonlinear ordinary differential equations. (Russian) *Proc. I. Vekua Inst. Appl. Math. Tbilis. St. Univ.* **31**(1988), 23-52.
5. M. Nagumo, Über die Differentialgleichung  $y'' = f(x, y, y')$ . *Proc. Phys.-Math. Soc. Japan* **19**(1937), 861-866.

6. Ph. Hartman, Ordinary differential equations. *John Wiley & Sons, New York*, 1964.

(Received 14.09.1993)

Author's address:  
Dept. of Applied Mathematics  
Georgian Technical University  
77, Kostava St., Tbilisi 380075  
Republic of Georgia