

Research Article

Applications of an HIDS Theorem to the Existence of Fixed Point, Abstract Equilibria and Optimization Problems

Wei-Shih Du

Department of Mathematics, National Kaohsiung Normal University, Kaohsiung 824, Taiwan

Correspondence should be addressed to Wei-Shih Du, wudu@nknuc.nknu.edu.tw

Received 16 June 2011; Accepted 18 July 2011

Academic Editor: Josip E. Pečarić

Copyright © 2011 Wei-Shih Du. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

By applying hybrid inclusion and disclusion systems (HIDS), we establish several vectorial variants of system of Ekeland's variational principle on topological vector spaces, some existence theorems of system of parametric vectorial quasi-equilibrium problem, and an existence theorem of system of the Stampacchia-type vectorial equilibrium problem. As an application, a vectorial minimization theorem is also given. Moreover, we discuss some equivalence relations between our vectorial variant of Ekeland's variational principle, common fixed point theorem, and maximal element theorem.

1. Introduction

Let X be a nonempty subset of a topological space (t.s., for short), and let $f : X \times X \rightarrow \mathbb{R}$ be a function with $f(x, x) \geq 0$ for all $x \in X$. Then the scalar equilibrium problem in the sense of Blum and Oettli [1] is to find $\bar{x} \in X$ such that $f(\bar{x}, y) \geq 0$ for all $y \in X$. The equilibrium problem was extensively investigated and generalized to the vectorial equilibrium problems for single-valued or multivalued maps and contains optimization problems, variational inequalities problems, saddle point problems, the Nash equilibrium problems, fixed point problems, complementary problems, bilevel problems, and semi-infinite problems as special cases and have some applications in mathematical program with equilibrium constraint; for detail one can refer to [1–4] and references therein.

The famous Ekeland's variational principle (EVP, for short) [5–7] is a forceful tool in various fields of applied mathematical analysis and nonlinear analysis. A number of generalizations in various different directions of these results for functions defined in metric (or quasimetric) spaces and more general in topological vector spaces have been investigated by several authors in the past; see [8–23] and references therein. It is wellknown that the original EVP is equivalent to Caristi's fixed point theorem, to Takahashi's nonconvex

minimization theorem; and to the flower petal theorem for detail, see [14, 16–18, 20] and references therein. EVPs were extended to the vector case by using scalarization method and were applied to the study of efficiency (or approximative efficiency) and others; see, for example, [3, 8, 9, 12, 13, 15, 23].

Recently, the author first studied the following mathematical model about hybrid inclusion and disclusion systems (HIDS, for short) [11]. Let I be any index set. For each $i \in I$, let Y_i be a nonempty closed convex subset of a Hausdorff topological vector space (t.v.s., for short) Y , $H_i \subseteq Y_i$, $Y = \prod_{i \in I} Y_i$, $A_i : Y \multimap Y_i$, and let $T_i : Y \multimap Y_i$ be multivalued maps. Hybrid inclusion and disclusion systems (HIDS) are defined as follows:

$$\begin{aligned} \text{(HIDS)} \quad & \text{Find } v = (v)_{i \in I} \in Y \quad \text{such that } v_i \in H_i, \\ & y_i \notin A_i(v), \quad \forall y_i \in T_i(v), \quad \forall i \in I. \end{aligned} \tag{1.1}$$

In fact, HIDS contains several important problems as special cases. Let X be a nonempty subset of a topological space E , and let $u \in X$ be given. For each $i \in I$, let U_i and Z_i be real t.v.s. with zero vector θ_{U_i} and θ_{Z_i} , respectively.

Example 1.1. For each $i \in I$, let $F_i : X \times Y_i \rightarrow \mathbb{R}$ and $G_i : Y \times Y_i \rightarrow \mathbb{R}$ be functions. If H_i and A_i are defined as follows:

$$\begin{aligned} H_i &= \{y_i \in Y_i : F_i(u, y_i) \leq 0\}, \\ A_i(y) &= \{z_i \in Y_i : G_i(y, z_i) \leq 0\}, \end{aligned} \tag{1.2}$$

then HIDS will reduce to the following system of hybrid scalar equilibrium problem (P_1):

$$(P_1) \quad \text{Find } v = (v)_{i \in I} \in Y \text{ such that } F_i(u, v_i) \leq 0 \text{ and } G_i(v, y_i) > 0 \text{ for all } y_i \in T_i(v) \text{ and for all } i \in I.$$

Example 1.2. For each $i \in I$, let $F_i : X \times Y_i \multimap U_i$ and $G_i : Y \times Y_i \multimap Z_i$ be multivalued maps with nonempty values, and let C_i and D_i be nonempty subsets of U_i and Z_i , respectively. If H_i and A_i are defined as follows:

$$\begin{aligned} H_i &= \{y_i \in Y_i : F_i(u, y_i) \cap (-C_i \setminus \{\theta_{U_i}\}) = \emptyset\}, \\ A_i(y) &= \{z_i \in Y_i : G_i(y, z_i) \cap (-D_i \setminus \{\theta_{Z_i}\}) = \emptyset\}, \end{aligned} \tag{1.3}$$

then HIDS will reduce to the following problem (P_2), which is an abstract equilibrium problem:

$$(P_2) \quad \text{Find } v = (v)_{i \in I} \in Y \text{ such that } F_i(u, v_i) \cap (-C_i \setminus \{\theta_{U_i}\}) = \emptyset \text{ and } G_i(v, y_i) \cap (-D_i \setminus \{\theta_{Z_i}\}) \neq \emptyset \text{ for all } y_i \in T_i(v) \text{ and for all } i \in I.$$

Example 1.3. For each $i \in I$, let $F_i : Y_i \multimap U_i$ and $G_i : Y \times Y_i \multimap Z_i$ be multivalued maps. If H_i and A_i are defined as follows:

$$\begin{aligned} H_i &= \{y_i \in Y_i : y_i \in F_i(y_i)\}, \\ A_i(y) &= \{z_i \in Y_i : y \notin G_i(y, z_i)\}, \end{aligned} \tag{1.4}$$

then HIDS will reduce to the following fixed point problem (P_3) :

(P_3) Find $v = (v)_{i \in I} \in Y$ such that $v_i \in F_i(v_i)$ and $v \in G_i(v, y_i)$ for all $y_i \in T_i(v)$ for all $i \in I$.

Example 1.4. For each $i \in I$, let $F_i : X \times Y_i \multimap U_i$ and $G_i : Y \times Y_i \multimap Z_i$ be multivalued maps with nonempty values. If H_i and A_i are defined as follows:

$$\begin{aligned} H_i &= \{y_i \in Y_i : \theta_{U_i} \notin F_i(u, y_i)\}, \\ A_i(y) &= \{z_i \in Y_i : \theta_{Z_i} \notin G_i(y, z_i)\}, \end{aligned} \tag{1.5}$$

then HIDS will reduce to the following system of mixed type of parametric variational inclusion and disclusion problem (P_4) :

(P_4) Find $v = (v)_{i \in I} \in Y$ such that $\theta_{U_i} \notin F_i(u, v_i)$ and $\theta_{Z_i} \in G_i(v, y_i)$ for all $y_i \in T_i(v)$ and for all $i \in I$.

In this paper, we study the existence theorems of system of parametric vectorial quasi-equilibrium problems, vectorial variants of system of Ekeland's variational principle (VSEVP, for short), and the existence theorems of system of the Stampacchia-type vectorial equilibrium problem by using an HIDS theorem established by the author [11]. Our results improve and generalize some theorems in [19] to the vector case. Till now, to my knowledge, there are extremely few results about Stampacchia-type vectorial equilibrium problem in the literature. The existence of VSEVP and the Stampacchia-type vectorial equilibrium problem are established by applying the HIDS theorem without the use of any scalarization method. So our results are completely different from [3, 8, 9, 12, 13, 15, 23]. As an application, a vectorial minimization theorem is also proved. Moreover, we prove some equivalence relations between our VSEVP, common fixed point theorem, and maximal element theorem.

2. Preliminaries

Let A and B be nonempty sets. A multivalued map $T : A \multimap B$ is a function from A to the power set 2^B of B . We denote $T(A) = \bigcup\{T(x) : x \in A\}$ and let $T^- : B \multimap A$ be defined by the condition that $x \in T^-(y)$ if and only if $y \in T(x)$. We recall that a point $x \in A$ is a maximal element of $T : A \rightarrow 2^B$ if $T(x) = \emptyset$. Let X be a linear space with zero vector θ . A nonempty subset C of X is called a *convex cone* if $C + C \subseteq C$ and $\lambda C \subseteq C$ for all $\lambda \geq 0$. A cone C in X is pointed if $C \cap (-C) = \{\theta\}$. Let Z be a real t.v.s., and let D be a proper convex cone in Z , and $A \subseteq Z$. A point $\bar{y} \in A$ is called a vectorial minimal point of A with respect to D if for any $y \in A$, $y - \bar{y} \notin -D \setminus \{\theta\}$. The set of vectorial minimal point of A is denoted by $\text{Min}_D A$. The convex hull of A and the closure of A are denoted by $\text{co } A$ and $\text{cl } A$, respectively.

Definition 2.1. Let X and Y be linear spaces, and let C be a proper convex cone in Y . A map $f : X \multimap Y$ is called *C-convex* if for any $x_1, x_2 \in X$ and $\lambda \in [0, 1]$, one has

$$\lambda f(x_1) + (1 - \lambda)f(x_2) - f(\lambda x_1 + (1 - \lambda)x_2) \subseteq C. \tag{2.1}$$

Clearly, if f_1 and f_2 are C -convex and $\alpha \geq 0$, then αf_1 and $f_1 + f_2$ are C -convex.

Definition 2.2. Let X be a nonempty convex subset of a vector space E , let Y be a nonempty convex subset of a vector space V , and let Z be a real t.v.s. Let $F : X \times Y \rightrightarrows Z$ and $C : X \rightrightarrows Z$ be multivalued maps such that for each $x \in X$, $C(x)$ is a nonempty closed convex cone. For each fixed $x \in X$, $y \rightrightarrows F(x, y)$ is called $C(x)$ -quasiconvex if for any $y_1, y_2 \in Y$ and $\lambda \in [0, 1]$, one has either

$$F(x, y_1) \subseteq F(x, \lambda y_1 + (1 - \lambda)y_2) + C(x), \quad (2.2)$$

or

$$F(x, y_2) \subseteq F(x, \lambda y_1 + (1 - \lambda)y_2) + C(x). \quad (2.3)$$

Now, we define the concept of vectorial upper and lower semicontinuous on t.v.s.

Definition 2.3. Let X be a t.s., let Y be a t.v.s. with zero vector θ , and let $C \neq \{\theta\}$ be a pointed convex cone in Y . A map $f : X \rightarrow Y$ is said to be

- (i) *vectorial lower semicontinuous with respect to C* (C-v.l.s.c., for short) at $x_0 \in X$ if for any $a \in C \setminus \{\theta\}$, there exists an open neighborhood $N(x_0)$ of x_0 such that $f(y) - f(x_0) + a \in C \setminus \{\theta\}$ for all $y \in N(x_0)$,
- (ii) *vectorial upper semicontinuous with respect to C* (C-v.u.s.c., for short) at $x_0 \in X$ if for any $a \in C \setminus \{\theta\}$, there exists an open neighborhood $N(x_0)$ of x_0 such that $f(x_0) - f(y) + a \in C \setminus \{\theta\}$ for all $y \in N(x_0)$.

The function f is called C-v.l.s.c. (resp., C-v.u.s.c.) on X if f is C-v.l.s.c. (resp., C-v.u.s.c.) at every point of X .

Proposition 2.4. *Let X be a t.s., let Y be a t.v.s with zero vector θ , and let $C \neq \{\theta\}$ be a pointed convex cone in Y . Let $f, g : X \rightarrow Y$ be maps and $\gamma \geq 0$. Then the following hold:*

- (a) f is C-v.u.s.c. (resp., C-v.l.s.c.) on $X \iff -f$ is C-v.l.s.c. (resp., C-v.u.s.c.) on X ;
- (b) if f and g are C-v.u.s.c. (resp., C-v.l.s.c.) on X , then γf and $f + g$ are C-v.u.s.c. (resp., C-v.l.s.c.) on X ;
- (c) if f is C-v.u.s.c. on X , then
 - (i) $\{x \in X : \lambda - f(x) \in C \setminus \{\theta\}\}$ is open in X for all $\lambda \in Y$,
 - (ii) $\{x \in X : \lambda - f(x) \notin C \setminus \{\theta\}\}$ is closed in X for all $\lambda \in Y$;
- (d) if f is C-v.l.s.c. on X , then
 - (iii) $\{x \in X : f(x) - \lambda \in C \setminus \{\theta\}\}$ is open in X for all $\lambda \in Y$,
 - (iv) $\{x \in X : f(x) - \lambda \notin C \setminus \{\theta\}\}$ is closed in X for all $\lambda \in Y$.

Proof. Clearly, (a) and (b) hold from definition. To prove (c), it suffices to show (i). Suppose that f is C-v.u.s.c. on X . Let $\lambda \in Y$ and $x_0 \in \{x \in X : \lambda - f(x) \in C \setminus \{\theta\}\}$. Then $\alpha := \lambda - f(x_0) \in C \setminus \{\theta\}$. Since f is C-v.u.s.c. at x_0 , there exists an open neighborhood $N(x_0)$ of x_0 such that

$$\lambda - f(y) = f(x_0) - f(y) + \alpha \in C \setminus \{\theta\} \quad (2.4)$$

for all $y \in N(x_0)$. Hence $\{x \in X : \lambda - f(x) \in C \setminus \{\theta\}\}$ is an open set in X and (i) is proved. Obviously, (ii) is immediate from (i). It is easy to see that conclusion (d) follows from (a) and (c). \square

Remark 2.5. Let X be a t.v.s. and $\alpha \in X$ with $\alpha \neq 0_X$, where 0_X is the origin of X . Let $f : X \rightarrow L_\alpha = \{\gamma\alpha : \gamma \in \mathbb{R}\}$ be a map. Hence in conclusion (c) (resp., (d)) of Proposition 2.4, we have

$$\begin{aligned} f \text{ is } C\text{-v.u.s.c. on } X &\iff (i) \iff (ii) \\ (\text{resp., } f \text{ is } C\text{-v.l.s.c. on } X &\iff (iii) \iff (iv)). \end{aligned} \quad (2.5)$$

In particular, if $X = \mathbb{R}$ (the set of real numbers) and $C = [0, \infty)$, then the C -v.l.s.c. (resp. C -v.u.s.c.) function $f : X \rightarrow \mathbb{R}$ is l.s.c. (resp. u.s.c.) in usual.

The concept of C -vectorial ℓ -function and C -vectorial quasi-distance on topological spaces are introduced as follows.

Definition 2.6. Let X and E be t.v.s., let θ be the zero vector of E , and let $C \neq \{\theta\}$, a pointed convex cone in E . A map $p : X \times X \rightarrow E$ is called

(a) a C -vectorial ℓ -function (ℓ_C -function, for short) if the following are satisfied:

- (VL1) $p(x, x) \in C$ for all $x \in X$;
- (VL2) for any $x \in X$, $p(x, \cdot)$ is C -convex;
- (VL3) for any $y \in X$, $p(\cdot, y)$ is C -v.u.s.c.

(b) a C -vectorial quasi-distance if the following are satisfied:

- (VQD1) $p(x, x) \in C$ for all $x \in X$;
- (VQD2) $p(x, y) + p(y, z) \in p(x, z) + C$ for any $x, y, z \in X$;
- (VQD3) for any $x \in X$, $p(x, \cdot)$ is C -convex and C -v.l.s.c.;
- (VQD4) for any $y \in X$, $p(\cdot, y)$ is C -v.u.s.c.

If $E = (-\infty, \infty]$ and letting $C = [0, \infty)$ be in (a) and (b), then the function $p : X \times X \rightarrow (-\infty, \infty]$ is called a ℓ -function and quasi-distance, respectively, introduced by Lin and Du [19]. For examples and results of ℓ -function and quasi-distance, one can see [19].

Remark 2.7. (a) Obviously, a C -vectorial quasi-distance is a ℓ_C -function, but the reverse is not true;

(b) if p_1 and p_2 are C -vectorial quasi-distances (resp., ℓ_C -functions) and $\alpha \geq 0$, then αp_1 and $p_1 + p_2$ are C -vectorial quasi-distances (resp., ℓ_C -functions);

(c) if $f : X \rightarrow E$ is a C -v.l.s.c. and C -convex function, then the function $p : X \times X \rightarrow E$ defined by $p(x, y) = f(y) - f(x)$ is a C -vectorial quasi-distance.

Lemma 2.8 (see [24, 25]). *Let X and Y be the Hausdorff topological spaces, and let $T : X \rightarrow Y$ be a multivalued map. Then T is l.s.c. at $x \in X$ if and only if for any $y \in T(x)$ and for any net $\{x_\alpha\}$ in X converging to x , there exists a subnet $\{x_{\phi(\lambda)}\}_{\lambda \in \Lambda}$ of $\{x_\alpha\}$ and a net $\{y_\lambda\}_{\lambda \in \Lambda}$ with $y_\lambda \rightarrow y$ such that $y_\lambda \in T(x_{\phi(\lambda)})$ for all $\lambda \in \Lambda$.*

3. The Existence of System of VSEVP and Abstract Equilibrium Problems

The following existence theorem for the solution of HIDS was established in [11].

Theorem 3.1 (HIDS theorem [11]). *Let I be any index set. For each $i \in I$, let Y_i be a nonempty closed convex subset of a Hausdorff t.v.s. V_i . Let X be a nonempty subset of a topological space E , $Y = \prod_{i \in I} Y_i$ and $u \in X$. For each $i \in I$, let H_i be a nonempty closed subset of Y_i , let $A_i : Y \multimap Y_i$ be a multivalued map, and let $T_i : Y \multimap Y_i$ be a multivalued map with nonempty values. For each $i \in I$, suppose that the following conditions are satisfied:*

- (i) for each $y = (y_i)_{i \in I} \in Y$, $y_i \notin A_i(y)$;
- (ii) for each $y \in Y$, $\text{co } T_i(y) \subseteq H_i$ and $A_i(y)$ is convex;
- (iii) for each $z_i \in Y_i$, $T_i^-(z_i)$ and $A_i^-(z_i)$ are open in Y ;
- (iv) there exist a nonempty compact subset K of Y and a nonempty compact convex subset M_i of Y_i for each $i \in I$ such that for each $y \in Y \setminus K$ there exist $j \in I$ and $z_j \in M_j \cap T_j(y) \cap A_j(y)$.

Then there exists $v = (v_i)_{i \in I} \in Y$ such that for each $i \in I$, $v_i \in H_i$ and $y_i \notin A_i(v)$ for all $y_i \in T_i(v)$.

Example 3.2. Let X and Y be Hausdorff t.v.s., let U be a real t.v.s. with its zero vector θ , and $u \in X$.

(a) Let $F : X \times Y \multimap U$ be a multivalued map with nonempty values such that there exists $w = w(u) \in Y$ such that $\theta \in F(u, w)$ and the map $y \multimap F(u, y)$ is closed. Then it is easy to see that $H = \{y \in Y : \theta \in F(u, y)\}$ is a nonempty closed subset of Y .

(b) Let $G : Y \times Y \multimap U$ be a multivalued map with nonempty values and W be a nonempty open set in U . Suppose that

- (i) for each $y \in Y$, $\theta \notin G(y, y) + W$,
- (ii) for each $y \in Y$, $G(y, \cdot)$ is $\{\theta\}$ -quasiconvex and for each $z \in Y$, $G(\cdot, z)$ is l.s.c.

Let $A : Y \multimap Y$ be defined by

$$A(y) = \{z \in Y : \theta \in G(y, z) + W\}. \quad (3.1)$$

Then $y \notin A(y)$ for each $y \in Y$. We claim that $A^-(z)$ is open in Y for each $z \in Y$. Indeed, let $z \in Y$ be given, and let $y \in \text{cl}(Y \setminus A^-(z))$. Then there exists a net $\{y_\alpha\}_{\alpha \in \Lambda}$ in $Y \setminus A^-(z)$ such that $y_\alpha \rightarrow y$. Thus we have $\theta \notin G(y_\alpha, z) + W$ or $G(y_\alpha, z) \subseteq U \setminus W$. Clearly, $y \in Y$. For any $w \in G(y, z)$, since $G(\cdot, z)$ is l.s.c. at y and $y_\alpha \rightarrow y$, by Lemma 2.8, there exists a net $\{w_\alpha\}$ with $w_\alpha \rightarrow w$ such that $w_\alpha \in G(y_\alpha, z) \subseteq U \setminus W$. Since $U \setminus W$ is closed, $w \in U \setminus W$. So $G(y, z) \subseteq U \setminus W$. It implies $\text{cl}(Y \setminus A^-(z)) = Y \setminus A^-(z)$, and hence $A^-(z)$ is open in Y . Next, we show that for each $y \in Y$, $A(y)$ is convex. Let $a, b \in A(y)$. Then $\theta \in G(y, a) + W$ and $\theta \in G(y, b) + W$. For any $\lambda \in [0, 1]$, let $e_\lambda := \lambda a + (1 - \lambda)b \in Y$. Suppose to the contrary that there exists $\lambda_0 \in (0, 1)$ such that $\theta \notin G(y, e_{\lambda_0}) + W$. By the $\{\theta\}$ -quasiconvexity of $G(y, \cdot)$, either

$$\theta \in G(y, a) + W \subseteq G(y, e_{\lambda_0}) + W \quad (3.2)$$

or

$$\theta \in G(y, b) + W \subseteq G(y, e_{\lambda_0}) + W, \quad (3.3)$$

which leads to a contradiction. Hence for each $y \in Y$, $A(y)$ is convex.

Applying Theorem 3.1, we establish the following existence theorem of system of parametric vectorial quasi-equilibrium problem.

Theorem 3.3. *Let I be any index set. For each $i \in I$, let X_i be a nonempty Hausdorff t.v.s., and let $C_i \neq \{\theta\}$ be a pointed convex cone in a t.v.s. E with zero vector θ . Let $X = \prod_{i \in I} X_i$. For each $i \in I$, let $p_i, q_i : X_i \times X_i \rightarrow E$ be C_i -vectorial quasi-distances and let $T_i : X \multimap X_i$ be a multivalued map with nonempty values. Let $u = (u_i)_{i \in I} \in X$ with $p_i(u_i, u_i) = q_i(u_i, u_i) = \theta$ for all $i \in I$. For each $i \in I$, suppose that the following conditions are satisfied:*

- (i) *for each $y \in X$, $\text{co}T_i(y) \subseteq \{x_i \in X_i : q_i(u_i, x_i) \notin C_i \setminus \{\theta\}\}$ and $T_i^-(z_i)$ is open for all $z_i \in X_i$;*
- (ii) *there exist a nonempty compact subset K of X and a nonempty compact convex subset M_i of X_i for each $i \in I$ such that for each $y \in X \setminus K$ there exist $j \in I$ and $z_j \in M_j \cap T_j(y)$ such that $p_j(y_j, z_j) \in -C_j \setminus \{\theta\}$.*

Then there exists $v = (v_i)_{i \in I} \in X$ such that for each $i \in I$, one has

- (a) $q_i(u_i, v_i) \notin C_i \setminus \{\theta\}$,
- (b) $p_i(v_i, x_i) \notin -C_i \setminus \{\theta\}$ for all $x_i \in T_i(v)$.

Proof. For each $i \in I$, define $F_i : X \times X_i \multimap E$ and $G_i : X \times X \times X_i \multimap E$ by

$$F_i(x, y_i) = -q_i(x_i, y_i) + C_i \setminus \{\theta\}, \quad \forall (x, y_i) \in X \times X_i, \quad (3.4)$$

$$G_i(t, x, y_i) = p_i(x_i, y_i) + C_i \setminus \{\theta\}, \quad \forall (t, x, y_i) \in X \times X \times X_i, \quad (3.5)$$

respectively. Thus, for each $i \in I$, we let

$$H_i := \{x_i \in X_i : \theta \notin F_i(u, x_i)\} = \{x_i \in X_i : q_i(u_i, x_i) \notin C_i \setminus \{\theta\}\}, \quad (3.6)$$

and let $A_i : X \multimap X_i$ be defined by

$$\begin{aligned} A_i(x) &= \{y_i \in X_i : \theta \in G_i(u, x, y_i)\} \\ &= \{y_i \in X_i : p_i(x_i, y_i) \in -C_i \setminus \{\theta\}\}, \quad \forall x = (x_i)_{i \in I} \in X. \end{aligned} \quad (3.7)$$

Clearly, for each $i \in I$, $\theta \notin G_i(u, x, x_i)$ for all $x = (x_i)_{i \in I} \in X$, and hence $x_i \notin A_i(x)$ for all $x = (x_i)_{i \in I} \in X$. By the C_i -vectorial lower semicontinuity of $q_i(u_i, \cdot)$, H_i is a nonempty closed subset of X_i . By (i), for each $x \in X$, $\text{co}T_i(x) \subseteq H_i$. For each $y_i \in X_i$, by the vectorial upper semicontinuity of $p_i(\cdot, y_i)$, the set $\{x_i \in X_i : p_i(x_i, y_i) \in -C_i \setminus \{\theta\}\}$ is open in X_i , and hence $A_i^-(y_i) = \{x_i \in X_i : q_i(x_i, y_i) \in -C_i \setminus \{\theta\}\} \times \prod_{j \neq i} X_j$ is open in X . For each $x \in X$, $A_i(x)$ is convex from the C_i -convexity of $p_i(x_i, \cdot)$ for all $i \in I$. By (ii), there exist a nonempty compact subset K of X and a nonempty compact convex subset M_i of X_i for each $i \in I$ such that for each $y \in X \setminus K$ there exist $j \in I$ and $z_j \in M_j \cap T_j(y) \cap A_j(y)$. Therefore all the conditions of Theorem 3.1 are satisfied. Applying Theorem 3.1, there exists $v = (v_i)_{i \in I} \in X$ such that:

- (1) $q_i(u_i, v_i) \notin C_i \setminus \{\theta\}$,
- (2) $p_i(v_i, x_i) \notin -C_i \setminus \{\theta\}$ for all $x_i \in T_i(v)$. □

Theorem 3.4. Let $I, X_i, X, C_i, E, \theta, T_i$, and p_i be the same as in Theorem 3.3. Let $u = (u_i)_{i \in I} \in X$ with $p_i(u_i, u_i) = \theta$ for all $i \in I$. For each $i \in I$, suppose that the following conditions are satisfied:

- (i) for each $y \in X$, $\text{co} T_i(y) \subseteq \{x_i \in X_i : p_i(u_i, x_i) \notin C_i \setminus \{\theta\}\}$ and $T_i^-(z_i)$ is open for all $z_i \in X_i$;
- (ii) there exist a nonempty compact subset K of X and a nonempty compact convex subset M_i of X_i for each $i \in I$ such that for each $y \in X \setminus K$ there exist $j \in I$ and $z_j \in M_j \cap T_j(y)$ such that $p_j(y_j, z_j) \in -C_j \setminus \{\theta\}$.

Then there exists $v = (v_i)_{i \in I} \in X$ such that for each $i \in I$, one has

- (a) $p_i(u_i, v_i) \notin C_i \setminus \{\theta\}$,
- (b) $p_i(v_i, x_i) \notin -C_i \setminus \{\theta\}$ for all $x_i \in T_i(v) \cup \{z_i \in X_i : p_i(u_i, z_i) \in C_i \setminus \{\theta\}\}$.

Proof. For each $i \in I$, let $q_i : X_i \times X_i \rightarrow E$ be defined by $q_i = p_i$. By Theorem 3.3, there exists $v = (v_i)_{i \in I} \in X$ such that for each $i \in I$, we have

- (1) $p_i(u_i, v_i) \notin C_i \setminus \{\theta\}$,
- (2) $p_i(v_i, x_i) \notin -C_i \setminus \{\theta\}$ for all $x_i \in T_i(v)$.

We claim that for each $i \in I$, $p_i(v_i, x_i) \notin -C_i \setminus \{\theta\}$ for all $x_i \in \{z_i \in X_i : p_i(u_i, z_i) \in C_i \setminus \{\theta\}\}$. For each $x_i \in \{z_i \in X_i : p_i(u_i, z_i) \in C_i \setminus \{\theta\}\}$, we have $p_i(u_i, x_i) \in C_i \setminus \{\theta\}$. If $p_i(v_i, x_i) \in -C_i \setminus \{\theta\}$, then, by (VQD2) (in Definition 2.6(b)), we obtain

$$p_i(u_i, v_i) \in p_i(u_i, x_i) - p_i(v_i, x_i) + C_i \subseteq C_i \setminus \{\theta\}, \quad (3.8)$$

which contradict with (1). Hence $p_i(v_i, x_i) \notin -C_i \setminus \{\theta\}$ for all $x_i \in T_i(v) \cup \{z_i \in X_i : p_i(u_i, z_i) \in C_i \setminus \{\theta\}\}$, and the proof is completed. \square

Remark 3.5. (i) Let X be a nonempty Hausdorff t.v.s., and let $C \neq \{\theta\}$ be a nonempty pointed convex cone in a t.v.s. E with zero vector θ . If a map $p : X \times X \rightarrow E$ satisfies that for each $x \in X$, $p(x, \cdot)$ is C -quasiconvex, then for each $u \in X$ with $p(u, u) = \theta$, $W_u := \{x \in X : p(u, x) \notin C \setminus \{\theta\}\}$ is a nonempty convex subset in X .

(ii) In Theorem 3.3, if for each $i \in I$, $W_i := \{x_i \in X_i : p_i(u_i, x_i) \notin C_i \setminus \{\theta\}\}$ is convex and $T_i(x) = W_i$ for all $x \in X$, then conclusion (b) can be replaced with conclusion (b)', where

$$(b)' \quad p_i(v_i, x_i) \notin -C_i \setminus \{\theta\}, \text{ for all } x_i \in X_i.$$

Since the sum of two C_i -vectorial quasi-distances is also a C_i -vectorial quasi-distance, the following results related with system of VSEVP for vectorial quasi-distances in a Hausdorff t.v.s. are immediate from Theorem 3.3.

Theorem 3.6. Let $I, X_i, X, C_i, E, \theta, T_i$, and p_i be the same as in Theorem 3.3. Let $u = (u_i)_{i \in I} \in X$ with $p_i(u_i, u_i) = \theta$ for all $i \in I$. For each $i \in I$, let $f_i : X_i \rightarrow E$ be a C_i -v.l.s.c. and C_i -convex function and suppose that the following conditions are satisfied:

- (i) for each $y \in X$, $\text{co} T_i(y) \subseteq \{x_i \in X_i : f_i(x_i) - f_i(u_i) \notin C_i \setminus \{\theta\}\}$ and $T_i^-(z_i)$ is open for all $z_i \in X_i$;
- (ii) there exist a nonempty compact subset K of X and a nonempty compact convex subset M_i of X_i for each $i \in I$ such that for each $y \in X \setminus K$ there exist $j \in I$ and $z_j \in M_j \cap T_j(y)$ such that $p_j(y_j, z_j) + f_j(z_j) - f_j(y_j) \in -C_j \setminus \{\theta\}$.

Then there exists $v = (v_i)_{i \in I} \in X$ such that for each $i \in I$, one has

- (a) $f_i(v_i) - f_i(u_i) \notin C_i \setminus \{\theta\}$,
- (b) $p_i(v_i, x_i) + f_i(x_i) - f_i(v_i) \notin -C_i \setminus \{\theta\}$ for all $x_i \in T_i(v)$.

In Theorem 3.6, if $E = (-\infty, \infty]$ and $C_i = [0, \infty)$ for all $i \in I$, then we have the following result.

Corollary 3.7. *Let I, X_i, X , and T_i be the same as in Theorem 3.3. For each $i \in I$, let $f_i : X_i \rightarrow (-\infty, \infty]$ be a l.s.c. and convex function and let $p_i : X_i \times X_i \rightarrow (-\infty, \infty]$ be a quasi-distance. Let $u = (u_i)_{i \in I} \in X$ with $p_i(u_i, u_i) = 0$ for all $i \in I$. Suppose that the following conditions are satisfied:*

- (i) for each $y \in X$, $\text{co} T_i(y) \subseteq \{x_i \in X_i : f_i(x_i) \leq f_i(u_i)\}$ and $T_i^-(z_i)$ is open for all $z_i \in X_i$;
- (ii) there exist a nonempty compact subset K of X and a nonempty compact convex subset M_i of X_i for each $i \in I$ such that for each $y \in X \setminus K$ there exist $j \in I$ and $z_j \in M_j \cap T_j(y)$ such that $p_j(y_j, z_j) < f_j(y_j) - f_j(z_j)$.

Then there exists $v = (v_i)_{i \in I} \in X$ such that for each $i \in I$, one has

- (a) $f_i(v_i) \leq f_i(u_i)$,
- (b) $p_i(v_i, x_i) \geq f_i(v_i) - f_i(x_i)$ for all $x_i \in T_i(v)$.

Theorem 3.8. *Let $I, X_i, X, C_i, E, \theta, T_i$, and p_i be the same as in Theorem 3.3. Let $u = (u_i)_{i \in I} \in X$ with $p_i(u_i, u_i) = \theta$ for all $i \in I$. For each $i \in I$, let $f_i : X_i \rightarrow E$ be a C_i -v.l.s.c. and C_i -convex function and suppose that the following conditions are satisfied:*

- (i) for each $y \in X$, $\text{co} T_i(y) \subseteq \{x_i \in X_i : p_i(u_i, x_i) + f_i(x_i) - f_i(u_i) \notin C_i \setminus \{\theta\}\}$ and $T_i^-(z_i)$ is open for all $z_i \in X_i$;
- (ii) there exist a nonempty compact subset K of X and a nonempty compact convex subset M_i of X_i for each $i \in I$ such that for each $y \in X \setminus K$ there exist $j \in I$ and $z_j \in M_j \cap T_j(y)$ such that $p_j(y_j, z_j) + f_j(z_j) - f_j(y_j) \in -C_j \setminus \{\theta\}$.

Then there exists $v = (v_i)_{i \in I} \in X$ such that for each $i \in I$, one has

- (a) $p_i(u_i, v_i) + f_i(v_i) - f_i(u_i) \notin C_i \setminus \{\theta\}$,
- (b) $p_i(v_i, x_i) + f_i(x_i) - f_i(v_i) \notin -C_i \setminus \{\theta\}$ for all $x_i \in T_i(v) \cup \{z_i \in X_i : p_i(u_i, z_i) + f_i(z_i) - f_i(u_i) \in C_i \setminus \{\theta\}\}$.

Corollary 3.9. *Let I, X_i, X , and T_i be the same as in Theorem 3.3. For each $i \in I$, let $f_i : X_i \rightarrow (-\infty, \infty]$ be a l.s.c. and convex function and $p_i : X_i \times X_i \rightarrow (-\infty, \infty]$ be a quasi-distance. Let $u = (u_i)_{i \in I} \in X$ with $p_i(u_i, u_i) = 0$ for all $i \in I$. Suppose that the following conditions are satisfied:*

- (i) for each $y \in X$, $\text{co} T_i(y) \subseteq \{x_i \in X_i : p_i(u_i, x_i) \leq f_i(u_i) - f_i(x_i)\}$ and $T_i^-(z_i)$ is open for all $z_i \in X_i$;
- (ii) there exist a nonempty compact subset K of X and a nonempty compact convex subset M_i of X_i for each $i \in I$ such that for each $y \in X \setminus K$ there exist $j \in I$ and $z_j \in M_j \cap T_j(y)$ such that $p_j(y_j, z_j) < f_j(y_j) - f_j(z_j)$.

Then there exists $v = (v_i)_{i \in I} \in X$ such that for each $i \in I$, one has

- (a) $p_i(u_i, v_i) \leq f_i(u_i) - f_i(v_i)$,
- (b) $p_i(v_i, x_i) \geq f_i(v_i) - f_i(x_i)$ for all $x_i \in T_i(v) \cup \{z_i \in X_i : p_i(u_i, z_i) > f_i(u_i) - f_i(z_i)\}$.

By using Theorem 3.1 again, we have the following result.

Theorem 3.10. *Let $I, X_i, X, C_i, E, \theta, u, p_i$, and q_i be the same as in Theorem 3.3. Suppose that there exist a nonempty compact subset K of X and a nonempty compact convex subset M_i of X_i for each $i \in I$ such that for each $y \in X \setminus K$ there exist $j \in I$ and $z_j \in M_j$ such that $q_j(u_j, z_j) \in -C_j$ and $p_j(y_j, z_j) \in -C_j \setminus \{\theta\}$. Then there exists $v = (v_i)_{i \in I} \in X$ such that for each $i \in I$, one has*

- (a) $q_i(u_i, v_i) \notin C_i \setminus \{\theta\}$,
- (b) $p_i(v_i, x_i) \notin -C_i \setminus \{\theta\}$ for all $x_i \in \{z_i \in X_i : q_i(u_i, z_i) \in -C_i\}$.

Proof. For each $i \in I$, let H_i and multivalued maps F_i, G_i , and A_i be defined as in the proof of Theorem 3.3. For each $i \in I$, define $T_i : X \multimap X_i$ by

$$\begin{aligned} T_i(x) &= L_i := \{z_i \in X_i : q_i(u_i, z_i) \in -C_i\}, \quad \forall x \in X \\ \iff T_i^-(z_i) &= \begin{cases} X & \text{if } z_i \in L_i, \\ \emptyset & \text{if } z_i \in X_i \setminus L_i. \end{cases} \end{aligned} \quad (3.9)$$

Clearly, for each $i \in I$, $T_i^-(z_i)$ is open in X for all $z_i \in X_i$. By the C_i -convexity of $q_i(u_i, \cdot)$, L_i is a nonempty convex subset of X_i for all $i \in I$. Since L_i is convex in X_i and $L_i \subseteq H_i$, we have $\text{co}T_i(x) \subseteq H_i$ for all $i \in I$. By our hypothesis, there exist a nonempty compact subset K of X and a nonempty compact convex subset M_i of X_i for each $i \in I$ such that for each $y \in X \setminus K$ there exist $j \in I$ and $z_j \in M_j \cap T_j(y) \cap A_j(y)$. Thus all the conditions of Theorem 3.1 are satisfied, and the conclusion follows from Theorem 3.1. \square

Remark 3.11. In Theorem 3.10, the multivalued map T_i and the condition “for each $y \in Y$, $\text{co}T_i(y) \subseteq \{x_i \in X_i : q_i(u_i, x_i) \notin C_i \setminus \{\theta\}\}$ ” are not assumed. So Theorems 3.10 and 3.3 are different.

Theorem 3.12. *Let $I, X_i, X, C_i, E, \theta$, and p_i be the same as in Theorem 3.3. Let $u = (u_i)_{i \in I} \in X$ with $p_i(u_i, u_i) = \theta$ for all $i \in I$. Suppose that there exist a nonempty compact subset K of X and a nonempty compact convex subset M_i of X_i for each $i \in I$ such that for each $y \in X \setminus K$ there exist $j \in I$ and $z_j \in M_j$ such that $p_j(u_j, z_j) \in -C_j$ and $p_j(y_j, z_j) \in -C_j \setminus \{\theta\}$. Then there exists $v = (v_i)_{i \in I} \in X$ such that for each $i \in I$, one has*

- (a) $p_i(u_i, v_i) \notin C_i \setminus \{\theta\}$,
- (b) $p_i(v_i, x_i) \notin -C_i \setminus \{\theta\}$ for all $x_i \in \{z_i \in X_i : p_i(u_i, z_i) \in -C_i\} \cup \{z_i \in X_i : p_i(u_i, z_i) \in C_i \setminus \{\theta\}\}$.

Proof. For each $i \in I$, let $q_i = p_i$. Applying Theorem 3.10 and following the same argument as in the proof of Theorem 3.4, one can prove the theorem. \square

Theorem 3.13. *Let $I, X_i, X, C_i, E, \theta$, and p_i be the same as in Theorem 3.3. Let $u = (u_i)_{i \in I} \in X$ with $p_i(u_i, u_i) = \theta$ for all $i \in I$. For each $i \in I$, let $f_i : X_i \rightarrow E$ be a C_i -v.l.s.c. and C_i -convex function.*

Suppose that there exist a nonempty compact subset K of X and a nonempty compact convex subset M_i of X_i for each $i \in I$ such that for each $y \in X \setminus K$ there exist $j \in I$ and $z_j \in M_j$ such that $f_j(z_j) - f_j(u_j) \in -C_j$ and $p_j(y_j, z_j) + f_j(z_j) - f_j(y_j) \in -C_j \setminus \{\theta\}$. Then there exists $v = (v_i)_{i \in I} \in X$ such that for each $i \in I$, one has

- (a) $f_i(v_i) - f_i(u_i) \notin C_i \setminus \{\theta\}$,
- (b) $p_i(v_i, x_i) + f_i(x_i) - f_i(v_i) \notin -C_i \setminus \{\theta\}$ for all $x_i \in \{z_i \in X_i : f_i(z_i) - f_i(u_i) \in -C_i\}$.

Corollary 3.14. Let I , X_i , and X be the same as in Theorem 3.3. For each $i \in I$, let $f_i : X_i \rightarrow (-\infty, \infty]$ be a l.s.c. and convex function and $p_i : X_i \times X_i \rightarrow (-\infty, \infty]$ be a quasi-distance. Let $u = (u_i)_{i \in I} \in X$ with $p_i(u_i, u_i) = 0$ for all $i \in I$. Suppose that there exist a nonempty compact subset K of X and a nonempty compact convex subset M_i of X_i for each $i \in I$ such that for each $y \in X \setminus K$ there exist $j \in I$ and $z_j \in M_j$ such that $f_j(z_j) \leq f_j(u_j)$ and $p_j(y_j, z_j) < f_j(y_j) - f_j(z_j)$. Then there exists $v = (v_i)_{i \in I} \in X$ such that for each $i \in I$, one has

- (a) $f_i(v_i) \leq f_i(u_i)$,
- (b) $p_i(v_i, x_i) \geq f_i(v_i) - f_i(x_i)$ for all $x_i \in \{z_i \in X_i : f_i(z_i) \leq f_i(u_i)\}$.

Remark 3.15. [19, Theorem 4.3] is a special case of Corollary 3.14.

Theorem 3.16. Let I , X_i , X , C_i , E , θ , and p_i be the same as in Theorem 3.3. Let $u = (u_i)_{i \in I} \in X$ with $p_i(u_i, u_i) = \theta$ for all $i \in I$. For each $i \in I$, let $f_i : X_i \rightarrow E$ be a C_i -v.l.s.c. and C_i -convex function, and suppose that there exist a nonempty compact subset K of X and a nonempty compact convex subset M_i of X_i for each $i \in I$ such that for each $y \in X \setminus K$ there exist $j \in I$ and $z_j \in M_j$ such that $p_j(u_j, z_j) + f_j(z_j) - f_j(u_j) \in -C_j$ and $p_j(y_j, z_j) + f_j(z_j) - f_j(y_j) \in -C_j \setminus \{\theta\}$. Then for each $u = (u_i)_{i \in I} \in X$ with $p_i(u_i, u_i) = \theta$ for all $i \in I$, there exists $v = (v_i)_{i \in I} \in X$ such that for each $i \in I$, one has

- (a) $p_i(u_i, v_i) + f_i(v_i) - f_i(u_i) \notin C_i \setminus \{\theta\}$,
- (b) $p_i(v_i, x_i) + f_i(x_i) - f_i(v_i) \notin -C_i \setminus \{\theta\}$ for all $x_i \in \{z_i \in X_i : p_i(u_i, z_i) + f_i(z_i) - f_i(u_i) \in -C_i\} \cup \{z_i \in X_i : p_i(u_i, z_i) + f_i(z_i) - f_i(u_i) \in C_i \setminus \{\theta\}\}$.

In Theorem 3.16, if $E = (-\infty, \infty]$ and $C_i = [0, \infty)$ for all $i \in I$, then we have the following system of Lin and Du's variant of system of Ekeland's variational principle in t.v.s.

Corollary 3.17. Let I , X_i , and X be the same as in Theorem 3.3. For each $i \in I$, let $f_i : X_i \rightarrow E$ be a l.s.c. and convex function and let $p_i : X_i \times X_i \rightarrow (-\infty, \infty]$ be a quasi-distance. Let $u = (u_i)_{i \in I} \in X$ with $p_i(u_i, u_i) = 0$ for all $i \in I$. Suppose that there exist a nonempty compact subset K of X and a nonempty compact convex subset M_i of X_i for each $i \in I$ such that for each $y \in X \setminus K$ there exist $j \in I$ and $z_j \in M_j$ such that $p_j(u_j, z_j) \leq f_j(u_j) - f_j(z_j)$ and $p_j(y_j, z_j) < f_j(y_j) - f_j(z_j)$. Then for each $u = (u_i)_{i \in I} \in X$ with $p_i(u_i, u_i) = 0$ for all $i \in I$, there exists $v = (v_i)_{i \in I} \in X$ such that for each $i \in I$, one has

- (a) $p_i(u_i, v_i) \leq f_i(u_i) - f_i(v_i)$,
- (b) $p_i(v_i, x_i) \geq f_i(v_i) - f_i(x_i)$ for all $x_i \in X_i$.

Remark 3.18. Corollary 3.17 generalizes [19, Theorem 4.1].

4. A Vectorial Minimization Theorem and Equivalent Formulations of VSEVP

Using Theorem 3.1 again, we also obtain an existence theorem of system of generalized vectorial equilibrium problem of the Stampacchia-type which can be regarded as a weak form of VSEVP for C -vectorial ℓ -function in a Hausdorff t.v.s.

Theorem 4.1. *Let I be any index set. For each $i \in I$, let X_i be a nonempty Hausdorff t.v.s. and $C_i \neq \{\theta\}$ be a nonempty pointed convex cone in a t.v.s. E with zero vector θ . Let $X = \prod_{i \in I} X_i$. For each $i \in I$, let $p_i, q_i : X_i \times X_i \rightarrow E$ be ℓ_{C_i} -function. Suppose that*

- (i) $\mathcal{L} = \{x = (x_i)_{i \in I} \in X : p_i(x_i, x_i) = q_i(x_i, x_i) = \theta \text{ for all } i \in I\} \neq \emptyset$,
- (ii) *there exist a nonempty compact subset K of X and a nonempty compact convex subset M_i of X_i for each $i \in I$ such that for each $y \in X \setminus K$ there exist $j \in I$ and $z_j \in M_j$ such that $p_j(y_j, z_j) + q_j(y_j, z_j) \in -C_i \setminus \{\theta\}$.*

Then there exists $v = (v_i)_{i \in I} \in X$ such that for each $i \in I$, one has $p_i(v_i, x_i) + q_i(v_i, x_i) \notin -C_i \setminus \{\theta\}$ for all $x_i \in X_i$.

Proof. Let $u = (u_i)_{i \in I} \in \mathcal{L}$ be given. For each $i \in I$, we define $F_i : X \times X_i \rightarrow E, G_i : X \times X \times X_i \rightarrow E$ and $A_i : X \rightarrow X_i$ by

$$F_i(x, y_i) = C_i \setminus \{\theta\}, \quad \forall (x, y_i) \in X \times X_i, \quad (4.1)$$

$$G_i(t, x, y_i) = p_i(x_i, y_i) + q_i(x_i, y_i) + C_i \setminus \{\theta\}, \quad \forall (t, x, y_i) \in X \times X \times X_i,$$

$$A_i(x) = \{y_i \in X_i : \theta \in G_i(u, x, y_i)\}, \quad \forall x = (x_i)_{i \in I} \in X, \quad (4.2)$$

respectively. Clearly,

$$H_i := \{x_i \in X_i : \theta \notin F_i(u, x_i)\} = X_i, \quad \forall i \in I. \quad (4.3)$$

Using the same argument in the proof of Theorem 3.3, one can verify that all the conditions of Theorem 3.1 are satisfied. By Theorem 3.1, there exists $v = (v_i)_{i \in I} \in X$ such that for each $i \in I$, we have $p_i(v_i, x_i) + q_i(v_i, x_i) \notin -C_i \setminus \{\theta\}$ for all $x_i \in X_i$. \square

An existence theorem of system of generalized vector equilibrium problem is immediate from Theorem 4.1 if $q_i \equiv 0$ (the zero map) for all $i \in I$.

Theorem 4.2. *Let I be any index set. For each $i \in I$, let X_i be a nonempty Hausdorff t.v.s., and let $C_i \neq \{\theta\}$ be a nonempty pointed convex cone in a t.v.s. E with zero vector θ . Let $X = \prod_{i \in I} X_i$. For each $i \in I$, let $p_i : X_i \times X_i \rightarrow E$ be a ℓ_{C_i} -function. Suppose that*

- (i) $\mathcal{L} = \{x = (x_i)_{i \in I} \in X : p_i(x_i, x_i) = \theta \text{ for all } i \in I\} \neq \emptyset$;
- (ii) *there exist a nonempty compact subset K of X and a nonempty compact convex subset M_i of X_i for each $i \in I$ such that for each $y \in X \setminus K$ there exist $j \in I$ and $z_j \in M_j$ such that $p_j(y_j, z_j) \in -C_i \setminus \{\theta\}$.*

Then there exists $v = (v_i)_{i \in I} \in X$ such that for each $i \in I$, one has $p_i(v_i, x_i) \notin -C_i \setminus \{\theta\}$ for all $x_i \in X_i$.

Remark 4.3. (a) Notice that Theorems 4.1 and 4.2 are indeed equivalent since the sum of two ℓ_{C_i} -functions is also a ℓ_{C_i} -function.

(b) In Theorem 4.1, the maps p_i and q_i are only assumed to be ℓ_{C_i} -functions which need not be C_i -vectorial quasi-distances. So Theorem 4.1 is different from any theorem in Section 3 and is not a special case of any theorem in Section 3.

Theorem 4.4. *Let I, X_i, X, C_i, E , and θ be the same as in Theorem 4.1. For each $i \in I$, let $p_i : X_i \times X_i \rightarrow E$ be a ℓ_{C_i} -function and let $f_i : X_i \rightarrow E$ be a C_i -v.l.s.c. and C_i -convex function. Suppose that*

- (i) $\mathcal{L} = \{x = (x_i)_{i \in I} \in X : p_i(x_i, x_i) = \theta \text{ for all } i \in I\} \neq \emptyset$,
- (ii) *there exist a nonempty compact subset K of X and a nonempty compact convex subset M_i of X_i for each $i \in I$ such that for each $y \in X \setminus K$ there exist $j \in I$ and $z_j \in M_j$ such that $p_j(y_j, z_j) + f_j(z_j) - f_j(y_j) \in -C_j \setminus \{\theta\}$.*

Then there exists $v = (v_i)_{i \in I} \in X$ such that for each $i \in I$, one has $p_i(v_i, x_i) + f_i(x_i) - f_i(v_i) \notin -C_i \setminus \{\theta\}$ for all $x_i \in X_i$.

Remark 4.5. [19, Theorem 4.2] is a special case of Theorem 4.4.

Applying Theorem 4.4, we obtain the following vectorial minimization theorem.

Theorem 4.6 (vectorial minimization theorem). *Let I, X_i, X, C_i, E , and θ be the same as in Theorem 4.1. For each $i \in I$, let $p_i : X_i \times X_i \rightarrow E$ be a ℓ_{C_i} -function and let $f_i : X_i \rightarrow E$ be a C_i -v.l.s.c. and C_i -convex function. Suppose that*

- (i) $\mathcal{L} = \{x = (x_i)_{i \in I} \in X : p_i(x_i, x_i) = \theta \text{ for all } i \in I\} \neq \emptyset$,
- (ii) *there exist a nonempty compact subset K of X and a nonempty compact convex subset M_i of X_i for each $i \in I$ such that for each $y \in X \setminus K$ there exist $j \in I$ and $z_j \in M_j$ such that $p_j(y_j, z_j) + f_j(z_j) - f_j(y_j) \in -C_j \setminus \{\theta\}$,*
- (iii) *for $i \in I$ and $x_i \in X_i$ with $f_i(x_i) \notin \text{Min}_{C_i} f_i(X_i)$, there exists $y_i \in X_i$ with $y_i \neq x_i$ such that $p_i(x_i, y_i) + f_i(y_i) - f_i(x_i) \in -C_i \setminus \{\theta\}$.*

Then there exists $\hat{x} = (\hat{x}_i)_{i \in I} \in X$ such that $f_i(\hat{x}_i) \in \text{Min}_{C_i} f_i(X_i)$ for all $i \in I$.

Proof. Applying Theorem 4.4, there exists $v = (v_i)_{i \in I} \in X$ such that for each $i \in I$, we have $p_i(v_i, x_i) + f_i(x_i) - f_i(v_i) \notin -C_i \setminus \{\theta\}$ for all $x_i \in X_i$. We claim that $f_i(v_i) \in \text{Min}_{C_i} f_i(X_i)$ for all $i \in I$. Suppose to the contrary that there exists $i_0 \in I$ such that $f_{i_0}(v_{i_0}) \notin \text{Min}_{C_{i_0}} f_{i_0}(X_{i_0})$. Then, by our assumption, there exists $y_{i_0} = y_{i_0}(v_{i_0}) \in X_{i_0}$ with $y_{i_0} \neq v_{i_0}$ such that $p_{i_0}(v_{i_0}, y_{i_0}) + f_{i_0}(y_{i_0}) - f_{i_0}(v_{i_0}) \in -C_{i_0} \setminus \{\theta\}$, which leads to a contradiction. Therefore $f_i(v_i) \in \text{Min}_{C_i} f_i(X_i)$ for all $i \in I$. \square

The following scalar minimization theorem follows from Theorem 4.6 immediately.

Corollary 4.7. *Let I, X_i , and X be the same as in Theorem 4.1. For each $i \in I$, let $f_i : X_i \rightarrow (-\infty, \infty]$ be a l.s.c. and convex function and let $p_i : X_i \times X_i \rightarrow (-\infty, \infty]$ be a ℓ -function. Suppose that*

- (i) $\mathfrak{D} = \{x = (x_i)_{i \in I} \in X : p_i(x_i, x_i) = 0 \text{ for all } i \in I\} \neq \emptyset$,
- (ii) *there exist a nonempty compact subset K of X and a nonempty compact convex subset M_i of X_i for each $i \in I$ such that for each $y \in X \setminus K$ there exist $j \in I$ and $z_j \in M_j$ such that $p_j(y_j, z_j) < f_j(y_j) - f_j(z_j)$,*

- (iii) for any $i \in I$ and $x_i \in X_i$ with $f_i(x_i) > \inf_{z_i \in X_i} f_i(z_i)$ there exists $y_i \in X_i$ with $y_i \neq x_i$ such that $p_i(x_i, y_i) < f_i(x_i) - f_i(y_i)$.

Then there exists $\hat{x} = (\hat{x}_i)_{i \in I} \in X$ such that $f_i(\hat{x}_i) = \inf_{z_i \in X_i} f_i(z_i)$ for all $i \in I$.

Remark 4.8. (a) [19, Theorem 5.5] is a special case of Corollary 4.7.

(b) Theorems 4.4 and 4.6 are equivalent if they further add the condition "for each $i \in I$, $p_i(x_i, y_i) \in C_i$ for all $x_i, y_i \in X_i$." Indeed, it suffices to show that Theorem 4.6 implies Theorem 4.4. Suppose that for each $x = (x_i)_{i \in I} \in X$, there exists $i_x \in I$ such that $p_{i_x}(x_{i_x}, y_{i_x}) + f_{i_x}(y_{i_x}) - f_{i_x}(x_{i_x}) \in -C_{i_x} \setminus \{\theta\}$ for some $y_{i_x} \in X_{i_x}$ with $y_{i_x} \neq x_{i_x}$. Then, by Theorem 4.6, there exists $v = (v_i)_{i \in I} \in X$ such that $f_i(v_i) \in \text{Min}_{C_i} f_i(X_i)$ or

$$f_i(x_i) - f_i(v_i) \notin -C_i \setminus \{\theta\}, \quad \forall x_i \in X_i, \forall i \in I. \quad (4.4)$$

But from our assumption, there exists $i_v \in I$ such that

$$p_{i_v}(v_{i_v}, w_{i_v}) + f_{i_v}(w_{i_v}) - f_{i_v}(v_{i_v}) \in -C_{i_v} \setminus \{\theta\} \quad (4.5)$$

for some $w_{i_v} \in X_{i_v}$ with $w_{i_v} \neq v_{i_v}$. It follows that

$$f_{i_v}(w_{i_v}) - f_{i_v}(v_{i_v}) \in -p_{i_v}(v_{i_v}, w_{i_v}) - C_{i_v} \setminus \{\theta\} \subseteq -C_{i_v} \setminus \{\theta\}, \quad (4.6)$$

which leads to a contradiction.

Now, we give some equivalent formulations of Theorem 4.1 (when I is a singleton) as follows.

Theorem 4.9. Let X be a Hausdorff t.v.s., let $C \neq \{\theta\}$ be a nonempty pointed convex cone in a t.v.s. E with zero vector θ , $p, q : X \times X \rightarrow E$ be \mathcal{L}_C -functions. Suppose that

- (1) $\mathcal{R} = \{x \in X : p(x, x) = q(x, x) = 0\} \neq \emptyset$,
- (2) there exist a nonempty compact subset K of X and a nonempty compact convex subset M of X such that for each $y \in X \setminus K$ there exists $z \in M$ such that $p(y, z) + q(y, z) \in -C \setminus \{\theta\}$.

Then the following statements are equivalent.

- (i) (Vectorial Variant of Ekeland's Variational Principle). There exists $v \in X$ such that $p(v, x) + q(v, x) \notin -C \setminus \{\theta\}$ for all $x \in X$.
- (ii) (Common Fixed Point Theorem for a Family of Multivalued Maps). Let Λ be an index set. For each $i \in \Lambda$, let $T_i : X \multimap X$ be a multivalued map with nonempty values such that for each $(i, x) \in \Lambda \times X$ with $x \notin T_i(x)$, there exists $y = y(x, i) \in X$ with $y \neq x$ such that

$$p(x, y) + q(x, y) \in -C \setminus \{\theta\}. \quad (4.7)$$

Then there exists $x_0 \in X$ such that $x_0 \in \bigcap_{i \in \Lambda} T_i(x_0)$. That is, $\{T_i\}_{i \in \Lambda}$ has a common fixed point in X .

- (iii) (*Common Fixed Point Theorem for a Family of Single-Valued Maps*). Let Λ be an index set. For each $i \in \Lambda$, suppose that $T_i : X \rightarrow X$ is a self-map satisfying

$$p(x, T_i x) + q(x, T_i x) \in -C \setminus \{\theta\}. \quad (4.8)$$

for all $x \neq T_i(x)$. Then there exists $x_0 \in X$ such that $T_i x_0 = x_0$ for all $i \in \Lambda$;

- (iv) (*Maximal Element Theorem for a Family of Multivalued Maps*). Let Λ be an index set. For each $i \in \Lambda$, let $T_i : X \multimap X$ be a multivalued map. Suppose that for each $(i, x) \in \Lambda \times X$ with $T_i(x) \neq \emptyset$, there exists $y = y(x, i) \in X$ with $y \neq x$ such that

$$p(x, y) + q(x, y) \in -C \setminus \{\theta\}. \quad (4.9)$$

Then there exists $x_0 \in X$ such that $T_i(x_0) = \emptyset$ for all $i \in \Lambda$.

Proof. By Theorem 4.1, conclusion (i) holds.

(a) “(i) \iff (ii)”

(\implies) By (i), there exists $v \in X$ such that $p(v, x) + q(v, x) \notin -C \setminus \{\theta\}$ for all $x \in X$. We claim that $v \in T_i(v)$ for all $i \in I$. If $v \notin T_{i_0} v$ for some $i_0 \in I$, then, by hypothesis, there exists $w(v, i_0) \in X$ with $w(v, i_0) \neq v$ such that

$$p(v, w(v, i_0)) + q(v, w(v, i_0)) \in -C \setminus \{\theta\}, \quad (4.10)$$

which leads to a contradiction. Hence $v \in \bigcap_{i \in I} T_i(v)$ and v is a common fixed point of $\{T_i\}_{i \in I}$.

(\impliedby) Suppose that for each $x \in X$, there exists $y \in X$ with $y \neq x$ such that $p(x, y) + q(x, y) \in -C \setminus \{\theta\}$. Then for each $x \in X$, we can define a multivalued map $T : X \multimap X \setminus \{\emptyset\}$ by

$$T(x) = \{y \in X : p(x, y) + q(x, y) \in -C \setminus \{\theta\}\}. \quad (4.11)$$

Clearly, $x \notin T(x)$ for all $x \in X$. But, by (ii), T has a fixed point v in X , a contradiction. So (i) is true.

(b) “(ii) \iff (iii)”

(\implies) Suppose (ii) holds. Under the assumption of (iii), for each $i \in I$, let $\varphi_i : X \multimap X$ be defined by $\varphi_i(x) = \{T_i(x)\}$. Then for each $(i, x) \in I \times X$ with $x \notin \varphi_i(x)$, we have $x \neq T_i(x)$. By hypothesis of (iii), $p(x, T_i x) + q(x, T_i x) \in -C \setminus \{\theta\}$. Therefore, by (ii), there exists $v \in X$ such that $v \in \bigcap_{i \in I} \varphi_i(v)$ or $T_i v = v$ for all $i \in I$, and hence (iii) is proved.

(\impliedby) Assume (iii) holds. Under the assumption of (ii), for each $(i, x) \in I \times X$ with $x \notin T_i(x)$, there exists $y(x, i) \in X$ with $y(x, i) \neq x$ such that

$$p(x, y(x, i)) + q(x, y(x, i)) \in -C \setminus \{\theta\}. \quad (4.12)$$

Define $\tau_i : X \rightarrow X$ by

$$\tau_i(x) = \begin{cases} x & \text{if } x \in T_i(x), \\ y(x, i) & \text{if } x \notin T_i(x). \end{cases} \quad (4.13)$$

Hence τ_i is a self-map of X into X satisfying $p(x, \tau_i(x)) + q(x, \tau_i(x)) \in -C \setminus \{\theta\}$ for all $x \neq \tau_i(x)$. By (iii), there exists $v \in X$ such that $v = \tau_i(v) \in T_i(v)$ for all $i \in I$. This shows that (iii) implies (ii).

(c) "(i) \iff (iv)"

(\implies) Using (i), there exists $v \in X$ such that $p(v, x) + q(v, x) \notin -C \setminus \{\theta\}$ for all $x \in X$. We want to show that $T_i(v) = \emptyset$ for all $i \in I$. Suppose to the contrary that there exists $i_0 \in I$ such that $T_{i_0}(v) \neq \emptyset$. By hypothesis of (iv), there exists $w = w(v, i_0) \in X$ with $w \neq v$ such that

$$p(v, w) + q(v, w) \in -C \setminus \{\theta\}, \quad (4.14)$$

which is a contradiction. Therefore $T_i(v) = \emptyset$ for all $i \in I$.

(\impliedby) Suppose that for each $x \in X$, there exists $y \in X$ with $y \neq x$ such that $p(x, y) + q(x, y) \in -C \setminus \{\theta\}$. For each $x \in X$, define a multivalued map $T : X \multimap X \setminus \{\emptyset\}$ by

$$T(x) = \{y \in X : p(x, y) + q(x, y) \in -C \setminus \{\theta\}\}. \quad (4.15)$$

Then $T(x) \neq \emptyset$ for all $x \in X$. But applying (iv), there exists $x_0 \in X$ such that $T(x_0) = \emptyset$, a contradiction. Hence (i) holds. \square

Remark 4.10. Theorem 4.9 improves and generalizes Theorems 4.2, 5.1, 5.2, 5.3, and 5.4 in [19].

Acknowledgment

This research was supported partially by grant no. NSC 99-2115-M-017-001 of the National Science Council of the Republic of China.

References

- [1] E. Blum and W. Oettli, "From optimization and variational inequalities to equilibrium problems," *The Mathematics Student*, vol. 63, no. 1–4, pp. 123–145, 1994.
- [2] Q. H. Ansari, S. Schaible, and J.-C. Yao, "The system of generalized vector equilibrium problems with applications," *Journal of Global Optimization*, vol. 22, no. 1–4, pp. 3–16, 2002.
- [3] G. Isac, V. A. Bulavsky, and V. V. Kalashnikov, *Complementarity, Equilibrium, Efficiency and Economics*, vol. 63 of *Nonconvex Optimization and Its Applications*, Kluwer Academic Publishers, Dordrecht, The Netherlands, 2002.
- [4] L.-J. Lin, "System of generalized vector quasi-equilibrium problems with applications to fixed point theorems for a family of nonexpansive multivalued mappings," *Journal of Global Optimization*, vol. 34, no. 1, pp. 15–32, 2006.
- [5] I. Ekeland, "Sur les problèmes variationnels," *Comptes Rendus de l'Académie des Sciences, Série A-B*, vol. 275, pp. A1057–A1059, 1972.

- [6] I. Ekeland, "On the variational principle," *Journal of Mathematical Analysis and Applications*, vol. 47, pp. 324–353, 1974.
- [7] I. Ekeland, "Nonconvex minimization problems," *Bulletin of the American Mathematical Society*, vol. 1, no. 3, pp. 443–474, 1979.
- [8] G.-Y. Chen, X. Huang, and X. Yang, *Vector Optimization*, vol. 541 of *Lecture Notes in Economics and Mathematical Systems*, Springer, Berlin, Germany, 2005.
- [9] G. Y. Chen, X. Q. Yang, and H. Yu, "A nonlinear scalarization function and generalized quasi-vector equilibrium problems," *Journal of Global Optimization*, vol. 32, no. 4, pp. 451–466, 2005.
- [10] W.-S. Du, "On some nonlinear problems induced by an abstract maximal element principle," *Journal of Mathematical Analysis and Applications*, vol. 347, no. 2, pp. 391–399, 2008.
- [11] W.-S. Du, "Hybrid inclusion and disclusion systems with applications to equilibria and parametric optimization," *Journal of Global Optimization*, vol. 47, no. 1, pp. 119–132, 2010.
- [12] A. Göpfert, H. Riahi, C. Tammer, and C. Zălinescu, *Variational Methods in Partially Ordered Spaces*, CMS Books in Mathematics/Ouvrages de Mathématiques de la SMC, 17, Springer, New York, NY, USA, 2003.
- [13] A. Göpfert, Chr. Tammer, and C. Zălinescu, "On the vectorial Ekeland's variational principle and minimal points in product spaces," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 39, no. 7, pp. 909–922, 2000.
- [14] A. H. Hamel, "Phelps' lemma, Daneš' drop theorem and Ekeland's principle in locally convex spaces," *Proceedings of the American Mathematical Society*, vol. 131, no. 10, pp. 3025–3038, 2003.
- [15] A. Hamel and A. Löhne, "A minimal point theorem in uniform spaces," in *Nonlinear Analysis and Applications: To V. Lakshmikantham on his 80th Birthday. Vol. 1, 2*, pp. 577–593, Kluwer Academic Publishers, Dordrecht, The Netherlands, 2003.
- [16] D. H. Hyers, G. Isac, and T. M. Rassias, *Topics in Nonlinear Analysis & Applications*, World Scientific, River Edge, NJ, USA, 1997.
- [17] L.-J. Lin and W.-S. Du, "Ekeland's variational principle, minimax theorems and existence of nonconvex equilibria in complete metric spaces," *Journal of Mathematical Analysis and Applications*, vol. 323, no. 1, pp. 360–370, 2006.
- [18] L.-J. Lin and W.-S. Du, "Some equivalent formulations of the generalized Ekeland's variational principle and their applications," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 67, no. 1, pp. 187–199, 2007.
- [19] L.-J. Lin and W.-S. Du, "Systems of equilibrium problems with applications to new variants of Ekeland's variational principle, fixed point theorems and parametric optimization problems," *Journal of Global Optimization*, vol. 40, no. 4, pp. 663–677, 2008.
- [20] L.-J. Lin and W.-S. Du, "On maximal element theorems, variants of Ekeland's variational principle and their applications," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 68, no. 5, pp. 1246–1262, 2008.
- [21] L.-J. Lin and C.-S. Chuang, "Existence theorems for variational inclusion problems and the set-valued vector Ekeland variational principle in a complete metric space," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 70, no. 7, pp. 2665–2672, 2009.
- [22] W. Oettli and M. Théra, "Equivalents of Ekeland's principle," *Bulletin of the Australian Mathematical Society*, vol. 48, no. 3, pp. 385–392, 1993.
- [23] Chr. Tammer, "A generalization of Ekeland's variational principle," *Optimization*, vol. 25, no. 2-3, pp. 129–141, 1992.
- [24] C. D. Aliprantis and K. C. Border, *Infinite-Dimensional Analysis*, Springer, Berlin, Germany, 2nd edition, 1999.
- [25] N. X. Tan, "Quasivariational inequalities in topological linear locally convex Hausdorff spaces," *Mathematische Nachrichten*, vol. 122, pp. 231–245, 1985.



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

