

## Research Article

# Approximate $n$ -Lie Homomorphisms and Jordan $n$ -Lie Homomorphisms on $n$ -Lie Algebras

M. Eshaghi Gordji<sup>1,2</sup> and G. H. Kim<sup>1,2</sup>

<sup>1</sup> Department of Mathematics, Semnan University, P.O. Box 35195-363, Semnan, Iran

<sup>2</sup> Department of Mathematics, Kangnam University, Yongin, Gyeonggi 446-702, Republic of Korea

Correspondence should be addressed to G. H. Kim, ghkim@kangnam.ac.kr

Received 18 October 2011; Accepted 5 January 2012

Academic Editor: Marcia Federson

Copyright © 2012 M. E. Gordji and G. H. Kim. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Using fixed point methods, we establish the stability of  $n$ -Lie homomorphisms and Jordan  $n$ -Lie homomorphisms on  $n$ -Lie algebras associated to the following generalized Jensen functional equation  $\mu f(\sum_{i=1}^n x_i/n) + \mu \sum_{j=2}^n f(\sum_{i=1, i \neq j}^n x_i - (n-1)x_j/n) = f(\mu x_1) (n \geq 2)$ .

## 1. Introduction

Let  $n$  be a natural number greater or equal to 3. The notion of an  $n$ -Lie algebra was introduced by Filippov in 1985 [1]. The Lie product is taken between  $n$  elements of the algebra instead of two. This new bracket is  $n$ -linear, antisymmetric and satisfies a generalization of the Jacobi identity. For  $n = 3$  this product is a special case of the Nambu bracket, well known in physics, which was introduced by Nambu [2] in 1973, as a generalization of the Poisson bracket in Hamiltonian mechanics.

An  $n$ -Lie algebra is a natural generalization of a Lie algebra. Namely, a vector space  $V$  together with a multilinear, antisymmetric  $n$ -ary operation  $[\cdot]: \Lambda^n V \rightarrow V$  is called an  $n$ -Lie algebra,  $n \geq 3$ , if the  $n$ -ary bracket is a derivation with respect to itself, that is,

$$[[x_1, \dots, x_n], x_{n+1}, \dots, x_{2n-1}] = \sum_{i=1}^n [x_1, \dots, x_{i-1} [x_i, x_{n+1}, \dots, x_{2n-1}], \dots, x_n], \quad (1.1)$$

where  $x_1, x_2, \dots, x_{2n-1} \in V$ . Equation (1.1) is called the generalized Jacobi identity. The meaning of this identity is similar to that of the usual Jacobi identity for a Lie algebra (which is a 2-Lie algebra).

In [1] and several subsequent papers, [3–5] a structure theory of finite-dimensional  $n$ -Lie algebras over a field  $\mathbb{F}$  of characteristic 0 was developed.

$n$ -ary algebras have been considered in physics in the context of Nambu mechanics [2, 6] and, recently (for  $n = 3$ ), in the search for the effective action of coincident  $M2$ -branes in  $M$ -theory initiated by the Bagger-Lambert-Gustavsson (BLG) model [7, 8] (further references on the physical applications of  $n$ -ary algebras are given in [9]).

From now on, we only consider  $n$ -Lie algebras over the field of complex numbers. An  $n$ -Lie algebra  $A$  is a normed  $n$ -Lie algebra if there exists a norm  $\|\cdot\|$  on  $A$  such that  $\|[x_1, x_2, \dots, x_n]\| \leq \|x_1\| \|x_2\| \cdots \|x_n\|$  for all  $x_1, x_2, \dots, x_n \in A$ . A normed  $n$ -Lie algebra  $A$  is called a Banach  $n$ -Lie algebra, if  $(A, \|\cdot\|)$  is a Banach space.

Let  $(A, [\cdot]_A)$  and  $(B, [\cdot]_B)$  be two Banach  $n$ -Lie algebras. A  $\mathbb{C}$ -linear mapping  $H : (A, [\cdot]_A) \rightarrow (B, [\cdot]_B)$  is called an  $n$ -Lie homomorphism if

$$H([x_1 x_2 \cdots x_n]_A) = [H(x_1)H(x_2) \cdots H(x_n)]_B \quad (1.2)$$

for all  $x_1, x_2, \dots, x_n \in A$ . A  $\mathbb{C}$ -linear mapping  $H : (A, [\cdot]_A) \rightarrow (B, [\cdot]_B)$  is called a Jordan  $n$ -Lie homomorphism if

$$H([x x \cdots x]_A) = [H(x)H(x) \cdots H(x)]_B \quad (1.3)$$

for all  $x \in A$ .

The study of stability problems had been formulated by Ulam [10] during a talk in 1940. Under what condition does there exist a homomorphism near an approximate homomorphism? In the following year, Hyers [11] answered affirmatively the question of Ulam for Banach spaces, which states that if  $\varepsilon > 0$  and  $f : X \rightarrow Y$  is a map with  $X$  a normed space,  $Y$  a Banach spaces such that

$$\|f(x + y) - f(x) - f(y)\| \leq \varepsilon \quad (1.4)$$

for all  $x, y \in X$ , then there exists a unique additive map  $T : X \rightarrow Y$  such that

$$\|f(x) - T(x)\| \leq \varepsilon \quad (1.5)$$

for all  $x \in X$ . A generalized version of the theorem of Hyers for approximately linear mappings was presented by Rassias [12] in 1978 by considering the case when inequality (1.4) is unbounded. Due to that fact, the additive functional equation  $f(x + y) = f(x) + f(y)$  is said to have the generalized Hyers-Ulam-Rassias stability property. A large list of references concerning the stability of functional equations can be found in [13–32].

In 1982–1994, Rassias (see [26–28]) solved the Ulam problem for different mappings and for many Euler-Lagrange type quadratic mappings, by involving a product of different powers of norms. In addition, Rassias considered the mixed product sum of powers of norms control function. For more details see [33–57].

In 2003 Cădariu and Radu applied the fixed-point method to the investigation of the Jensen functional equation [58]. They could present a short and a simple proof (different of the “*direct method*”, initiated by Hyers in 1941) for the generalized Hyers-Ulam stability of Jensen functional equation [58] and for quadratic functional equation.

Park and Rassias [59] proved the stability of homomorphisms in  $C^*$ -algebras and Lie  $C^*$ -algebras and also of derivations on  $C^*$ -algebras and Lie  $C^*$ -algebras for the Jensen-type functional equation

$$\mu f\left(\frac{x+y}{2}\right) + \mu f\left(\frac{x-y}{2}\right) - f(\mu x) = 0 \tag{1.6}$$

for all  $\mu \in \mathbb{T}^1 := \{\lambda \in \mathbb{C}; |\lambda| = 1\}$ .

In this paper, by using the fixed-point methods, we establish the stability of  $n$ -Lie homomorphisms and Jordan  $n$ -Lie homomorphisms on  $n$ -Lie Banach algebras associated to the following generalized Jensen type functional equation:

$$\mu f\left(\frac{\sum_{i=1}^n x_i}{n}\right) + \mu \sum_{j=2}^n f\left(\frac{\sum_{i=1, i \neq j}^n x_i - (n-1)x_j}{n}\right) - f(\mu x_1) = 0 \tag{1.7}$$

for all  $\mu \in (\mathbb{T}_{1/n_0}^1 := \{e^{i\theta}; 0 \leq \theta \leq 2\pi/n_0\} \cup \{1\})$ , where  $n \geq 2$ .

Throughout this paper, assume that  $(A, [\ ]_A), (B, [\ ]_B)$  are two  $n$ -Lie Banach algebras.

## 2. Main Results

Before proceeding to the main results, we recall a fundamental result in fixed point theory.

**Theorem 2.1** (see [60]). *Let  $(\Omega, d)$  be a complete generalized metric space, and let  $T : \Omega \rightarrow \Omega$  be a strictly contractive function with Lipschitz constant  $L$ . Then for each given  $x \in \Omega$ , either*

$$d(T^m x, T^{m+1} x) = \infty \quad \forall m \geq 0, \tag{2.1}$$

or other exists a natural number  $m_0$  such that

- (i)  $d(T^m x, T^{m+1} x) < \infty$  for all  $m \geq m_0$ ;
- (ii) the sequence  $\{T^m x\}$  is convergent to a fixed point  $y^*$  of  $T$ ;
- (iii)  $y^*$  is the unique fixed point of  $T$  in  $\Lambda = \{y \in \Omega : d(T^{m_0} x, y) < \infty\}$ ;
- (iv)  $d(y, y^*) \leq (1/(1-L))d(y, Ty)$  for all  $y \in \Lambda$ .

We start our work with the main theorem of the our paper.

**Theorem 2.2.** *Let  $n_0 \in \mathbb{N}$  be a fixed positive integer number. Let  $f : A \rightarrow B$  be a function for which there exists a function  $\phi : A^n \rightarrow [0, \infty)$  such that*

$$\left\| \mu f\left(\frac{\sum_{i=1}^n x_i}{n}\right) + \mu \sum_{j=2}^n f\left(\frac{\sum_{i=1, i \neq j}^n x_i - (n-1)x_j}{n}\right) - f(\mu x_1) \right\|_B \leq \phi(x_1, x_2, \dots, x_n) \tag{2.2}$$

for all  $\mu \in (T_{1/n_0}^1 := \{e^{i\theta}; 0 \leq \theta \leq 2\pi/n_0\} \cup \{1\})$  and all  $x_1, \dots, x_n \in A$ , and that

$$\|f([x_1 x_2 \cdots x_n]_A) - [f(x_1)f(x_2) \cdots f(x_n)]_B\|_B \leq \phi(x_1, x_2, \dots, x_n) \quad (2.3)$$

for all  $x_1, \dots, x_n \in A$ . If there exists an  $L < 1$  such that

$$\phi(x_1, x_2, \dots, x_n) \leq nL\phi\left(\frac{x_1}{n}, \frac{x_2}{n}, \dots, \frac{x_n}{n}\right) \quad (2.4)$$

for all  $x_1, \dots, x_n \in A$ , then there exists a unique  $n$ -Lie homomorphism  $H : A \rightarrow B$  such that

$$\|f(x) - H(x)\| \leq \frac{L}{1-L}\phi(x, 0, 0, \dots, 0) \quad (2.5)$$

for all  $x \in A$ .

*Proof.* Let  $\Omega$  be the set of all functions from  $A$  into  $B$  and let

$$d(g, h) := \inf\{C \in \mathbb{R}^+ : \|g(x) - h(x)\|_B \leq C\phi(x, 0, \dots, 0), \forall x \in A\}. \quad (2.6)$$

It is easy to show that  $(\Omega, d)$  is a generalized complete metric space [61].

Now we define the mapping  $J : \Omega \rightarrow \Omega$  by  $J(h)(x) = (1/n)h(nx)$  for all  $x \in A$ .

Note that for all  $g, h \in \Omega$ ,

$$\begin{aligned} d(g, h) < C &\implies \|g(x) - h(x)\| \leq C\phi(x, 0, \dots, 0), \quad \forall x \in A, \\ &\implies \left\| \frac{1}{n}g(nx) - \frac{1}{n}h(nx) \right\| \leq \frac{1}{|n|^e} C\phi(nx, 0, \dots, 0), \quad \forall x \in A, \\ &\implies \left\| \frac{1}{n}g(nx) - \frac{1}{n}h(nx) \right\| \leq LC\phi(x, 0, \dots, 0), \quad \forall x \in A, \\ &\implies d(J(g), J(h)) \leq LC. \end{aligned} \quad (2.7)$$

Hence we see that

$$d(J(g), J(h)) \leq Ld(g, h) \quad (2.8)$$

for all  $g, h \in \Omega$ . It follows from (2.4) that

$$\lim_{m \rightarrow \infty} \frac{1}{n^m} \phi(n^m x_1, n^m x_2, \dots, n^m x_n) = 0 \quad (2.9)$$

for all  $x_1, \dots, x_n \in A$ . Putting  $\mu = 1$ ,  $x_1 = x$ , and  $x_j = 0$  ( $j = 2, \dots, n$ ) in (2.2), we obtain

$$\left\| n f\left(\frac{x}{n}\right) - f(x) \right\|_B \leq \phi(x, 0, \dots, 0) \quad (2.10)$$

for all  $x \in A$ . Thus by using (2.4), we obtain that

$$\left\| \frac{1}{n} f(nx) - f(x) \right\|_B \leq \frac{1}{n} \phi(nx, 0, \dots, 0) \leq L \phi(x, 0, \dots, 0) \quad (2.11)$$

for all  $x \in A$ , that is,

$$d(f, J(f)) \leq L < \infty. \quad (2.12)$$

By Theorem 2.1,  $J$  has a unique fixed point in the set  $X_1 := \{h \in \Omega : d(f, h) < \infty\}$ . Let  $H$  be the fixed point of  $J$ .  $H$  is the unique mapping with

$$H(nx) = nH(x) \quad (2.13)$$

for all  $x \in A$ , such that there exists  $C \in (0, \infty)$  satisfying

$$\|f(x) - H(x)\|_B \leq C \phi(x, 0, \dots, 0) \quad (2.14)$$

for all  $x \in A$ . On the other hand we have  $\lim_{m \rightarrow \infty} d(J^m(f), H) = 0$ , so

$$\lim_{m \rightarrow \infty} \frac{1}{n^m} f(n^m x) = H(x) \quad (2.15)$$

for all  $x \in A$ . Also by Theorem 2.1, we have

$$d(f, H) \leq \frac{1}{1-L} d(f, J(f)). \quad (2.16)$$

It follows from (2.12) and (2.16) that

$$d(f, H) \leq \frac{L}{1-L}. \quad (2.17)$$

This implies the inequality (2.5). By (2.21), we have

$$\begin{aligned} & \|H([x_1 x_2 \cdots x_n]_A) - [H(x_1)H(x_2)H(x_3) \cdots H(x_n)]_B\|_B \\ &= \lim_{m \rightarrow \infty} \left\| \frac{1}{n^{nm}} H([n^m x_1 n^m x_2 \cdots n^m x_n]_A) - \frac{1}{n^{nm}} ([H(n^m x_1)H(n^m x_2)H(n^m x_3) \cdots H(n^m x_n)]_B) \right\| \\ &\leq \lim_{m \rightarrow \infty} \frac{1}{n^{nm}} \phi(n^m x_1, n^m x_2, \dots, n^m x_n) = 0 \end{aligned} \quad (2.18)$$

for all  $x_1, \dots, x_n \in A$ . Hence

$$H([x_1 x_2 \cdots x_n]_A) = [H(x_1)H(x_2)H(x_3) \cdots H(x_n)]_B \quad (2.19)$$

for all  $x_1, \dots, x_n \in A$ .

On the other hand, it follows from (2.2), (2.9), and (2.15) that

$$\begin{aligned} & \left\| H\left(\frac{\sum_{i=1}^n x_i}{n}\right) + \sum_{j=2}^n H\left(\frac{\sum_{i=1, i \neq j}^n x_i - (n-1)x_j}{n}\right) - H(x_1) \right\|_B \\ &= \lim_{m \rightarrow \infty} \frac{1}{n^m} \left\| f\left(n^{m-1} \sum_{i=1}^n x_i\right) + \sum_{j=2}^n \left(f\left(n^{m-1} \left(\sum_{i=1, i \neq j}^n x_i - (n-1)x_j\right)\right)\right) - f(n^m x_1) \right\|_B \\ &\leq \lim_{m \rightarrow \infty} \frac{1}{n^m} \phi(n^m x_1, n^m x_2, \dots, n^m x_n) = 0 \end{aligned} \quad (2.20)$$

for all  $x_1, \dots, x_n \in A$ . Then

$$H\left(\frac{\sum_{i=1}^n x_i}{n}\right) + \sum_{j=2}^n H\left(\frac{\sum_{i=1, i \neq j}^n x_i - (n-1)x_j}{n}\right) = H(x_1) \quad (2.21)$$

for all  $x_1, \dots, x_n \in A$ . Putting  $s_1 = \sum_{i=1}^n x_i/n$  and  $s_j = \sum_{i=1, i \neq j}^n x_i - (n-1)x_j/n$  ( $j = 2, 3, \dots, n$ ) in (2.21), we obtain

$$H\left(\sum_{j=1}^n s_j\right) = \sum_{j=1}^n H(s_j) \quad (2.22)$$

for all  $s_1, \dots, s_n \in A$ . Setting  $s_j = 0$  ( $j = 3, 4, \dots, n$ ) in (2.22) to get

$$H(s_1 + s_2) = H(s_1) + H(s_2) \quad (2.23)$$

hence  $H$  is cauchy additive. Letting  $x_i = x$  for all  $i = 1, 2, \dots, n$  in (2.2), we obtain

$$\|\mu f(x) - f(\mu x)\|_B \leq \phi(x, x, \dots, x) \quad (2.24)$$

for all  $x \in A$ . It follows that

$$\begin{aligned} \|H(\mu x) - \mu H(x)\| &= \lim_{m \rightarrow \infty} \frac{1}{n^m} \|f(\mu n^m x) - \mu f(n^m x)\|_B \\ &\leq \lim_{m \rightarrow \infty} \frac{1}{n^m} \phi(n^m x, n^m x, \dots, n^m x) = 0 \end{aligned} \quad (2.25)$$

for all  $\mu \in \mathbb{T}_{1/n_0}^1$ , and all  $x \in A$ . One can show that the mapping  $H : A \rightarrow B$  is  $\mathbb{C}$ -linear. Hence,  $H : A \rightarrow B$  is an  $n$ -Lie homomorphism satisfying (2.5), as desired.  $\square$

**Corollary 2.3.** *Let  $\theta$  and  $p$  be nonnegative real numbers such that  $p < 1$ . Suppose that a function  $f : A \rightarrow B$  satisfies*

$$\left\| \mu f\left(\frac{\sum_{i=1}^n x_i}{n}\right) + \mu \sum_{j=2}^n f\left(\frac{\sum_{i=1, i \neq j}^n x_i - (n-1)x_j}{n}\right) - f(\mu x_1) \right\|_B \leq \theta \sum_{i=1}^n (\|x_i\|_A^p) \quad (2.26)$$

for all  $\mu \in \mathbb{T}^1$  and all  $x_1, \dots, x_n \in A$  and

$$\|f([x_1 x_2 \cdots x_n]_A) - [f(x_1) f(x_2) \cdots f(x_n)]_B\|_B \leq \theta \sum_{i=1}^n (\|x_i\|_A^p) \quad (2.27)$$

for all  $x_1, \dots, x_n \in A$ . Then there exists a unique  $n$ -Lie homomorphism  $H : A \rightarrow B$  such that

$$\|f(x) - H(x)\|_B \leq \frac{2^p}{\varrho(2-2^p)} \theta \|x\|_A^p \quad (2.28)$$

for all  $x \in A$ .

*Proof.* Put  $\phi(x_1, x_2, \dots, x_n) := \theta \sum_{i=1}^n (\|x_i\|_A^p)$  for all  $x_1, \dots, x_n \in A$  in Theorem 2.2. Then (2.9) holds for  $p < 1$ , and (2.28) holds when  $L = 2^{(p-1)}$ .  $\square$

**Theorem 2.4.** *Let  $n_0 \in \mathbb{N}$  be a fixed positive integer number. Let  $f : A \rightarrow B$  be a function for which there exists a function  $\phi : A^n \rightarrow [0, \infty)$  such that*

$$\left\| \mu f\left(\frac{\sum_{i=1}^n x_i}{n}\right) + \mu \sum_{j=2}^n f\left(\frac{\sum_{i=1, i \neq j}^n x_i - (n-1)x_j}{n}\right) - f(\mu x_1) \right\|_B \leq \phi(x_1, x_2, \dots, x_n) \quad (2.29)$$

for all  $\mu \in (\mathbb{T}_{1/n_0}^1 := \{e^{i\theta}; 0 \leq \theta \leq 2\pi/n_0\} \cup \{1\})$  and all  $x_1, \dots, x_n \in A$ , and that

$$\|f([x x \cdots x]_A) - [f(x) f(x) \cdots f(x)]_B\|_B \leq \phi(x, x, \dots, x) \quad (2.30)$$

for all  $x \in A$ . If there exists an  $L < 1$  such that

$$\phi(x_1, x_2, \dots, x_n) \leq nL\phi\left(\frac{x_1}{n}, \frac{x_2}{n}, \dots, \frac{x_n}{n}\right) \quad (2.31)$$

for all  $x_1, \dots, x_n \in A$ , then there exists a unique Jordan  $n$ -Lie homomorphism  $H : A \rightarrow B$  such that

$$\|f(x) - H(x)\|_B \leq \frac{L}{1-L} \phi(x, 0, 0, \dots, 0) \quad (2.32)$$

for all  $x \in A$ .

*Proof.* By the same reasoning as the proof of Theorem 2.2, we can define the mapping

$$H(x) = \lim_{m \rightarrow \infty} \frac{1}{n^{m\ell}} f(n^{m\ell} x) \quad (2.33)$$

for all  $x \in A$ . Moreover, we can show that  $H$  is  $\mathbb{C}$ -linear. The inequality (2.30) follows that

$$\begin{aligned} & \|H([xx \cdots x]_A) - [H(x)H(x) \cdots H(x)]_B\|_B \\ &= \lim_{m \rightarrow \infty} \left\| \frac{1}{n^{nm}} H([n^m x \cdots n^m x]_A) - \frac{1}{n^{nm}} ([H(n^m x)H(n^m x) \cdots H(n^m x)]_B) \right\|_B \\ &\leq \lim_{m \rightarrow \infty} \frac{1}{n^{nm}} \phi(n^m x, n^m x, \dots, n^m x) = 0 \end{aligned} \quad (2.34)$$

for all  $x \in A$ . So

$$H([xx \cdots x]_A) = [H(x)H(x) \cdots H(x)]_B \quad (2.35)$$

for all  $x \in A$ . Hence  $H : A \rightarrow B$  is a Jordan  $n$ -Lie homomorphism satisfying (2.32).  $\square$

**Corollary 2.5.** *Let  $\theta$  and  $p$  be nonnegative real numbers such that  $p < 1$ . Suppose that a function  $f : A \rightarrow B$  satisfies*

$$\left\| \mu f\left(\frac{\sum_{i=1}^n x_i}{n}\right) + \mu \sum_{j=2}^n f\left(\frac{\sum_{i=1, i \neq j}^n x_i - (n-1)x_j}{n}\right) - f(\mu x_1) \right\|_B \leq \theta \sum_{i=1}^n (\|x_i\|_A^p) \quad (2.36)$$

for all  $\mu \in \mathbb{T}^1$  and all  $x_1, \dots, x_n \in A$  and

$$\|f([xx \cdots x]_A) - [f(x)f(x) \cdots f(x)]_B\|_B \leq n\theta (\|x\|_A^p) \quad (2.37)$$

for all  $x \in A$ . Then there exists a unique Jordan  $n$ -Lie homomorphism  $H : A \rightarrow B$  such that

$$\|f(x) - H(x)\|_B \leq \frac{2^p}{\varrho(2-2^p)} \theta \|x\|_A^p \quad (2.38)$$

for all  $x \in A$ .

*Proof.* It follows by Theorem 2.4 by putting  $\phi(x_1, x_2, \dots, x_n) := \theta \sum_{i=1}^n (\|x_i\|_A^p)$  for all  $x_1, \dots, x_n \in A$  and  $L = 2^{(p-1)}$ .  $\square$

## References

- [1] V. T. Filippov, "n-Lie algebras," *Sibirskiĭ Matematicheskiĭ Zhurnal*, vol. 26, no. 6, pp. 126–140, 1985.
- [2] Y. Nambu, "Generalized Hamiltonian dynamics," *Physical Review D*, vol. 7, pp. 2405–2412, 1973.



- [3] V. T. Filippov, "On the  $n$ -Lie algebra of Jacobians," *Sibirskit Matematicheskii Zhurnal*, vol. 39, no. 3, pp. 660–669, 1998.
- [4] Sh. M. Kasymov, "On a theory of  $n$ -Lie algebras," *Algebra i Logika*, vol. 26, no. 3, pp. 277–297, 1987.
- [5] Sh. M. Kasymov, "Nil-elements and nil-subsets in  $n$ -Lie algebras," *Sibirskit Matematicheskii Zhurnal*, vol. 32, no. 6, pp. 77–80, 1991.
- [6] L. Takhtajan, "On foundation of the generalized Nambu mechanics," *Communications in Mathematical Physics*, vol. 160, no. 2, pp. 295–315, 1994.
- [7] J. Bagger and N. Lambert, "Comments on multiple M2-branes," *Journal of High Energy Physics*, no. 2, article 105, 2008.
- [8] A. Gustavsson, "One-loop corrections to Bagger-Lambert theory," *Nuclear Physics B*, vol. 807, no. 1-2, pp. 315–333, 2009.
- [9] J. A. de Azcarraga and J. M. Izquierdo, " $n$ -ary algebras: a review with applications," *Journal of Physics A*, vol. 43, no. 29, Article ID 293001, 2010.
- [10] S. M. Ulam, *Problems in Modern Mathematics*, John Wiley & Sons, New York, NY, USA, 1940.
- [11] D. H. Hyers, "On the stability of the linear functional equation," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 27, pp. 222–224, 1941.
- [12] T. M. Rassias, "On the stability of the linear mapping in Banach spaces," *Proceedings of the American Mathematical Society*, vol. 72, no. 2, pp. 297–300, 1978.
- [13] M. B. Savadkouhi, M. E. Gordji, J. M. Rassias, and N. Ghobadipour, "Approximate ternary Jordan derivations on Banach ternary algebras," *Journal of Mathematical Physics*, vol. 50, no. 4, Article ID 042303, 9 pages, 2009.
- [14] St. Czerwik, "On the stability of the quadratic mapping in normed spaces," *Abhandlungen aus dem Mathematischen Seminar der Universität Hamburg*, vol. 62, pp. 59–64, 1992.
- [15] A. Ebadian, N. Ghobadipour, and M. Eshaghi Gordji, "A fixed point method for perturbation of bimultipliers and Jordan bimultipliers in  $C^*$ -ternary algebras," *Journal of Mathematical Physics*, vol. 51, no. 10, Article ID 103508, 10 pages, 2010.
- [16] M. E. Gordji, M. B. Savadkouhi, M. Bidkham, C. Park, and J. R. Lee, "Nearly partial derivations on Banach ternary algebras," *Journal of Mathematics and Statistics*, vol. 6, no. 4, pp. 454–461, 2010.
- [17] M. E. Gordji and N. Ghobadipour, "Stability of  $\alpha, \beta, \gamma$ -derivations on Lie  $C^*$ -algebras," *International Journal of Geometric Methods in Modern Physics*, vol. 7, no. 7, pp. 1093–1102, 2010.
- [18] M. E. Gordji, R. Khodabakhsh, and H. Khodaei, "On approximate  $n$ -ary derivations," *International Journal of Geometric Methods in Modern Physics*, vol. 8, no. 3, pp. 485–500, 2011.
- [19] M. E. Gordji and H. Khodaei, *Stability of Functional Equations*, LAP LAMBERT Academic Publishing, 2010.
- [20] Z. Gajda, "On stability of additive mappings," *International Journal of Mathematics and Mathematical Sciences*, vol. 14, no. 3, pp. 431–434, 1991.
- [21] R. Farokhzad and S. A. R. Hosseinioun, "Perturbations of Jordan higher derivations in Banach ternary algebras: an alternative fixed point approach," *International Journal of Nonlinear Analysis and Applications*, vol. 1, no. 1, pp. 42–53, 2010.
- [22] M. E. Gordji, S. K. Gharetapeh, E. Rashidi, T. Karimi, and M. Aghaei, "Ternary Jordan derivations in  $C^*$ -ternary algebras," *Journal of Computational Analysis and Applications*, vol. 12, no. 2, pp. 463–470, 2010.
- [23] S.-M. Jung, "Hyers-Ulam-Rassias stability of Jensen's equation and its application," *Proceedings of the American Mathematical Society*, vol. 126, no. 11, pp. 3137–3143, 1998.
- [24] S.-M. Jung, *Hyers-Ulam-Rassias Stability of Functional Equations in Nonlinear Analysis*, vol. 48 of *Springer Optimization and Its Applications*, Springer, New York, NY, USA, 2011.
- [25] H. Khodaei and Th. M. Rassias, "Approximately generalized additive functions in several variables," *International Journal of Nonlinear Analysis and Applications*, vol. 1, pp. 22–41, 2010.
- [26] J. M. Rassias, "On approximation of approximately linear mappings by linear mappings," *Bulletin des Sciences Mathématiques. 2e Série*, vol. 108, no. 4, pp. 445–446, 1984.
- [27] J. M. Rassias, "On approximation of approximately linear mappings by linear mappings," *Journal of Functional Analysis*, vol. 46, no. 1, pp. 126–130, 1982.
- [28] J. M. Rassias, "On the stability of the non-linear Euler-Lagrange functional equation in real normed linear spaces," *Journal of Mathematical and Physical Sciences*, vol. 28, no. 5, pp. 231–235, 1994.
- [29] Th. M. Rassias and J. Brzdek, Eds., *Functional Equations in Mathematical Analysis*, Springer, New York, NY, USA, 2012.
- [30] T. M. Rassias, "On the stability of functional equations in Banach spaces," *Journal of Mathematical Analysis and Applications*, vol. 251, no. 1, pp. 264–284, 2000.

- [31] T. M. Rassias, "On the stability of functional equations and a problem of Ulam," *Acta Applicandae Mathematicae*, vol. 62, no. 1, pp. 23–130, 2000.
- [32] T. M. Rassias and P. Šemrl, "On the behavior of mappings which do not satisfy Hyers-Ulam stability," *Proceedings of the American Mathematical Society*, vol. 114, no. 4, pp. 989–993, 1992.
- [33] P. Kannappan, *Functional Equations and Inequalities with Applications*, Springer Monographs in Mathematics, Springer, New York, NY, USA, 2009.
- [34] H.-M. Kim, J. M. Rassias, and Y.-S. Cho, "Stability problem of Ulam for Euler-Lagrange quadratic mappings," *Journal of Inequalities and Applications*, vol. 2007, Article ID 10725, 15 pages, 2007.
- [35] J. M. Rassias, "Complete solution of the multi-dimensional problem of Ulam," *Discussiones Mathematicae*, vol. 14, pp. 101–107, 1994.
- [36] P. Gavruta, "An answer to a question of John M. Rassias concerning the stability of Cauchy equation," in *Advances in Equations and Inequalities*, Hadronic Math. Ser., pp. 67–71, Hadronic Press, 1999.
- [37] A. Pietrzyk, "Stability of the Euler-Lagrange-Rassias functional equation," *Demonstratio Mathematica*, vol. 39, no. 3, pp. 523–530, 2006.
- [38] Y.-S. Lee and S.-Y. Chung, "Stability of an Euler-Lagrange-Rassias equation in the spaces of generalized functions," *Applied Mathematics Letters*, vol. 21, no. 7, pp. 694–700, 2008.
- [39] P. Nakmahachalasint, "On the generalized Ulam-Gavruta-Rassias stability of mixed-type linear and Euler-Lagrange-Rassias functional equations," *International Journal of Mathematics and Mathematical Sciences*, vol. 2007, Article ID 63239, 10 pages, 2007.
- [40] K.-W. Jun, H.-M. Kim, and J. M. Rassias, "Extended Hyers-Ulam stability for Cauchy-Jensen mappings," *Journal of Difference Equations and Applications*, vol. 13, no. 12, pp. 1139–1153, 2007.
- [41] J. M. Rassias and M. J. Rassias, "On some approximately quadratic mappings being exactly quadratic," *The Journal of the Indian Mathematical Society*, vol. 69, no. 1–4, pp. 155–160, 2002.
- [42] V. Fatziev and J. M. Rassias, "Stability of generalized additive equations on Banach spaces and groups," *Journal of Nonlinear Functional Analysis and Differential Equations*, vol. 1, no. 2, pp. 153–173, 2007.
- [43] M. Eshaghi Gordji, J. M. Rassias, and N. Ghobadipour, "Generalized hyers-ulam stability of generalized  $(N,K)$ -derivations," *Abstract and Applied Analysis*, vol. 2009, Article ID 437931, 8 pages, 2009.
- [44] M. Eshaghi Gordji, S. Kaboli Gharetapeh, J. M. Rassias, and S. Zolfaghari, "Solution and stability of a mixed type additive, quadratic, and cubic functional equation," *Advances in Difference Equations*, vol. 2009, Article ID 826130, 17 pages, 2009.
- [45] J. M. Rassias, J. Lee, and H. M. Kim, "Refined Hyers-Ulam stability for Jensen type mappings," *Journal of the Chungcheong Mathematical Society*, vol. 22, no. 1, pp. 101–116, 2009.
- [46] M. E. Gordji, J. M. Rassias, and M. B. Savadkouhi, "Approximation of the quadratic and cubic functional equations in RN-spaces," *European Journal of Pure and Applied Mathematics*, vol. 2, no. 4, pp. 494–507, 2009.
- [47] J. M. Rassias, "Solution of the Ulam stability problem for quartic mappings," *Glasnik Matematički. Serija III*, vol. 34(54), no. 2, pp. 243–252, 1999.
- [48] B. Bouikhalene, E. Elqorachi, and J. M. Rassias, "The superstability of d'Alembert's functional equation on the Heisenberg group," *Applied Mathematics Letters*, vol. 23, no. 1, pp. 105–109, 2010.
- [49] M. Eshaghi Gordji, S. Zolfaghari, J. M. Rassias, and M. B. Savadkouhi, "Solution and stability of a mixed type cubic and quartic functional equation in quasi-Banach spaces," *Abstract and Applied Analysis*, vol. 2009, Article ID 417473, 14 pages, 2009.
- [50] M. E. Gordji and H. Khodaei, "On the generalized Hyers-Ulam-Rassias stability of quadratic functional equations," *Abstract and Applied Analysis*, vol. 2009, Article ID 923476, 11 pages, 2009.
- [51] K. Ravi and M. Arunkumar, "On the Ulam-Gavruta-Rassias stability of the orthogonally Euler-Lagrange type functional equation," *International Journal of Applied Mathematics & Statistics*, vol. 7, no. Fe07, pp. 143–156, 2007.
- [52] P. Nakmahachalasint, "Hyers-Ulam-Rassias and Ulam-Gavruta-Rassias stabilities of an additive functional equation in several variables," *International Journal of Mathematics and Mathematical Sciences*, vol. 2007, Article ID 13437, 6 pages, 2007.
- [53] M. A. SiBaha, B. Bouikhalene, and E. Elqorachi, "Ulam-Găvrută-Rassias stability of a linear functional equation," *International Journal of Applied Mathematics & Statistics*, vol. 7, no. Fe07, pp. 157–166, 2007.
- [54] B. Bouikhalene and E. Elqorachi, "Ulam-Găvruta-Rassias stability of the Pexider functional equation," *International Journal of Applied Mathematics & Statistics*, vol. 7, no. Fe07, pp. 27–39, 2007.

- [55] K. Ravi, M. Arunkumar, and J. M. Rassias, "Ulam stability for the orthogonally general Euler-Lagrange type functional equation," *International Journal of Mathematics and Statistics*, vol. 3, no. A08, pp. 36–46, 2008.
- [56] H.-X. Cao, J.-R. Lv, and J. M. Rassias, "Superstability for generalized module left derivations and generalized module derivations on a Banach module. I," *Journal of Inequalities and Applications*, vol. 2009, Article ID 718020, 10 pages, 2009.
- [57] H.-X. Cao, J.-R. Lv, and J. M. Rassias, "Superstability for generalized module left derivations and generalized module derivations on a Banach module. II," *Journal of Inequalities in Pure and Applied Mathematics*, vol. 10, no. 3, article 85, 2009.
- [58] L. Cădariu and V. Radu, "Fixed points and the stability of Jensen's functional equation," *Journal of Inequalities in Pure and Applied Mathematics*, vol. 4, no. 1, article 4, 2003.
- [59] C. Park and J. M. Rassias, "Stability of the Jensen-type functional equation in  $C^*$ -algebras: a fixed point approach," *Abstract and Applied Analysis*, vol. 2009, Article ID 360432, 17 pages, 2009.
- [60] J. B. Diaz and B. Margolis, "A fixed point theorem of the alternative, for contractions on a generalized complete metric space," *Bulletin of the American Mathematical Society*, vol. 74, pp. 305–309, 1968.
- [61] L. Cădariu and V. Radu, "On the stability of the Cauchy functional equation: a fixed point approach," in *Iteration Theory*, vol. 346 of *Grazer Math. Ber.*, pp. 43–52, Karl-Franzens-Univ. Graz, Graz, Austria, 2004.



# Hindawi

Submit your manuscripts at  
<http://www.hindawi.com>

