

Research Article

Iterative Schemes for Fixed Point Computation of Nonexpansive Mappings

Rudong Chen

Department of Mathematics, Tianjin Polytechnic University, Tianjin 300387, China

Correspondence should be addressed to Rudong Chen, chenrd@tjpu.edu.cn

Received 18 December 2011; Accepted 3 February 2012

Academic Editor: Muhammad Aslam Noor

Copyright © 2012 Rudong Chen. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Fixed point (especially, the minimum norm fixed point) computation is an interesting topic due to its practical applications in natural science. The purpose of the paper is devoted to finding the common fixed points of an infinite family of nonexpansive mappings. We introduce an iterative algorithm and prove that suggested scheme converges strongly to the common fixed points of an infinite family of nonexpansive mappings under some mild conditions. As a special case, we can find the minimum norm common fixed point of an infinite family of nonexpansive mappings.

1. Introduction

In many problems, it is needed to find a solution with minimum norm. In an abstract way, we may formulate such problems as finding a point x^\dagger with the property

$$x^\dagger \in C, \quad \|x^\dagger\|^2 = \min_{x \in C} \|x\|^2, \quad (1.1)$$

where C is a nonempty closed convex subset of a real Hilbert space H . In other words, x^\dagger is the (nearest point or metric) projection of the origin onto C ,

$$x^\dagger = P_C(0), \quad (1.2)$$

where P_C is the metric (or nearest point) projection from H onto C .

A typical example is the least-squares solution to the constrained linear inverse problem [1]

$$Ax = b, \quad x \in C, \quad (1.3)$$

where A is a bounded linear operator from H to another real Hilbert space H_1 and b is a given point in H_1 . The least-squares solution to (1.3) is the least-norm minimizer of the minimization problem

$$\min_{x \in C} \|Ax - b\|^2. \quad (1.4)$$

Recently, some authors consider the minimum norm solution problem by using the iterative algorithm. For some related works, please refer to [2–4]. Yao and Xu [3] introduced the following algorithm:

$$x_{n+1} = P_C(\alpha_n f(x_n) + (1 - \alpha_n)Tx_n), \quad n \geq 0. \quad (1.5)$$

They proved that the sequence $\{x_n\}$ converges in norm to the unique solution \tilde{x} of VI $\langle (I - f)\tilde{x}, x - \tilde{x} \rangle \geq 0, x \in \text{Fix}(T)$. Particularly, the sequence $\{x_n\}$ defined by

$$x_{n+1} = P_C((1 - \alpha_n)Tx_n), \quad n \geq 0, \quad (1.6)$$

converges to the minimum norm fixed point of T . We note that the authors added an additional assumption, that is, $\lim_{n \rightarrow \infty} \alpha_{n+1}/\alpha_n = 1$. Iterative algorithm for finding the fixed points of nonexpansive mappings has been considered by many authors, see [5–20].

The purpose of this paper is to extend Yao and Xu's result to an infinite family of nonexpansive mappings $\{T_n\}_{n=0}^{\infty}$. We suggest a new algorithm. Particularly, we drop the above additional assumption and prove the suggested algorithm converges strongly to the common fixed points of $\{T_n\}_{n=0}^{\infty}$. As a special case, we can find the minimum norm fixed point of $\{T_n\}_{n=0}^{\infty}$.

2. Preliminaries

Let H be a real Hilbert space with inner product $\langle \cdot, \cdot \rangle$ and norm $\|\cdot\|$, respectively, and let C be a nonempty closed convex subset of H . We call $f : C \rightarrow H$ a κ -contraction if there exists a constant $\kappa \in [0, 1)$ such that $\|f(x) - f(y)\| \leq \kappa\|x - y\|$ for all $x, y \in C$. A bounded linear operator B is said to be strongly positive on H if there exists a constant $\alpha > 0$ such that

$$\langle Bx, x \rangle \geq \alpha\|x\|^2, \quad \forall x \in H. \quad (2.1)$$

Recall that the (nearest point or metric) projection from H onto C , denoted by P_C , is defined in such a way that, for each $x \in H$, $P_C x$ is the unique point in C with the property

$$\|x - P_C x\| = \min\{\|x - y\| : y \in C\}. \quad (2.2)$$

It is known that P_C satisfies

$$\langle x - y, P_C x - P_C y \rangle \geq \|P_C x - P_C y\|^2, \quad \forall x, y \in H. \quad (2.3)$$

Moreover, P_C is characterized by the following properties:

$$\begin{aligned} \langle x - P_Cx, y - P_Cx \rangle &\leq 0, \\ \|x - y\|^2 &\geq \|x - P_Cx\|^2 + \|y - P_Cx\|^2, \end{aligned} \tag{2.4}$$

for all $x \in H$ and $y \in C$.

We also need other sorts of nonlinear operators which are introduced below. Let $T : H \rightarrow H$ be a nonlinear operator.

- (a) T is nonexpansive if $\|Tx - Ty\| \leq \|x - y\|$ for all $x, y \in H$.
- (b) T is firmly nonexpansive if $2T - I$ is nonexpansive. Equivalently, $T = (I + S)/2$, where $S : H \rightarrow H$ is nonexpansive. Alternatively, T is firmly nonexpansive if and only if

$$\|Tx - Ty\|^2 \leq \langle Tx - Ty, x - y \rangle, \quad x, y \in H. \tag{2.5}$$

- (c) T is averaged if $T = (1 - \tau)I + \tau S$, where $\tau \in (0, 1)$ and $S : H \rightarrow H$ is nonexpansive. In this case, we also say that T is τ -averaged. A firmly nonexpansive mapping is $1/2$ -averaged.

It is well known that both P_C and $I - P_C$ are firmly nonexpansive. We will need to use the following notation:

- (i) $\text{Fix}(T)$ stands for the set of fixed points of T ;
- (ii) $x_n \rightharpoonup x$ stands for the weak convergence of $\{x_n\}$ to x ;
- (iii) $x_n \rightarrow x$ stands for the strong convergence of $\{x_n\}$ to x .

Let T_1, T_2, \dots be infinite mappings of C into itself, and let ξ_1, ξ_2, \dots be real numbers such that $0 \leq \xi_i \leq 1$ for every $i \in \mathbf{N}$. For any $n \in \mathbf{N}$, define a mapping W_n of C into itself as follows:

$$\begin{aligned} U_{n,n+1} &= I, \\ U_{n,n} &= \xi_n T_n U_{n,n+1} + (1 - \xi_n)I, \\ U_{n,n-1} &= \xi_{n-1} T_{n-1} U_{n,n} + (1 - \xi_{n-1})I, \\ &\vdots \\ U_{n,k} &= \xi_k T_k U_{n,k+1} + (1 - \xi_k)I, \\ U_{n,k-1} &= \xi_{k-1} T_{k-1} U_{n,k} + (1 - \xi_{k-1})I, \\ &\vdots \\ U_{n,2} &= \xi_2 T_2 U_{n,3} + (1 - \xi_2)I, \\ W_n = U_{n,1} &= \xi_1 T_1 U_{n,2} + (1 - \xi_1)I. \end{aligned} \tag{2.6}$$

Such W_n is called the W -mapping generated by $T_n, T_{n-1}, \dots, T_2, T_1$ and $\xi_n, \xi_{n-1}, \dots, \xi_2, \xi_1$. For the iterative algorithm for a finite family of nonexpansive mappings, we refer the reader to [21].

We have the following crucial lemmas concerning W_n which can be found in [22].

Lemma 2.1. *Let C be a nonempty closed convex subset of a real Hilbert space H . Let T_1, T_2, \dots be nonexpansive mappings of C into itself such that $\bigcap_{n=1}^{\infty} F(T_n)$ is nonempty, and let ξ_1, ξ_2, \dots be real numbers such that $0 < \xi_i \leq b < 1$ for any $i \in \mathbf{N}$. Then, for every $x \in C$ and $k \in \mathbf{N}$, the limit $\lim_{n \rightarrow \infty} U_{n,k}x$ exists.*

Lemma 2.2. *Let C be a nonempty closed convex subset of a real Hilbert space H . Let T_1, T_2, \dots be nonexpansive mappings of C into itself such that $\bigcap_{n=1}^{\infty} F(T_n)$ is nonempty, and let ξ_1, ξ_2, \dots be real numbers such that $0 < \xi_i \leq b < 1$ for any $i \in \mathbf{N}$. Then, $F(W) = \bigcap_{n=1}^{\infty} F(T_n)$.*

The following remark [23] is important to prove our main results.

Remark 2.3. Using Lemma 2.1, one can define a mapping W of C into itself as $Wx = \lim_{n \rightarrow \infty} W_n x = \lim_{n \rightarrow \infty} U_{n,1}x$, for every $x \in C$. If $\{x_n\}$ is a bounded sequence in C , then one has

$$\lim_{n \rightarrow \infty} \|Wx_n - W_n x_n\| = 0. \quad (2.7)$$

Throughout this paper, we will assume that $0 < \xi_i \leq b < 1$ for every $i \in \mathbf{N}$.

Lemma 2.4 (see [24]). *Let K be a nonempty closed convex subset of a real Hilbert space H . Let $T : K \rightarrow K$ be a nonexpansive mapping with $\text{Fix}(T) \neq \emptyset$. Then T is demiclosed on K , that is, if $x_n \rightharpoonup x \in K$ weakly and $x_n - Tx_n \rightarrow 0$, then $x = Tx$.*

Lemma 2.5 (see [25]). *Let $\{x_n\}$ and $\{z_n\}$ be bounded sequences in a Banach space X , and let $\{\beta_n\}$ be a sequence in $[0, 1]$ with $0 < \liminf_{n \rightarrow \infty} \beta_n \leq \limsup_{n \rightarrow \infty} \beta_n < 1$. Suppose $x_{n+1} = (1 - \beta_n)z_n + \beta_n x_n$ for all integers $n \geq 0$ and $\limsup_{n \rightarrow \infty} (\|z_{n+1} - z_n\| - \|x_{n+1} - x_n\|) \leq 0$. Then, $\lim_{n \rightarrow \infty} \|z_n - x_n\| = 0$.*

Lemma 2.6 (see [26]). *Assume $\{a_n\}$ is a sequence of nonnegative real numbers such that $a_{n+1} \leq (1 - \gamma_n)a_n + \delta_n$ where $\{\gamma_n\}$ is a sequence in $(0, 1)$ and $\{\delta_n\}$ is a sequence such that*

- (1) $\sum_{n=1}^{\infty} \gamma_n = \infty$;
- (2) $\limsup_{n \rightarrow \infty} \delta_n / \gamma_n \leq 0$ or $\sum_{n=1}^{\infty} |\delta_n| < \infty$.

Then $\lim_{n \rightarrow \infty} a_n = 0$.

3. Main Result

In this section, we introduce our algorithm and prove its strong convergence.

Theorem 3.1. *Let C be a nonempty closed convex subset of a real Hilbert space H . Let $\{T_n\}_{n=1}^{\infty}$ be a sequence of nonexpansive mappings from C to C such that the common fixed point set $F := \bigcap_{n=1}^{\infty} F(T_n) \neq \emptyset$. Let $f : C \rightarrow H$ be a κ -contraction and $B : H \rightarrow H$ be a self-adjoint, strongly*

positive bounded linear operator with coefficient $\alpha > 0$. Let σ be a constant such that $0 < \sigma\kappa < \alpha$. For an arbitrary initial point x_0 belonging to C , one defines a sequence $\{x_n\}_{n \geq 0}$ iteratively

$$x_{n+1} = P_C [\alpha_n \sigma f(x_n) + (I - \alpha_n B) W_n x_n], \quad \forall n \geq 0, \quad (3.1)$$

where $\{\alpha_n\}$ is a real sequence in $[0, 1]$. Assume the sequence $\{\alpha_n\}$ satisfies the following conditions:

$$(C1) \lim_{n \rightarrow \infty} \alpha_n = 0;$$

$$(C2) \sum_{n=0}^{\infty} \alpha_n = \infty.$$

Then the sequence $\{x_n\}$ generated by (3.1) converges in norm to the unique solution x^* which solves the following variational inequality:

$$x^* \in F \text{ such that } \langle \sigma f(x^*) - Bx^*, \tilde{x} - x^* \rangle \leq 0, \quad \forall \tilde{x} \in F. \quad (3.2)$$

Proof. Let $\tilde{x} \in F$. From (3.1), we have

$$\begin{aligned} \|x_{n+1} - \tilde{x}\| &= \|P_C [\alpha_n \sigma f(x_n) + (I - \alpha_n B) W_n x_n] - \tilde{x}\| \\ &\leq \|\alpha_n \sigma f(x_n) + (I - \alpha_n B) W_n x_n - \tilde{x}\| \\ &\leq \alpha_n \sigma \|f(x_n) - f(\tilde{x})\| + \|I - \alpha_n B\| \|W_n x_n - \tilde{x}\| + \alpha_n \|\sigma f(\tilde{x}) - B\tilde{x}\| \\ &\leq \alpha_n \sigma \kappa \|x_n - \tilde{x}\| + (1 - \alpha_n \alpha) \|x_n - \tilde{x}\| + \alpha_n \|\sigma f(\tilde{x}) - B\tilde{x}\| \\ &= [1 - (\alpha - \sigma\kappa)\alpha_n] \|x_n - \tilde{x}\| + \frac{(\alpha - \sigma\kappa)\alpha_n \|f(\tilde{x}) - B\tilde{x}\|}{(\alpha - \sigma\kappa)}. \end{aligned} \quad (3.3)$$

It follows by induction that

$$\begin{aligned} \|x_{n+1} - \tilde{x}\| &\leq \max \left\{ \|x_n - \tilde{x}\|, \frac{\|f(\tilde{x}) - B\tilde{x}\|}{(\alpha - \sigma\kappa)} \right\} \\ &\leq \max \left\{ \|x_0 - \tilde{x}\|, \frac{\|f(\tilde{x}) - B\tilde{x}\|}{(\alpha - \sigma\kappa)} \right\}. \end{aligned} \quad (3.4)$$

This indicates that $\{x_n\}$ is bounded. It is easy to deduce that $\{f(x_n)\}$, $\{W_n x_n\}$, and $\{B W_n x_n\}$ are also bounded.

Set $S = 2P_C - I$. It is known that S is nonexpansive. Note that $W_n = \xi_1 T_1 U_{n,2} x_n + (1 - \xi_1) x_n$. Then, we can rewrite (3.1) as

$$\begin{aligned}
 x_{n+1} &= \frac{I+S}{2} [\alpha_n \sigma f(x_n) + (I - \alpha_n B) W_n x_n] \\
 &= \frac{1 - \alpha_n}{2} W_n x_n + \frac{\alpha_n}{2} (\sigma f(x_n) - B W_n x_n + W_n x_n) \\
 &\quad + \frac{1}{2} S [\alpha_n \sigma f(x_n) + (I - \alpha_n B) W_n x_n] \\
 &= \frac{1 - \alpha_n}{2} [(1 - \xi) I + \xi T_1 U_{n,2}] x_n + \frac{\alpha_n}{2} (\sigma f(x_n) - B W_n x_n + W_n x_n) \\
 &\quad + \frac{1}{2} S [\alpha_n \sigma f(x_n) + (I - \alpha_n B) W_n x_n] \\
 &= \frac{(1 - \xi)(1 - \alpha_n)}{2} x_n + \frac{\xi(1 - \alpha_n)}{2} T_1 U_{n,2} x_n + \frac{\alpha_n}{2} (\sigma f(x_n) - B W_n x_n + W_n x_n) \\
 &\quad + \frac{1}{2} S [\alpha_n \sigma f(x_n) + (I - \alpha_n B) W_n x_n].
 \end{aligned} \tag{3.5}$$

Note that

$$\begin{aligned}
 0 < \lim_{n \rightarrow \infty} \frac{(1 - \xi)(1 - \alpha_n)}{2} &= \frac{1 - \xi}{2} < 1, \\
 \frac{\xi(1 - \alpha_n)}{2} + \frac{1}{2} &= \frac{1 + \xi}{2} - \frac{\xi}{2} \alpha_n.
 \end{aligned} \tag{3.6}$$

From (3.5), we have

$$\begin{aligned}
 x_{n+1} &= \left[1 - \left(\frac{1 + \xi}{2} + \frac{1 - \xi}{2} \alpha_n \right) \right] x_n + \left(\frac{1 + \xi}{2} + \frac{1 - \xi}{2} \alpha_n \right) \\
 &\quad \times \frac{(\xi(1 - \alpha_n)/2) T_1 U_{n,2} x_n + (\alpha_n/2) (\sigma f(x_n) - B W_n x_n + W_n x_n)}{((1 + \xi)/2) + ((1 - \xi)/2) \alpha_n} \\
 &\quad + \frac{(1/2) S [\alpha_n \sigma f(x_n) + (I - \alpha_n B) W_n x_n]}{((1 + \xi)/2) + ((1 - \xi)/2) \alpha_n} \\
 &= \left[1 - \left(\frac{1 + \xi}{2} + \frac{1 - \xi}{2} \alpha_n \right) \right] x_n + \left(\frac{1 + \xi}{2} + \frac{1 - \xi}{2} \alpha_n \right) y_n,
 \end{aligned} \tag{3.7}$$

where

$$\begin{aligned}
 y_n &= \frac{(\xi(1-\alpha_n)/2)T_1U_{n,2}x_n + (\alpha_n/2)(\sigma f(x_n) - BW_nx_n + W_nx_n)}{((1+\xi)/2) + ((1-\xi)/2)\alpha_n} \\
 &\quad + \frac{(1/2)S[\alpha_n\sigma f(x_n) + (I - \alpha_nB)W_nx_n]}{((1+\xi)/2) + ((1-\xi)/2)\alpha_n} \\
 &= \frac{\xi(1-\alpha_n)T_1U_{n,2}x_n + \alpha_n(\sigma f(x_n) - BW_nx_n + W_nx_n) + S[\alpha_n\sigma f(x_n) + (I - \alpha_nB)W_nx_n]}{1 + \xi + (1-\xi)\alpha_n}.
 \end{aligned}
 \tag{3.8}$$

Set $z_n = \sigma f(x_n) - BW_nx_n + W_nx_n$ and $\tilde{z}_n = \alpha_n\sigma f(x_n) + (I - \alpha_nB)W_nx_n$ for all n . Then

$$y_n = \frac{\xi(1-\alpha_n)T_1U_{n,2}x_n + \alpha_nz_n + S\tilde{z}_n}{1 + \xi + (1-\xi)\alpha_n}, \quad \forall n \geq 0.
 \tag{3.9}$$

It follows that

$$\begin{aligned}
 y_{n+1} - y_n &= \frac{\xi(1-\alpha_{n+1})T_1U_{n+1,2}x_{n+1} + \alpha_{n+1}z_{n+1} + S\tilde{z}_{n+1}}{1 + \xi + (1-\xi)\alpha_{n+1}} \\
 &\quad - \frac{\xi(1-\alpha_n)T_1U_{n,2}x_n + \alpha_nz_n + S\tilde{z}_n}{1 + \xi + (1-\xi)\alpha_n} \\
 &= \frac{\xi(1-\alpha_{n+1})}{1 + \xi + (1-\xi)\alpha_{n+1}}(T_1U_{n+1,2}x_{n+1} - T_1U_{n,2}x_n) \\
 &\quad + \left(\frac{\xi(1-\alpha_{n+1})}{1 + \xi + (1-\xi)\alpha_{n+1}} - \frac{\xi(1-\alpha_n)}{1 + \xi + (1-\xi)\alpha_n} \right) T_1U_{n,2}x_n \\
 &\quad + \frac{\alpha_{n+1}z_{n+1}}{1 + \xi + (1-\xi)\alpha_{n+1}} - \frac{\alpha_nz_n}{1 + \xi + (1-\xi)\alpha_n} \\
 &\quad + \frac{S\tilde{z}_{n+1} - S\tilde{z}_n}{1 + \xi + (1-\xi)\alpha_{n+1}} + \left(\frac{1}{1 + \xi + (1-\xi)\alpha_{n+1}} - \frac{1}{1 + \xi + (1-\xi)\alpha_n} \right) S\tilde{z}_n.
 \end{aligned}
 \tag{3.10}$$

Thus,

$$\begin{aligned}
 \|y_{n+1} - y_n\| &\leq \frac{\xi(1-\alpha_{n+1})}{1 + \xi + (1-\xi)\alpha_{n+1}} \|T_1U_{n+1,2}x_{n+1} - T_1U_{n,2}x_n\| \\
 &\quad + \left| \frac{\xi(1-\alpha_{n+1})}{1 + \xi + (1-\xi)\alpha_{n+1}} - \frac{\xi(1-\alpha_n)}{1 + \xi + (1-\xi)\alpha_n} \right| \|T_1U_{n,2}x_n\|
 \end{aligned}$$

$$\begin{aligned}
& + \frac{\alpha_{n+1}}{1 + \xi + (1 - \xi)\alpha_{n+1}} \|z_{n+1}\| + \frac{\alpha_n}{1 + \xi + (1 - \xi)\alpha_n} \|z_n\| \\
& + \frac{1}{1 + \xi + (1 - \xi)\alpha_{n+1}} \|S\tilde{z}_{n+1} - S\tilde{z}_n\| \\
& + \left| \frac{1}{1 + \xi + (1 - \xi)\alpha_{n+1}} - \frac{1}{1 + \xi + (1 - \xi)\alpha_n} \right| \|S\tilde{z}_n\|.
\end{aligned} \tag{3.11}$$

From the nonexpansivity of S , we get

$$\begin{aligned}
\|S\tilde{z}_{n+1} - S\tilde{z}_n\| & \leq \|\tilde{z}_{n+1} - \tilde{z}_n\| \\
& = \|\alpha_{n+1}\sigma f(x_{n+1}) + (I - \alpha_{n+1}B)W_{n+1}x_{n+1} - (\alpha_n\sigma f(x_n) + (I - \alpha_nB)W_nx_n)\| \\
& \leq \alpha_{n+1}\|\sigma f(x_{n+1}) - BW_{n+1}x_{n+1}\| + \alpha_n\|\sigma f(x_n) - BW_nx_n\| + \|W_{n+1}x_{n+1} - W_nx_n\| \\
& \leq \alpha_{n+1}\|\sigma f(x_{n+1}) - BW_{n+1}x_{n+1}\| + \alpha_n\|\sigma f(x_n) - BW_nx_n\| \\
& \quad + \|W_{n+1}x_{n+1} - W_{n+1}x_n\| + \|W_{n+1}x_n - W_nx_n\| \\
& \leq \alpha_{n+1}\|\sigma f(x_{n+1}) - BW_{n+1}x_{n+1}\| + \alpha_n\|\sigma f(x_n) - BW_nx_n\| + \|x_{n+1} - x_n\| \\
& \quad + \|W_{n+1}x_n - W_nx_n\|.
\end{aligned} \tag{3.12}$$

Since T_i and $U_{n,i}$ are nonexpansive, we have

$$\begin{aligned}
\|T_1U_{n+1,2}x_n - T_1U_{n,2}x_n\| & \leq \|U_{n+1,2}x_n - U_{n,2}x_n\| \\
& = \|\xi_2T_2U_{n+1,3}x_n - \xi_2T_2U_{n,3}x_n\| \\
& \leq \xi_2\|U_{n+1,3}x_n - U_{n,3}x_n\| \\
& \leq \dots \\
& \leq \xi_2 \cdots \xi_n \|U_{n+1,n+1}x_n - U_{n,n+1}x_n\| \\
& \leq M \prod_{i=2}^n \xi_i,
\end{aligned} \tag{3.13}$$

where $M > 0$ is a constant such that $\|U_{n+1,n+1}x_n - U_{n,n+1}x_n\| \leq M$ for all $n \geq 0$. So,

$$\begin{aligned}
\|T_1U_{n+1,2}x_{n+1} - T_1U_{n,2}x_n\| & \leq \|T_1U_{n+1,2}x_{n+1} - T_1U_{n+1,2}x_n\| + \|T_1U_{n+1,2}x_n - T_1U_{n,2}x_n\| \\
& \leq \|x_{n+1} - x_n\| + M \prod_{i=2}^n \xi_i.
\end{aligned} \tag{3.14}$$

Hence,

$$\begin{aligned}
 \|y_{n+1} - y_n\| &\leq \frac{\xi(1 - \alpha_{n+1})}{1 + \xi + (1 - \xi)\alpha_{n+1}} \|x_{n+1} - x_n\| + M \prod_{i=2}^n \xi_i \\
 &+ \left| \frac{\xi(1 - \alpha_{n+1})}{1 + \xi + (1 - \xi)\alpha_{n+1}} - \frac{\xi(1 - \alpha_n)}{1 + \xi + (1 - \xi)\alpha_n} \right| \|T_1 U_{n,2} x_n\| \\
 &+ \frac{\alpha_{n+1}}{1 + \xi + (1 - \xi)\alpha_{n+1}} \|z_{n+1}\| + \frac{\alpha_n}{1 + \xi + (1 - \xi)\alpha_n} \|z_n\| \\
 &+ \frac{1}{1 + \xi + (1 - \xi)\alpha_{n+1}} \\
 &\times (\alpha_{n+1} \|\sigma f(x_{n+1}) - BW_{n+1} x_{n+1}\| + \alpha_n \|\sigma f(x_n) - BW_n x_n\| + \|x_{n+1} - x_n\|) \\
 &+ \|W_{n+1} x_n - W_n x_n\| + \left| \frac{1}{1 + \xi + (1 - \xi)\alpha_{n+1}} - \frac{1}{1 + \xi + (1 - \xi)\alpha_n} \right| \|S\tilde{z}_n\| \\
 &= \frac{1 + \xi - \xi\alpha_{n+1}}{1 + \xi + (1 - \xi)\alpha_{n+1}} \|x_{n+1} - x_n\| \\
 &+ \left| \frac{\xi(1 - \alpha_{n+1})}{1 + \xi + (1 - \xi)\alpha_{n+1}} - \frac{\xi(1 - \alpha_n)}{1 + \xi + (1 - \xi)\alpha_n} \right| \|T_1 U_{n,2} x_n\| \\
 &+ \frac{\alpha_{n+1}}{1 + \xi + (1 - \xi)\alpha_{n+1}} \|z_{n+1}\| + \frac{\alpha_n}{1 + \xi + (1 - \xi)\alpha_n} \|z_n\| \\
 &+ \frac{1}{1 + \xi + (1 - \xi)\alpha_{n+1}} \\
 &\times (\alpha_{n+1} \|\sigma f(x_{n+1}) - BW_{n+1} x_{n+1}\| + \alpha_n \|\sigma f(x_n) - BW_n x_n\|) \\
 &+ \|W_{n+1} x_n - W_n x_n\| + \left| \frac{1}{1 + \xi + (1 - \xi)\alpha_{n+1}} - \frac{1}{1 + \xi + (1 - \xi)\alpha_n} \right| \|S\tilde{z}_n\|.
 \end{aligned} \tag{3.15}$$

Since $\alpha_n \rightarrow 0$, we have $\xi(1 - \alpha_{n+1})/(1 + \xi + (1 - \xi)\alpha_{n+1}) - \xi(1 - \alpha_n)/(1 + \xi + (1 - \xi)\alpha_n) \rightarrow 0$, $\|W_{n+1} x_n - W_n x_n\| \rightarrow 0$, and $1/(1 + \xi + (1 - \xi)\alpha_{n+1}) - 1/(1 + \xi + (1 - \xi)\alpha_n) \rightarrow 0$. Therefore,

$$\limsup_{n \rightarrow \infty} (\|y_{n+1} - y_n\| - \|x_{n+1} - x_n\|) \leq 0. \tag{3.16}$$

By Lemma 2.5, we get

$$\lim_{n \rightarrow \infty} \|y_n - x_n\| = 0. \tag{3.17}$$

Hence, from (3.7), we deduce

$$\lim_{n \rightarrow \infty} \|x_{n+1} - x_n\| = \lim_{n \rightarrow \infty} \left(\frac{1 + \xi}{2} + \frac{1 - \xi}{2} \alpha_n \right) \|y_n - x_n\| = 0. \tag{3.18}$$

Observe that

$$\begin{aligned} \|x_{n+1} - x_n\| &= \|P_C[\alpha_n \sigma f(x_n) + (I - \alpha_n B)W_n]x_n - Wx_n\| \\ &\leq \alpha_n \|\sigma f(x_n) - BW_n x_n\| + \|W_n x_n - Wx_n\| \rightarrow 0. \end{aligned} \quad (3.19)$$

From (3.18) and (3.19), we deduce

$$\lim_{k \rightarrow \infty} \|Wx_n - x_n\| = 0. \quad (3.20)$$

Next we prove

$$\limsup_{k \rightarrow \infty} \langle \sigma f(x^*) - Bx^*, x^k - x^* \rangle \leq 0, \quad (3.21)$$

where x^* is the unique solution of VI (3.2).

Indeed, we can choose a subsequence $\{x_{n_i}\}$ of $\{x_n\}$ such that

$$\limsup_{n \rightarrow \infty} \langle \sigma f(x^*) - Bx^*, x_n - x^* \rangle = \lim_{i \rightarrow \infty} \langle \sigma f(x^*) - Bx^*, x_{n_i} - x^* \rangle. \quad (3.22)$$

Since $\{x_{n_i}\}$ is bounded, there exists a subsequence of $\{x_{n_i}\}$ which converges weakly to a point \tilde{x} . Without loss of generality, we may assume that $\{x_{n_i}\}$ converges weakly to \tilde{x} . Therefore, from (3.20) and Lemma 2.4, we have $x_{n_i} \rightharpoonup \tilde{x} \in \text{Fix}(W) = F$. Therefore,

$$\begin{aligned} \limsup_{n \rightarrow \infty} \langle \sigma f(x^*) - Bx^*, x_n - x^* \rangle &= \lim_{i \rightarrow \infty} \langle \sigma f(x^*) - Bx^*, x_{n_i} - x^* \rangle \\ &= \langle \sigma f(x^*) - Bx^*, \tilde{x} - x^* \rangle \leq 0. \end{aligned} \quad (3.23)$$

Finally, we show that $x_n \rightarrow x^*$. We observe that

$$\|x_{n+1} - x^*\|^2 = \langle x_{n+1} - \tilde{z}_n, x_{n+1} - x^* \rangle + \langle \tilde{z}_n - x^*, x_{n+1} - x^* \rangle. \quad (3.24)$$

Since $\langle x_{n+1} - \tilde{z}_n, x_{n+1} - x^* \rangle \leq 0$, we get

$$\begin{aligned}
 \|x_{n+1} - x^*\|^2 &\leq \langle \tilde{z}_n - x^*, x_{n+1} - x^* \rangle \\
 &= \langle \alpha_n \sigma (f(x_n) - f(x^*)) + (I - \alpha_n B)(W_n x_n - x^*), x_{n+1} - x^* \rangle \\
 &\quad + \alpha_n \langle \sigma f(x^*) - Bx^*, x_{n+1} - x^* \rangle \\
 &\leq (\alpha_n \sigma \|f(x_n) - f(x^*)\| + \|I - \alpha_n B\| \|W_n x_n - x^*\|) \|x_{n+1} - x^*\| \\
 &\quad + \alpha_n \langle \sigma f(x^*) - Bx^*, x_{n+1} - x^* \rangle \\
 &\leq (1 - \alpha_n(\alpha - \sigma\kappa)) \|x_n - x^*\| \|x_{n+1} - x^*\| \\
 &\quad + \alpha_n \langle \sigma f(x^*) - Bx^*, x_{n+1} - x^* \rangle \\
 &\leq \frac{[1 - \alpha_n(\alpha - \sigma\kappa)]^2}{2} \|x_n - x^*\|^2 + \frac{1}{2} \|x_{n+1} - x^*\|^2 \\
 &\quad + \alpha_n \langle \sigma f(x^*) - Bx^*, x_{n+1} - x^* \rangle.
 \end{aligned} \tag{3.25}$$

It follows that

$$\begin{aligned}
 \|x_{n+1} - x^*\|^2 &\leq [1 - \alpha_n(\alpha - \sigma\kappa)] \|x_n - x^*\|^2 \\
 &\quad + 2\alpha_n \langle \sigma f(x^*) - Bx^*, x_{n+1} - x^* \rangle.
 \end{aligned} \tag{3.26}$$

Hence, all conditions of Lemma 2.6 are satisfied. Therefore, we immediately deduce that $x_k \rightarrow x^*$. This completes the proof. \square

From (3.1) and Theorem 3.1, we can deduce easily the following result.

Corollary 3.2. *Let C be a nonempty closed convex subset of a real Hilbert space H . Let $\{T_n\}_{n=1}^\infty$ be a sequence of nonexpansive mappings from C to C such that the common fixed point set $F := \bigcap_{n=1}^\infty F(T_n) \neq \emptyset$. For an arbitrary initial point x_0 , one defines a sequence $\{x_n\}_{n \geq 0}$ iteratively*

$$x_{n+1} = P_C[(1 - \alpha_n)W_n x_n], \quad n \geq 0, \tag{3.27}$$

where $\{\alpha_n\}$ is a real sequence in $[0, 1]$. Assume the sequence $\{\alpha_n\}$ satisfies the following conditions:

- (C1) $\lim_{n \rightarrow \infty} \alpha_n = 0$;
- (C2) $\sum_{n=0}^\infty \alpha_n = \infty$.

Then the sequence $\{x_n\}$ generated by (3.27) converges to the minimum norm common fixed point x^* of $\{T_n\}$.

Acknowledgment

The paper was supported by NSFC 11071279.

References

- [1] A. Sabharwal and L. C. Potter, "Convexly constrained linear inverse problems: iterative least-squares and regularization," *IEEE Transactions on Signal Processing*, vol. 46, no. 9, pp. 2345–2352, 1998.
- [2] Y.-L. Cui and X. Liu, "Notes on Browder's and Halpern's methods for nonexpansive mappings," *Fixed Point Theory*, vol. 10, no. 1, pp. 89–98, 2009.
- [3] Y. Yao and H.-K. Xu, "Iterative methods for finding minimum-norm fixed points of nonexpansive mappings with applications," *Optimization*, vol. 60, no. 6, pp. 645–658, 2011.
- [4] Y. Yao, R. Chen, and H.-K. Xu, "Schemes for finding minimum-norm solutions of variational inequalities," *Nonlinear Analysis. Theory, Methods & Applications*, vol. 72, no. 7-8, pp. 3447–3456, 2010.
- [5] H. H. Bauschke, "The approximation of fixed points of compositions of nonexpansive mappings in Hilbert space," *Journal of Mathematical Analysis and Applications*, vol. 202, no. 1, pp. 150–159, 1996.
- [6] F. E. Browder, "Convergence of approximants to fixed points of nonexpansive non-linear mappings in Banach spaces," *Archive for Rational Mechanics and Analysis*, vol. 24, pp. 82–90, 1967.
- [7] F. E. Browder and W. V. Petryshyn, "Construction of fixed points of nonlinear mappings in Hilbert space," *Journal of Mathematical Analysis and Applications*, vol. 20, pp. 197–228, 1967.
- [8] H.-K. Xu, "Viscosity approximation methods for nonexpansive mappings," *Journal of Mathematical Analysis and Applications*, vol. 298, no. 1, pp. 279–291, 2004.
- [9] S. A. Hirstoaga, "Iterative selection methods for common fixed point problems," *Journal of Mathematical Analysis and Applications*, vol. 324, no. 2, pp. 1020–1035, 2006.
- [10] P.-E. Mainge, "Approximation methods for common fixed points of nonexpansive mappings in Hilbert spaces," *Journal of Mathematical Analysis and Applications*, vol. 325, no. 1, pp. 469–479, 2007.
- [11] Y. Yao, R. Chen, and J.-C. Yao, "Strong convergence and certain control conditions for modified Mann iteration," *Nonlinear Analysis. Theory, Methods & Applications*, vol. 68, no. 6, pp. 1687–1693, 2008.
- [12] M. Kikkawa and W. Takahashi, "Approximating fixed points of infinite nonexpansive mappings by the hybrid method," *Journal of Optimization Theory and Applications*, vol. 117, no. 1, pp. 93–101, 2003.
- [13] H. Zegeye and N. Shahzad, "Viscosity approximation methods for a common fixed point of finite family of nonexpansive mappings," *Applied Mathematics and Computation*, vol. 191, no. 1, pp. 155–163, 2007.
- [14] Y. Yao, Y. C. Liou, and G. Marino, "Strong convergence of two iterative algorithms for nonexpansive mappings in Hilbert spaces," *Fixed Point Theory and Applications*, Article ID 279058, 7 pages, 2009.
- [15] Y. Yao, Y. J. Cho, and Y.-C. Liou, "Algorithms of common solutions for variational inclusions, mixed equilibrium problems and fixed point problems," *European Journal of Operational Research*, vol. 212, no. 2, pp. 242–250, 2011.
- [16] S.-S. Chang, "Viscosity approximation methods for a finite family of nonexpansive mappings in Banach spaces," *Journal of Mathematical Analysis and Applications*, vol. 323, no. 2, pp. 1402–1416, 2006.
- [17] M. A. Noor, "Some developments in general variational inequalities," *Applied Mathematics and Computation*, vol. 152, no. 1, pp. 199–277, 2004.
- [18] M. A. Noor, K. I. Noor, and E. Al-Said, "Iterative methods for solving nonconvex equilibrium variational inequalities," *Applied Mathematics & Information Sciences*, vol. 6, no. 1, pp. 65–69, 2012.
- [19] M. A. Noor, "Extended general variational inequalities," *Applied Mathematics Letters*, vol. 22, no. 2, pp. 182–186, 2009.
- [20] M. A. Noor, "Some aspects of extended general variational inequalities," *Abstract and Applied Analysis*, vol. 2012, Article ID 303569, 16 pages, 2012.
- [21] P. K. F. Kuhfittig, "Common fixed points of nonexpansive mappings by iteration," *Pacific Journal of Mathematics*, vol. 97, no. 1, pp. 137–139, 1981.
- [22] K. Shimoji and W. Takahashi, "Strong convergence to common fixed points of infinite nonexpansive mappings and applications," *Taiwanese Journal of Mathematics*, vol. 5, no. 2, pp. 387–404, 2001.
- [23] Y. Yao, Y.-C. Liou, and J.-C. Yao, "Convergence theorem for equilibrium problems and fixed point problems of infinite family of nonexpansive mappings," *Fixed Point Theory and Applications*, Article ID 64363, 12 pages, 2007.
- [24] K. Goebel and W. A. Kirk, *Topics in Metric Fixed Point Theory*, vol. 28 of *Cambridge Studies in Advanced Mathematics*, Cambridge University Press, Cambridge, UK, 1990.
- [25] T. Suzuki, "Strong convergence theorems for infinite families of nonexpansive mappings in general Banach spaces," *Fixed Point Theory and Applications*, no. 1, pp. 103–123, 2005.
- [26] H.-K. Xu, "Iterative algorithms for nonlinear operators," *Journal of the London Mathematical Society*, vol. 66, no. 1, pp. 240–256, 2002.



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

