

## Research Article

# A Fixed Point Approach to the Stability of an $n$ -Dimensional Mixed-Type Additive and Quadratic Functional Equation

Yang-Hi Lee<sup>1</sup> and Soon-Mo Jung<sup>2</sup>

<sup>1</sup> Department of Mathematics Education, Gongju National University of Education, Gongju 314-711, Republic of Korea

<sup>2</sup> Mathematics Section, College of Science and Technology, Hongik University, Jochiwon 339-701, Republic of Korea

Correspondence should be addressed to Soon-Mo Jung, smjung@hongik.ac.kr

Received 8 October 2011; Accepted 3 December 2011

Academic Editor: Alberto d'Onofrio

Copyright © 2012 Y.-H. Lee and S.-M. Jung. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We investigate the stability problems for a functional equation  $2f(\sum_{j=1}^n x_j) + \sum_{1 \leq i, j \leq n, i \neq j} f(x_i - x_j) = (n+1) \sum_{j=1}^n f(x_j) + (n-1) \sum_{j=1}^n f(-x_j)$  by using the fixed point method.

## 1. Introduction

In 1940, Ulam [1] gave a wide-ranging talk before a Mathematical Colloquium at the University of Wisconsin in which he discussed a number of important unsolved problems. Among those was the following question concerning the stability of homomorphisms.

*Let  $G_1$  be a group and let  $G_2$  be a metric group with a metric  $d(\cdot, \cdot)$ . Given  $\varepsilon > 0$ , does there exist a  $\delta > 0$  such that if a function  $h : G_1 \rightarrow G_2$  satisfies the inequality  $d(h(xy), h(x)h(y)) < \delta$  for all  $x, y \in G_1$ , then there is a homomorphism  $H : G_1 \rightarrow G_2$  with  $d(h(x), H(x)) < \varepsilon$  for all  $x \in G_1$ ?*

If the answer is affirmative, we say that the functional equation for homomorphisms is stable. Hyers [2] was the first mathematician to present the result concerning the stability of functional equations. He answered the question of Ulam for the case where  $G_1$  and  $G_2$  are assumed to be Banach spaces. This result of Hyers is stated as follows.

*Let  $f : E_1 \rightarrow E_2$  be a function between Banach spaces such that*

$$\|f(x+y) - f(x) - f(y)\| \leq \delta \tag{1.1}$$

for some  $\delta > 0$  and for all  $x, y \in E_1$ . Then the limit  $A(x) = \lim_{n \rightarrow \infty} 2^{-n} f(2^n x)$  exists for each  $x \in E_1$ , and  $A : E_1 \rightarrow E_2$  is the unique additive function such that  $\|f(x) - A(x)\| \leq \delta$  for every  $x \in E_1$ . Moreover, if  $f(tx)$  is continuous in  $t$  for each fixed  $x \in E_1$ , then the function  $A$  is linear.

We remark that the additive function  $A$  is directly constructed from the given function  $f$  and this method is called the *direct method*. The direct method is a very powerful method for studying the stability problems of various functional equations. Taking this famous result into consideration, the additive Cauchy equation  $f(x + y) = f(x) + f(y)$  is said to have the *Hyers-Ulam stability* on  $(E_1, E_2)$  if, for every function  $f : E_1 \rightarrow E_2$  satisfying the inequality (1.1) for some  $\delta \geq 0$  and for all  $x, y \in E_1$ , there exists an additive function  $A : E_1 \rightarrow E_2$  such that  $f - A$  is bounded on  $E_1$ .

In 1950, Aoki [3] generalized the theorem of Hyers for additive functions, and in the following year, Bourgin [4] extended the theorem without proof. Unfortunately, it seems that their results failed to receive attention from mathematicians at that time. No one has made use of these results for a long time.

In 1978, Rassias [5] actually rediscovered the result of Aoki; he proved the following theorem of Hyers for linear functions.

Let  $f : E_1 \rightarrow E_2$  be a function between Banach spaces. If  $f$  satisfies the functional inequality

$$\|f(x + y) - f(x) - f(y)\| \leq \theta(\|x\|^p + \|y\|^p) \quad (1.2)$$

for some  $\theta \geq 0$ ,  $p$  with  $0 \leq p < 1$  and for all  $x, y \in E_1$ , then there exists a unique additive function  $A : E_1 \rightarrow E_2$  such that  $\|f(x) - A(x)\| \leq (2\theta/(2 - 2^p))\|x\|^p$  for each  $x \in E_1$ . If, in addition,  $f(tx)$  is continuous in  $t$  for each fixed  $x \in E_1$ , then the function  $A$  is linear.

This result of Rassias attracted a number of mathematicians who began to be stimulated to investigate the stability problems of functional equations. By regarding a large influence of Ulam, Hyers, and Rassias on the study of stability problems of functional equations, the stability phenomenon proved by Aoki and Rassias is called the *Hyers-Ulam-Rassias stability*. For the last thirty years many results concerning the Hyers-Ulam-Rassias stability of various functional equations have been obtained (see [6–16]). For some discussion of possible definitions of stability for functional equations, see [17].

The following functional equation

$$2f\left(\sum_{j=1}^n x_j\right) + \sum_{1 \leq i, j \leq n, i \neq j} f(x_i - x_j) = (n + 1) \sum_{j=1}^n f(x_j) + (n - 1) \sum_{j=1}^n f(-x_j) \quad (1.3)$$

is called an *n-dimensional mixed-type additive and quadratic functional equation*, and each solution of (1.3) is called a *quadratic-additive function*. Recently, Towanlong and Nakmahachalasint [18] solved the *n-dimensional mixed-type additive and quadratic functional equation*.

Let  $E_1$  and  $E_2$  be vector spaces. A function  $f : E_1 \rightarrow E_2$  satisfies (1.3) for all  $x_1, x_2, \dots, x_n \in E_1$  if and only if there exist an additive function  $a : E_1 \rightarrow E_2$  and a quadratic function  $q : E_1 \rightarrow E_2$  such that  $f(x) = a(x) + q(x)$  for all  $x \in E_1$ .

Moreover, they also investigated the Hyers-Ulam-Rassias stability of (1.3) by using the direct method (see [18]). Indeed, they tried to approximate the even and odd parts of each solution of a perturbed inequality by the even and odd parts of an “exact” solution of (1.3), respectively.

In Theorems 3.1 and 3.3 of this paper, we will apply the fixed point method and prove the Hyers-Ulam-Rassias stability of the  $n$ -dimensional mixed-type additive and quadratic functional equation. The advantage of this paper, in comparison with [18], is to approximate each solution of a perturbed inequality by an “exact” solution of (1.3), and we obtain sharper estimations in consequence of this advantage.

Throughout this paper, let  $V$  be a (real or complex) vector space,  $Y$  a Banach space, and  $n$  an integer larger than 1.

## 2. Preliminaries

Let  $X$  be a nonempty set. A function  $d : X^2 \rightarrow [0, \infty]$  is called a *generalized metric* on  $X$  if and only if  $d$  satisfies the following:

- (M<sub>1</sub>)  $d(x, y) = 0$  if and only if  $x = y$ ;
- (M<sub>2</sub>)  $d(x, y) = d(y, x)$  for all  $x, y \in X$ ;
- (M<sub>3</sub>)  $d(x, z) \leq d(x, y) + d(y, z)$  for all  $x, y, z \in X$ .

We remark that the only difference between the generalized metric and the usual metric is that the range of the former is permitted to include the infinity.

We now introduce one of the fundamental results of the fixed point theory. For the proof, we refer to [19].

**Theorem 2.1.** *Let  $(X, d)$  be a complete generalized metric space. Assume that  $\Lambda : X \rightarrow X$  is a strict contraction with the Lipschitz constant  $L < 1$ . If there exists a nonnegative integer  $n_0$  such that  $d(\Lambda^{n_0+1}x, \Lambda^{n_0}x) < \infty$  for some  $x \in X$ , then the following statements are true.*

- (i) *The sequence  $\{\Lambda^n x\}$  converges to a fixed point  $x^*$  of  $\Lambda$ .*
- (ii)  *$x^*$  is the unique fixed point of  $\Lambda$  in  $X^* = \{y \in X \mid d(\Lambda^{n_0}x, y) < \infty\}$ .*
- (iii) *If  $y \in X^*$ , then*

$$d(y, x^*) \leq \frac{1}{1-L} d(\Lambda y, y). \quad (2.1)$$

In 1991, Baker applied the fixed point method to prove the Hyers-Ulam stability of a nonlinear functional equation (see [20]). Thereafter, Radu noticed that many theorems concerning the Hyers-Ulam stability of various functional equations follow from the fixed point alternative (Theorem 2.1). Indeed, he applied the fixed point method to prove the existence of a solution of the inequality (1.1) and investigated the Hyers-Ulam stability of the additive Cauchy equation (see [21] and also [22–26]). For a somewhat different fixed point approach to stability of functional equations, see [27, 28].

### 3. Hyers-Ulam-Rassias Stability

Let  $V$  be a (real or complex) vector space and let  $Y$  be a Banach space. For a given function  $f : V \rightarrow Y$ , we use the following abbreviation:

$$Df(x_1, x_2, \dots, x_n) := 2f\left(\sum_{j=1}^n x_j\right) + \sum_{1 \leq i, j \leq n, i \neq j} f(x_i - x_j) - (n+1) \sum_{j=1}^n f(x_j) - (n-1) \sum_{j=1}^n f(-x_j) \quad (3.1)$$

for all  $x_1, x_2, \dots, x_n \in V$ .

In the following theorem, we prove the stability of the functional equation (1.3) by using the fixed point method.

**Theorem 3.1.** *Let  $\varphi : (V \setminus \{0\})^n \rightarrow [0, \infty)$  be a given function. Assume that a function  $f : V \rightarrow Y$  satisfies  $f(0) = 0$  and the inequality*

$$\|Df(x_1, x_2, \dots, x_n)\| \leq \varphi(x_1, x_2, \dots, x_n) \quad (3.2)$$

for all  $x_1, x_2, \dots, x_n \in V \setminus \{0\}$ . If there exists a constant  $0 < L < 1$  such that  $\varphi$  has the property

$$\varphi(nx_1, nx_2, \dots, nx_n) \leq nL\varphi(x_1, x_2, \dots, x_n) \quad (3.3)$$

for all  $x_1, x_2, \dots, x_n \in V \setminus \{0\}$ , then there exists a unique function  $F : V \rightarrow Y$  such that  $DF(x_1, x_2, \dots, x_n) = 0$  for all  $x_1, x_2, \dots, x_n \in V \setminus \{0\}$  and

$$\|f(x) - F(x)\| \leq \frac{n+1}{4n^2(1-L)}\varphi(x) \quad (3.4)$$

for all  $x \in V \setminus \{0\}$ , where the function  $\varphi : V \setminus \{0\} \rightarrow Y$  is defined by

$$\varphi(x) := \varphi(x, x, \dots, x) + \varphi(-x, -x, \dots, -x) \quad (3.5)$$

for all  $x \in V \setminus \{0\}$ . In particular,  $F$  is represented by

$$F(x) = \lim_{m \rightarrow \infty} \left( \frac{f(n^m x) - f(-n^m x)}{2n^m} + \frac{f(n^m x) + f(-n^m x)}{2n^{2m}} \right) \quad (3.6)$$

for all  $x \in V$ .

*Proof.* Let  $S$  be the set of all functions  $g : V \rightarrow Y$  with  $g(0) = 0$ . We introduce a generalized metric on  $S$  by

$$d(g, h) := \inf\{K \in [0, \infty) \mid \|g(x) - h(x)\| \leq K\varphi(x) \forall x \in V \setminus \{0\}\}, \quad (3.7)$$

where  $\psi$  is given in (3.5). It is not difficult to show that  $(S, d)$  is a complete generalized metric space (see [29] or [30, 31]).

Now we consider the operator  $J : S \rightarrow S$  defined by

$$Jg(x) := \frac{g(nx) - g(-nx)}{2n} + \frac{g(nx) + g(-nx)}{2n^2} \tag{3.8}$$

for all  $x \in V$ . We can apply induction on  $m$  to prove

$$J^m g(x) = \frac{g(n^m x) - g(-n^m x)}{2n^m} + \frac{g(n^m x) + g(-n^m x)}{2n^{2m}} \tag{3.9}$$

for all  $x \in V$  and  $m \in \mathbb{N}$ .

Let  $g, h \in S$  and let  $K \in [0, \infty]$  be an arbitrary constant with  $d(g, h) \leq K$ . By the definition of  $d$ , (3.3), and (3.5), we have

$$\begin{aligned} \|Jg(x) - Jh(x)\| &\leq \frac{(n+1)\|g(nx) - h(nx)\|}{2n^2} + \frac{(n-1)\|g(-nx) - h(-nx)\|}{2n^2} \leq K \frac{\psi(nx)}{n} \\ &\leq KL\psi(x) \end{aligned} \tag{3.10}$$

for all  $x \in V \setminus \{0\}$ , which implies that

$$d(Jg, Jh) \leq Ld(g, h) \tag{3.11}$$

for any  $g, h \in S$ . That is,  $J$  is a strict contraction with the Lipschitz constant  $L$ .

Moreover, by (3.2), we see that

$$\|f(x) - Jf(x)\| = \left\| \frac{n-1}{4n^2} Df(-x, -x, \dots, -x) - \frac{n+1}{4n^2} Df(x, x, \dots, x) \right\| \leq \frac{n+1}{4n^2} \psi(x) \tag{3.12}$$

for all  $x \in V \setminus \{0\}$ , that is,  $d(f, Jf) \leq (n+1)/4n^2 < \infty$  (see the definition of  $d$ ). Therefore, according to Theorem 2.1, the sequence  $\{J^m f\}$  converges to the unique "fixed point"  $F : V \rightarrow Y$  of  $J$  in the set  $T = \{g \in S \mid d(f, g) < \infty\}$  and  $F$  is represented by (3.6) for all  $x \in V$ . Notice that  $F(0) = 0$ . By Theorem 2.1, we have

$$d(f, F) \leq \frac{1}{1-L} d(f, Jf) \leq \frac{n+1}{4n^2(1-L)}, \tag{3.13}$$

which implies the validity of (3.4).

By a somewhat tedious manipulation, it follows from (3.2), (3.3), and (3.6) that

$$\begin{aligned} \|DF(x_1, x_2, \dots, x_n)\| &= \lim_{m \rightarrow \infty} \left\| \frac{Df(n^m x_1, n^m x_2, \dots, n^m x_n) - Df(-n^m x_1, -n^m x_2, \dots, -n^m x_n)}{2n^m} \right. \\ &\quad \left. + \frac{Df(n^m x_1, n^m x_2, \dots, n^m x_n) + Df(-n^m x_1, -n^m x_2, \dots, -n^m x_n)}{2n^{2m}} \right\| \\ &\leq \lim_{m \rightarrow \infty} \frac{n^m + 1}{2n^{2m}} [\varphi(n^m x_1, \dots, n^m x_n) + \varphi(-n^m x_1, \dots, -n^m x_n)] \\ &= 0 \end{aligned} \tag{3.14}$$

for all  $x_1, x_2, \dots, x_n \in V \setminus \{0\}$ . □

We can prove the following corollary by a way similar to that presented in the proof of the preceding theorem.

**Corollary 3.2.** *Let  $\varphi : V^n \rightarrow [0, \infty)$  be a given function such that  $\varphi(x_1, x_2, \dots, x_n) = \varphi(-x_1, -x_2, \dots, -x_n)$  for all  $x_1, x_2, \dots, x_n \in V$ . Assume that a function  $f : V \rightarrow Y$  satisfies  $f(0) = 0$  and the inequality (3.2) for all  $x_1, x_2, \dots, x_n \in V$ . If there exists a constant  $0 < L < 1$  such that  $\varphi$  has the property (3.3) for all  $x_1, x_2, \dots, x_n \in V$ , then there exists a unique quadratic-additive function  $F : V \rightarrow Y$  such that*

$$\|f(x) - F(x)\| \leq \frac{\varphi(x, x, \dots, x)}{2n(1-L)} \tag{3.15}$$

for all  $x \in V$ .

*Proof.* Notice that

$$\|f(x) - Jf(x)\| = \left\| \frac{n-1}{4n^2} Df(-x, -x, \dots, -x) - \frac{n+1}{4n^2} Df(x, x, \dots, x) \right\| \leq \frac{\varphi(x, x, \dots, x)}{2n} \tag{3.16}$$

for all  $x \in V$  (see the proof of Theorem 3.1). By a similar method used in the proof of Theorem 3.1, we can show that there exists a unique quadratic-additive function  $F : V \rightarrow Y$  satisfying (3.15). □

In the following theorem, we prove the Hyers-Ulam-Rassias stability of (1.3) under the condition (3.17) instead of (3.3).

**Theorem 3.3.** *Given  $\varphi : (V \setminus \{0\})^n \rightarrow [0, \infty)$ , assume that a function  $f : V \rightarrow Y$  satisfies the condition  $f(0) = 0$  and the inequality (3.2) for all  $x_1, x_2, \dots, x_n \in V \setminus \{0\}$ . If there exists a constant  $0 < L < 1$  such that*

$$L\varphi(nx_1, nx_2, \dots, nx_n) \geq n^2\varphi(x_1, x_2, \dots, x_n) \tag{3.17}$$

for all  $x_1, x_2, \dots, x_n \in V \setminus \{0\}$ , then there exists a unique function  $F : V \rightarrow Y$  such that  $DF(x_1, x_2, \dots, x_n) = 0$  for all  $x_1, x_2, \dots, x_n \in V \setminus \{0\}$  and

$$\|f(x) - F(x)\| \leq \frac{L}{2n^2(1-L)}\varphi(x) \quad (3.18)$$

for all  $x \in V \setminus \{0\}$ , where  $\varphi$  is defined by (3.5). In particular,  $F$  is represented by

$$F(x) = \lim_{m \rightarrow \infty} \left( \frac{n^m}{2} \left( f\left(\frac{x}{n^m}\right) - f\left(-\frac{x}{n^m}\right) \right) + \frac{n^{2m}}{2} \left( f\left(\frac{x}{n^m}\right) + f\left(-\frac{x}{n^m}\right) \right) \right) \quad (3.19)$$

for all  $x \in V$ .

*Proof.* Let  $S$  and  $d$  be defined as in the proof of Theorem 3.1. Then  $(S, d)$  is a complete generalized metric space. We now consider the operator  $J : S \rightarrow S$  defined by

$$Jg(x) := \frac{n}{2} \left( g\left(\frac{x}{n}\right) - g\left(-\frac{x}{n}\right) \right) + \frac{n^2}{2} \left( g\left(\frac{x}{n}\right) + g\left(-\frac{x}{n}\right) \right) \quad (3.20)$$

for all  $g \in S$  and  $x \in V$ .

Notice that

$$J^m g(x) = \frac{n^m}{2} \left( g\left(\frac{x}{n^m}\right) - g\left(-\frac{x}{n^m}\right) \right) + \frac{n^{2m}}{2} \left( g\left(\frac{x}{n^m}\right) + g\left(-\frac{x}{n^m}\right) \right) \quad (3.21)$$

for all  $x \in V$  and  $m \in \mathbb{N}$ .

Let  $g, h \in S$  and let  $K \in [0, \infty]$  be an arbitrary constant with  $d(g, h) \leq K$ . By the definition of  $d$ , together with (3.5) and (3.17), we have

$$\begin{aligned} \|Jg(x) - Jh(x)\| &\leq \frac{n^2 + n}{2} \left\| g\left(\frac{x}{n}\right) - h\left(\frac{x}{n}\right) \right\| + \frac{n^2 - n}{2} \left\| g\left(-\frac{x}{n}\right) - h\left(-\frac{x}{n}\right) \right\| \\ &\leq n^2 K \varphi\left(\frac{x}{n}\right) \\ &\leq LK\varphi(x) \end{aligned} \quad (3.22)$$

for all  $x \in V \setminus \{0\}$ . Thus, we get

$$d(Jg, Jh) \leq Ld(g, h) \quad (3.23)$$

for any  $g, h \in S$ . That is,  $J$  is a strict contraction with the Lipschitz constant  $L$ .

Moreover, it follows from (3.2), (3.5), and (3.17) that

$$\begin{aligned}
\|f(x) - Jf(x)\| &= \left\| \frac{1}{2} Df\left(\frac{x}{n}, \frac{x}{n}, \dots, \frac{x}{n}\right) \right\| \\
&\leq \frac{1}{2} \varphi\left(\frac{x}{n}, \frac{x}{n}, \dots, \frac{x}{n}\right) \\
&\leq \frac{L}{2n^2} \varphi(x, x, \dots, x) \\
&\leq \frac{L}{2n^2} \varphi(x)
\end{aligned} \tag{3.24}$$

for all  $x \in V \setminus \{0\}$ , which implies that  $d(f, Jf) \leq L/2n^2 < \infty$ . Therefore, according to Theorem 2.1, the sequence  $\{J^m f\}$  converges to the unique "fixed point"  $F$  of  $J$  in the set  $T = \{g \in S \mid d(f, g) < \infty\}$  and  $F$  is represented by (3.19). In view of Theorem 2.1, we have

$$d(f, F) \leq \frac{1}{1-L} d(f, Jf) \leq \frac{L}{2n^2(1-L)}. \tag{3.25}$$

Hence, the inequality (3.18) is true.

In a similar way presented in the proof of Theorem 3.1, it follows from (3.2), (3.17), and (3.19) that

$$\begin{aligned}
\|DF(x_1, x_2, \dots, x_n)\| &= \lim_{m \rightarrow \infty} \left\| \frac{n^m}{2} \left( Df\left(\frac{x_1}{n^m}, \frac{x_2}{n^m}, \dots, \frac{x_n}{n^m}\right) - Df\left(-\frac{x_1}{n^m}, -\frac{x_2}{n^m}, \dots, -\frac{x_n}{n^m}\right) \right) \right. \\
&\quad \left. + \frac{n^{2m}}{2} \left( Df\left(\frac{x_1}{n^m}, \frac{x_2}{n^m}, \dots, \frac{x_n}{n^m}\right) + Df\left(-\frac{x_1}{n^m}, -\frac{x_2}{n^m}, \dots, -\frac{x_n}{n^m}\right) \right) \right\| \\
&\leq \lim_{m \rightarrow \infty} \frac{n^{2m} + n^m}{2} \left( \varphi\left(\frac{x_1}{n^m}, \frac{x_2}{n^m}, \dots, \frac{x_n}{n^m}\right) + \varphi\left(-\frac{x_1}{n^m}, -\frac{x_2}{n^m}, \dots, -\frac{x_n}{n^m}\right) \right) \\
&= 0
\end{aligned} \tag{3.26}$$

for all  $x_1, x_2, \dots, x_n \in V \setminus \{0\}$ . □

We can prove the following corollary by a similar way as we did in the proof of Corollary 3.2. We omit the proof.

**Corollary 3.4.** *Let  $\varphi : V^n \rightarrow [0, \infty)$  be a given function such that  $\varphi(x_1, x_2, \dots, x_n) = \varphi(-x_1, -x_2, \dots, -x_n)$  for all  $x_1, x_2, \dots, x_n \in V$ . Assume that a function  $f : V \rightarrow Y$  satisfies  $f(0) = 0$  and the inequality (3.2) for all  $x_1, x_2, \dots, x_n \in V$ . If there exists a constant  $0 < L < 1$  such that  $\varphi$  has*



the property (3.17) for all  $x_1, x_2, \dots, x_n \in V$ , then there exists a unique quadratic-additive function  $F : V \rightarrow Y$  such that

$$\|f(x) - F(x)\| \leq \frac{L}{2n^2(1-L)} \varphi(x, x, \dots, x) \quad (3.27)$$

for all  $x \in V$ .

#### 4. Applications

For a given function  $f : V \rightarrow Y$ , we will use the following abbreviation:

$$Af(x, y) := f(x + y) - f(x) - f(y) \quad (4.1)$$

for all  $x, y \in V$ .

**Corollary 4.1.** Let  $f_k : V \rightarrow Y$  be functions for which  $f_k(0) = 0$  and there exist functions  $\phi_k : V^2 \rightarrow [0, \infty)$  such that

$$\|Af_k(x, y)\| \leq \phi_k(x, y) \quad (4.2)$$

for all  $x, y \in V$  and  $k \in \{1, 2\}$ . If there exists a constant  $0 < L < 1$  such that

$$\phi_1(nx, ny) \leq nL\phi_1(x, y), \quad (4.3)$$

$$n^2\phi_2(x, y) \leq L\phi_2(nx, ny), \quad (4.4)$$

and  $\phi_k(x, y) = \phi_k(-x, -y)$  for all  $x, y \in V$  and  $k \in \{1, 2\}$ , then there exist additive functions  $F_k : V \rightarrow Y$  such that

$$\|f_1(x) - F_1(x)\| \leq \frac{\Phi_1(x)}{2n(1-L)}, \quad (4.5)$$

$$\|f_2(x) - F_2(x)\| \leq \frac{L\Phi_2(x)}{2n^2(1-L)}, \quad (4.6)$$

for all  $x \in V \setminus \{0\}$ , where  $\Phi_k : V \rightarrow Y$  are defined by

$$\Phi_k(x) := n(n-1)\phi_k(x, -x) + 2 \sum_{j=1}^{n-1} \phi_k(jx, x) \quad (4.7)$$

for all  $x \in V$  and  $k \in \{1, 2\}$ . In particular, the functions  $F_1$  and  $F_2$  are represented by

$$F_1(x) = \lim_{m \rightarrow \infty} \frac{f_1(n^m x)}{n^m}, \quad (4.8)$$

$$F_2(x) = \lim_{m \rightarrow \infty} n^m f_2\left(\frac{x}{n^m}\right) \quad (4.9)$$

for all  $x \in V$ .

*Proof.* By a long manipulation, we obtain

$$Df_k(x_1, x_2, \dots, x_n) = \sum_{1 \leq i, j \leq n, i \neq j} Af_k(x_i, -x_j) + 2 \sum_{j=1}^{n-1} Af_k\left(\sum_{i=1}^j x_i, x_{j+1}\right) \quad (4.10)$$

for all  $x_1, x_2, \dots, x_n \in V$  and  $k \in \{1, 2\}$ . If we put

$$\varphi_k(x_1, x_2, \dots, x_n) := \sum_{1 \leq i, j \leq n, i \neq j} \phi_k(x_i, -x_j) + 2 \sum_{j=1}^{n-1} \phi_k\left(\sum_{i=1}^j x_i, x_{j+1}\right) \quad (4.11)$$

for all  $x_1, x_2, \dots, x_n \in V$  and  $k \in \{1, 2\}$ , then it follows from (4.2) and (4.10) that

$$\|Df_k(x_1, x_2, \dots, x_n)\| \leq \varphi_k(x_1, x_2, \dots, x_n) \quad (4.12)$$

for all  $x_1, x_2, \dots, x_n \in V$  and  $k \in \{1, 2\}$ . By (4.3) and (4.4), we know that  $\varphi_1$  and  $\varphi_2$  satisfy (3.3) and (3.17), respectively. Furthermore,  $\varphi_k(x_1, x_2, \dots, x_n) = \varphi_k(-x_1, -x_2, \dots, -x_n)$  for all  $x_1, x_2, \dots, x_n \in V$  and  $k \in \{1, 2\}$ .

Therefore, according to Corollary 3.2, there exists a unique quadratic-additive function  $F_1 : V \rightarrow Y$  satisfying (4.5) and  $F_1$  is represented by (3.6). By (4.2) and (4.3), we see

$$\begin{aligned} \lim_{m \rightarrow \infty} \left\| \frac{f_1(n^m x) + f_1(-n^m x)}{2n^m} \right\| &= \lim_{m \rightarrow \infty} \left\| \frac{f_1(n^m x) + f_1(-n^m x) - f_1(0)}{2n^m} \right\| \\ &= \lim_{m \rightarrow \infty} \frac{1}{2n^m} \|Af_1(n^m x, -n^m x)\| \\ &\leq \lim_{m \rightarrow \infty} \frac{1}{2n^m} \phi_1(n^m x, -n^m x) \\ &\leq \lim_{m \rightarrow \infty} \frac{L^m}{2} \phi_1(x, -x) \\ &= 0, \\ \lim_{m \rightarrow \infty} \left\| \frac{f_1(n^m x) + f_1(-n^m x)}{2n^{2m}} \right\| &\leq \lim_{m \rightarrow \infty} \frac{L^m}{2n^m} \phi_1(x, -x) = 0 \end{aligned} \quad (4.13)$$

for all  $x \in V$ . From these and (3.6), we get (4.8).

Moreover, we have

$$\left\| \frac{Af_1(n^m x, n^m y)}{n^m} \right\| \leq \frac{\phi_1(n^m x, n^m y)}{n^m} \leq L^m \phi_1(x, y) \quad (4.14)$$

for all  $x, y \in V$ . Taking the limit as  $m \rightarrow \infty$  in the aforementioned inequality, we get

$$AF_1(x, y) = 0 \quad (4.15)$$

for all  $x, y \in V$ .

Now, according to Corollary 3.4, there exists a unique function  $F_2 : V \rightarrow Y$  satisfying (4.6) and  $F_2$  is represented by (3.19). Due to (4.2) and (4.4), we get

$$\begin{aligned} \lim_{m \rightarrow \infty} \frac{n^{2m}}{2} \left\| f_2\left(\frac{x}{n^m}\right) + f_2\left(-\frac{x}{n^m}\right) \right\| &= \lim_{m \rightarrow \infty} \frac{n^{2m}}{2} \left\| Af_2\left(\frac{x}{n^m}, -\frac{x}{n^m}\right) \right\| \\ &\leq \lim_{m \rightarrow \infty} \frac{n^{2m}}{2} \phi_2\left(\frac{x}{n^m}, -\frac{x}{n^m}\right) \\ &\leq \lim_{m \rightarrow \infty} \frac{L^m}{2} \phi_2(x, -x) \\ &= 0 \end{aligned} \quad (4.16)$$

as well as

$$\lim_{m \rightarrow \infty} \frac{n^m}{2} \left\| f_2\left(\frac{x}{n^m}\right) + f_2\left(-\frac{x}{n^m}\right) \right\| \leq \lim_{m \rightarrow \infty} \frac{L^m}{2n^m} \phi_2(x, -x) = 0 \quad (4.17)$$

for all  $x \in V$ . From these and (3.19), we get (4.9).

Moreover, we have

$$\left\| n^m Af_2\left(\frac{x}{n^m}, \frac{y}{n^m}\right) \right\| \leq n^m \phi_2\left(\frac{x}{n^m}, \frac{y}{n^m}\right) \leq \frac{L^m}{n^m} \phi_2(x, y) \quad (4.18)$$

for all  $x, y \in V$ . Taking the limit as  $m \rightarrow \infty$  in the aforementioned inequality, we get

$$AF_2(x, y) = 0 \quad (4.19)$$

for all  $x, y \in V$ . □

**Corollary 4.2.** *Let  $X$  be a normed space and  $p \in (-\infty, 1) \cup (2, \infty)$ . If a function  $f : X \rightarrow Y$  satisfies  $f(0) = 0$  and the inequality*

$$\|Df(x_1, x_2, \dots, x_n)\| \leq \theta(\|x_1\|^p + \|x_2\|^p + \dots + \|x_n\|^p) \quad (4.20)$$

for all  $x_1, x_2, \dots, x_n \in X \setminus \{0\}$  and for some  $\theta \geq 0$ , then there exists a unique function  $F : X \rightarrow Y$  such that  $DF(x_1, x_2, \dots, x_n) = 0$  for all  $x_1, x_2, \dots, x_n \in X \setminus \{0\}$  and

$$\|f(x) - F(x)\| \leq \begin{cases} \frac{(n+1)\theta\|x\|^p}{2(n-n^p)} & \text{if } p < 1, \\ \frac{n\theta\|x\|^p}{n^p - n^2} & \text{if } p > 2 \end{cases} \quad (4.21)$$

for all  $x \in X \setminus \{0\}$ .

*Proof.* If we put

$$\varphi(x_1, x_2, \dots, x_n) := \theta(\|x_1\|^p + \|x_2\|^p + \dots + \|x_n\|^p) \quad (4.22)$$

for all  $x_1, x_2, \dots, x_n \in X \setminus \{0\}$  and

$$L := \begin{cases} n^{p-1} & \text{if } p < 1, \\ n^{2-p} & \text{if } p > 2, \end{cases} \quad (4.23)$$

then our assertions follow from Theorems 3.1 and 3.3.  $\square$

**Corollary 4.3.** *Let  $X$  be a normed space. If a function  $f : X \rightarrow Y$  satisfies  $f(0) = 0$  and the inequality*

$$\|Df(x_1, x_2, \dots, x_n)\| \leq \theta \sum_{1 \leq i \leq n, x_i \neq 0} \|x_i\|^p \quad (4.24)$$

for all  $x_1, x_2, \dots, x_n \in X$  and for some  $\theta \geq 0$  and  $p \in (-\infty, 1) \cup (2, \infty)$ , then there exists a unique quadratic-additive function  $F : X \rightarrow Y$  such that

$$\|f(x) - F(x)\| \leq \begin{cases} \frac{n\theta\|x\|^p}{2(n-n^p)} & \text{if } p < 1, \\ \frac{n\theta\|x\|^p}{2(n^p - n^2)} & \text{if } p > 2 \end{cases} \quad (4.25)$$

for all  $x \in X \setminus \{0\}$ . In particular, if  $p < 0$ , then  $f$  is a quadratic-additive function.

*Proof.* This corollary follows from Corollaries 3.2 and 3.4 by putting

$$\varphi(x_1, x_2, \dots, x_n) := \theta \sum_{1 \leq i \leq n, x_i \neq 0} \|x_i\|^p \quad (4.26)$$

for all  $x_1, x_2, \dots, x_n \in X$  with  $L$  defined in (4.23).  $\square$

**Corollary 4.4.** Let  $X$  be a normed space and let  $p_1, p_2, \dots, p_n$  be real constants with  $p_1 + p_2 + \dots + p_n \in (-\infty, 1) \cup (2, \infty)$ . If a function  $f : X \rightarrow Y$  satisfies  $f(0) = 0$  and the inequality

$$\|Df(x_1, x_2, \dots, x_n)\| \leq \theta \|x_1\|^{p_1} \|x_2\|^{p_2} \cdots \|x_n\|^{p_n} \quad (4.27)$$

for all  $x_1, x_2, \dots, x_n \in X \setminus \{0\}$ , then there exists a unique function  $F : X \rightarrow Y$  such that  $DF(x_1, x_2, \dots, x_n) = 0$  for all  $x_1, x_2, \dots, x_n \in X \setminus \{0\}$  and

$$\|f(x) - F(x)\| \leq \begin{cases} \frac{(n+1)\theta \|x\|^p}{2n(n-n^p)} & \text{if } p < 1, \\ \frac{\theta \|x\|^p}{n^p - n^2} & \text{if } p > 2 \end{cases} \quad (4.28)$$

for all  $x \in X \setminus \{0\}$ , where  $p := p_1 + p_2 + \dots + p_n$ .

*Proof.* If we put

$$\varphi(x_1, x_2, \dots, x_n) := \theta \|x_1\|^{p_1} \|x_2\|^{p_2} \cdots \|x_n\|^{p_n} \quad (4.29)$$

for all  $x_1, x_2, \dots, x_n \in X \setminus \{0\}$  and  $L$  is defined in (4.23), then our assertion follows from Theorems 3.1 and 3.3.  $\square$



## Acknowledgment

The second author was supported by the Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education, Science and Technology (no. 2011-0004919).

## References

- [1] S. M. Ulam, *A Collection of Mathematical Problems*, Interscience Tracts in Pure and Applied Mathematics No. 8, Interscience, New York, NY, USA, 1960.
- [2] D. H. Hyers, "On the stability of the linear functional equation," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 27, pp. 222–224, 1941.
- [3] T. Aoki, "On the stability of the linear transformation in Banach spaces," *Journal of the Mathematical Society of Japan*, vol. 2, pp. 64–66, 1950.
- [4] D. G. Bourgin, "Classes of transformations and bordering transformations," *Bulletin of the American Mathematical Society*, vol. 57, pp. 223–237, 1951.
- [5] T. M. Rassias, "On the stability of the linear mapping in Banach spaces," *Proceedings of the American Mathematical Society*, vol. 72, no. 2, pp. 297–300, 1978.
- [6] S. Czerwik, *Functional Equations and Inequalities in Several Variables*, World Scientific, River Edge, NJ, USA, 2002.
- [7] P. Găvruta, "A generalization of the Hyers-Ulam-Rassias stability of approximately additive mappings," *Journal of Mathematical Analysis and Applications*, vol. 184, no. 3, pp. 431–436, 1994.
- [8] D. H. Hyers, G. Isac, and T. M. Rassias, *Stability of functional equations in several variables*, Progress in Nonlinear Differential Equations and Their Applications, 34, Birkhäuser, Boston, Mass, USA, 1998.
- [9] S.-M. Jung, *Hyers-Ulam-Rassias Stability of Functional Equations in Nonlinear Analysis*, vol. 48 of Springer Optimization and Its Applications, Springer, New York, NY, USA, 2011.

- [10] G. H. Kim, "On the stability of functional equations with square-symmetric operation," *Mathematical Inequalities & Applications*, vol. 4, no. 2, pp. 257–266, 2001.
- [11] H.-M. Kim, "On the stability problem for a mixed type of quartic and quadratic functional equation," *Journal of Mathematical Analysis and Applications*, vol. 324, no. 1, pp. 358–372, 2006.
- [12] Y.-H. Lee, "On the stability of the monomial functional equation," *Bulletin of the Korean Mathematical Society*, vol. 45, no. 2, pp. 397–403, 2008.
- [13] Y.-H. Lee and K.-W. Jun, "A generalization of the Hyers-Ulam-Rassias stability of Jensen's equation," *Journal of Mathematical Analysis and Applications*, vol. 238, no. 1, pp. 305–315, 1999.
- [14] Y.-H. Lee and K.-W. Jun, "A generalization of the Hyers-Ulam-Rassias stability of the Pexider equation," *Journal of Mathematical Analysis and Applications*, vol. 246, no. 2, pp. 627–638, 2000.
- [15] Y.-H. Lee and K.-W. Jun, "A note on the Hyers-Ulam-Rassias stability of Pexider equation," *Journal of the Korean Mathematical Society*, vol. 37, no. 1, pp. 111–124, 2000.
- [16] Y.-H. Lee and K.-W. Jun, "On the stability of approximately additive mappings," *Proceedings of the American Mathematical Society*, vol. 128, no. 5, pp. 1361–1369, 2000.
- [17] Z. Moszner, "On the stability of functional equations," *Aequationes Mathematicae*, vol. 77, no. 1-2, pp. 33–88, 2009.
- [18] W. Towanlong and P. Nakmahachalasint, "An n-dimensional mixed-type additive and quadratic functional equation and its stability," *ScienceAsia*, vol. 35, pp. 381–385, 2009.
- [19] B. Margolis and J. B. Diaz, "A fixed point theorem of the alternative, for contractions on a generalized complete metric space," *Bulletin of the American Mathematical Society*, vol. 74, pp. 305–309, 1968.
- [20] J. A. Baker, "The stability of certain functional equations," *Proceedings of the American Mathematical Society*, vol. 112, no. 3, pp. 729–732, 1991.
- [21] V. Radu, "The fixed point alternative and the stability of functional equations," *Fixed Point Theory*, vol. 4, no. 1, pp. 91–96, 2003.
- [22] L. Cădariu and V. Radu, "Fixed points and the stability of Jensen's functional equation," *JIPAM: Journal of Inequalities in Pure and Applied Mathematics*, vol. 4, no. 1, article 4, 2003.
- [23] L. Cădariu and V. Radu, "Fixed points and the stability of quadratic functional equations," *Analele Universitatii Din Timisoara (Seria Matematica-Informatica)*, vol. 41, no. 1, pp. 25–48, 2003.
- [24] L. Cădariu and V. Radu, "On the stability of the Cauchy functional equation: a fixed point approach," in *Iteration Theory*, vol. 346 of *Grazer Mathematische Berichte*, pp. 43–52, Karl-Franzens-Universitaet Graz, Graz, Austria, 2004.
- [25] L. Cădariu and V. Radu, "Fixed point methods for the generalized stability of functional equations in a single variable," *Fixed Point Theory and Applications*, vol. 2008, Article ID 749392, 15 pages, 2008.
- [26] M. Mirzavaziri and M. S. Moslehian, "A fixed point approach to stability of a quadratic equation," *Bulletin of the Brazilian Mathematical Society New Series*, vol. 37, no. 3, pp. 361–376, 2006.
- [27] J. Brzdęk, J. Chudziak, and Z. Páles, "A fixed point approach to stability of functional equations," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 74, no. 17, pp. 6728–6732, 2011.
- [28] J. Brzdęk and K. Ciepliski, "A fixed point approach to the stability of functional equations in non-Archimedean metric spaces," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 74, no. 18, pp. 6861–6867, 2011.
- [29] S.-M. Jung, "A fixed point approach to the stability of a Volterra integral equation," *Fixed Point Theory and Applications*, vol. 2007, Article ID 57064, 9 pages, 2007.
- [30] S.-M. Jung and T.-S. Kim, "A fixed point approach to the stability of the cubic functional equation," *Sociedad Matemática Mexicana Boletín Tercera Serie*, vol. 12, no. 1, pp. 51–57, 2006.
- [31] S.-M. Jung, T.-S. Kim, and K.-S. Lee, "A fixed point approach to the stability of quadratic functional equation," *Bulletin of the Korean Mathematical Society*, vol. 43, no. 3, pp. 531–541, 2006.



# Hindawi

Submit your manuscripts at  
<http://www.hindawi.com>

