

Research Article

Bounds of Double Integral Dynamic Inequalities in Two Independent Variables on Time Scales

S. H. Saker

College of Science Research Centre, King Saud University, P. O. Box 2455, Riyadh 11451, Saudi Arabia

Correspondence should be addressed to S. H. Saker, shsaker@mans.edu.eg

Received 4 October 2011; Revised 29 October 2011; Accepted 31 October 2011

Academic Editor: Delfim F. M. Torres

Copyright © 2011 S. H. Saker. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Our aim in this paper is to establish some explicit bounds of the unknown function in a certain class of nonlinear dynamic inequalities in two independent variables on time scales which are unbounded above. These on the one hand generalize and on the other hand furnish a handy tool for the study of qualitative as well as quantitative properties of solutions of partial dynamic equations on time scales. Some examples are considered to demonstrate the applications of the results.

1. Introduction

During the past decade, a number of dynamic inequalities have been established by some authors which are motivated by some applications, for example, when studying the behavior of solutions of certain classes of dynamic equations, the bounds provided by earlier inequalities are inadequate in applications and some new and specific type of dynamic inequalities on time scales are required. The general idea is to prove a result for a dynamic inequality where the domain of the unknown function is a so-called time scale \mathbb{T} , which may be an arbitrary closed subset of the real numbers \mathbb{R} . In [1, Theorem 6.1], it is proved that if u , f , and $p \in C_{rd}$ and $p \in \mathcal{R}^+$, then

$$u^\Delta(t) \leq f(t) + p(t)u(t), \quad \forall t \in [t_0, \infty)_{\mathbb{T}}, \quad (1.1)$$

implies

$$u(t) \leq u(t_0)e_p(t, t_0) + \int_{t_0}^t e_p(t, \sigma(s))f(s)\Delta s, \quad \forall t \in [t_0, \infty)_{\mathbb{T}}, \quad (1.2)$$

where $\mathcal{R}^+ := \{f \in \mathcal{R} : 1 + \mu(t)f(t) > 0, t \in \mathbb{T}\}$ and \mathcal{R} is the class of rd-continuous and regressive functions. A function $f : \mathbb{T} \rightarrow \mathbb{R}$ is said to be right-dense continuous (rd-continuous) provided f is continuous at right-dense points and at left-dense points in \mathbb{T} , left-hand limits exist and are finite. The set of all such rd-continuous functions is denoted by $C_{\text{rd}}(\mathbb{T})$. The graininess function μ for a time scale \mathbb{T} is defined by $\mu(t) := \sigma(t) - t$, and, for any function $f : \mathbb{T} \rightarrow \mathbb{R}$, the notation $f^\sigma(t)$ denotes $f(\sigma(t))$, where $\sigma(t)$ is the forward jump operator defined by $\sigma(t) := \inf\{s \in \mathbb{T} : s > t\}$. We say that a function $f : \mathbb{T} \rightarrow \mathbb{R}$ is regressive provided $1 + \mu(t)f(t) \neq 0, t \in \mathbb{T}$. The set of all regressive functions on a time scale \mathbb{T} forms an Abelian group under the addition \oplus defined by $p \oplus q := p + q + \mu pq$. Throughout this paper, we will assume that $\sup \mathbb{T} = \infty$ and define the time scale interval $[t_0, \infty)_{\mathbb{T}}$ by $[t_0, \infty)_{\mathbb{T}} := [t_0, \infty) \cap \mathbb{T}$. The exponential function $e_p(t, s)$ on time scales is defined by

$$e_p(t, s) = \exp\left(\int_s^t \xi_{\mu(\tau)}(p(\tau)) \Delta\tau\right), \quad \text{for } t, s \in \mathbb{T}, \quad (1.3)$$

where $\xi_h(z)$ is the cylinder transformation, which is given by

$$\xi_h(z) = \begin{cases} \frac{\log(1 + hz)}{h}, & h \neq 0, \\ z, & h = 0. \end{cases} \quad (1.4)$$

Alternatively, for $p \in \mathcal{R}$, one can define the exponential function $e_p(\cdot, t_0)$ to be the unique solution of the IVP $x^\Delta = p(t)x$, with $x(t_0) = 1$. If $p \in \mathcal{R}$, then $e_p(t, s)$ is real-valued and nonzero on \mathbb{T} . If $p \in \mathcal{R}^+$, then $e_p(t, t_0)$ is always positive, $e_p(t, t) = 1$, and $e_0(t, s) = 1$. Note that

$$\begin{aligned} e_p(t, t_0) &= \exp\left(\int_{t_0}^t p(s) ds\right), & \text{if } \mathbb{T} = \mathbb{R}, \\ e_p(t, t_0) &= \prod_{s=t_0}^{t-1} (1 + p(s)), & \text{if } \mathbb{T} = \mathbb{N}, \\ e_p(t, t_0) &= \prod_{s=t_0}^{t-1} (1 + (q-1)sp(s)), & \text{if } \mathbb{T} = q^{\mathbb{N}_0}. \end{aligned} \quad (1.5)$$

The book on the subject of time scales by Bohner and Peterson [1] summarizes and organizes much of time scale calculus. The three most popular examples of calculus on time scales are differential calculus, difference calculus, and quantum calculus (see [2]), that is, when $\mathbb{T} = \mathbb{R}$, $\mathbb{T} = \mathbb{N}$, and $\mathbb{T} = q^{\mathbb{N}_0} = \{q^t : t \in \mathbb{N}_0\}$, where $q > 1$.

In this paper, we will refer to the (delta) integral which we can define as follows: If $G^\Delta(t) = g(t)$, then the Cauchy (delta) integral of g is defined by $\int_a^t g(s) \Delta s := G(t) - G(a)$. It can be shown (see [1]) that if $g \in C_{\text{rd}}(\mathbb{T})$, then the Cauchy integral $G(t) := \int_{t_0}^t g(s) \Delta s$ exists, $t_0 \in \mathbb{T}$, and satisfies $G^\Delta(t) = g(t), t \in \mathbb{T}$. There are applications of dynamic equations on time scales to quantum mechanics, electrical engineering, neural networks, heat transfer,

and combinatorics. A recent cover story article in New Scientist [3] discusses several possible applications. Also, in [1, Theorem 6.4], it is proved that if u, a , and $p \in C_{rd}$ and $p \in \mathcal{R}^+$, then

$$u(t) \leq a(t) + \int_{t_0}^t p(s)u(s)\Delta s, \quad \forall t \in [t_0, \infty)_{\mathbb{T}}, \tag{1.6}$$

implies that

$$u(t) \leq a(t) + \int_{t_0}^t e_p(t, \sigma(s))a(s)p(s)\Delta s, \quad \forall t \in [t_0, \infty)_{\mathbb{T}}. \tag{1.7}$$

Since (1.7) provides an explicit bound to the unknown function $u(t)$ and a tool to the study of many qualitative as well as quantitative properties of solutions of dynamic equations, it has become one of the very few classic and most influential results in the theory and applications of dynamic inequalities. Because of its fundamental importance, over the years, many generalizations and analogous results of (1.7) have been established. Since the discovery of the inequalities (1.1)–(1.7), much work has been done, and many papers which deal with various generalizations and extensions have appeared in the literature, we refer the reader to [4–9] and the references cited therein. On the other hand, a few authors have focused on the theory of partial dynamic equations on time scales [10–15]. However, to the best of author’s knowledge, only [16–19] have studied integral inequalities useful in the theory of partial dynamic equations on time scales. Before, we give a brief summary of some of the results of dynamic inequalities in two independent variables, we present some basic definitions about calculus in two variables on time scales (for more details, we refer to [12]).

Let \mathbb{T}_1 and \mathbb{T}_2 be two time scales with at least two points, and consider the time scale intervals $\Omega_1 = [t_0, \infty) \cap \mathbb{T}_1$ and $\Omega_2 = [s_0, \infty) \cap \mathbb{T}_2$ for $t_0 \in \mathbb{T}_1$ and $s_0 \in \mathbb{T}_2$. Let $\sigma_1, \rho_1, \Delta_1$ and $\sigma_2, \rho_2, \Delta_2$ denote the forward jump operators, backward jump operators, and the delta differentiation operator, respectively, on \mathbb{T}_1 and \mathbb{T}_2 . We say that a real valued function f on $\mathbb{T}_1 \times \mathbb{T}_2$ at $(t, s) \in \Omega \equiv \Omega_1 \times \Omega_2$ has a Δ_1 partial derivative $f^{\Delta_1}(t, s)$ with respect to t if for each $\epsilon > 0$ there exists a neighborhood U_t of t such that

$$\left| f(\sigma_1(t), s) - f(\eta, s) - f^{\Delta_1}(t, s)[\sigma_1(t) - \eta] \right| \leq \epsilon |\sigma(t) - \eta|, \quad \forall \eta \in U_t. \tag{1.8}$$

In this case, we say $f^{\Delta_1}(t, s)$ is the (partial delta) derivative of $f(t, s)$ at t . We say that a real valued function f on $\mathbb{T}_1 \times \mathbb{T}_2$ at $(t, s) \in \Omega_1 \times \Omega_2$ has a Δ_2 partial derivative $f^{\Delta_2}(t, s)$ with respect to s if for each $\epsilon > 0$ there exists a neighborhood U_s of s such that

$$\left| f(t, \sigma_2(s)) - f(t, \xi) - f^{\Delta_2}(t, s)[\sigma_2(s) - \xi] \right| \leq \epsilon |\sigma(s) - \xi|, \quad \forall \xi \in U_s. \tag{1.9}$$

In this case, we say $f^{\Delta_2}(t, s)$ is the (partial delta) derivative of $f(t, s)$ at s . The function f is called rd-continuous in t if for every $\alpha_2 \in \mathbb{T}_2$ the function $f(t, \alpha_2)$ is rd-continuous on \mathbb{T}_1 . The function f is called rd-continuous in s if for every $\alpha_1 \in \mathbb{T}_1$ the function $f(\alpha_1, s)$ is rd-continuous on \mathbb{T}_2 . Now, we are ready to present some results for dynamic inequalities in two independent variables on times scales which are related to the main results in our paper.

In [18], the authors proved that if a , f , and u are positive rd-continuous functions and a is nonnegative and nondecreasing in each of its variables, then

$$u(x, y) \leq a(x, y) + \int_{x_0}^x \int_{y_0}^y f(s, t) u(s, t) \Delta t \Delta s, \quad (1.10)$$

for all $(x, y) \in [x_0, \infty)_{\mathbb{T}} \times [y_0, \infty)_{\mathbb{T}}$, implies

$$u(x, y) \leq a(x, y) e_F(x, x_0), \quad \text{where } F = \int_{y_0}^y f(x, t) \Delta t. \quad (1.11)$$

In [19], the author proved that if a, b, g, h , and u are positive continuous real functions defined on $\mathbb{T} \times \mathbb{T}$ and $\gamma > 1$ is a real constant, then

$$u^\gamma(x, y) \leq a(x, y) + b(x, y) \int_{x_0}^x \int_{y_0}^y [g(s, t) u^\gamma(s, t) + h(s, t) u(s, t)] \Delta t \Delta s, \quad (1.12)$$

implies

$$u(x, y) \leq [a(x, y) + b(x, y) m(x, y) e_G(t, t_0)]^{1/\gamma}, \quad (1.13)$$

for all $(x, y) \in [x_0, \infty)_{\mathbb{T}} \times [y_0, \infty)_{\mathbb{T}}$, where

$$\begin{aligned} m(x, y) &= \int_{x_0}^x \int_{y_0}^y \left[a(s, t) g(s, t) + \left(\frac{\gamma - 1}{\gamma} + \frac{a(s, t)}{\gamma} \right) h(s, t) \right] \Delta t \Delta s, \\ G(s, t) &= \int_{y_0}^y \left[g(x, t) + \frac{h(x, t)}{\gamma} \right] b(x, t) \Delta t. \end{aligned} \quad (1.14)$$

In this paper, we are concerned with bounds of the double integral nonlinear dynamic inequality in two independent variables

$$u^\gamma(t, s) \leq a(t, s) + b(t, s) \int_{t_0}^t \int_{s_0}^s [f(\tau, \eta) u^\delta(\tau, \eta) + g(\tau, \eta) u^\alpha(\tau, \eta)]^\lambda \Delta \eta \Delta \tau, \quad (1.15)$$

for all $(t, s) \in [t_0, \infty)_{\mathbb{T}} \times [s_0, \infty)_{\mathbb{T}}$. For (1.15), we will assume the following hypotheses:

$$(H) \begin{cases} u, a, b, f, \text{ and } g \text{ are rd-continuous positive functions on } \Omega_1 \times \Omega_2, \\ \alpha, \delta, \lambda, \text{ and } \gamma \text{ are positive constants.} \end{cases} \quad (1.16)$$

The main aim in this paper is to establish some explicit bounds of the unknown function $u(t, s)$ of the inequality (1.15). Our results not only complement the results in [18, 19] but also improve the results in [19], in the sense that the results can be applied in the cases

when $\gamma \leq 1$. The main results will be proved by employing the Bernoulli inequality [20, Bernoulli's inequality]:

$$(1+x)^\gamma \leq 1+\gamma x, \quad \text{for } 0 < \gamma \leq 1, x > -1, \quad (1.17)$$

the Young inequality [20]:

$$ab \leq \frac{a^p}{p} + \frac{b^q}{q}, \quad \text{where } a, b \geq 0, p > 1, \frac{1}{p} + \frac{1}{q} = 1, \quad (1.18)$$

and the algebraic inequalities [20]:

$$(a+b)^\lambda \leq 2^{\lambda-1}(a^\lambda + b^\lambda), \quad \text{for } a, b \geq 0, \lambda \geq 1, \quad (1.19)$$

$$(a+b)^\lambda \leq a^\lambda + b^\lambda, \quad \text{for } a, b \geq 0, 0 \leq \lambda \leq 1. \quad (1.20)$$

Some examples are considered to illustrate the main results.

2. Main Results

Before, we stated and proved the main results and we proved some Lemmas which play important roles in the proofs of the main results. We will assume that the equations or the inequalities possess such nontrivial solutions.

Lemma 2.1. *Let \mathbb{T} be an unbounded time scale with (t_0, s_0) and $(t, s) \in \mathbb{T} \times \mathbb{T}$. Let $g_i : \mathbb{R} \rightarrow \mathbb{R}$ for $i = 1, 2, \dots, n$ be functions with $g_i(x_1(t, s)) \leq g_i(x_2(t, s))$ for $i = 1, 2, \dots, n$, where $x_i(t, s) : \mathbb{T} \times \mathbb{T} \rightarrow \mathbb{R}$ for $i = 1, 2$, whenever $x_1 \leq x_2$. Let $v, w : \mathbb{T} \times \mathbb{T} \rightarrow \mathbb{R}$ be differentiable with*

$$v^{\Delta_t \Delta_s}(t, s) \leq \sum_{i=1}^n a_i(t, s) g_i(v(t, s)), \quad w^{\Delta_t \Delta_s}(t, s) \geq \sum_{i=1}^n a_i(t, s) g_i(w(t, s)), \quad (2.1)$$

for all $(t, s) \in \mathbb{T} \times \mathbb{T}$. Then, $v(t_0, s_0) < w(t_0, s_0)$ implies $v(t, s) \leq w(t, s)$ for all $(t, s) \in [t_0, \infty)_{\mathbb{T}} \times [s_0, \infty)_{\mathbb{T}}$.

Proof. The proof is by induction and similar to the proof of Theorem 6.9 in [1] and hence is omitted. \square

Lemma 2.2. *Let \mathbb{T} be an unbounded time scale with (t_0, s_0) and $(t, s) \in \mathbb{T} \times \mathbb{T}$. Suppose that $g_i : \mathbb{R} \rightarrow \mathbb{R}$ is nondecreasing for $i = 1, 2, \dots, n$ and $\mathbf{y} : \mathbb{T} \times \mathbb{T} \rightarrow \mathbb{R}^+$ is such that $g_i(\mathbf{y})$ is rd-continuous. Let p_i be rd-continuous and positive for $i = 1, 2, \dots, n$ and $f : \mathbb{T} \times \mathbb{T} \rightarrow \mathbb{R}^+$ differentiable. Then,*

$$\mathbf{y}(t, s) \leq f(t, s) + \sum_{i=1}^n \int_{t_0}^t \int_{s_0}^s p_i(\eta, \tau) g_i(\mathbf{y}(\eta, s)) \Delta \eta \Delta \tau, \quad \forall (t, s) \in \mathbb{T} \times \mathbb{T}, \quad (2.2)$$

implies $y(t, s) \leq x(t, s)$ for all $(t, s) \in [t_0, \infty)_{\mathbb{T}} \times [s_0, \infty)_{\mathbb{T}}$, where $x(t, s)$ solves the initial value problem

$$\begin{aligned} x^{\Delta_t \Delta_s}(t, s) &= f^{\Delta_t \Delta_s}(t, s) + \sum_{i=1}^n p_i(t, s) g_i(x(t, s)), \\ x(t_0, s_0) &> f(t_0, s_0) > 0. \end{aligned} \quad (2.3)$$

Proof. Let

$$v(t, s) := f(t, s) + \sum_{i=1}^n \int_{t_0}^t \int_{s_0}^s p_i(\tau, \eta) g_i(y(\tau, \eta)) \Delta \eta \Delta \tau, \quad (2.4)$$

for all $(t, s) \in [t_0, \infty)_{\mathbb{T}} \times [s_0, \infty)_{\mathbb{T}}$. Then,

$$v^{\Delta_t \Delta_s}(t, s) := f^{\Delta_t \Delta_s}(t, s) + \sum_{i=1}^n p_i(t, s) g_i(y(t, s)), \quad (2.5)$$

for all $(t, s) \in [t_0, \infty)_{\mathbb{T}} \times [s_0, \infty)_{\mathbb{T}}$ and $y(t, s) \leq v(t, s)$ so that

$$v^{\Delta_t \Delta_s}(t, s) \leq f^{\Delta_t \Delta_s}(t, s) + \sum_{i=1}^n p_i(t, s) g_i(v(t, s)), \quad (2.6)$$

for all $(t, s) \in [t_0, \infty)_{\mathbb{T}} \times [s_0, \infty)_{\mathbb{T}}$. Since $v(t_0, s_0) = f(t_0, s_0) < x_0 = x(t_0, s_0)$, the comparison Lemma 2.1 yields $v(t, s) \leq x(t, s)$ for all $(t, s) \geq (t_0, s_0)$, where $x(t, s)$ solves the initial value problem (2.3). Hence, since $y(t, s) \leq v(t, s)$, we obtain $y(t, s) \leq x(t, s)$. The proof is complete. \square

Now, we are ready to state and prove the main results in this paper. First, we consider the case when $\lambda \geq 1$ and $\alpha, \delta \leq \gamma$. For simplicity, we introduce the following notations:

$$\begin{aligned} F(t, s) &:= 2^{2(\lambda-1)} \int_{t_0}^t \int_{s_0}^s \left[f^\lambda(\tau, \eta) \left[a^{\delta/\gamma}(\tau, \eta) \right]^\lambda \right] \Delta \eta \Delta \tau \\ &\quad + 2^{2(\lambda-1)} \int_{t_0}^t \int_{s_0}^s \left[g^\lambda(\tau, \eta) \left[a^{\alpha/\gamma}(\tau, \eta) \right]^\lambda \right] \Delta \eta \Delta \tau, \\ G(t, s) &:= 2^{2(\lambda-1)} \left(f^\lambda(t, s) \left[\frac{\delta}{\gamma} a^{(\delta/\gamma)-1}(t, s) \right]^\lambda + g^\lambda(t, s) \left[\frac{\alpha}{\gamma} a^{(\alpha/\gamma)-1}(t, s) \right]^\lambda \right). \end{aligned} \quad (2.7)$$

Theorem 2.3. *Let \mathbb{T} be an unbounded time scale with $(t_0, s_0) \in \mathbb{T} \times \mathbb{T}$. Assume that (H) holds, $\lambda, \gamma \geq 1$ and $\alpha, \delta \leq \gamma$. Then,*

$$u^\gamma(t, s) \leq a(t, s) + b(t, s) \int_{t_0}^t \int_{s_0}^s \left[f(\tau, \eta) u^\delta(\tau, \eta) + g(\tau, \eta) u^\alpha(\tau, \eta) \right]^\lambda \Delta \eta \Delta \tau, \quad (2.8)$$

for $(t, s) \in \Omega$, implies that

$$u(t, s) \leq a^{1/\gamma}(t, s) + \frac{1}{\gamma} a^{(1/\gamma)-1}(t, s) b(t, s) w(t, s), \quad \forall (t, s) \in \Omega, \quad (2.9)$$

where $w(t)$ solves the initial value problem

$$w^{\Delta_t \Delta_s}(t, s) = F^{\Delta_t \Delta_s}(t, s) + b^\lambda(t, s) G(t, s) w^\lambda(t, s), \quad w(t_0, s_0) > 0. \quad (2.10)$$

Proof. Define a function $y(t, s)$ by

$$y(t, s) := \int_{t_0}^t \int_{t_0}^s [f(\tau, \eta) u^\delta(\tau, \eta) + g(\tau, \eta) u^\alpha(\tau, \eta)]^\lambda \Delta \eta \Delta \tau. \quad (2.11)$$

This reduces (2.8) to

$$u^\gamma(t, s) \leq a(t, s) + b(t, s) y(t, s), \quad \text{for } (t, s) \in \Omega. \quad (2.12)$$

This implies that

$$u(t, s) \leq (a(t, s) + b(t, s) y(t, s))^{1/\gamma}, \quad \text{for } (t, s) \in \Omega. \quad (2.13)$$

Applying the inequality (1.17) (noting that $1/\gamma \leq 1$), we see that

$$u(t, s) \leq a^{1/\gamma}(t, s) + \frac{1}{\gamma} a^{(1/\gamma)-1}(t, s) b(t, s) y(t, s), \quad \text{for } (t, s) \in \Omega. \quad (2.14)$$

From (2.13), we obtain

$$u^\alpha(t, s) \leq a^{\alpha/\gamma}(t, s) \left[1 + \frac{b(t, s) y(t, s)}{a(t, s)} \right]^{\alpha/\gamma}, \quad \text{for } (t, s) \in \Omega. \quad (2.15)$$

Applying inequality (1.17) on (2.15) (where $\alpha \leq \gamma$), we obtain for $(t, s) \in \Omega$ that

$$\begin{aligned} u^\alpha(t, s) &\leq a^{\alpha/\gamma}(t, s) \left[1 + \frac{\alpha b(t, s)}{\gamma a(t, s)} y(t, s) \right] \\ &= a^{\alpha/\gamma}(t, s) + \frac{\alpha}{\gamma} a^{(\alpha/\gamma)-1}(t, s) b(t, s) y(t, s). \end{aligned} \quad (2.16)$$

Also, from (2.13), we obtain

$$u^\delta(t, s) \leq a^{\delta/\gamma}(t, s) \left[1 + \frac{b(t, s)y(t, s)}{a(t, s)} \right]^{\delta/\gamma}, \quad \text{for } (t, s) \in \Omega. \quad (2.17)$$

Applying inequality (1.17) on (2.17) (where $\delta \leq \gamma$), we have for $(t, s) \in \Omega$ that

$$\begin{aligned} u^\delta(t, s) &\leq a^{\delta/\gamma}(t, s) \left[1 + \frac{\delta}{\gamma} \frac{b(t, s)}{a(t, s)} y(t, s) \right] \\ &= a^{\delta/\gamma}(t, s) + \frac{\delta}{\gamma} a^{(\delta/\gamma)-1}(t, s) b(t, s) y(t, s). \end{aligned} \quad (2.18)$$

Combining (2.11), (2.16), and (2.18) and applying the inequality (1.19) (noting that $\lambda \geq 1$), we have

$$\begin{aligned} y(t, s) &= \int_{t_0}^t \int_{t_0}^s \left[f(\tau, \eta) u^\delta(\tau, \eta) + g(\tau, \eta) u^\alpha(\tau, \eta) \right]^\lambda \Delta\eta \Delta\tau \\ &\leq 2^{\lambda-1} \int_{t_0}^t \int_{t_0}^s \left[f(\tau, \eta) u^\delta(\tau, \eta) \right]^\lambda \Delta\eta \Delta\tau \\ &\quad + 2^{\lambda-1} \int_{t_0}^t \int_{t_0}^s \left[g(\tau, \eta) u^\alpha(\tau, \eta) \right]^\lambda \Delta\eta \Delta\tau \\ &\leq 2^{\lambda-1} \int_{t_0}^t \int_{t_0}^s f^\lambda(\tau, \eta) \left[a^{\delta/\gamma}(\tau, \eta) + \frac{\delta}{\gamma} a^{(\delta/\gamma)-1}(\tau, \eta) b(\tau, \eta) y(\tau, \eta) \right]^\lambda \Delta\eta \Delta\tau \\ &\quad + 2^{\lambda-1} \int_{t_0}^t \int_{t_0}^s g^\lambda(\tau, \eta) \left[a^{\alpha/\gamma}(\tau, \eta) + \frac{\alpha}{\gamma} a^{(\alpha/\gamma)-1}(\tau, \eta) b(\tau, \eta) y(\tau, \eta) \right]^\lambda \Delta\eta \Delta\tau. \end{aligned} \quad (2.19)$$

This implies that

$$\begin{aligned} y(t, s) &\leq 2^{2(\lambda-1)} \int_{t_0}^t \int_{t_0}^s f^\lambda(\tau, \eta) \left[a^{(\delta/\gamma)}(\tau, \eta) \right]^\lambda \Delta\eta \Delta\tau \\ &\quad + 2^{2(\lambda-1)} \int_{t_0}^t \int_{t_0}^s f^\lambda(\tau, \eta) \left[\frac{\delta}{\gamma} a^{(\delta/\gamma)-1}(\tau, \eta) b(\tau, \eta) \right]^\lambda y^\lambda(\tau, \eta) \Delta\eta \Delta\tau \\ &\quad + 2^{2(\lambda-1)} \int_{t_0}^t \int_{t_0}^s g^\lambda(\tau, \eta) \left[a^{\alpha/\gamma}(\tau, \eta) \right]^\lambda \Delta\eta \Delta\tau \\ &\quad + 2^{2(\lambda-1)} \int_{t_0}^t \int_{t_0}^s g^\lambda(\tau, \eta) \left[\frac{\alpha}{\gamma} a^{(\alpha/\gamma)-1}(\tau, \eta) b(\tau, \eta) \right]^\lambda y^\lambda(\tau, \eta) \Delta\eta \Delta\tau \\ &= F(t, s) + \int_{t_0}^t \int_{t_0}^s G(\tau, \eta) y^\lambda(\tau, \eta) \Delta\eta \Delta\tau, \quad \text{for } (t, s) \in \Omega. \end{aligned} \quad (2.20)$$

Now an application of Lemma 2.2 (with $n = 1$ and $g(y) = y^\lambda$) gives that

$$y(t, s) < w(t, s), \quad \text{for } (t, s) \in \Omega, \quad (2.21)$$

where $w(t, s)$ solves the initial value problem (2.10). Substituting (2.21) into (2.14), we obtain the desired inequality (2.9). The proof is complete. \square

Theorem 2.4. *Let \mathbb{T} be an unbounded time scale with $(t_0, s_0) \in \mathbb{T} \times \mathbb{T}$. Assume that (H) holds, $\lambda, \gamma \geq 1$ and $\alpha, \delta \leq \gamma$. Then, (2.8) implies*

$$u(t, s) \leq a^{1/\gamma}(t, s) + b^{1/\gamma}(t, s)w^{1/\gamma}(t, s), \quad \forall (t, s) \in \Omega, \quad (2.22)$$

where $w(t)$ solves the initial value problem

$$\begin{aligned} w^{\Delta_t \Delta_s}(t, s) &= F^{\Delta_t \Delta_s}(t, s) + G_1(t, s)w^{\lambda(\delta/\gamma)}(t, s) + G_2(t, s)w^{\lambda(\alpha/\gamma)}(t, s), \\ w(t_0, s_0) &> 0, \end{aligned} \quad (2.23)$$

where $F(t)$ is defined as in (2.7) and

$$G_1(t, s) := 2^{2(\lambda-1)} f^\lambda(t, s) b^{\lambda\delta/\gamma}(t, s), \quad G_2 := 2^{2(\lambda-1)} g^\lambda(t, s) b^{\lambda\alpha/\gamma}(t, s). \quad (2.24)$$

Proof. Define a function $y(t, s)$ by (2.11) and proceed as in the proof of Theorem 2.3 to obtain

$$u(t, s) \leq (a(t, s) + b(t, s)y(t, s))^{1/\gamma}, \quad \text{for } (t, s) \in \Omega. \quad (2.25)$$

Applying the inequality (1.20), we see that

$$u(t, s) \leq a^{1/\gamma}(t, s) + b^{1/\gamma}(t, s)y^{1/\gamma}(t, s), \quad \text{for } (t, s) \in \Omega. \quad (2.26)$$

From (2.25), we obtain

$$u^\alpha(t, s) \leq (a(t, s) + b(t, s)y(t, s))^{\alpha/\gamma}, \quad \text{for } (t, s) \in \Omega. \quad (2.27)$$

Applying inequality (1.20) on (2.27) (where $\alpha \leq \gamma$), we obtain for $(t, s) \in \Omega$ that

$$u^\alpha(t, s) \leq a^{\alpha/\gamma}(t, s) + b^{\alpha/\gamma}(t, s)y^{\alpha/\gamma}(t, s). \quad (2.28)$$

Also, from (2.25), we have by (1.20) that

$$u^\delta(t, s) \leq a^{\delta/\gamma}(t, s) + b^{\delta/\gamma}(t, s)y^{\delta/\gamma}(t, s), \quad \text{for } (t, s) \in \Omega. \quad (2.29)$$

Combining (2.11), (2.28), and (2.29) and applying the inequality (1.19) (noting that $\lambda \geq 1$), we have

$$\begin{aligned}
y(t, s) &= \int_{t_0}^t \int_{t_0}^s [f(\tau, \eta)u^\delta(\tau, \eta) + g(\tau, \eta)u^\alpha(\tau, \eta)]^\lambda \Delta\eta\Delta\tau \\
&\leq 2^{\lambda-1} \int_{t_0}^t \int_{t_0}^s [f(\tau, \eta)u^\delta(\tau, \eta)]^\lambda \Delta\eta\Delta\tau \\
&\quad + 2^{\lambda-1} \int_{t_0}^t \int_{t_0}^s [g(\tau, \eta)u^\alpha(\tau, \eta)]^\lambda \Delta\eta\Delta\tau \\
&\leq 2^{\lambda-1} \int_{t_0}^t \int_{t_0}^s f^\lambda(\tau, \eta) [a^{\delta/\gamma}(\tau, \eta) + b^{\delta/\gamma}(\tau, \eta)y^{\delta/\gamma}(\tau, \eta)]^\lambda \Delta\eta\Delta\tau \\
&\quad + 2^{\lambda-1} \int_{t_0}^t \int_{t_0}^s g^\lambda(\tau, \eta) [a^{\alpha/\gamma}(\tau, \eta) + b^{\alpha/\gamma}(\tau, \eta)y^{\alpha/\gamma}(\tau, \eta)]^\lambda \Delta\eta\Delta\tau.
\end{aligned} \tag{2.30}$$

This implies that

$$\begin{aligned}
y(t, s) &\leq 2^{2(\lambda-1)} \int_{t_0}^t \int_{t_0}^s f^\lambda(\tau, \eta) [a^{\delta/\gamma}(\tau, \eta)]^\lambda \Delta\eta\Delta\tau \\
&\quad + 2^{2(\lambda-1)} \int_{t_0}^t \int_{t_0}^s g^\lambda(\tau, \eta) [a^{\alpha/\gamma}(\tau, \eta)]^\lambda \Delta\eta\Delta\tau \\
&\quad + 2^{2(\lambda-1)} \int_{t_0}^t \int_{t_0}^s f^\lambda(\tau, \eta) [b^{\delta/\gamma}(\tau, \eta)]^\lambda y^{\lambda(\delta/\gamma)}(\tau, \eta) \Delta\eta\Delta\tau \\
&\quad + 2^{2(\lambda-1)} \int_{t_0}^t \int_{t_0}^s g^\lambda(\tau, \eta) [b^{\alpha/\gamma}(\tau, \eta)]^\lambda y^{\lambda(\alpha/\gamma)}(\tau, \eta) \Delta\eta\Delta\tau \\
&= F(t, s) + \int_{t_0}^t \int_{t_0}^s [G_1(\tau, \eta)y^{\lambda(\delta/\gamma)}(\tau, \eta) + G_2(\tau, \eta)y^{\lambda(\alpha/\gamma)}(\tau, \eta)] \Delta\eta\Delta\tau,
\end{aligned} \tag{2.31}$$

for $(t, s) \in \Omega$. Now, an application of Lemma 2.2 (with $n = 2$, $g_1(y) = y^{\lambda(\delta/\gamma)}$, and $g_2(y) = y^{\lambda(\alpha/\gamma)}$) gives that

$$y(t, s) < w(t, s), \quad \text{for } (t, s) \in \Omega, \tag{2.32}$$

where $w(t, s)$ solves the initial value problem (2.23). Substituting (2.32) into (2.26), we obtain the desired inequality (2.22). The proof is complete. \square

As in the proof of Theorem 2.3 by employing the inequality (1.20) instead of the inequality (1.19), we can obtain an explicit bound for $u(t)$ when $0 \leq \lambda \leq 1$. This will be presented below in Theorem 2.5 without proof since the proof is similar to the proof of Theorem 2.3. For simplicity, we introduce the following notations:

$$\begin{aligned}
 F_1(t, s) &:= \int_{t_0}^t \int_{s_0}^s \left[f^\lambda(\tau, \eta) a^{\lambda(\delta/\gamma)}(\tau, \eta) \right] \Delta\eta \Delta\tau \\
 &\quad + \int_{t_0}^t \int_{s_0}^s \left[g^\lambda(\tau, \eta) a^{\lambda(\alpha/\gamma)}(\tau, \eta) \right] \Delta\eta \Delta\tau, \\
 G_3(t, s) &:= \left(f^\lambda(t, s) \left[\frac{\delta}{\gamma} a^{(\delta/\gamma)-1}(t, s) \right]^\lambda + g^\lambda(t, s) \left[\frac{\alpha}{\gamma} a^{(\alpha/\gamma)-1}(t, s) \right]^\lambda \right).
 \end{aligned} \tag{2.33}$$

Theorem 2.5. *Let \mathbb{T} be an unbounded time scale with $(t_0, s_0) \in \mathbb{T} \times \mathbb{T}$. Assume that (H) holds, $\gamma \geq 1$, $0 < \lambda \leq 1$, $\delta \leq \gamma$, and $\alpha \leq \gamma$. Then, (2.8) implies that*

$$u(t, s) \leq a^{1/\gamma}(t, s) + \frac{1}{\gamma} a^{(1/\gamma)-1}(t, s) b(t, s) z(t, s), \quad \text{for } (t, s) \in \Omega, \tag{2.34}$$

where $z(t)$ solves the initial value problem

$$z^{\Delta_t \Delta_s}(t) = F_1^{\Delta_t \Delta_s}(t) + G_3(t, s) b^\lambda(t, s) z^\lambda(t, s), \quad z(t_0, s_0) > 0. \tag{2.35}$$

In the following, we apply the Young inequality (1.18) to find a new explicit upper bound for $u(t)$ of (2.8) when $\lambda \geq 1$ and $0 \leq \lambda \leq 1$. First, we consider the case when $\lambda \geq 1$ and assume that $\lambda(\alpha/\gamma) < 1$ and $\lambda(\delta/\gamma) < 1$.

Theorem 2.6. *Let \mathbb{T} be an unbounded time scale with $(t_0, s_0) \in \mathbb{T} \times \mathbb{T}$. Assume that (H) holds, $\gamma, \lambda \geq 1$ and $\alpha, \delta \leq \gamma$ such that $(\lambda\alpha/\gamma) < 1$ and $(\lambda\delta/\gamma) < 1$. Then, (2.8) implies that*

$$u(t, s) \leq a^{1/\gamma}(t, s) + b^{1/\gamma}(t, s) F_3^{1/\gamma}(t, s), \quad \forall (t, s) \in \Omega, \tag{2.36}$$

where

$$\begin{aligned}
 F_3(t, s) &:= F_0(t, s) + e_{\beta(s-s_0)}(t, t_0), \quad \beta = \lambda \left[\frac{\alpha}{\gamma} + \frac{\delta}{\gamma} \right], \\
 F_0(t, s) &:= F(t, s) + \frac{(\gamma - \lambda\delta)}{\gamma} \int_{t_0}^t (G_1(\tau, \eta))^{\gamma/(\gamma-\lambda\delta)} \Delta\eta \Delta\tau \\
 &\quad + \frac{(\gamma - \lambda\alpha)}{\gamma} \int_{t_0}^t (G_2(\tau, \eta))^{\gamma/(\gamma-\lambda\alpha)} \Delta\eta \Delta\tau,
 \end{aligned} \tag{2.37}$$

and F , G_1 , and G_2 are defined as in (2.7) and (2.24).

Proof. Define a function $y(t, s)$ by (2.11) and proceed as in the proof of Theorem 2.4 to obtain

$$u(t, s) \leq a^{1/\gamma}(t, s) + b^{1/\gamma}(t, s)y^{1/\gamma}(t, s), \quad \text{for } (t, s) \in \Omega, \quad (2.38)$$

$$y(t, s) \leq F(t, s) + \int_{t_0}^t \int_{t_0}^s \left[G_1(\tau, \eta) y^{\lambda(\delta/\gamma)}(\tau, \eta) + G_2(\tau, \eta) y^{\lambda(\alpha/\gamma)}(\tau, \eta) \right] \Delta\eta \Delta\tau, \quad (2.39)$$

where F , G_1 , and G_2 are defined as in (2.7) and (2.24). Applying the Young inequality (1.18) on the term $G_1 y^{\lambda(\delta/\gamma)}$ with $q = \gamma/\lambda\delta > 1$ and $p = \gamma/(\gamma - \lambda\delta) > 1$, we see that

$$G_1 y^{\lambda(\delta/\gamma)} \leq \frac{(\gamma - \lambda\delta)}{\gamma} (G_1)^{\gamma/(\gamma - \lambda\delta)} + \left(\frac{\lambda\delta}{\gamma} \right) y. \quad (2.40)$$

Again applying the Young inequality (1.18) on the term $G_2 y^{\lambda(\alpha/\gamma)}$ with $q = \gamma/\lambda\alpha > 1$ and $p = \gamma/(\gamma - \lambda\alpha) > 1$, we see that

$$G_2 y^{\lambda(\alpha/\gamma)} \leq \frac{(\gamma - \lambda\alpha)}{\gamma} (G_2)^{\gamma/(\gamma - \lambda\alpha)} + \left(\frac{\lambda\alpha}{\gamma} \right) y. \quad (2.41)$$

Substituting (2.40) and (2.41) into (2.39), we have

$$\begin{aligned} y(t, s) &\leq F(t, s) + \frac{(\gamma - \lambda\delta)}{\gamma} \int_{t_0}^t \int_{t_0}^s (G_1(\tau, \eta))^{\gamma/(\gamma - \lambda\delta)} \Delta\eta \Delta\tau \\ &\quad + \frac{(\gamma - \lambda\alpha)}{\gamma} \int_{t_0}^t \int_{t_0}^s (G_2(\tau, \eta))^{\gamma/(\gamma - \lambda\alpha)} \Delta\eta \Delta\tau \\ &\quad + \left[\frac{\lambda\alpha}{\gamma} + \frac{\lambda\delta}{\gamma} \right] \int_{t_0}^t \int_{s_0}^s y(\tau, \eta) \Delta\eta \Delta\tau, \quad \forall (t, s) \in \Omega. \end{aligned} \quad (2.42)$$

From the definitions of $F_0(t)$ and β , we get that

$$y(t, s) \leq F_0(t, s) + \beta \int_{t_0}^t \int_{t_0}^s y(\tau, \eta) \Delta\eta \Delta\tau, \quad \text{for } (t, s) \in \Omega. \quad (2.43)$$

Applying the inequality (1.10) with $f(t, s) = \beta$, we have

$$y(t, s) < F_0(t, s) + e_{\beta(s-s_0)}(t, t_0), \quad \forall (t, s) \in \Omega. \quad (2.44)$$

Substituting (2.44) into (2.38), we get the desired inequality (2.36). The proof is complete. \square

Theorem 2.7. Let \mathbb{T} be an unbounded time scale with $(t_0, s_0) \in \mathbb{T} \times \mathbb{T}$. Assume that (H) holds, $\gamma \geq 1$, $0 < \lambda \leq 1$, and $\alpha, \delta \leq \gamma$. Then, (2.8) implies that

$$u(t, s) \leq a^{1/\gamma}(t, s) + \frac{1}{\gamma} a^{(1/\gamma)-1}(t, s) b(t, s) F_4(t, s), \quad \text{for } (t, s) \in \Omega, \quad (2.45)$$

where

$$\begin{aligned} F_4(t, s) &:= F_2(t, s) + e_{\lambda(s-s_0)}(t, t_0), \\ F_2(t, s) &:= F_1(t, s) + (1 - \lambda) \int_{t_0}^t \int_{s_0}^s (G_3(\tau, \eta))^{1/(1-\lambda)} \Delta\eta \Delta\tau, \end{aligned} \quad (2.46)$$

and F_1 and G_3 are defined as in (2.33).

Proof. Define a function $y(t, s)$ by (2.11) and proceed as in the proof of Theorem 2.3 to obtain

$$u(t, s) \leq a^{1/\gamma}(t, s) + \frac{1}{\gamma} a^{(1/\gamma)-1}(t, s) b(t, s) y(t, s), \quad \text{for } (t, s) \in \Omega, \quad (2.47)$$

$$y(t, s) \leq F_1(t, s) + \int_{t_0}^t \int_{s_0}^s G_3(s) y^\lambda(\tau, \eta) \Delta\eta \Delta\tau, \quad \text{for } (t, s) \in \Omega, \quad (2.48)$$

where F_1 and G_3 are defined in (2.33). Applying the Young inequality (1.18) on the term $G_3 y^\lambda$ with $q = 1/\lambda > 1$ and $p = 1/(1 - \lambda) > 1$, we see that

$$G_3 y^\lambda \leq (1 - \lambda)(G_3)^{1/1-\lambda} + \lambda y. \quad (2.49)$$

This and (2.48) imply that

$$\begin{aligned} y(t, s) &\leq F_1(t, s) + (1 - \lambda) \int_{t_0}^t \int_{s_0}^s (G_3(\tau, \eta))^{1/(1-\lambda)} \Delta\eta \Delta\tau \\ &\quad + \int_{t_0}^t \int_{s_0}^s \lambda y(\tau, \eta) \Delta\eta \Delta\tau. \end{aligned} \quad (2.50)$$

Using the definition of $F_2(t, s)$, we get that

$$y(t, s) \leq F_2(t, s) + \lambda \int_{t_0}^t \int_{s_0}^s y(\tau, \eta) \Delta\eta \Delta\tau, \quad \text{for } (t, s) \in \Omega. \quad (2.51)$$

Applying the inequality (1.10) with $f(t, s) = \lambda$, we have

$$y(t, s) < F_2(t, s) + e_{\lambda(s-s_0)}(t, t_0), \quad \forall (t, s) \in \Omega. \quad (2.52)$$

Substituting (2.52) into (2.47), we get the desired inequality (2.45). The proof is complete. \square

Next in the following, we consider the case when $\gamma \leq 1$ and establish some new explicit bounds of the unknown function $u(t, s)$ of (2.8).

Theorem 2.8. *Let \mathbb{T} be an unbounded time scale with $(t_0, s_0) \in \mathbb{T} \times \mathbb{T}$. Assume that (H) holds, $\lambda \leq 1$, $\gamma \leq 1$, and $\alpha\lambda, \delta\lambda \leq \gamma$. Then, (2.8) implies that*

$$u^\gamma(t, s) \leq a(t, s) + b(t, s) [H(t, s) + e_{\beta_1(s-s_0)}(t, t_0)], \quad \text{for } (t, s) \in \Omega, \quad (2.53)$$

where $\beta_1 = (\lambda\alpha/\gamma) + (\lambda\delta/\gamma)$ and

$$\begin{aligned} H(t, s) &= 2^{\lambda\delta((1/\gamma)-1)} \int_{t_0}^t \int_{t_0}^s f^\lambda(\tau, \eta) a^{\lambda\delta/\gamma}(\tau, \eta) \Delta\eta \Delta\tau \\ &+ 2^{\lambda\alpha((1/\gamma)-1)} \int_{t_0}^t \int_{t_0}^s g^\lambda(\tau, \eta) a^{\lambda\alpha/\gamma}(\tau, \eta) \Delta\eta \Delta\tau \\ &+ 2^{\lambda\delta((1/\gamma)-1)} \frac{(\gamma - \lambda\delta)}{\gamma} \int_{t_0}^t \int_{t_0}^s \left(f^\lambda(\tau, \eta) b^{\lambda\delta/\gamma}(\tau, \eta) \right)^{\gamma/(\gamma-\lambda\delta)} \Delta\eta \Delta\tau \\ &+ 2^{\lambda\alpha((1/\gamma)-1)} \frac{(\gamma - \lambda\alpha)}{\gamma} \int_{t_0}^t \int_{t_0}^s \left(g^\lambda(\tau, \eta) b^{\lambda\alpha/\gamma}(\tau, \eta) \right)^{\gamma/(\gamma-\lambda\alpha)} \Delta\eta \Delta\tau. \end{aligned} \quad (2.54)$$

Proof. Define a function $y(t, s)$ by

$$y(t, s) := \int_{t_0}^t \int_{t_0}^s \left[f(\tau, \eta) u^\delta(\tau, \eta) + g(\tau, \eta) u^\alpha(\tau, \eta) \right]^\lambda \Delta\eta \Delta\tau. \quad (2.55)$$

This reduces (2.8) to

$$u^\gamma(t, s) \leq a(t, s) + b(t, s) y(t, s), \quad \text{for } (t, s) \in \Omega. \quad (2.56)$$

This implies that

$$u(t, s) \leq (a(t, s) + b(t, s) y(t, s))^{1/\gamma}, \quad \text{for } (t, s) \in \Omega. \quad (2.57)$$

Applying the inequality (1.19) (noting that $\gamma \leq 1$), we see that

$$u(t, s) \leq 2^{(1/\gamma)-1} \left[a^{1/\gamma}(t, s) + b^{1/\gamma}(t, s) y^{1/\gamma}(t, s) \right], \quad \text{for } (t, s) \in \Omega. \quad (2.58)$$

From (2.58), we obtain

$$u^\alpha(t, s) \leq 2^{\alpha((1/\gamma)-1)} \left[a^{1/\gamma}(t, s) + b^{1/\gamma}(t, s) y^{1/\gamma}(t, s) \right]^\alpha, \quad \text{for } (t, s) \in \Omega. \quad (2.59)$$

Also, from (2.58), we obtain

$$u^\delta(t, s) \leq 2^{\delta((1/\gamma)-1)} \left[a^{1/\gamma}(t, s) + b^{1/\gamma}(t, s) y^{1/\gamma}(t, s) \right]^\delta, \quad \text{for } (t, s) \in \Omega. \quad (2.60)$$

Combining (2.55), (2.59), and (2.60) and applying the inequality (1.19) (noting that $\lambda \leq 1$), we have

$$\begin{aligned} y(t, s) &= \int_{t_0}^t \int_{t_0}^s \left[f(\tau, \eta) u^\delta(\tau, \eta) + g(\tau, \eta) u^\alpha(\tau, \eta) \right]^\lambda \Delta\eta \Delta\tau \\ &\leq \int_{t_0}^t \int_{t_0}^s \left[f(\tau, \eta) u^\delta(\tau, \eta) \right]^\lambda \Delta\eta \Delta\tau \\ &\quad + \int_{t_0}^t \int_{t_0}^s \left[g(\tau, \eta) u^\alpha(\tau, \eta) \right]^\lambda \Delta\eta \Delta\tau \\ &\leq 2^{\lambda\delta((1/\gamma)-1)} \int_{t_0}^t \int_{t_0}^s f^\lambda(\tau, \eta) \left[a^{1/\gamma}(t, s) + b^{1/\gamma}(t, s) y^{1/\gamma}(t, s) \right]^{\lambda\delta} \Delta\eta \Delta\tau \\ &\quad + 2^{\lambda\alpha((1/\gamma)-1)} \int_{t_0}^t \int_{t_0}^s g^\lambda(\tau, \eta) \left[a^{1/\gamma}(t, s) + b^{1/\gamma}(t, s) y^{1/\gamma}(t, s) \right]^{\lambda\alpha} \Delta\eta \Delta\tau. \end{aligned} \quad (2.61)$$

This implies (noting that $\alpha\lambda \leq 1$ and $\delta\lambda \leq 1$) that

$$\begin{aligned} y(t, s) &\leq 2^{\lambda\delta((1/\gamma)-1)} \int_{t_0}^t \int_{t_0}^s f^\lambda(\tau, \eta) a^{\lambda\delta/\gamma}(\tau, \eta) \Delta\eta \Delta\tau \\ &\quad + 2^{\lambda\delta((1/\gamma)-1)} \int_{t_0}^t \int_{t_0}^s f^\lambda(\tau, \eta) b^{\lambda\delta/\gamma}(\tau, \eta) y^{\lambda\delta/\gamma}(\tau, \eta) \Delta\eta \Delta\tau \\ &\quad + 2^{\lambda\alpha((1/\gamma)-1)} \int_{t_0}^t \int_{t_0}^s g^\lambda(\tau, \eta) a^{\lambda\alpha/\gamma}(\tau, \eta) \Delta\eta \Delta\tau \\ &\quad + 2^{\lambda\alpha((1/\gamma)-1)} \int_{t_0}^t \int_{t_0}^s g^\lambda(\tau, \eta) b^{\lambda\alpha/\gamma}(\tau, \eta) y^{\lambda\alpha/\gamma}(\tau, \eta) \Delta\eta \Delta\tau. \end{aligned} \quad (2.62)$$

Applying the Young inequality (1.18) on the term $H_1 y^{\lambda(\delta/\gamma)}$ with $q = \gamma/\lambda\delta > 1$ and $p = \gamma/(\gamma - \lambda\delta) > 1$, we see that

$$H_1 y^{\lambda(\delta/\gamma)} \leq \frac{(\gamma - \lambda\delta)}{\gamma} (H_1)^{\gamma/(\gamma - \lambda\delta)} + \left(\frac{\lambda\delta}{\gamma} \right) y, \quad (2.63)$$

where $H_1 = f^\lambda b^{\lambda\delta/\gamma}$. Again applying the Young inequality (1.18) on the term $H_2 y^{\lambda(\alpha/\gamma)}$ with $q = \gamma/\lambda\alpha > 1$ and $p = \gamma/(\gamma - \lambda\alpha) > 1$, we see that

$$H_2 y^{\lambda(\alpha/\gamma)} \leq \frac{(\gamma - \lambda\alpha)}{\gamma} (H_2)^{\gamma/(\gamma - \lambda\alpha)} + \left(\frac{\lambda\alpha}{\gamma} \right) y, \quad (2.64)$$

where $H_2 = g^\lambda b^{\lambda\alpha/\gamma}$. Combining (2.62)–(2.64), we have

$$y(t, s) \leq H(t, s) + \beta_1 \int_{t_0}^t \int_{t_0}^s y(\tau, \eta) \Delta\eta \Delta\tau. \quad (2.65)$$

Applying the inequality (1.10) on (2.65) with $f(t, s) = \beta_1$, we have

$$y(t, s) < H(t, s) + e_{\beta_1(s-s_0)}(t, t_0), \quad \forall (t, s) \in \Omega. \quad (2.66)$$

Substituting (2.66) into (2.56), we get the desired inequality (2.45). The proof is complete. \square

Remark 2.9. The above results can be extended to the general inequality

$$u^\gamma(t, s) \leq a(t, s) + b(t, s) \int_{t_0}^t \int_{s_0}^s L(\tau, \eta, u(\tau, \eta)) \Delta\eta \Delta\tau, \quad (2.67)$$

when

$$L(\tau, \eta, u) - L(\tau, \eta, v) \leq \left[L_1(\tau, \eta, v)(u - v)^\delta + L_2(\tau, \eta, v)(u - v)^\alpha \right]^\lambda. \quad (2.68)$$

In fact, by using the new substitution

$$y(t, s) = \int_{t_0}^t \int_{s_0}^s L(\tau, \eta, u(\tau, \eta)) \Delta\eta \Delta\tau, \quad (2.69)$$

the inequality (2.67) can be written as

$$u^\gamma(t, s) \leq a(t, s) + b(t, s)y(t, s), \quad (2.70)$$

and then, since $\gamma \geq 1$, we have

$$u(t, s) \leq A(t, s) + B(t, s)y(t, s), \quad (2.71)$$

where $A(t, s) = a^{1/\gamma}(t, s)$ and $B(t, s) = a^{(1/\gamma)-1}(t, s)b(t, s)$. This implies that

$$\begin{aligned} y(t, s) &= \int_{t_0}^t \int_{s_0}^s L(\tau, \eta, u(\tau, \eta)) \Delta\eta \Delta\tau \\ &\leq \int_{t_0}^t \int_{s_0}^s [L(\tau, \eta, A(\tau, \eta) + B(\tau, \eta)y(\tau, \eta)) - L(\tau, \eta, A(\tau, \eta))] \Delta\eta \Delta\tau \\ &\quad + \int_{t_0}^t \int_{s_0}^s L(\tau, \eta, A(\tau, \eta)) \Delta\eta \Delta\tau. \end{aligned} \quad (2.72)$$

Using (2.68) in the last inequality, we get an inequality similar to the inequality (2.8) and then follow the proof of the above results to find some explicit bounds of (2.8). The details are left to the reader.

3. Applications

In this section, we present some applications of our main results.

Example 3.1. Consider the partial dynamic equation on time scales

$$(u^\gamma(t, s))^{\Delta_t \Delta_s} = H(t, s, u(t, s)) + h(t, s), \quad (t, s) \in \Omega \equiv [t_0, \infty)_{\mathbb{T}} \times [s_0, \infty)_{\mathbb{T}}, \quad (3.1)$$

with initial boundary conditions

$$u(t, s_0) = a(t) >, \quad u(t_0, s) = b(s) >, \quad a(t_0) = b(s_0) = 0, \quad (3.2)$$

where $\gamma \geq 1$ is a constant and H and h are rd-continuous functions on Ω , $a : [t_0, \infty)_{\mathbb{T}} \rightarrow \mathbb{R}^+$ and $b : [t_0, \infty)_{\mathbb{T}} \rightarrow \mathbb{R}^+$ are rd-continuous functions. Assume that

$$|H(t, s, u)| \leq f(t, s)|u(t, s)|^\delta + g(t, s)|u(t, s)|^\alpha, \quad (3.3)$$

where $f(t, s)$ and $g(t, s)$ are nonnegative rd-continuous functions for $(t, s) \in \Omega$ and $\alpha, \delta < \gamma$. If $u(t, s)$ is a solution of (3.1)-(3.2), then $u(t, s)$ satisfies

$$|u(t, s)| \leq a^{1/\gamma}(t, s) + A^{1/\gamma}(t, s), \quad \forall (t, s) \in \Omega, \quad (3.4)$$

where

$$\begin{aligned} a(t, s) &= a^\gamma(t) + b^\gamma(s) + \int_{t_0}^t \int_{s_0}^s |h(\tau, \eta)| \Delta\eta \Delta\tau, \\ A(t, s) &:= H_0(t, s) + e_{\beta(s-s_0)}(t, t_0), \quad \beta = \left[\frac{\alpha}{\gamma} + \frac{\delta}{\gamma} \right], \\ H_0(t, s) &= \int_{t_0}^t \int_{s_0}^s \left[f(\tau, \eta) a^{\delta/\gamma}(\tau, \eta) \right] \Delta\eta \Delta\tau \\ &\quad + \int_{t_0}^t \int_{s_0}^s \left[g(\tau, \eta) a^{\alpha/\gamma}(\tau, \eta) \right] \Delta\eta \Delta\tau \\ &\quad + \frac{(\gamma - \delta)}{\gamma} \int_{t_0}^t \int_{s_0}^s (f(\tau, s))^{\gamma/(\gamma-\delta)} \Delta\eta \Delta\tau \\ &\quad + \frac{(\gamma - \alpha)}{\gamma} \int_{t_0}^t \int_{s_0}^s (g(\tau, \eta))^{\gamma/(\gamma-\alpha)} \Delta\eta \Delta\tau. \end{aligned} \quad (3.5)$$

In fact, the solution of (3.1)-(3.2) satisfies

$$|u(t, s)|^\gamma = a^\gamma(t) + b^\gamma(t) + \int_{t_0}^t \int_{s_0}^s h(\tau, \eta) \Delta\eta \Delta\tau + \int_{t_0}^t \int_{s_0}^s H(\tau, \eta, u(\tau, \eta)) \Delta\eta \Delta\tau, \quad (3.6)$$

for $(t, s) \in \Omega$. Therefore,

$$|u(t, s)|^\gamma \leq a(t, s) + \int_{t_0}^t \int_{s_0}^s |H(\tau, \eta, u(\tau, \eta))| \Delta\eta \Delta\tau, \quad \text{for } (t, s) \in \Omega. \quad (3.7)$$

It follows from (3.3) and (3.7) that

$$|u(t, s)|^\gamma \leq a(t, s) + \int_{t_0}^t \int_{s_0}^s f(\tau, s) |u(\tau, \eta)|^\delta + g(\tau, \eta) |u(\tau, \eta)|^\alpha \Delta\eta \Delta\tau, \quad (3.8)$$

for $(t, s) \in \Omega$. Applying Theorem 2.6 on (3.8) with $\lambda = 1$ and $b(t, s) = 1$, we obtain (3.4).

Example 3.2. Consider the equation

$$u^\gamma(t, s) = H(t, s, u(t, s)) + h(t, s), \quad (t, s) \in \Omega \equiv [t_0, \infty)_{\mathbb{T}} \times [s_0, \infty)_{\mathbb{T}}, \quad (3.9)$$

where $\gamma \geq 1$ is a constant and H and h are rd-continuous on Ω , $a : [t_0, \infty)_{\mathbb{T}} \rightarrow \mathbb{R}^+$ and $b : [t_0, \infty)_{\mathbb{T}} \rightarrow \mathbb{R}^+$ are rd-continuous functions. Assume that

$$|H(t, s, u)| \leq f(t, s) |u(t, s)|^\delta + g(t, s) |u(t, s)|^\alpha, \quad (3.10)$$

where $f(t, s)$ and $g(t, s)$ are nonnegative rd-continuous functions for $(t, s) \in \Omega$ and $\alpha, \delta < \gamma$. If $u(t, s)$ is a solution of (3.1)-(3.2), then $u(t, s)$ satisfies

$$|u(t, s)| \leq |h(t, s)|^{1/\gamma} + \frac{1}{\gamma} |h(t, s)|^{(1/\gamma)-1} B^{1/\gamma}(t, s), \quad \forall (t, s) \in \Omega, \quad (3.11)$$

where

$$\begin{aligned} B(t, s) &:= F^*(t, s) + (1 - \lambda) \int_{t_0}^t \int_{s_0}^s (G^*(\tau, \eta))^{1/(1-\lambda)} \Delta\eta \Delta\tau + e_{\lambda(s-s_0)}(t, t_0), \\ F^*(t, s) &:= \int_{t_0}^t \int_{s_0}^s [f^\lambda(\tau, \eta) a^{\lambda(\delta/\gamma)}(\tau, \eta) + g^\lambda(\tau, \eta) a^{\lambda(\alpha/\gamma)}(\tau, \eta)] \Delta\eta \Delta\tau, \\ G^*(t, s) &:= \left(f^\lambda(t, s) \left[\frac{\delta}{\gamma} a^{(\delta/\gamma)-1}(t, s) \right]^\lambda + g^\lambda(t, s) \left[\frac{\alpha}{\gamma} a^{(\alpha/\gamma)-1}(t, s) \right]^\lambda \right). \end{aligned} \quad (3.12)$$

In fact, the solution of (3.9) satisfies

$$|u(t, s)|^\gamma \leq |h(t, s)| + \int_{t_0}^t \int_{s_0}^s |H(\tau, \eta, u(\tau, \eta))| \Delta\eta \Delta\tau, \quad \text{for } (t, s) \in \Omega. \quad (3.13)$$

It follows from (3.10) and (3.13) that

$$|u(t, s)|^\gamma \leq |h(t, s)| + \int_{t_0}^t \int_{s_0}^s [f(\tau, s)|u(\tau, \eta)|^\delta + g(\tau, \eta)|u(\tau, \eta)|^\alpha]^\lambda \Delta\eta \Delta\tau, \quad (3.14)$$

for $(t, s) \in \Omega$. Applying Theorem 2.7 on (3.14) with $b(t, s) = 1$, we obtain (3.11).

Example 3.3. Assume that $\mathbb{T} = \mathbb{R}$ and consider the partial differential equation

$$\frac{\partial}{\partial s} \left(u^{\gamma-1}(t, s) \frac{\partial}{\partial t} u(t, s) \right) + H(t, s, u(t, s)) = h(t, s), \quad (t, s) \in \Omega^*, \quad (3.15)$$

where $\Omega^* = [0, \infty) \times [0, \infty)$, with initial boundary conditions

$$u(t, 0) = a(t) > 0, \quad u(0, s) = b(s) > 0, \quad a(0) = b(0) = 0. \quad (3.16)$$

Assume that $\gamma \geq 1$ is a constant and $H : [0, \infty) \times [0, \infty) \times \mathbb{R} \rightarrow \mathbb{R}$ and $h : [0, \infty)_{\mathbb{R}} \times [0, \infty)_{\mathbb{R}} \rightarrow \mathbb{R}$, $a : \mathbb{R} \rightarrow \mathbb{R}^+$ and $b : \mathbb{R} \rightarrow \mathbb{R}^+$ are continuous functions. Also, we assume that

$$|H(t, s, u)| \leq f(t, s)|u(t, s)|^\delta + g(t, s)|u(t, s)|^\alpha, \quad (3.17)$$

where $f(t, s)$ and $g(t, s)$ are nonnegative continuous functions for $(t, s) \in \Omega^*$ and $\alpha, \delta < \gamma$. If $u(t, s)$ is a solution of (3.1)-(3.2), then $u(t, s)$ satisfies

$$|u(t, s)| \leq a^{1/\gamma}(t, s) + \gamma^{1/\gamma} B^{1/\gamma}(t, s), \quad \forall (t, s) \in \Omega^*, \quad (3.18)$$

where

$$\begin{aligned} a(t, s) &= a^\gamma(t) + b^\gamma(s) + \gamma \int_{t_0}^t \int_{s_0}^s |h(\tau, \eta)| \Delta\eta \Delta\tau, \\ B(t, s) &:= H_0(t, s) + e_{\beta(s-s_0)}(t, t_0), \quad \beta = \left[\frac{\alpha}{\gamma} + \frac{\delta}{\gamma} \right], \\ H_0(t, s) &= \int_{t_0}^t \int_{s_0}^s [f(\tau, \eta) a^{\delta/\gamma}(\tau, \eta)] \Delta\eta \Delta\tau + \int_{t_0}^t \int_{s_0}^s [g(\tau, \eta) a^{\alpha/\gamma}(\tau, \eta)] \Delta\eta \Delta\tau \\ &\quad + \frac{(\gamma - \delta)}{\gamma} \int_{t_0}^t \int_{s_0}^s (f(\tau, s))^{\gamma/(\gamma-\delta)} \Delta\eta \Delta\tau \\ &\quad + \frac{(\gamma - \alpha)}{\gamma} \int_{t_0}^t \int_{s_0}^s (g(\tau, \eta))^{\gamma/(\gamma-\alpha)} \Delta\eta \Delta\tau. \end{aligned} \quad (3.19)$$

In fact, the solution formula of (3.15)-(3.16), after integration twice, is given by

$$\begin{aligned} |u(t, s)|^\gamma - a^\gamma(t) - b^\gamma(t) + \gamma \int_{t_0}^t \int_{s_0}^s H(\tau, \eta, u(\tau, \eta)) \Delta\eta \Delta\tau \\ = \gamma \int_{t_0}^t \int_{s_0}^s h(\tau, \eta) \Delta\eta \Delta\tau, \quad \text{for } (t, s) \in \Omega^*. \end{aligned} \quad (3.20)$$

Therefore,

$$|u(t, s)|^\gamma \leq a(t, s) + \gamma \int_{t_0}^t \int_{s_0}^s |H(\tau, \eta, u(\tau, \eta))| \Delta\eta \Delta\tau, \quad \text{for } (t, s) \in \Omega^*. \quad (3.21)$$

It follows from (3.17) and (3.21) that

$$|u(t, s)|^\gamma \leq a(t, s) + \gamma \int_{t_0}^t \int_{s_0}^s [f(\tau, s)|u(\tau, \eta)|^\delta + g(\tau, \eta)|u(\tau, \eta)|^\alpha] \Delta\eta \Delta\tau, \quad (3.22)$$

for $(t, s) \in \Omega^*$. Applying Theorem 2.6 on (3.22) with $\lambda = 1$ and $b(t, s) = \gamma$, we obtain (3.18).


Acknowledgments

The author is very grateful to the referees for their valuable remarks and comments which significantly contributed to the quality of the paper. This project was supported by King Saud University, Dean-ship of Scientific Research, College of Science Research Centre.

References

- [1] M. Bohner and A. Peterson, *Dynamic Equations on Time Scales*, Birkhäuser, Boston, Mass, USA, 2001, An Introduction with Application.
- [2] V. Kac and P. Cheung, *Quantum Calculus*, Springer, New York, NY, USA, 2001.
- [3] V. Spedding, "Taming nature's numbers," *New Scientist*, pp. 28–31, 2003.
- [4] D. R. Anderson, "Nonlinear dynamic integral inequalities in two independent variables on time scale pairs," *Advances in Dynamical Systems and Applications*, vol. 3, no. 1, pp. 1–13, 2008.
- [5] E. Akin-Bohner, M. Bohner, and F. Akin, "Pachpatte inequalities on time scales," *Journal of Inequalities in Pure and Applied Mathematics*, vol. 6, no. 1, Article 6, pp. 1–23, 2005.
- [6] W. N. Li, "Some Pachpatte type inequalities on time scales," *Computers & Mathematics with Applications*, vol. 57, no. 2, pp. 275–282, 2009.
- [7] W. N. Li, "Some new dynamic inequalities on time scales," *Journal of Mathematical Analysis and Applications*, vol. 319, no. 2, pp. 802–814, 2007.
- [8] S. H. Saker, "Some nonlinear dynamic inequalities on time scales and applications," *Journal of Mathematical Inequalities*, vol. 4, no. 4, pp. 561–579, 2010.
- [9] S. H. Saker, "Some nonlinear dynamic inequalities on time scales," *Mathematical Inequalities and Applications*, vol. 14, pp. 633–645, 2011.
- [10] C. D. Ahlbrandt and C. Morian, "Partial differential equations on time scales," *Journal of Computational and Applied Mathematics*, vol. 141, no. 1-2, pp. 35–55, 2002, Dynamic equations on time scale.
- [11] J. Hoffacker, "Basic partial dynamic equations on time scales," *Journal of Difference Equations and Applications*, vol. 8, no. 4, pp. 307–319, 2002.
- [12] B. Jackson, "Partial dynamic equations on time scales," *Journal of Computational and Applied Mathematics*, vol. 186, no. 2, pp. 391–415, 2006.

- [13] M. Bohner and G. Sh. Guseinov, "Partial differentiation on time scales," *Dynamic Systems and Applications*, vol. 13, no. 3-4, pp. 351–379, 2004.
- [14] M. Bohner and G. Sh. Guseinov, "Double integral calculus of variations on time scales," *Computers & Mathematics with Applications*, vol. 54, no. 1, pp. 45–57, 2007.
- [15] P. Wang and P. Li, "Monotone iterative technique for partial dynamic equations of first order on time scales," *Discrete Dynamics in Nature and Society*, vol. 2008, Article ID 265609, 7 pages, 2008.
- [16] D. R. Anderson, "Young's integral inequality on time scales revisited," *Journal of Inequalities in Pure and Applied Mathematics*, vol. 8, no. 3, Article 64, p. 5, 2007.
- [17] D. R. Anderson, "Dynamic double integral inequalities in two independent variables on time scales," *Journal of Mathematical Inequalities*, vol. 2, no. 2, pp. 163–184, 2008.
- [18] R. A. C. Ferreira and D. F. M. Torres, "Some linear and nonlinear integral inequalities on time scales in two independent variables," *Nonlinear Dynamics and Systems Theory*, vol. 9, no. 2, pp. 161–169, 2009.
- [19] W. N. Li, "Nonlinear integral inequalities in two independent variables on time scales," *Advances in Difference Equations*, vol. 2011, Article ID 283926, 11 pages, 2011.
- [20] D. S. Mitrinović, J. E. Pečarić, and A. M. Fink, *Classical and New Inequalities in Analysis*, vol. 61, Kluwer Academic, Dodrecht, The Netherlands, 1993.



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

