

Research Article

An Iteration Method for Nonexpansive Mappings in Hilbert Spaces

Lin Wang

Received 22 August 2006; Revised 2 November 2006; Accepted 2 November 2006

Recommended by Nan-Jing Huang

In real Hilbert space H , from an arbitrary initial point $x_0 \in H$, an explicit iteration scheme is defined as follows: $x_{n+1} = \alpha_n x_n + (1 - \alpha_n) T^{\lambda_{n+1}} x_n, n \geq 0$, where $T^{\lambda_{n+1}} x_n = T x_n - \lambda_{n+1} \mu F(T x_n)$, $T : H \rightarrow H$ is a nonexpansive mapping such that $F(T) = \{x \in K : Tx = x\}$ is nonempty, $F : H \rightarrow H$ is a η -strongly monotone and k -Lipschitzian mapping, $\{\alpha_n\} \subset (0, 1)$, and $\{\lambda_n\} \subset [0, 1)$. Under some suitable conditions, the sequence $\{x_n\}$ is shown to converge strongly to a fixed point of T and the necessary and sufficient conditions that $\{x_n\}$ converges strongly to a fixed point of T are obtained.

Copyright © 2007 Lin Wang. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction

Let H be a Hilbert space with inner product $\langle \cdot, \cdot \rangle$ and norm $\| \cdot \|$. A mapping $T : H \rightarrow H$ is said to be nonexpansive if $\|Tx - Ty\| \leq \|x - y\|$ for any $x, y \in H$. A mapping $F : H \rightarrow H$ is said to be η -strongly monotone if there exists constant $\eta > 0$ such that $\langle Fx - Fy, x - y \rangle \geq \eta \|x - y\|^2$ for any $x, y \in H$. $F : H \rightarrow H$ is said to be k -Lipschitzian if there exists constant $k > 0$ such that $\|Fx - Fy\| \leq k \|x - y\|$ for any $x, y \in H$.

The interest and importance of construction of fixed points of nonexpansive mappings stem mainly from the fact that it may be applied in many areas, such as image recovery and signal processing (see, e.g., [1–3]). Iterative techniques for approximating fixed points of nonexpansive mappings have been studied by various authors (see, e.g., [1, 4–10], etc.), using famous Mann iteration method, Ishikawa iteration method, and many other iteration methods such as, viscosity approximation method [6] and CQ method [7].

Let $F : H \rightarrow H$ be a nonlinear mapping and K nonempty closed convex subset of H . The variational inequality problem is formulated as finding a point $u^* \in K$ such

2 Fixed Point Theory and Applications

that

$$(\text{VI}(F, K)) \langle F(u^*), v - u^* \rangle \geq 0, \quad \forall v \in K. \quad (1.1)$$

The variational inequalities were initially studied by Kinderlehrer and Stampacchia [11], and ever since have been widely studied. It is well known that the $\text{VI}(F, K)$ is equivalent to the fixed point equation

$$u^* = P_K(u^* - \mu F(u^*)), \quad (1.2)$$

where P_K is the projection from H onto K and μ is an arbitrarily fixed constant. In fact, when F is an η -strongly monotone and Lipschitzian mapping on K and $\mu > 0$ small enough, then the mapping defined by the right-hand side of (1.2) is a contraction.

For reducing the complexity of computation caused by the projection P_K , Yamada [12] proposed an iteration method to solve the variational inequalities $\text{VI}(F, K)$. For arbitrary $u_0 \in H$,

$$u_{n+1} = Tu_n - \lambda_{n+1}\mu F(T(u_n)), \quad n \geq 0, \quad (1.3)$$

where T is a nonexpansive mapping from H into itself, K is the fixed point set of T , F is an η -strongly monotone and k -Lipschitzian mapping on K , $\{\lambda_n\}$ is a real sequence in $[0, 1)$, and $0 < \mu < 2\eta/k^2$. Then Yamada [12] proved that $\{u_n\}$ converges strongly to the unique solution of the $\text{VI}(F, K)$ as $\{\lambda_n\}$ satisfies the following conditions:

- (1) $\lim_{n \rightarrow \infty} \lambda_n = 0$,
- (2) $\sum_{n=0}^{\infty} \lambda_n = \infty$,
- (3) $\lim_{n \rightarrow \infty} (\lambda_n - \lambda_{n+1})/\lambda_{n+1}^2 = 0$.

Motivated by the above work, we propose a new explicit iteration scheme with mapping F to approximate the fixed point of nonexpansive mapping T in Hilbert space. The strong and weak convergence theorems to a fixed point of T are obtained. The necessary and sufficient conditions for strong convergence of this iteration scheme are obtained, too.

2. Preliminaries

Let T be a nonexpansive mapping from H into itself, $F : H \rightarrow H$ an η -strongly monotone and k -Lipschitzian mapping, $\{\lambda_n\} \subset (0, 1)$, $\{\lambda_n\} \subset [0, 1)$, and μ a fixed constant in $(0, 2\eta/k^2)$. Starting with an initial point $x_0 \in H$, the explicit iteration scheme with mapping F is defined as follows:

$$x_{n+1} = \alpha_n x_n + (1 - \alpha_n)(Tx_n - \lambda_{n+1}\mu F(Tx_n)), \quad n \geq 0. \quad (2.1)$$

For simplicity, we define a mapping $T^\lambda : H \rightarrow H$ by

$$T^\lambda x = Tx - \lambda\mu F(Tx), \quad \forall x \in H. \quad (2.2)$$

Then (2.1) may be written as follows:

$$x_{n+1} = \alpha_n x_n + (1 - \alpha_n)T^{\lambda_{n+1}} x_n, \quad n \geq 0. \quad (2.3)$$

In fact, as $\lambda_n = 0$, $n \geq 1$, then the iteration scheme (2.3) reduces to the famous Mann iteration scheme.

A Banach space E is said to satisfy Opial's condition if for any sequence $\{x_n\}$ in E , $x_n \rightharpoonup x$ implies that $\limsup_{n \rightarrow \infty} \|x_n - x\| < \limsup_{n \rightarrow \infty} \|x_n - y\|$ for all $y \in E$ with $y \neq x$, where $x_n \rightharpoonup x$ denotes that $\{x_n\}$ converges weakly to x . It is well known that every Hilbert space satisfies Opial's condition.

A mapping $T : K \rightarrow E$ is said to be semicompact if, for any sequence $\{x_n\}$ in K such that $\|x_n - Tx_n\| \rightarrow 0$ ($n \rightarrow \infty$), there exists subsequence $\{x_{n_j}\}$ of $\{x_n\}$ such that $\{x_{n_j}\}$ converges strongly to $x^* \in K$.

A mapping T with domain $D(T)$ and range $R(T)$ in E is said to be demiclosed at p ; if whenever $\{x_n\}$ is a sequence in $D(T)$ such that $\{x_n\}$ converges weakly to $x^* \in D(T)$ and $\{Tx_n\}$ converges strongly to p , then $Tx^* = p$.

LEMMA 2.1 [13]. *Let $\{\alpha_n\}$ and $\{t_n\}$ be two nonnegative sequences satisfying*

$$\alpha_{n+1} \leq (1 + a_n)\alpha_n + b_n, \quad \forall n \geq 1. \tag{2.4}$$

If $\sum_{n=1}^{\infty} a_n < \infty$ and $\sum_{n=1}^{\infty} b_n < \infty$, then $\lim_{n \rightarrow \infty} \alpha_n$ exists.

LEMMA 2.2 [12]. *Let $T^\lambda x = Tx - \lambda\mu F(Tx)$, where $T : H \rightarrow H$ is a nonexpansive mapping from H into itself and F is an η -strongly monotone and k -Lipschitzian mapping from H into itself. If $0 \leq \lambda < 1$ and $0 < \mu < 2\eta/k^2$, then T^λ is a contraction and satisfies*

$$\|T^\lambda x - T^\lambda y\| \leq (1 - \lambda\tau)\|x - y\|, \quad \forall x, y \in H, \tag{2.5}$$

where $\tau = 1 - \sqrt{1 - \mu(2\eta - \mu k^2)}$.

LEMMA 2.3 [14]. *Let K be a nonempty closed convex subset of a real Hilbert space H and T a nonexpansive mapping from K into itself. If T has a fixed point, then $I - T$ is demiclosed at zero, where I is the identity mapping of H , that is, whenever $\{x_n\}$ is a sequence in K weakly converging to some $x \in K$ and the sequence $\{(I - T)x_n\}$ strongly converges to some y , it follows that $(I - T)x = y$.*

3. Main results

LEMMA 3.1. *Let H be a Hilbert space, $T : H \rightarrow H$ a nonexpansive mapping with $F(T) \neq \emptyset$, and $F : H \rightarrow H$ an η -strongly monotone and k -Lipschitzian mapping. For any given $x_0 \in H$, $\{x_n\}$ is defined by*

$$x_{n+1} = \alpha_n x_n + (1 - \alpha_n) T^{\lambda_{n+1}} x_n, \quad n \geq 0, \tag{3.1}$$

where $\{\alpha_n\}$ and $\{\lambda_n\} \subset [0, 1)$ satisfy the following conditions:

- (1) $\alpha \leq \alpha_n \leq \beta$ for some $\alpha, \beta \in (0, 1)$;
- (2) $\sum_{n=1}^{\infty} \lambda_n < \infty$;
- (3) $0 < \mu < 2\eta/k^2$.

Then,

- (1) $\lim_{n \rightarrow \infty} \|x_n - q\|$ exists for each $q \in F(T)$;
- (2) $\lim_{n \rightarrow \infty} \|x_n - Tx_n\| = 0$.

4 Fixed Point Theory and Applications

Proof. (1) For any $q \in F(T)$, we have

$$\begin{aligned} \|x_{n+1} - q\|^2 &= \|\alpha_n(x_n - q) + (1 - \alpha_n)(T^{\lambda_{n+1}}x_n - q)\|^2 \\ &= \alpha_n\|x_n - q\|^2 + (1 - \alpha_n)\|T^{\lambda_{n+1}}x_n - q\|^2 - \alpha_n(1 - \alpha_n)\|x_n - T^{\lambda_{n+1}}x_n\|^2, \end{aligned} \quad (3.2)$$

where (by Lemma 2.2)

$$\begin{aligned} \|T^{\lambda_{n+1}}x_n - q\| &= \|T^{\lambda_{n+1}}x_n - T^{\lambda_{n+1}}q + T^{\lambda_{n+1}}q - q\| \\ &\leq \|T^{\lambda_{n+1}}x_n - T^{\lambda_{n+1}}q\| + \|T^{\lambda_{n+1}}q - q\| \\ &\leq (1 - \lambda_{n+1}\tau)\|x_n - q\| + \lambda_{n+1}\mu\|F(q)\|. \end{aligned} \quad (3.3)$$

Furthermore,

$$\|T^{\lambda_{n+1}}x_n - q\|^2 \leq (1 - \lambda_{n+1}\tau)\|x_n - q\|^2 + \frac{\lambda_{n+1}\mu^2}{\tau}\|F(q)\|^2. \quad (3.4)$$

Thus,

$$\begin{aligned} \|x_{n+1} - q\|^2 &\leq \alpha_n\|x_n - q\|^2 + (1 - \alpha_n)(1 - \lambda_{n+1}\tau)\|x_n - q\|^2 \\ &\quad + (1 - \alpha_n)\frac{\lambda_{n+1}\mu^2}{\tau}\|F(q)\|^2 - \alpha_n(1 - \alpha_n)\|x_n - T^{\lambda_{n+1}}x_n\|^2 \\ &\leq \alpha_n\|x_n - q\|^2 + (1 - \alpha_n)(1 - \lambda_{n+1}\tau)\|x_n - q\|^2 \\ &\quad + (1 - \alpha_n)\frac{\lambda_{n+1}\mu^2}{\tau}\|F(q)\|^2 - \alpha_n\|x_{n+1} - x_n\|^2 \\ &\leq \|x_n - q\|^2 + \frac{\lambda_{n+1}\mu^2}{\tau}\|F(q)\|^2 - \alpha_n\|x_{n+1} - x_n\|^2. \end{aligned} \quad (3.5)$$

Since $\sum_{n=1}^{\infty} \lambda_n < \infty$, it follows from Lemma 2.1 that $\lim_{n \rightarrow \infty} \|x_n - q\|$ exists for each $q \in F(T)$. It also implies that $\{x_n\}$ is bounded.

(2) From (3.5), we have

$$\alpha\|x_{n+1} - x_n\|^2 \leq \alpha_n\|x_{n+1} - x_n\|^2 \leq \|x_n - q\|^2 - \|x_{n+1} - q\|^2 + \frac{\lambda_{n+1}\mu^2}{\tau}\|F(q)\|^2. \quad (3.6)$$

Therefore, $\lim_{n \rightarrow \infty} \|x_{n+1} - x_n\| = 0$. In addition,

$$(1 - \beta)\|x_n - T^{\lambda_{n+1}}x_n\| \leq (1 - \alpha_n)\|x_n - T^{\lambda_{n+1}}x_n\| = \|x_{n+1} - x_n\|. \quad (3.7)$$

Hence, $\lim_{n \rightarrow \infty} \|x_n - T^{\lambda_{n+1}}x_n\| = 0$. Thus,

$$\begin{aligned} \|x_n - Tx_n\| &= \|x_n - T^{\lambda_{n+1}}x_n + T^{\lambda_{n+1}}x_n - Tx_n\| \\ &\leq \|x_n - T^{\lambda_{n+1}}x_n\| + \lambda_{n+1}\mu\|F(Tx_n)\|. \end{aligned} \quad (3.8)$$

Since $\{x_n\}$ is bounded, then $\{Tx_n\}$ and $\{F(Tx_n)\}$ are bounded, as well. Therefore, $\lim_{n \rightarrow \infty} \|x_n - Tx_n\| = 0$. The proof is completed. \square

THEOREM 3.2. *Let H be a Hilbert space, $T : H \rightarrow H$ a nonexpansive mapping with $F(T) \neq \emptyset$, and $F : H \rightarrow H$ an η -strongly monotone and k -Lipschitzian mapping. For any given $x_0 \in H$, $\{x_n\}$ is defined by*

$$x_{n+1} = \alpha_n x_n + (1 - \alpha_n) T^{\lambda_{n+1}} x_n, \quad n \geq 0, \quad (3.9)$$

where $\{\alpha_n\}$ and $\{\lambda_n\} \subset [0, 1)$ satisfy the following conditions:

- (1) $\alpha \leq \alpha_n \leq \beta$ for some $\alpha, \beta \in (0, 1)$;
- (2) $\sum_{n=1}^{\infty} \lambda_n < \infty$;
- (3) $0 < \mu < 2\eta/k^2$.

Then,

- (1) $\{x_n\}$ converges weakly to a fixed point of T ;
- (2) $\{x_n\}$ converges strongly to a fixed point of T if and only if $\liminf_{n \rightarrow \infty} d(x_n, F(T)) = 0$.

Proof. (1) It follows from Lemma 3.1 that $\{x_n\}$ is bounded. Thus, let q_1 and q_2 be weak limits of subsequences $\{x_{n_k}\}$ and $\{x_{n_j}\}$ of $\{x_n\}$, respectively. It follows from Lemmas 2.3 and 3.1 that $q_1, q_2 \in F(T)$. Assume $q_1 \neq q_2$, then by Opial's condition, we obtain

$$\begin{aligned} \lim_{n \rightarrow \infty} \|x_n - q_1\| &= \lim_{k \rightarrow \infty} \|x_{n_k} - q_1\| < \lim_{k \rightarrow \infty} \|x_{n_k} - q_2\| \\ &= \lim_{j \rightarrow \infty} \|x_{n_j} - q_2\| < \lim_{k \rightarrow \infty} \|x_{n_k} - q_1\| = \lim_{n \rightarrow \infty} \|x_n - q_1\|, \end{aligned} \quad (3.10)$$

which is a contradiction; hence, $q_1 = q_2$. Then, $\{x_n\}$ converges weakly to a common fixed point of T .

(2) Suppose that $\{x_n\}$ converges strongly to a fixed point q of T , then $\lim_{n \rightarrow \infty} \|x_n - q\| = 0$. Since $0 \leq d(x_n, F(T)) \leq \|x_n - q\|$, we have $\liminf_{n \rightarrow \infty} d(x_n, F(T)) = 0$.

Conversely, suppose that $\liminf_{n \rightarrow \infty} d(x_n, F(T)) = 0$. For any $p \in F(T)$, $\|F(p)\| \leq \|F(p) - F(x_n)\| + \|F(x_n)\| \leq k\|x_n - p\| + \|F(x_n)\|$. Since $\{x_n\}$ and $\{F(x_n)\}$ are bounded, $\|F(p)\|$ is bounded for any $p \in F(T)$, that is, there exists constant $M > 0$ such that $\|F(p)\| \leq M$ for all $p \in F(T)$. In addition, it follows from (3.5) that

$$\|x_{n+1} - p\|^2 \leq \|x_n - p\|^2 + \frac{\lambda_{n+1}\mu^2}{\tau} \|F(p)\|^2. \quad (3.11)$$

So,

$$\begin{aligned} \|x_{n+1} - p\|^2 &\leq \|x_n - p\|^2 + \frac{\lambda_{n+1}\mu^2}{\tau} (2k^2\|x_n - p\|^2 + 2\|F(x_n)\|^2) \\ &= \left(1 + 2k^2 \frac{\lambda_{n+1}\mu^2}{\tau}\right) \|x_n - p\|^2 + 2 \frac{\lambda_{n+1}\mu^2}{\tau} \|F(x_n)\|^2. \end{aligned} \quad (3.12)$$

Thus,

$$[d(x_{n+1}, F(T))]^2 \leq \left(1 + 2k^2 \frac{\lambda_{n+1}\mu^2}{\tau}\right) [d(x_n, F(T))]^2 + 2 \frac{\lambda_{n+1}\mu^2}{\tau} \|F(x_n)\|^2. \quad (3.13)$$

In addition, we obtain that $\sum_{n=1}^{\infty} 2k^2(\lambda_{n+1}\mu^2/\tau) < \infty$ and $\sum_{n=1}^{\infty} 2(\lambda_{n+1}\mu^2/\tau)\|F(x_n)\|^2 < \infty$ since $\sum_{n=1}^{\infty} \lambda_n < \infty$ and $\{F(x_n)\}$ is bounded. It follows from Lemma 2.1 that

6 Fixed Point Theory and Applications

$\lim_{n \rightarrow \infty} d(x_n, F(T))$ exists. Furthermore, since $\liminf_{n \rightarrow \infty} d(x_n, F(T)) = 0$, we have $\lim_{n \rightarrow \infty} d(x_n, F(T)) = 0$. We now prove that $\{x_n\}$ is a Cauchy sequence.

Taking $M_1 = \max\{2e^{(2\mu^2 k^2/\tau)\sum_{i=1}^{\infty} \lambda_i}, 4(\mu^2 M^2/\tau)e^{(2\mu^2 k^2/\tau)\sum_{i=1}^{\infty} \lambda_i}\}$, for any $\epsilon > 0$, there exists positive integer N such that $d(x_n, F(T)) < \sqrt{\epsilon/4M_1}$ and $\sum_{i=n}^{\infty} \lambda_i < \epsilon/4M_1$ as $n \geq N$. Taking $q \in F(T)$, for any $n, m \geq N$, it follows from (3.12) that

$$\begin{aligned}
 \frac{\|x_n - x_m\|^2}{2} &\leq \|x_n - q\|^2 + \|x_m - q\|^2 \\
 &\leq \left(1 + 2k^2 \frac{\lambda_n \mu^2}{\tau}\right) \|x_{n-1} - q\|^2 + 2 \frac{\lambda_n \mu^2}{\tau} \|F(x_{n-1})\|^2 \\
 &\quad + \left(1 + 2k^2 \frac{\lambda_m \mu^2}{\tau}\right) \|x_{m-1} - q\|^2 + 2 \frac{\lambda_m \mu^2}{\tau} \|F(x_{m-1})\|^2 \\
 &\leq \left(1 + 2k^2 \frac{\lambda_n \mu^2}{\tau}\right) \|x_{n-1} - q\|^2 + 2 \frac{\lambda_n \mu^2}{\tau} M^2 \\
 &\quad + \left(1 + 2k^2 \frac{\lambda_m \mu^2}{\tau}\right) \|x_{m-1} - q\|^2 + 2 \frac{\lambda_m \mu^2}{\tau} M^2 \\
 &\leq \prod_{i=N+1}^n \left(1 + 2k^2 \frac{\lambda_i \mu^2}{\tau}\right) \|x_N - q\|^2 + \sum_{i=N+1}^{n-1} 2 \frac{\lambda_i \mu^2}{\tau} M^2 \prod_{j=i+1}^n \left(1 + 2k^2 \frac{\lambda_j \mu^2}{\tau}\right) \\
 &\quad + 2 \frac{\lambda_n \mu^2}{\tau} M^2 + \prod_{i=N+1}^m \left(1 + 2k^2 \frac{\lambda_i \mu^2}{\tau}\right) \|x_N - q\|^2 \\
 &\quad + \sum_{i=N+1}^{m-1} 2 \frac{\lambda_i \mu^2}{\tau} M^2 \prod_{j=i+1}^m \left(1 + 2k^2 \frac{\lambda_j \mu^2}{\tau}\right) + 2 \frac{\lambda_m \mu^2}{\tau} M^2 \\
 &\leq 2e^{(2\mu^2 k^2/\tau)\sum_{i=N+1}^{\infty} \lambda_i} \|x_N - q\|^2 + 4 \frac{\mu^2 M^2}{\tau} e^{(2\mu^2 k^2/\tau)\sum_{i=N+1}^{\infty} \lambda_i} \sum_{i=N+1}^{\infty} \lambda_i.
 \end{aligned} \tag{3.14}$$

Thus,

$$\|x_n - x_m\|^2 \leq 2M_1 \|x_N - q\|^2 + 2M_1 \sum_{i=N+1}^{\infty} \lambda_i. \tag{3.15}$$

Taking the infimum for all $q \in F(T)$, we have

$$\|x_n - x_m\|^2 \leq 2M_1 [d(x_N, F(T))]^2 + 2M_1 \sum_{i=N+1}^{\infty} \lambda_i < \epsilon. \tag{3.16}$$

This implies that $\{x_n\}$ is a Cauchy sequence. Therefore, there exists $p \in H$ such that $\{x_n\}$ converges strongly to p . It follows from Lemma 3.1 that

$$\|p - Tp\| \leq \|p - x_n\| + \|x_n - Tx_n\| \rightarrow 0, \quad \text{as } n \rightarrow \infty. \tag{3.17}$$

Hence, $p \in F(T)$. The proof is completed. \square

COROLLARY 3.3. *Under the conditions of Lemma 3.1, if T is completely continuous, then $\{x_n\}$ converges strongly to a fixed point of T .*

Proof. By Lemma 3.1, $\{x_n\}$ is bounded and $\lim_{n \rightarrow \infty} \|x_n - Tx_n\| = 0$, then $\{Tx_n\}$ is also bounded. Since T is completely continuous, there exists subsequence $\{Tx_{n_j}\}$ of $\{Tx_n\}$ such that $Tx_{n_j} \rightarrow p$ as $j \rightarrow \infty$. It follows from Lemma 3.1 that $\lim_{j \rightarrow \infty} \|x_{n_j} - Tx_{n_j}\| = 0$. So by the continuity of T and Lemma 2.3, we have $\lim_{j \rightarrow \infty} \|x_{n_j} - p\| = 0$ and $p \in F(T)$. Furthermore, by Lemma 3.1, we get that $\lim_{n \rightarrow \infty} \|x_n - p\|$ exists. Thus, $\lim_{n \rightarrow \infty} \|x_n - p\| = 0$. The proof is completed. \square

COROLLARY 3.4. *Under the conditions of Lemma 3.1, if T is demicompact, then $\{x_n\}$ converges strongly to a fixed point of T .*

Proof. Since T is demicompact, $\{x_n\}$ is bounded and $\lim_{n \rightarrow \infty} \|x_n - Tx_n\| = 0$, then there exists subsequence $\{x_{n_j}\}$ of $\{x_n\}$ such that $\{x_{n_j}\}$ converges strongly to $q \in H$. It follows from Lemma 2.3 that $q \in F(T)$. Thus, $\lim_{n \rightarrow \infty} \|x_n - q\|$ exists by Lemma 3.1. Since the subsequence $\{x_{n_j}\}$ of $\{x_n\}$ such that $\{x_{n_j}\}$ converges strongly to q , then $\{x_n\}$ converges strongly to the common fixed point $q \in F(T)$. The proof is completed. \square

For studying the strong convergence of fixed points of a nonexpansive mapping, Senter and Dotson [9] introduced Condition (A). Later on, Maiti and Ghosh [5] well as Tan and Xu [10] studied Condition (A) and pointed out that Condition (A) is weaker than the requirement of demicompactness for nonexpansive mappings. A mapping $T : K \rightarrow K$ with $F(T) = \{x \in K : Tx = x\} \neq \emptyset$ is said to satisfy condition (A) if there exists a non-decreasing function $f : [0, \infty) \rightarrow [0, \infty)$ with $f(0) = 0$ and $f(t) > 0$ for all $t \in (0, \infty)$ such that $\|x - Tx\| \geq f(d(x, F(T)))$ for all $x \in K$, where $d(x, F(T)) = \inf\{\|x - q\| : q \in F(T)\}$.

THEOREM 3.5. *Under the conditions of Lemma 3.1, if T satisfies condition (A), then $\{x_n\}$ converges strongly to a fixed point of T .*

Proof. Since T satisfies condition (A), then $f(d(x_n, F(T))) \leq \|x_n - Tx_n\|$. It follows from Lemma 3.1 that $\liminf_{n \rightarrow \infty} d(x_n, F(T)) = 0$. Thus, it follows from Theorem 3.2 that $\{x_n\}$ converges strongly to a fixed point of T . The proof is completed. \square

References

- [1] F. E. Browder and W. V. Petryshyn, "Construction of fixed points of nonlinear mappings in Hilbert space," *Journal of Mathematical Analysis and Applications*, vol. 20, no. 2, pp. 197–228, 1967.
- [2] C. Byrne, "A unified treatment of some iterative algorithms in signal processing and image reconstruction," *Inverse Problems*, vol. 20, no. 1, pp. 103–120, 2004.
- [3] C. I. Podilchuk and R. J. Mammone, "Image recovery by convex projections using a least-squares constraint," *Journal of the Optical Society of America. A*, vol. 7, no. 3, pp. 517–521, 1990.
- [4] G. Marino and H.-K. Xu, "A general iterative method for nonexpansive mappings in Hilbert spaces," *Journal of Mathematical Analysis and Applications*, vol. 318, no. 1, pp. 43–52, 2006.
- [5] M. Maiti and M. K. Ghosh, "Approximating fixed points by Ishikawa iterates," *Bulletin of the Australian Mathematical Society*, vol. 40, no. 1, pp. 113–117, 1989.
- [6] A. Moudafi, "Viscosity approximation methods for fixed-points problems," *Journal of Mathematical Analysis and Applications*, vol. 241, no. 1, pp. 46–55, 2000.

8 Fixed Point Theory and Applications

- [7] K. Nakajo and W. Takahashi, “Strong convergence theorems for nonexpansive mappings and nonexpansive semigroups,” *Journal of Mathematical Analysis and Applications*, vol. 279, no. 2, pp. 372–379, 2003.
- [8] K. Nakajo, K. Shimoji, and W. Takahashi, “Strong convergence theorems by the hybrid method for families of nonexpansive mappings in Hilbert spaces,” *Taiwanese Journal of Mathematics*, vol. 10, no. 2, pp. 339–360, 2006.
- [9] H. F. Senter and W. G. Dotson Jr., “Approximating fixed points of nonexpansive mappings,” *Proceedings of the American Mathematical Society*, vol. 44, no. 2, pp. 375–380, 1974.
- [10] K.-K. Tan and H.-K. Xu, “Approximating fixed points of nonexpansive mappings by the Ishikawa iteration process,” *Journal of Mathematical Analysis and Applications*, vol. 178, no. 2, pp. 301–308, 1993.
- [11] D. Kinderlehrer and G. Stampacchia, *An Introduction to Variational Inequalities and Their Applications*, vol. 88 of *Pure and Applied Mathematics*, Academic Press, New York, NY, USA, 1980.
- [12] I. Yamada, “The hybrid steepest descent method for the variational inequality problem over the intersection of fixed point sets of nonexpansive mappings,” in *Inherently Parallel Algorithms in Feasibility and Optimization and Their Applications (Haifa, 2000)*, D. Butnariu, Y. Censor, and S. Reich, Eds., vol. 8 of *Stud. Comput. Math.*, pp. 473–504, North-Holland, Amsterdam, The Netherlands, 2001.
- [13] M. O. Osilike, S. C. Aniagbosor, and B. G. Akuchu, “Fixed points of asymptotically demicontractive mappings in arbitrary Banach spaces,” *Panamerican Mathematical Journal*, vol. 12, no. 2, pp. 77–88, 2002.
- [14] K. Geobel and W. A. Kirk, *Topics in Metric Fixed Point Theory*, vol. 28 of *Cambridge Studies in Advanced Mathematics*, Cambridge University Press, Cambridge, UK, 1990.

Lin Wang: Department of Mathematics, Kunming Teachers College, Kunming,
Yunnan 650031, China
Email address: wl64mail@yahoo.com.cn