

Research Article

Mann Type Implicit Iteration Approximation for Multivalued Mappings in Banach Spaces

Huimin He,¹ Sanyang Liu,¹ and Rudong Chen²

¹ Department of Mathematics, Xidian University, Xi'an 710071, China

² Department of Mathematics, Tianjin Polytechnic University, Tianjin 300160, China

Correspondence should be addressed to Huimin He, huiminhe@126.com

Received 16 March 2010; Accepted 5 July 2010

Academic Editor: Mohamed Amine Khamsi

Copyright © 2010 Huimin He et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Let K be a nonempty compact convex subset of a uniformly convex Banach space E and let T be a multivalued nonexpansive mapping. For the implicit iterates $x_0 \in K$, $x_n = \alpha_n x_{n-1} + (1 - \alpha_n)y_n$, $y_n \in Tx_n$, $n \geq 1$. We proved that $\{x_n\}$ converges strongly to a fixed point of T under some suitable conditions. Our results extended corresponding ones and revised a gap in the work of Panyanak (2007).

1. Introduction

Let K be a nonempty subset of a Banach space E . We will denote 2^E by the family of all subsets of E , $CB(E)$ the family of nonempty closed and bounded subsets of E , $C(E)$ the family of nonempty compact subsets of E . Let $CK(E)$ symbolize the family of nonempty compact convex subsets of E . Let $H(\cdot, \cdot)$ be Hausdorff metric on $CB(E)$; that is,

$$H(A, B) = \max \left\{ \sup_{x \in A} d(x, B), \sup_{x \in B} d(x, A) \right\}, \quad \forall A, B \in CB(E), \quad (1.1)$$

where $d(x, B) = \inf\{\|x - y\| : y \in B\}$. A multivalued mapping $T : K \rightarrow CB(E)$ is called nonexpansive (resp., contractive), if for any $x, y \in K$, there holds

$$H(Tx, Ty) \leq \|x - y\|, \quad (1.2)$$

(resp., $H(Tx, Ty) \leq k\|x - y\|$, for some $k \in (0, 1)$).

A point x is called a fixed point of T if $x \in Tx$. In this paper, $F(T)$ stands for the fixed point set of a mapping T .

The fixed point theory of multivalued nonexpansive mappings is much more complicated and difficult than the corresponding theory of single-valued nonexpansive mappings. However, some classical fixed point theorems for single-valued nonexpansive mappings have already been extended to multivalued mappings.

In 1968, Markin [1] firstly established the nonexpansive multivalued convergence results in Hilbert space. Banach's Contraction Principle was extended to a multivalued contraction in 1969. (Below is stated in a Banach space setting.)

Theorem 1.1 (see [2]). *Let K be a nonempty closed subset of a Banach space E and $T : K \rightarrow CB(K)$ a multivalued contraction. Then T has a fixed point.*

In 1974, one breakthrough was achieved by Lim using Edelstein's method of asymptotic centers [3].

Theorem 1.2 (see Lim [3]). *Let K be a nonempty closed bounded convex subset of a uniformly convex Banach space E and $T : K \rightarrow C(E)$ a multivalued nonexpansive mapping. Then T has a fixed point.*

In 1990, Kirk and Massa [4] obtained another important result for multivalued nonexpansive mappings.

Theorem 1.3 (see Kirk and Massa [4]). *Let K be a nonempty closed bounded convex subset of a Banach space E and $T : K \rightarrow CK(E)$ a multivalued nonexpansive mapping. Suppose that the asymptotic center in E of each bounded sequence of X is nonempty and compact. Then T has a fixed point.*

In 1999, Sahu [5] obtained the strong convergence theorems of the nonexpansive type and nonself multivalued mappings for the following (1.3) iteration process:

$$x_n = t_n u + (1 - t_n) y_n, \quad n \geq 0, \quad (1.3)$$

where $y_n \in Tx_n$, $u \in K$, $t_n \in (0, 1)$ and $\lim_{n \rightarrow \infty} t_n = 0$. He proved that $\{x_n\}$ converges strongly to some fixed points of T . Xu [6] extended Theorem 1.3 to a multivalued nonexpansive nonself mapping and obtained the fixed theorem in 2001. The recent fixed point results for nonexpansive mappings can be found in [7–12] and references therein.

Recently, Panyanak [13] studied the Mann iteration (1.4) and Ishikawa iterative processes (1.5) for $x_0 \in K$ as follows:

$$x_{n+1} = \alpha_n x_n + (1 - \alpha_n) y_n, \quad n \geq 0, \quad (1.4)$$

where $\alpha_n \in [0, 1]$, $y_n \in Tx_n$, and fixed $p \in F(T)$ are such that $\|y_n - p\| \leq d(p, Tx_n)$,

$$\begin{aligned} y_n &= (1 - \beta_n) x_n + \beta_n z_n, \\ x_{n+1} &= (1 - \alpha_n) x_n + \alpha_n z'_n, \quad n \geq 0, \end{aligned} \quad (1.5)$$

where $\alpha_n \in [0, 1]$, $\beta_n \in [0, 1]$, $z_n \in Tx_n$, $z'_n \in Ty_n$, and fixed $p \in F(T)$ are such that $\|z_n - p\| \leq d(p, Tx_n)$ and $\|z'_n - p\| \leq d(p, Ty_n)$ and proved the strong convergence theorems for multivalued nonexpansive mappings T in Banach spaces.

In this paper, motivated by Panyanak [13] and the previous results, we will study the following iteration process (1.6). Let K be a nonempty convex subset of E , $\alpha_n \in [0, 1]$,

$$\begin{aligned} x_0 &\in K, \\ x_n &= \alpha_n x_{n-1} + (1 - \alpha_n) y_n, \quad y_n \in Tx_n, \quad n \geq 1, \end{aligned} \tag{1.6}$$

and we prove some strong convergence theorems of the sequence $\{x_n\}$ defined by (1.6) for nonexpansive multivalued mappings in Banach spaces. The results presented in this paper establish a new type iteration convergence theorems for multivalued nonexpansive mappings in Banach spaces and extend the corresponding results of Panyanak [13].

2. Preliminaries

Let E be a real Banach space and let J denote the normalized duality mapping from E to 2^{E^*} defined by

$$J(x) = \{f \in E^*, \langle x, f \rangle = \|x\| \|f\|, \|x\| = \|f\|\}, \quad \forall x \in E, \tag{2.1}$$

where E^* denotes the dual space of E and $\langle \cdot, \cdot \rangle$ denotes the generalized duality pair. It is well known that if E^* is strictly convex, then J is single valued. And if Banach space E admits sequentially continuous duality mapping J from weak topology to weak star topology, then, by [14, Lemma 1], we know that the duality mapping J is also single valued. In this case, the duality mapping J is also said to be weakly sequentially continuous; that is, if $\{x_n\}$ is a subject of E with $x_n \rightharpoonup x$, then $J(x_n) \xrightarrow{*} J(x)$. By Theorem 1 of [14], we know that if E admits a weakly sequentially continuous duality mapping, then E satisfies Opial's condition, and E is smooth; for the details, see [14]. In the sequel, we will denote the single-valued duality mapping by j .

Throughout this paper, we write $x_n \rightharpoonup x$ (resp., $x_n \xrightarrow{*} x$) to indicate that the sequence x_n weakly (resp., weak $*$) converges to x , as usual $x_n \rightarrow x$ will symbolize strong convergence. In order to show our main results, the following concepts and lemmas are needed.

A Banach space E is called uniformly convex if for each $\epsilon > 0$ there is a $\delta > 0$ such that for $x, y \in E$ with $\|x\|, \|y\| \leq 1$ and $\|x - y\| \geq \epsilon$, $\|x + y\| \leq 2(1 - \delta)$ holds. The modulus of convexity of E is defined by

$$\delta_E(\epsilon) = \inf \left\{ 1 - \left\| \frac{1}{2}(x + y) \right\| : \|x\|, \|y\| \leq 1, \|x - y\| \geq \epsilon \right\}, \tag{2.2}$$

for all $\epsilon \in [0, 2]$. E is said to be uniformly convex if $\delta_E(0) = 0$, and $\delta(\epsilon) > 0$ for all $0 < \epsilon \leq 2$.

Lemma 2.1 (see [10]). *In Banach space E , the following result (subdifferential inequality) is well known: for all $x, y \in E$, for all $j(x) \in J(x)$, for all $j(x+y) \in J(x+y)$,*

$$\|x\|^2 + 2\langle y, j(x) \rangle \leq \|x+y\|^2 \leq \|x\|^2 + 2\langle y, j(x+y) \rangle. \quad (2.3)$$

Definition 2.2. A Banach space E is said to satisfy *Opial's condition* if for any sequence $\{x_n\}$ in E , $x_n \rightharpoonup x$ ($n \rightarrow \infty$) implies

$$\limsup_{n \rightarrow \infty} \|x_n - x\| < \limsup_{n \rightarrow \infty} \|x_n - y\|, \quad \forall y \in E \text{ with } x \neq y. \quad (2.4)$$

We know that Hilbert spaces, l^p ($1 < p < \infty$), and Banach space with weakly sequentially continuous duality mappings satisfy Opial's condition; for the details, see [14, 15].

Definition 2.3. A multivalued mapping $T : K \rightarrow CB(K)$ is said to satisfy *Condition I* if there is a nondecreasing function $f : [0, \infty) \rightarrow [0, \infty)$ with $f(0) = 0$, $f(r) > 0$ for $r \in (0, \infty)$ such that

$$d(x, Tx) \geq f(d(x, F(T))), \quad \forall x \in K. \quad (2.5)$$

Example of mappings that satisfy *Condition I* can be founded in [13].

3. Main Results

Now, we prove our results.

Theorem 3.1. *Let K be a nonempty compact convex subset of a uniformly convex Banach space E and let $T : K \rightarrow CB(K)$ be a multivalued nonexpansive mapping, where $\alpha_n \in (0, 1)$ and $\lim_{n \rightarrow \infty} \alpha_n = 0$, the sequence $\{x_n\}_{n=1}^{\infty}$ is generated by (1.6).*

Then,

- (i) *by the compactness of K , there exists a subsequence $\{x_{n_i}\}$ of $\{x_n\}$ such that $x_{n_i} \rightarrow p$ for some $p \in K$. In addition if $\|y_n - p\| \leq d(p, Tx_n)$, then*
- (ii) *p is a fixed point of T and the sequence $\{x_n\}$ converges strongly to p .*

Proof. Part (i) is trivial. And part (ii) remains to be proved.

Due to the compactness of K and boundness of $CB(K)$, there exists a real number $M > 0$ such that

$$\|x_{n-1} - y_n\| \leq M. \quad (3.1)$$

It follows from (1.6), that

$$\|x_n - y_n\| = \alpha_n \|x_{n-1} - y_n\| \leq \alpha_n M, \quad (3.2)$$

thus

$$\|x_n - y_n\| \rightarrow 0, \quad \text{as } n \rightarrow \infty, \quad (3.3)$$

therefore

$$\begin{aligned} d(p, Tp) &\leq \|p - x_n\| + d(x_n, Tx_n) + H(Tx_n, Tp) \\ &\leq 2\|p - x_n\| + d(x_n, Tx_n) \\ &\leq 2\|p - x_n\| + \|x_n - y_n\|, \end{aligned} \quad (3.4)$$

so

$$d(p, Tp) \rightarrow 0, \quad \text{as } n \rightarrow \infty. \quad (3.5)$$

Hence, p is a fixed point of T .

Next we show that $\lim_{n \rightarrow \infty} \|x_n - p\|$ exists.

For all $n \geq 1$, there exist $j(x_n - p) \in J(x_n - p)$, using Lemma 2.1, we obtain

$$\begin{aligned} \|x_n - p\|^2 &= \langle \alpha_n x_{n-1} + (1 - \alpha_n) y_n - p, j(x_n - p) \rangle \\ &= (1 - \alpha_n) \langle y_n - p, j(x_n - p) \rangle + \alpha_n \langle x_{n-1} - p, j(x_n - p) \rangle \\ &\leq (1 - \alpha_n) \|y_n - p\| \cdot \|x_n - p\| + \alpha_n \|x_{n-1} - p\| \cdot \|x_n - p\| \\ &\leq (1 - \alpha_n) H(Tx_n, Tp) \cdot \|x_n - p\| + \alpha_n \|x_{n-1} - p\| \cdot \|x_n - p\| \\ &\leq (1 - \alpha_n) \|x_n - p\|^2 + \alpha_n \|x_{n-1} - p\| \cdot \|x_n - p\|, \end{aligned} \quad (3.6)$$

so

$$\|x_n - p\|^2 \leq \|x_{n-1} - p\| \cdot \|x_n - p\|. \quad (3.7)$$

If $\|x_n - p\| = 0$, then $\lim_{n \rightarrow \infty} \|x_n - p\| = 0$ apparently holds.

Let $\|x_n - p\| > 0$, from (3.7), we have

$$\|x_n - p\| \leq \|x_{n-1} - p\|. \quad (3.8)$$

We get that $\{\|x_n - p\|\}$ is a decreasing sequence, so

$$\lim_{n \rightarrow \infty} \|x_n - p\| \text{ exists.} \quad (3.9)$$

So the desired conclusion follows.

The proof is completed. \square

Remark 3.2. The above result modified the gap in the proof of Theorem 3.1 in [13] by a new method; the gap discovered by Song and Wang [16] is as follows.

Panyanak [13] introduced the Ishikawa iterates (1.5) of a multivalued mapping T . It is obvious that x_n depends on p and T . For $p \in F(T)$, we have

$$\begin{aligned}\|z_n - p\| &= d(p, Tx_n) \leq H(Tp, Tx_n) \leq \|x_n - p\|, \\ \|z'_n - p\| &= d(p, Ty_n) \leq H(Tp, Ty_n) \leq \|y_n - p\|.\end{aligned}\tag{3.10}$$

Clearly, if $q \in F(T)$ and $q \neq p$, then the above inequalities cannot be assured. Indeed, from the monotone decreasing sequence of $\{\|x_n - p\|\}$ in the proof of (Theorem 3.1 [13]), we cannot obtain that $\{\|x_n - q\|\}$ is a decreasing sequence. Hence, the conclusion of Theorem 3.1 in [13] cannot be achieved.

Theorem 3.3. *Let E be a Banach space satisfying Opial's condition and let K be a nonempty weakly compact convex subset of E . Suppose that $T : K \rightarrow CB(K)$ is a multivalued nonexpansive mapping, where $\alpha_n \in (0, 1)$ and $\lim_{n \rightarrow \infty} \alpha_n = 0$, the sequence $\{x_n\}_{n=1}^{\infty}$ is generated by (1.6).*

Then,

- (i) *by the weak compactness of K , there exists a subsequence $\{x_{n_i}\}$ of $\{x_n\}$ such that $x_{n_i} \rightharpoonup p$ for some $p \in K$. In addition if, $\|y_n - p\| \leq d(p, Tx_n)$, then*
- (ii) *p is a fixed point of T and the sequence $\{x_n\}$ converges weakly to p .*

Proof. Part (i) is trivial. Now we prove part (ii).

It follows from (3.3) of Theorem 3.1 that

$$\lim_{n \rightarrow \infty} d(x_n, Tx_n) = 0.\tag{3.11}$$

Since K is weakly compact, from part (i), there exists a subsequence $\{x_{n_i}\}$ of $\{x_n\}$ such that

$$x_{n_i} \rightharpoonup p, \quad \text{for some } p \in K.\tag{3.12}$$

Suppose that p does not belong to Tp . By the compactness of Tp , for any given x_{n_i} , there exist $z_i \in Tp$ such that $\|x_{n_i} - z_i\| = d(x_{n_i}, Tp)$ and $z_i \rightarrow z \in Tp$.

Thus $p \neq z$, from

$$\begin{aligned}\limsup_{i \rightarrow \infty} \|x_{n_i} - z\| &\leq \limsup_{i \rightarrow \infty} [\|x_{n_i} - z_i\| + \|z_i - z\|] \\ &= \limsup_{i \rightarrow \infty} \|x_{n_i} - z_i\| \\ &\leq \limsup_{i \rightarrow \infty} [d(x_{n_i}, Tx_{n_i}) + H(Tx_{n_i}, Tp)] \\ &\leq \limsup_{i \rightarrow \infty} \|x_{n_i} - p\| < \limsup_{i \rightarrow \infty} \|x_{n_i} - z\|.\end{aligned}\tag{3.13}$$

This is a contradiction by satisfying Opial's condition.

Hence, p is a fixed point of T .

It follows from (3.7) of Theorem 3.1 that

$$\lim_{n \rightarrow \infty} \|x_n - p\| \text{ exists.} \quad (3.14)$$

Next we show $x_n \rightarrow p$. Suppose not. There exists another subsequence $\{x_{n_k}\}$ of $\{x_n\}$ such that $x_{n_k} \rightarrow q \neq p$.

Then, we also obtain $q \in Tq$. From Opial's condition, we have

$$\begin{aligned} \lim_{i \rightarrow \infty} \|x_n - p\| &= \limsup_{i \rightarrow \infty} \|x_{n_i} - p\| \\ &< \limsup_{i \rightarrow \infty} \|x_{n_i} - q\| = \limsup_{k \rightarrow \infty} \|x_{n_k} - q\| \\ &< \limsup_{k \rightarrow \infty} \|x_{n_k} - p\| = \lim_{i \rightarrow \infty} \|x_n - p\|. \end{aligned} \quad (3.15)$$

Which is a contradiction, so the conclusion of the theorem follows.

The proof is completed. \square

Corollary 3.4. *Let E be a reflexive Banach space which admits a weakly sequentially continuous duality mapping J from E to E^* , and let K be a nonempty weakly compact convex subset of E . Suppose that $T : K \rightarrow CB(K)$ is a multivalued nonexpansive mapping, where $\alpha_n \in (0, 1)$ and $\lim_{n \rightarrow \infty} \alpha_n = 0$, the sequence $\{x_n\}_{n=1}^{\infty}$ is generated by (1.6).*

Then,

(i) *by the weak compactness of K , there exists a subsequence $\{x_{n_i}\}$ of $\{x_n\}$ such that $x_{n_i} \rightarrow p$ for some $p \in K$. In addition if, $\|y_n - p\| \leq d(p, Tx_n)$, then*

(ii) *p is a fixed point of T and the sequence $\{x_n\}$ converges weakly to p .*

Proposition 3.5. *Let K be a nonempty compact convex subset of a uniformly convex Banach space E and let $T : K \rightarrow CB(K)$ be a multivalued nonexpansive mapping. Then $F(T)$ is a closed subset of K .*

Proof. Suppose $q_n \in F(T)$, $n \geq 1$, such that $\lim_{n \rightarrow \infty} q_n = q$, then we have

$$\begin{aligned} d(q, Tq) &\leq \|p - q_n\| + d(q_n, Tq_n) + H(Tq_n, Tq) \\ &\leq 2\|q - q_n\| + d(q_n, Tq_n), \end{aligned} \quad (3.16)$$

so

$$d(q, Tq) \rightarrow 0, \quad \text{as } n \rightarrow \infty. \quad (3.17)$$

Hence, q is a fixed point of T .

Thus, $F(T)$ is a closed subset of K .

The proof is completed. \square

Theorem 3.6. *Let K be a nonempty compact convex subset of a uniformly convex Banach space E and let $T : K \rightarrow CB(K)$ be a multivalued nonexpansive mapping satisfying Condition I, where $\alpha_n \in (0, 1)$ and $\lim_{n \rightarrow \infty} \alpha_n = 0$, then the sequence $\{x_n\}_{n=1}^{\infty}$ generated by (1.6) converges strongly to a fixed point.*

Proof. It follows from (3.3) of Theorem 3.1 that

$$\lim_{n \rightarrow \infty} d(x_n, Tx_n) = 0. \quad (3.18)$$

The proof of remained part is omitted because it is similar to the proof of Theorem 3.8 in [13]. \square

Acknowledgment

The work was supported by the Fundamental Research Funds for the Central Universities, No. JY10000970006, and National Nature Science Foundation, No. 60974082.

References

- [1] J. T. Markin, "A fixed point theorem for set valued mappings," *Bulletin of the American Mathematical Society*, vol. 74, pp. 639–640, 1968.
- [2] S. B. Nadler Jr., "Multi-valued contraction mappings," *Pacific Journal of Mathematics*, vol. 30, pp. 475–488, 1969.
- [3] T. C. Lim, "A fixed point theorem for multivalued nonexpansive mappings in a uniformly convex Banach space," *Bulletin of the American Mathematical Society*, vol. 80, pp. 1123–1126, 1974.
- [4] W. A. Kirk and S. Massa, "Remarks on asymptotic and Chebyshev centers," *Houston Journal of Mathematics*, vol. 16, no. 3, pp. 357–364, 1990.
- [5] D. R. Sahu, "Strong convergence theorems for nonexpansive type and non-self-multi-valued mappings," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 37, no. 3, pp. 401–407, 1999.
- [6] H.-K. Xu, "Multivalued nonexpansive mappings in Banach spaces," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 43, no. 6, pp. 693–706, 2001.
- [7] Y. Feng and S. Liu, "Fixed point theorems for multi-valued contractive mappings and multi-valued Caristi type mappings," *Journal of Mathematical Analysis and Applications*, vol. 317, no. 1, pp. 103–112, 2006.
- [8] J. S. Jung, "Strong convergence theorems for multivalued nonexpansive nonself-mappings in Banach spaces," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 66, no. 11, pp. 2345–2354, 2007.
- [9] D. Turkoglu and I. Altun, "A fixed point theorem for multi-valued mappings and its applications to integral inclusions," *Applied Mathematics Letters*, vol. 20, no. 5, pp. 563–570, 2007.
- [10] T. Domínguez Benavides and B. Gavira, "The fixed point property for multivalued nonexpansive mappings," *Journal of Mathematical Analysis and Applications*, vol. 328, no. 2, pp. 1471–1483, 2007.
- [11] R. Chen and H. He, "Viscosity approximation of common fixed points of nonexpansive semigroups in Banach space," *Applied Mathematics Letters*, vol. 20, no. 7, pp. 751–757, 2007.
- [12] H. He, X. Wang, R. Chen, and N. Cakic, "Strong convergence theorems for the implicit iteration process for a finite family of hemicontractive mappings in Banach space," *Applied Mathematics Letters*, vol. 22, no. 7, pp. 990–993, 2009.
- [13] B. Panyanak, "Mann and Ishikawa iterative processes for multivalued mappings in Banach spaces," *Computers & Mathematics with Applications*, vol. 54, no. 6, pp. 872–877, 2007.
- [14] J.-P. Gossez and E. Lami Dozo, "Some geometric properties related to the fixed point theory for nonexpansive mappings," *Pacific Journal of Mathematics*, vol. 40, pp. 565–573, 1972.
- [15] K. Yanagi, "On some fixed point theorems for multivalued mappings," *Pacific Journal of Mathematics*, vol. 87, no. 1, pp. 233–240, 1980.
- [16] Y. Song and H. Wang, "Convergence of iterative algorithms for multivalued mappings in Banach spaces," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 70, no. 4, pp. 1547–1556, 2009.