

Research Article

Regularity Criterion for Weak Solution to the 3D Micropolar Fluid Equations

Yu-Zhu Wang and Zigao Chen

School of Mathematics and Information Sciences, North China University of Water Resources and Electric Power, Zhengzhou 450011, China

Correspondence should be addressed to Yu-Zhu Wang, yuzhu108@163.com

Received 20 March 2011; Accepted 26 June 2011

Academic Editor: Ch Tsitouras

Copyright © 2011 Y.-Z. Wang and Z. Chen. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Regularity criterion for the 3D micropolar fluid equations is investigated. We prove that, for some $T > 0$, if $\int_0^T \|v_{x_3}\|_{L^q}^p dt < \infty$, where $3/\varrho + 2/\rho \leq 1$ and $\varrho \geq 3$, then the solution (v, w) can be extended smoothly beyond $t = T$. The derivative v_{x_3} can be substituted with any directional derivative of v .

1. Introduction

In the paper, we investigate the initial value problem for the micropolar fluid equations in \mathbb{R}^3 :

$$\begin{aligned} \partial_t v - (\nu + \kappa)\Delta v + v \cdot \nabla v + \nabla \pi - 2\kappa \nabla \times w &= 0, \\ \partial_t w - \gamma \Delta w - (\alpha + \beta) \nabla \nabla \cdot w + 4\kappa w + v \cdot \nabla w - 2\kappa \nabla \times v &= 0, \\ \nabla \cdot v &= 0 \end{aligned} \quad (1.1)$$

with the initial value

$$t = 0 : \quad v = v_0(x), \quad w = w_0(x), \quad (1.2)$$

where $v(t, x)$, $w(t, x)$, and $\pi(t, x)$ stand for the divergence free velocity field, nondivergence free microrotation field (angular velocity of the rotation of the particles of the fluid), the scalar pressure, respectively $\nu > 0$ is the Newtonian kinetic viscosity, $\kappa > 0$ is the dynamics microrotation viscosity, and $\alpha, \beta, \gamma > 0$ are the angular viscosity (see, e.g., Lukaszewicz [1]).

The micropolar fluid equations was first proposed by Eringen [2]. It is a type of fluids which exhibits the microrotational effects and microrotational inertia and can be viewed as a non-Newtonian fluid. Physically, micropolar fluid may represent fluids that consists of rigid, randomly oriented (or spherical) particles suspended in a viscous medium, where the deformation of fluid particles is ignored. It can describe many phenomena appeared in a large number of complex fluids such as the suspensions, animal blood, and liquid crystals which cannot be characterized appropriately by the Navier-Stokes equations, and that is important to the scientists working with the hydrodynamic fluid problems and phenomena. For more background, we refer to [1] and references therein. Besides their physical applications, the micropolar fluid equations are also mathematically significant. The existences of weak and strong solutions for micropolar fluid equations were treated by Galdi and Rionero [3] and Yamaguchi [4], respectively. The convergence of weak solutions of the micropolar fluids in bounded domains of \mathbb{R}^n was investigated (see [5]). When the viscosities tend to zero, in the limit, a fluid governed by an Euler-like system was found. Fundamental mathematical issues such as the global regularity of their solutions have generated extensive research, and many interesting results have been obtained (see [6–8]). A Beale-Kato-Madja criterion (see [9]) of smooth solutions to a related model with (1.1) was established in [10].

If $\kappa = 0$ and $w = 0$, then (1.1) reduces to be the Navier-Stokes equations. Besides its physical applications, the Navier-Stokes equations are also mathematically significant. In the last century, Leray [11] and Hopf [12] constructed weak solutions to the Navier-Stokes equations. The solution is called the Leray-Hopf weak solution. Later on, much effort has been devoted to establish the global existence and uniqueness of smooth solutions to the Navier-Stokes equations. Different criteria for regularity of the weak solutions have been proposed, and many interesting results are established (see [13–31]).

The purpose of this paper is to establish the regularity criteria of weak solutions to (1.1), (1.2) via the derivative of the velocity in one direction. It is proved that if $\int_0^T \|v_{x_3}\|_{L^q}^\rho dt < \infty$ with

$$\frac{3}{q} + \frac{2}{\rho} \leq 1, \quad q \geq 3, \quad (1.3)$$

then the solution (v, w) can be extended smoothly beyond $t = T$.

The paper is organized as follows. We first state some important inequalities in Section 2, which play an important roles in the proof of our main result. Then, we give definition of weak solution and state main results in Section 3 and then prove main result in Section 4.

2. Preliminaries

In order to prove our main result, we need the following Lemma, which may be found in [32] (see also [33, 34]). For the convenience of the readers, the proof of the Lemmas are provided.

Lemma 2.1. *Assume that $\mu, \lambda, \iota \in \mathbb{R}$ and satisfy*

$$1 \leq \mu, \lambda < \infty, \quad \frac{1}{\mu} + \frac{2}{\lambda} > 1, \quad 1 + \frac{3}{\iota} = \frac{1}{\mu} + \frac{2}{\lambda}. \quad (2.1)$$

Assume that $f \in H^1(\mathbb{R}^3)$, $f_{x_1}, f_{x_2} \in L^\lambda(\mathbb{R}^3)$, and $f_{x_3} \in L^\mu(\mathbb{R}^3)$. Then, there exists a positive constant such that

$$\|f\|_{L^t} \leq C \|f_{x_1}\|_{L^\lambda}^{1/3} \|f_{x_2}\|_{L^\lambda}^{1/3} \|f_{x_3}\|_{L^\mu}^{1/3}. \quad (2.2)$$

Especially, when $\lambda = 2$, there exists a positive constant $C = C(\mu)$ such that

$$\|f\|_{L^{3\mu}} \leq C \|f_{x_1}\|_{L^2}^{1/3} \|f_{x_2}\|_{L^2}^{1/3} \|f_{x_3}\|_{L^\mu}^{1/3}, \quad (2.3)$$

which holds for any $f \in H^1(\mathbb{R}^3)$ and $f_{x_3} \in L^\mu(\mathbb{R}^3)$ with $1 \leq \mu < \infty$.

Proof. It is not difficult to find

$$\begin{aligned} |f(x_1, x_2, x_3)|^{1+(1-1/\lambda)t} &\leq C \int_{-\infty}^{x_1} |f(x_1, x_2, x_3)|^{(1-(1/\lambda))t} |\partial_\tau f(\tau, x_2, x_3)| d\tau, \\ |f(x_1, x_2, x_3)|^{1+(1-1/\lambda)t} &\leq C \int_{-\infty}^{x_2} |f(x_1, x_2, x_3)|^{(1-(1/\lambda))t} |\partial_\tau f(x_1, \tau, x_3)| d\tau, \\ |f(x_1, x_2, x_3)|^{1+(1-1/\mu)t} &\leq C \int_{-\infty}^{x_3} |f(x_1, x_2, x_3)|^{(1-(1/\mu))t} |\partial_\tau f(x_1, x_2, \tau)| d\tau. \end{aligned} \quad (2.4)$$

Then, we obtain

$$\begin{aligned} |f(x_1, x_2, x_3)|^t &\leq C \left[\int_{-\infty}^{\infty} |f(x_1, x_2, x_3)|^{(1-1/\lambda)t} |\partial_{x_1} f(x_1, x_2, x_3)| dx_1 \right]^{1/2} \\ &\quad \times \left[\int_{-\infty}^{\infty} |f(x_1, x_2, x_3)|^{(1-1/\lambda)t} |\partial_{x_2} f(x_1, x_2, x_3)| dx_2 \right]^{1/2} \\ &\quad \times \left[\int_{-\infty}^{\infty} |f(x_1, x_2, x_3)|^{(1-1/\mu)t} |\partial_{x_3} f(x_1, x_2, x_3)| dx_3 \right]^{1/2}. \end{aligned} \quad (2.5)$$

Integrating with respect to x_1 and using Hölder inequality, we have

$$\begin{aligned} \int_{-\infty}^{\infty} |f(x_1, x_2, x_3)|^t dx_1 &\leq C \left[\int_{-\infty}^{\infty} |f(x_1, x_2, x_3)|^{(1-1/\lambda)t} |\partial_{x_1} f(x_1, x_2, x_3)| dx_1 \right]^{1/2} \\ &\quad \times \left[\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} |f(x_1, x_2, x_3)|^{(1-1/\lambda)t} |\partial_{x_2} f(x_1, x_2, x_3)| dx_2 dx_1 \right]^{1/2} \\ &\quad \times \left[\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} |f(x_1, x_2, x_3)|^{(1-1/\mu)t} |\partial_{x_3} f(x_1, x_2, x_3)| dx_3 dx_1 \right]^{1/2}. \end{aligned} \quad (2.6)$$

Integrating with respect to x_2, x_3 and using Hölder inequality, we obtain

$$\begin{aligned} \int_{\mathbb{R}^3} |f(x_1, x_2, x_3)|^t dx &\leq C \left[\int_{-\infty}^{\infty} |f(x_1, x_2, x_3)|^{(1-1/\lambda)t} |\partial_{x_1} f(x_1, x_2, x_3)| dx \right]^{1/2} \\ &\quad \times \left[\int_{\mathbb{R}^3} |f(x_1, x_2, x_3)|^{(1-1/\lambda)t} |\partial_{x_2} f(x_1, x_2, x_3)| dx \right]^{1/2} \\ &\quad \times \left[\int_{\mathbb{R}^3} |f(x_1, x_2, x_3)|^{(1-1/\mu)t} |\partial_{x_3} f(x_1, x_2, x_3)| dx \right]^{1/2}. \end{aligned} \quad (2.7)$$

It follows from Hölder inequality that

$$\|f\|_{L^t}^t \leq C \|f\|_{L^t}^{(1-1/\lambda)t/2} \|\partial_{x_1} f\|_{L^\lambda}^{1/2} \|f\|_{L^t}^{(1-1/\lambda)t/2} \|\partial_{x_2} f\|_{L^\lambda}^{1/2} \|f\|_{L^t}^{(1-1/\mu)t/2} \|\partial_{x_3} f\|_{L^\mu}^{1/2}. \quad (2.8)$$

By the above inequality, we get (2.2). \square

Lemma 2.2. *Let $2 \leq q \leq 6$ and assume that $f \in H^1(\mathbb{R}^3)$. Then, there exists a positive constant $C = C(q)$ such that*

$$\|f\|_{L^q} \leq C \|f\|_{L^2}^{(6-q)/2q} \|\partial_{x_1} f\|_{L^2}^{(q-2)/2q} \|\partial_{x_2} f\|_{L^2}^{(q-2)/2q} \|\partial_{x_3} f\|_{L^2}^{(q-2)/2q}. \quad (2.9)$$

Proof. Using the interpolating inequality, we obtain

$$\|f\|_{L^q} \leq C \|f\|_{L^2}^{(6-q)/2q} \|f\|_{L^6}^{(3q-6)/2q}. \quad (2.10)$$

By (2.3) with $\mu = 2$, we have

$$\|f\|_{L^6} \leq C \|\partial_{x_1} f\|_{L^2}^{1/3} \|\partial_{x_2} f\|_{L^2}^{1/3} \|\partial_{x_3} f\|_{L^2}^{1/3}. \quad (2.11)$$

Combining (2.10) and (2.11) yields (2.9). \square

3. Main Results

Before stating our main results, we introduce some function spaces. Let

$$C_{0,\sigma}^\infty(\mathbb{R}^3) = \left\{ \varphi \in (C^\infty(\mathbb{R}^3))^3 : \nabla \cdot \varphi = 0 \right\} \subset (C^\infty(\mathbb{R}^3))^3. \quad (3.1)$$

The subspace

$$L_\sigma^2 = \overline{C_{0,\sigma}^\infty(\mathbb{R}^3)}^{\|\cdot\|_{L^2}} = \left\{ \varphi \in L^2(\mathbb{R}^3) : \nabla \cdot \varphi = 0 \right\} \quad (3.2)$$

is obtained as the closure of $C_{0,\sigma}^\infty$ with respect to L^2 -norm $\|\cdot\|_{L^2}$. H_σ^r is the closure of $C_{0,\sigma}^\infty$ with respect to the H^r -norm

$$\|\varphi\|_{H^r} = \|(I - \Delta)^{r/2} \varphi\|_{L^2}, \quad r \geq 0. \quad (3.3)$$

Before stating our main results, we give the definition of weak solution to (1.1), (1.2) (see [6]).

Definition 3.1 (Weak solutions). Let $T > 0$, $v_0 \in L_\sigma^2(\mathbb{R}^3)$, and $w_0 \in L^2(\mathbb{R}^3)$. A measurable \mathbb{R}^3 -valued triple (v, w) is said to be a weak solution to (1.1), (1.2) on $[0, T]$ if the following conditions hold the following.

(1)

$$\begin{aligned} v &\in L^\infty(0, T; L_\sigma^2(\mathbb{R}^3)) \cap L^2(0, T; H_\sigma^1(\mathbb{R}^3)), \\ w &\in L^\infty(0, T; L^2(\mathbb{R}^3)) \cap L^2(0, T; H^1(\mathbb{R}^3)). \end{aligned} \quad (3.4)$$

(2) Equations (1.1), (1.2) are satisfied in the sense of distributions; that is, for every $\varphi \in H^1((0, T); H_\sigma^1)$ and $\psi \in H^1((0, T); H^1)$ with $\varphi(T) = \psi(T) = 0$, hold

$$\begin{aligned} \int_0^T \{-\langle v, \partial_\tau \varphi \rangle + \langle v \cdot \nabla v, \varphi \rangle + (v + \kappa) \langle \nabla v, \nabla \varphi \rangle\} d\tau - \int_0^T \{2\kappa \langle \nabla \times w, \varphi \rangle\} d\tau \\ = \langle v_0, \varphi(0) \rangle, \end{aligned} \quad (3.5)$$

$$\begin{aligned} \int_0^T \{-\langle w, \partial_\tau \psi \rangle + \gamma \langle \nabla w, \nabla \psi \rangle + (\alpha + \beta) \langle \nabla \cdot w, \nabla \psi \rangle + 4\kappa \langle w, \psi \rangle\} d\tau \\ + \int_0^T \{\langle v \cdot \nabla w, \psi \rangle - 2\kappa \langle \nabla \times v, \psi \rangle\} d\tau \\ = \langle w_0, \psi(0) \rangle. \end{aligned} \quad (3.6)$$

(3) The energy inequality, that is,

$$\begin{aligned} \|v(t)\|_{L^2}^2 + \|w(t)\|_{L^2}^2 + 2 \int_0^t \left(\nu \|\nabla v(\tau)\|_{L^2}^2 + \gamma \|\nabla w(\tau)\|_{L^2}^2 \right) d\tau + 2(\alpha + \beta) \int_0^t \|\nabla \cdot w(\tau)\|_{L^2}^2 d\tau \\ \leq \|v_0\|_{L^2}^2 + \|w_0\|_{L^2}^2. \end{aligned} \quad (3.7)$$

Theorem 3.2. Let $v_0 \in H_\sigma^1(\mathbb{R}^3)$ with $w_0 \in H^1(\mathbb{R}^3)$. Assume that (v, w) is a weak solution to (1.1), (1.2) on some interval $[0, T]$. If

$$\Theta(T) \equiv \int_0^T \|v_{x_3}\|_{L^q}^p dt < \infty, \quad (3.8)$$

where

$$\frac{3}{\varrho} + \frac{2}{\rho} \leq 1, \quad \varrho \geq 3, \quad (3.9)$$

then the solution (v, w) can be extended smoothly beyond $t = T$.

4. Proof of Theorem 3.2

Proof. Multiplying the first equation of (1.1) by v and integrating with respect to x on \mathbb{R}^3 , using integration by parts, we obtain

$$\frac{1}{2} \frac{d}{dt} \|v(t)\|_{L^2}^2 + (\nu + \kappa) \|\nabla v(t)\|_{L^2}^2 = 2\kappa \int_{\mathbb{R}^3} (\nabla \times w) \cdot v dx. \quad (4.1)$$

Similarly, we get

$$\frac{1}{2} \frac{d}{dt} \|w(t)\|_{L^2}^2 + \gamma \|\nabla w(t)\|_{L^2}^2 + (\alpha + \beta) \|\nabla \cdot w\|_{L^2}^2 + 4\kappa \|w\|_{L^2}^2 = 2\kappa \int_{\mathbb{R}^3} (\nabla \times v) \cdot w dx. \quad (4.2)$$

Summing up (4.1)-(4.2), we deduce that

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \left(\|v(t)\|_{L^2}^2 + \|w(t)\|_{L^2}^2 \right) + (\nu + \kappa) \|\nabla v(t)\|_{L^2}^2 \\ & \quad + \gamma \|\nabla w(t)\|_{L^2}^2 + (\alpha + \beta) \|\nabla \cdot w\|_{L^2}^2 + 4\kappa \|w\|_{L^2}^2 \\ & = 2\kappa \int_{\mathbb{R}^3} (\nabla \times w) \cdot v dx + 2\kappa \int_{\mathbb{R}^3} (\nabla \times v) \cdot w dx. \end{aligned} \quad (4.3)$$

By integration by parts and Cauchy inequality, we obtain

$$2\kappa \int_{\mathbb{R}^3} (\nabla \times w) \cdot v dx + 2\kappa \int_{\mathbb{R}^3} (\nabla \times v) \cdot w dx \leq \kappa \|\nabla v\|_{L^2}^2 + 4\kappa \|w\|_{L^2}^2. \quad (4.4)$$

Combining (4.3)-(4.4) yields

$$\frac{1}{2} \frac{d}{dt} \left(\|v(t)\|_{L^2}^2 + \|w(t)\|_{L^2}^2 \right) + \nu \|\nabla v(t)\|_{L^2}^2 + \gamma \|\nabla w(t)\|_{L^2}^2 + (\alpha + \beta) \|\nabla \cdot w\|_{L^2}^2 \leq 0. \quad (4.5)$$

Integrating with respect to t , we have

$$\begin{aligned} & \|v(t)\|_{L^2}^2 + \|w(t)\|_{L^2}^2 + 2 \int_0^t \left(\nu \|\nabla v(\tau)\|_{L^2}^2 + \gamma \|\nabla w(\tau)\|_{L^2}^2 \right) d\tau + 2(\alpha + \beta) \int_0^t \|\nabla \cdot w(\tau)\|_{L^2}^2 d\tau \\ & \leq \|v_0\|_{L^2}^2 + \|w_0\|_{L^2}^2. \end{aligned} \quad (4.6)$$

Differentiating (1.1) with respect to x_3 , we obtain

$$\begin{aligned} \partial_t v_{x_3} - (\nu + \kappa) \Delta v_{x_3} + v_{x_3} \cdot \nabla v + v \cdot \nabla v_{x_3} + \nabla \pi_{x_3} - 2\kappa \nabla \times w_{x_3} &= 0, \\ \partial_t w_{x_3} - \gamma \Delta w_{x_3} - (\alpha + \beta) \nabla \cdot \nabla w_{x_3} + 4\kappa w_{x_3} + v_{x_3} \cdot \nabla w + v \cdot \nabla w_{x_3} - 2\kappa \nabla \times v_{x_3} &= 0. \end{aligned} \quad (4.7)$$

Taking the inner product of v_{x_3} with the first equation of (4.7) and using integration by parts yields

$$\frac{1}{2} \frac{d}{dt} \|v_{x_3}(t)\|_{L^2}^2 + (\nu + \kappa) \|\nabla v_{x_3}(t)\|_{L^2}^2 = - \int_{\mathbb{R}^3} v_{x_3} \cdot \nabla v \cdot v_{x_3} dx + 2\kappa \int_{\mathbb{R}^3} (\nabla \times w_{x_3}) \cdot v_{x_3} dx. \quad (4.8)$$

Similarly, we get

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \|w_{x_3}(t)\|_{L^2}^2 + \gamma \|\nabla w_{x_3}(t)\|_{L^2}^2 + (\alpha + \beta) \|\nabla \cdot w_{x_3}\|_{L^2}^2 + 4\kappa \|w_{x_3}\|_{L^2}^2 \\ = - \int_{\mathbb{R}^3} v_{x_3} \cdot \nabla w \cdot w_{x_3} dx + 2\kappa \int_{\mathbb{R}^3} (\nabla \times v_{x_3}) \cdot w_{x_3} dx. \end{aligned} \quad (4.9)$$

Combining (4.8)–(4.9) yields

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \left(\|v_{x_3}(t)\|_{L^2}^2 + \|w_{x_3}(t)\|_{L^2}^2 \right) + (\nu + \kappa) \|\nabla v_{x_3}(t)\|_{L^2}^2 \\ + \gamma \|\nabla w_{x_3}(t)\|_{L^2}^2 + (\alpha + \beta) \|\nabla \cdot w_{x_3}\|_{L^2}^2 + 4\kappa \|w_{x_3}\|_{L^2}^2 \\ = - \int_{\mathbb{R}^3} v_{x_3} \cdot \nabla v \cdot v_{x_3} dx + 2\kappa \int_{\mathbb{R}^3} (\nabla \times w_{x_3}) \cdot v_{x_3} dx \\ - \int_{\mathbb{R}^3} v_{x_3} \cdot \nabla w \cdot w_{x_3} dx + 2\kappa \int_{\mathbb{R}^3} (\nabla \times v_{x_3}) \cdot w_{x_3} dx. \end{aligned} \quad (4.10)$$

Using integration by parts and Cauchy inequality, we obtain

$$2\kappa \int_{\mathbb{R}^3} (\nabla \times w_{x_3}) \cdot v_{x_3} dx + 2\kappa \int_{\mathbb{R}^3} (\nabla \times v_{x_3}) \cdot w_{x_3} dx \leq \kappa \|\nabla v_{x_3}\|_{L^2}^2 + 4\kappa \|w_{x_3}\|_{L^2}^2. \quad (4.11)$$

Combining (4.10)–(4.11) yields

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \left(\|v_{x_3}(t)\|_{L^2}^2 + \|w_{x_3}(t)\|_{L^2}^2 \right) + \nu \|\nabla v_{x_3}(t)\|_{L^2}^2 \\ + \gamma \|\nabla w_{x_3}(t)\|_{L^2}^2 + (\alpha + \beta) \|\nabla \cdot w_{x_3}\|_{L^2}^2 \\ \leq - \int_{\mathbb{R}^3} v_{x_3} \cdot \nabla v \cdot v_{x_3} dx - \int_{\mathbb{R}^3} v_{x_3} \cdot \nabla w \cdot w_{x_3} dx \\ \triangleq I_1 + I_2. \end{aligned} \quad (4.12)$$

In what follows, we estimate I_j ($j = 1, 2, \dots, 5$). By integration by parts and Hölder inequality, we obtain

$$I_1 \leq C \|\nabla v_{x_3}\|_{L^2} \|v_{x_3}\|_{L^\sigma} \|v\|_{L^{3\varrho}}, \quad (4.13)$$

where

$$\frac{1}{\sigma} + \frac{1}{3\varrho} = \frac{1}{2}, \quad 2 \leq \sigma \leq 6. \quad (4.14)$$

It follows from the interpolating inequality that

$$\|v_{x_3}\|_{L^\sigma} \leq C \|v_{x_3}\|_{L^2}^{1-3(1/2-1/\sigma)} \|\nabla v_{x_3}\|_{L^2}^{3(1/2-1/\sigma)}. \quad (4.15)$$

From (2.3), we get

$$\begin{aligned} I_1 &\leq C \|\nabla v_{x_3}\|_{L^2} \|v_{x_3}\|_{L^2}^{1-3(1/2-1/\sigma)} \|\nabla v_{x_3}\|_{L^2}^{3(1/2-1/\sigma)} \|\nabla v\|_{L^2}^{2/3} \|v_{x_3}\|_{L^\varrho}^{1/3} \\ &\leq C \|\nabla v_{x_3}\|_{L^2}^{1+3(1/2-1/\sigma)} \|v_{x_3}\|_{L^2}^{1-3(1/2-1/\sigma)} \|\nabla v\|_{L^2}^{2/3} \|v_{x_3}\|_{L^\varrho}^{1/3} \\ &\leq \frac{\nu}{2} \|\nabla v_{x_3}\|_{L^2}^2 + C \|v_{x_3}\|_{L^2}^2 \|\nabla v\|_{L^2}^{2q} \|v_{x_3}\|_{L^2}^q, \end{aligned} \quad (4.16)$$

where

$$q = \frac{2}{3-9(1/2-1/\sigma)} = \frac{2}{3(1-1/\varrho)}. \quad (4.17)$$

When $\varrho \geq 3$, we have $2q \leq 2$ and application of Young inequality yields

$$I_1 \leq \frac{\nu}{2} \|\nabla v_{x_3}\|_{L^2}^2 + C \|v_{x_3}\|_{L^2}^2 \left(\|\nabla v\|_{L^2}^2 + \|v_{x_3}\|_{L^\varrho}^\delta \right), \quad (4.18)$$

where

$$\frac{3}{\varrho} + \frac{2}{\delta} = 1. \quad (4.19)$$

From Hölder inequality, we obtain

$$\begin{aligned} I_2 &\leq C \|\nabla w\|_{L^2} \|w_{x_3}\|_{L^{2\varrho/(\varrho-2)}} \|v_{x_3}\|_{L^\varrho} \\ &\leq C \|\nabla w\|_{L^2} \|v_{x_3}\|_{L^\varrho} \|w_{x_3}\|_{L^2}^{1-3/\varrho} \|\nabla w_{x_3}\|_{L^2}^{3/\varrho} \\ &\leq C \|\nabla w_{x_3}\|_{L^2}^2 + \|\nabla w\|_{L^2}^{2\varrho/(2\varrho-3)} \|v_{x_3}\|_{L^\varrho}^{2\varrho/(2\varrho-3)} \|w_{x_3}\|_{L^2}^{(2\varrho-6)/(2\varrho-3)} \\ &\leq \frac{\gamma}{2} \|\nabla w_{x_3}\|_{L^2}^2 + C \left(\|\nabla w\|_{L^2}^2 + \|v_{x_3}\|_{L^\varrho}^\delta \right) \|w_{x_3}\|_{L^2}^{(2\varrho-6)/(2\varrho-3)}, \end{aligned} \quad (4.20)$$

where

$$\frac{3}{\varrho} + \frac{2}{\delta} = 1. \quad (4.21)$$

Combining (4.12)–(4.20) yields

$$\begin{aligned} & \frac{d}{dt} \left(\|v_{x_3}\|_{L^2}^2 + \|w_{x_3}\|_{L^2}^2 \right) + \nu \|\nabla v_{x_3}\|_{L^2}^2 + \gamma \|\nabla w_{x_3}\|_{L^2}^2 + (\alpha + \beta) \|\nabla \cdot w_{x_3}\|_{L^2}^2 \\ & \leq C \|v_{x_3}\|_{L^2}^2 \left(\|\nabla v\|_{L^2}^2 + \|v_{x_3}\|_{L^\varrho}^\delta \right) + C \left(\|\nabla w\|_{L^2}^2 + \|v_{x_3}\|_{L^\varrho}^\delta \right) \|w_{x_3}\|_{L^2}^{(2\varrho-6)/(2\varrho-3)}. \end{aligned} \quad (4.22)$$

From Gronwall inequality, we get

$$\begin{aligned} & \|v_{x_3}\|_{L^2}^2 + \|w_{x_3}\|_{L^2}^2 + \nu \int_0^t \|\nabla v_{x_3}\|_{L^2}^2 d\tau + \int_0^t \left(\gamma \|\nabla w_{x_3}\|_{L^2}^2 + (\alpha + \beta) \|\nabla \cdot w_{x_3}\|_{L^2}^2 \right) d\tau \\ & \leq C e^{(\|v_0\|_{L^2}^2 + \|w_0\|_{L^2}^2)} e^{\Theta(t)} \left[\|v_0\|_{H^1}^2 + \|w_0\|_{H^1}^2 + C \left(\|v_0\|_{L^2}^2 + \|w_0\|_{L^2}^2 + \Theta(t) \right)^{2\varrho-3/\varrho} \right]. \end{aligned} \quad (4.23)$$

Multiplying the first equation of (1.1) by $-\Delta v$ and integrating with respect to x on \mathbb{R}^3 , then using integration by parts, we obtain

$$\frac{1}{2} \frac{d}{dt} \|\nabla v(t)\|_{L^2}^2 + (\nu + \kappa) \|\Delta v\|_{L^2}^2 = \int_{\mathbb{R}^3} v \cdot \nabla v \cdot \Delta v dx - 2\kappa \int_{\mathbb{R}^3} (\nabla \times w) \cdot \Delta v dx. \quad (4.24)$$

Similarly, we get

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \|\nabla w(t)\|_{L^2}^2 + \gamma \|\Delta w\|_{L^2}^2 + (\alpha + \beta) \|\nabla \nabla \cdot w\|_{L^2}^2 + 4\kappa \|\nabla w\|_{L^2}^2 \\ & = \int_{\mathbb{R}^3} v \cdot \nabla w \cdot \Delta w dx - 2\kappa \int_{\mathbb{R}^3} (\nabla \times v) \cdot \Delta w dx. \end{aligned} \quad (4.25)$$

Collecting (4.24) and (4.25) yields

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \left(\|\nabla v(t)\|_{L^2}^2 + \|\nabla w(t)\|_{L^2}^2 \right) + (\nu + \kappa) \|\Delta v\|_{L^2}^2 \\ & \quad + \gamma \|\Delta w\|_{L^2}^2 + (\alpha + \beta) \|\nabla \nabla \cdot w\|_{L^2}^2 + 4\kappa \|\nabla w\|_{L^2}^2 \\ & = \int_{\mathbb{R}^3} v \cdot \nabla v \cdot \Delta v dx - 2\kappa \int_{\mathbb{R}^3} (\nabla \times w) \cdot \Delta v dx \\ & \quad + \int_{\mathbb{R}^3} v \cdot \nabla w \cdot \Delta w dx - 2\kappa \int_{\mathbb{R}^3} (\nabla \times v) \cdot \Delta w dx. \end{aligned} \quad (4.26)$$

Thanks to integration by parts and Cauchy inequality, we get

$$-2\kappa \int_{\mathbb{R}^3} (\nabla \times w) \cdot \Delta v dx - 2\kappa \int_{\mathbb{R}^3} (\nabla \times v) \cdot \Delta w dx \leq \kappa \|\Delta v\|_{L^2}^2 + 4\kappa \|\nabla w\|_{L^2}^2. \quad (4.27)$$

It follows from (4.26)-(4.27) and integration by parts that

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \left(\|\nabla v(t)\|_{L^2}^2 + \|\nabla w(t)\|_{L^2}^2 \right) + \nu \|\Delta v\|_{L^2}^2 + \gamma \|\Delta w\|_{L^2}^2 + (\alpha + \beta) \|\nabla \nabla \cdot w\|_{L^2}^2 \\ & \leq - \int_{\mathbb{R}^3} \nabla v \cdot \nabla v \cdot \nabla v dx - \int_{\mathbb{R}^3} \nabla v \cdot \nabla w \cdot \nabla w dx \\ & \triangleq J_1 + J_2. \end{aligned} \quad (4.28)$$

In what follows, we estimate $J_i (i = 1, 2)$.

By (2.9) and Young inequality, we deduce that

$$\begin{aligned} J_1 & \leq C \|\nabla v\|_{L^3}^3 \\ & \leq C \|\nabla v\|_{L^2}^{3/2} \|\nabla_{\bar{x}} \nabla v\|_{L^2} \|\nabla v_{x_3}\|_{L^2}^{1/2} \\ & \leq \frac{\nu}{4} \|\nabla_{\bar{x}} \nabla v\|_{L^2}^2 + C \|\nabla v\|_{L^2}^3 \|\nabla v_{x_3}\|_{L^2} \\ & \leq \frac{\nu}{4} \|\nabla_{\bar{x}} \nabla v\|_{L^2}^2 + C \left(\|\nabla v\|_{L^2}^2 + \|\nabla v_{x_3}\|_{L^2}^2 \right) \|\nabla v\|_{L^2}^2, \end{aligned} \quad (4.29)$$

where $\nabla_{\bar{x}} = (\partial_{x_1}, \partial_{x_2})$.

By (2.9) and Young inequality, we have

$$\begin{aligned} J_2 & \leq \|\nabla v\|_{L^3} \|\nabla w\|_{L^3}^2 \\ & \leq C \|\nabla v\|_{L^2}^{1/2} \|\nabla_{\bar{x}} \nabla v\|_{L^2}^{1/3} \|\nabla v_{x_3}\|_{L^2}^{1/6} \|\nabla w\|_{L^2} \|\nabla_{\bar{x}} \nabla w\|_{L^2}^{2/3} \|\nabla w_{x_3}\|_{L^2}^{1/3} \\ & \leq \frac{\nu}{4} \|\nabla_{\bar{x}} \nabla v\|_{L^2}^2 + C \|\nabla v\|_{L^2}^{3/5} \|\nabla v_{x_3}\|_{L^2}^{1/5} \|\nabla w\|_{L^2}^{6/5} \|\nabla_{\bar{x}} \nabla w\|_{L^2}^{4/5} \|\nabla w_{x_3}\|_{L^2}^{2/5} \\ & \leq \frac{\nu}{4} \|\nabla_{\bar{x}} \nabla v\|_{L^2}^2 + \frac{\gamma}{2} \|\nabla_{\bar{x}} \nabla w\|_{L^2}^2 + C \|\nabla v\|_{L^2} \|\nabla v_{x_3}\|_{L^2}^{1/3} \|\nabla w\|_{L^2}^2 \|\nabla w_{x_3}\|_{L^2}^{2/3} \\ & \leq \frac{\nu}{4} \|\nabla_{\bar{x}} \nabla v\|_{L^2}^2 + \frac{\gamma}{2} \|\nabla_{\bar{x}} \nabla w\|_{L^2}^2 + C \|\nabla w\|_{L^2}^2 \left(\|\nabla v\|_{L^2}^2 + \|\nabla v_{x_3}\|_{L^2}^2 + \|\nabla w_{x_3}\|_{L^2}^2 \right), \end{aligned} \quad (4.30)$$

where $\nabla_{\bar{x}} = (\partial_{x_1}, \partial_{x_2})$.

Combining (4.28)-(4.30) yields

$$\begin{aligned} & \frac{d}{dt} \left(\|\nabla v(t)\|_{L^2}^2 + \|\nabla w(t)\|_{L^2}^2 \right) + \nu \|\Delta v\|_{L^2}^2 + \gamma \|\Delta w\|_{L^2}^2 + (\alpha + \beta) \|\nabla \nabla \cdot w\|_{L^2}^2 \\ & \leq C \left(\|\nabla v\|_{L^2}^2 + \|\nabla w\|_{L^2}^2 \right) \left(\|\nabla v\|_{L^2}^2 + \|\nabla v_{x_3}\|_{L^2}^2 + \|\nabla w_{x_3}\|_{L^2}^2 \right). \end{aligned} \quad (4.31)$$

From (4.31), Gronwall inequality, (4.6), and (4.23), we know that $(v, w) \in L^\infty(0, T; H^1(\mathbb{R}^3))$. Thus, (v, w) can be extended smoothly beyond $t = T$. We have completed the proof of Theorem 3.2. \square

Acknowledgments

This work was supported in part by the NNSF of China (Grant no. 10971190) and the Research Initiation Project for High-level Talents (201031) of the North China University of Water Resources and Electric Power.

References

- [1] G. Łukaszewicz, *Micropolar Fluids. Theory and Applications*, Modeling and Simulation in Science, Engineering and Technology, Birkhäuser Boston, Boston, Mass, USA, 1999.
- [2] A. C. Eringen, "Theory of micropolar fluids," *Journal of Mathematics and Mechanics*, vol. 16, pp. 1–18, 1966.
- [3] G. P. Galdi and S. Rionero, "A note on the existence and uniqueness of solutions of the micropolar fluid equations," *International Journal of Engineering Science*, vol. 15, no. 2, pp. 105–108, 1977.
- [4] N. Yamaguchi, "Existence of global strong solution to the micropolar fluid system in a bounded domain," *Mathematical Methods in the Applied Sciences*, vol. 28, no. 13, pp. 1507–1526, 2005.
- [5] E. Ortega-Torres, M. A. Rojas-Medar, and E. J. Villamizar-Roa, "Micropolar fluids with vanishing viscosity," *Abstract and Applied Analysis*, vol. 2010, Article ID 843692, 18 pages, 2010.
- [6] B.-Q. Dong and Z.-M. Chen, "Regularity criteria of weak solutions to the three-dimensional micropolar flows," *Journal of Mathematical Physics*, vol. 50, no. 10, p. 103525-1-13, 2009.
- [7] B.-Q. Dong and Z. Zhang, "Global regularity of the 2D micropolar fluid flows with zero angular viscosity," *Journal of Differential Equations*, vol. 249, no. 1, pp. 200–213, 2010.
- [8] E. Ortega-Torres and M. Rojas-Medar, "On the regularity for solutions of the micropolar fluid equations," *Rendiconti del Seminario Matematico della Università di Padova*, vol. 122, pp. 27–37, 2009.
- [9] J. T. Beale, T. Kato, and A. Majda, "Remarks on the breakdown of smooth solutions for the 3-D Euler equations," *Communications in Mathematical Physics*, vol. 94, no. 1, pp. 61–66, 1984.
- [10] Y. Z. Wang, L. Hu, and Y. X. Wang, "A Beale-Kato-Madja criterion for magneto-micropolar fluid equations with partial viscosity," *Boundary Value Problems*, vol. 2011, Article ID 128614, 14 pages, 2011.
- [11] J. Leray, "Sur le mouvement d'un liquide visqueux emplissant l'espace," *Acta Mathematica*, vol. 63, no. 1, pp. 193–248, 1934.
- [12] E. Hopf, "Über die Anfangswertaufgabe für die hydrodynamischen Grundgleichungen," *Mathematische Nachrichten*, vol. 4, pp. 213–231, 1951.
- [13] D. Chae, "On the regularity conditions for the Navier-Stokes and related equations," *Revista Matemática Iberoamericana*, vol. 23, no. 1, pp. 371–384, 2007.
- [14] C. H. Chan and A. Vasseur, "Log improvement of the Prodi-Serrin criteria for Navier-Stokes equations," *Methods and Applications of Analysis*, vol. 14, no. 2, pp. 197–212, 2007.
- [15] J. Fan and T. Ozawa, "Regularity criterion for weak solutions to the Navier-Stokes equations in terms of the gradient of the pressure," *Journal of Inequalities and Applications*, vol. 2008, Article ID 412678, 6 pages, 2008.
- [16] J. Fan, S. Jiang, G. Nakamura, and Y. Zhou, "Logarithmically improved regularity criteria for the navier-stokes and MHD equations," *Journal of Mathematical Fluid Mechanics*. In press.
- [17] C. He, "New sufficient conditions for regularity of solutions to the Navier-Stokes equations," *Advances in Mathematical Sciences and Applications*, vol. 12, no. 2, pp. 535–548, 2002.
- [18] H. Kozono, T. Ogawa, and Y. Taniuchi, "The critical Sobolev inequalities in Besov spaces and regularity criterion to some semi-linear evolution equations," *Mathematische Zeitschrift*, vol. 242, no. 2, pp. 251–278, 2002.
- [19] H. Kozono and N. Yatsu, "Extension criterion via two-components of vorticity on strong solutions to the 3D Navier-Stokes equations," *Mathematische Zeitschrift*, vol. 246, no. 1-2, pp. 55–68, 2004.
- [20] I. Kukavica and M. Ziane, "One component regularity for the Navier-Stokes equations," *Nonlinearity*, vol. 19, no. 2, pp. 453–469, 2006.

- [21] I. Kukavica and M. Ziane, "Navier-Stokes equations with regularity in one direction," *Journal of Mathematical Physics*, vol. 48, no. 6, p. 065203-1-10, 2007.
- [22] J. Serrin, "On the interior regularity of weak solutions of the Navier-Stokes equations," *Archive for Rational Mechanics and Analysis*, vol. 9, pp. 187–195, 1962.
- [23] Z. Zhifei and C. Qionglei, "Regularity criterion via two components of vorticity on weak solutions to the Navier-Stokes equations in \mathbb{R}^3 ," *Journal of Differential Equations*, vol. 216, no. 2, pp. 470–481, 2005.
- [24] Y. Zhou, "Regularity criteria in terms of pressure for the 3-D Navier-Stokes equations in a generic domain," *Mathematische Annalen*, vol. 328, no. 1-2, pp. 173–192, 2004.
- [25] Y. Zhou, "On regularity criteria in terms of pressure for the Navier-Stokes equations in \mathbb{R}^3 ," *Proceedings of the American Mathematical Society*, vol. 134, no. 1, pp. 149–156, 2006.
- [26] Y. Zhou, "On a regularity criterion in terms of the gradient of pressure for the Navier-Stokes equations in \mathbb{R}^N ," *Zeitschrift für Angewandte Mathematik und Physik*, vol. 57, no. 3, pp. 384–392, 2006.
- [27] Y. Zhou and S. Gala, "Logarithmically improved regularity criteria for the Navier-Stokes equations in multiplier spaces," *Journal of Mathematical Analysis and Applications*, vol. 356, no. 2, pp. 498–501, 2009.
- [28] Y. Zhou, "A new regularity criterion for the Navier-Stokes equations in terms of the gradient of one velocity component," *Methods and Applications of Analysis*, vol. 9, no. 4, pp. 563–578, 2002.
- [29] Y. Zhou, "A new regularity criterion for weak solutions to the Navier-Stokes equations," *Journal de Mathématiques Pures et Appliquées*, vol. 84, no. 11, pp. 1496–1514, 2005.
- [30] Y. Zhou and M. Pokorný, "On a regularity criterion for the Navier-Stokes equations involving gradient of one velocity component," *Journal of Mathematical Physics*, vol. 50, no. 12, pp. 123514-1–123514-11, 2009.
- [31] Y. Zhou and M. Pokorný, "On the regularity of the solutions of the Navier-Stokes equations via one velocity component," *Nonlinearity*, vol. 23, no. 5, pp. 1097–1107, 2010.
- [32] R. A. Adams, *Sobolev Spaces*, Academic Press, New York, NY, USA, 1975, Pure and Applied Mathematics.
- [33] G. P. Galdi, *An introduction to the Mathematical Theory of the Navier-Stokes Equations. Vol. I*, vol. I, II of *Springer Tracts in Natural Philosophy*, Springer, New York, NY, USA, 1994.
- [34] O. A. Ladyzhenskaya, *The Mathematical Theory of Viscous Incompressible Flow*, Second English edition, revised and enlarged. Translated from the Russian by Richard A. Silverman and John Chu. *Mathematics and its Applications, Vol. 2*, Gordon and Breach Science Publishers, New York, 1969.



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

