

Research Article

Left and Right Inverse Eigenpairs Problem for κ -Hermitian Matrices

Fan-Liang Li,¹ Xi-Yan Hu,² and Lei Zhang²

¹ Institute of Mathematics and Physics, School of Sciences, Central South University of Forestry and Technology, Changsha 410004, China

² College of Mathematics and Econometrics, Hunan University, Changsha 410082, China

Correspondence should be addressed to Fan-Liang Li; lfl302@tom.com

Received 6 December 2012; Accepted 20 March 2013

Academic Editor: Panayiotis J. Psarrakos

Copyright © 2013 Fan-Liang Li et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Left and right inverse eigenpairs problem for κ -hermitian matrices and its optimal approximate problem are considered. Based on the special properties of κ -hermitian matrices, the equivalent problem is obtained. Combining a new inner product of matrices, the necessary and sufficient conditions for the solvability of the problem and its general solutions are derived. Furthermore, the optimal approximate solution and a calculation procedure to obtain the optimal approximate solution are provided.

1. Introduction

Throughout this paper we use some notations as follows. Let $C^{n \times m}$ be the set of all $n \times m$ complex matrices, $UC^{n \times n}$, $HC^{n \times n}$, $SHC^{n \times n}$ denote the set of all $n \times n$ unitary matrices, hermitian matrices, skew-hermitian matrices, respectively. Let \bar{A} , A^H , and A^+ be the conjugate, conjugate transpose, and the Moore-Penrose generalized inverse of A , respectively. For $A, B \in C^{n \times m}$, $\langle A, B \rangle = \text{re}(\text{tr}(B^H A))$, where $\text{re}(\text{tr}(B^H A))$ denotes the real part of $\text{tr}(B^H A)$, the inner product of matrices A and B . The induced matrix norm is called Frobenius norm. That is, $\|A\| = \langle A, A \rangle^{1/2} = (\text{tr}(A^H A))^{1/2}$.

Left and right inverse eigenpairs problem is a special inverse eigenvalue problem. That is, giving partial left and right eigenpairs (eigenvalue and corresponding eigenvector), (λ_i, x_i) , $i = 1, \dots, h$; (μ_j, y_j) , $j = 1, \dots, l$, a special matrix set S , finding a matrix $A \in S$ such that

$$\begin{aligned} Ax_i &= \lambda_i x_i, & i &= 1, \dots, h, \\ y_j^T A &= \mu_j y_j^T, & j &= 1, \dots, l. \end{aligned} \quad (1)$$

This problem, which usually arises in perturbation analysis of matrix eigenvalues and in recursive matters, has profound application background [1–6]. When the matrix set S is different, it is easy to obtain different left and right inverse

eigenpairs problem. For example, we studied the left and right inverse eigenpairs problem of skew-centrosymmetric matrices and generalized centrosymmetric matrices, respectively [5, 6]. Based on the special properties of left and right eigenpairs of these matrices, we derived the solvability conditions of the problem and its general solutions. In this paper, combining the special properties of κ -hermitian matrices and a new inner product of matrices, we first obtain the equivalent problem, then derive the necessary and sufficient conditions for the solvability of the problem and its general solutions.

Hill and Waters [7] introduced the following matrices.

Definition 1. Let κ be a fixed product of disjoint transpositions, and let K be the associated permutation matrix, that is, $\bar{K} = K^H = K$, $K^2 = I_n$, a matrix $A \in C^{n \times n}$ is said to be κ -hermitian matrices (skew κ -hermitian matrices) if and only if $a_{ij} = \bar{a}_{k(j)k(i)}$ ($a_{ij} = -\bar{a}_{k(j)k(i)}$), $i, j = 1, \dots, n$. We denote the set of κ -hermitian matrices (skew κ -hermitian matrices) by $KHC^{n \times n}$ ($SKHC^{n \times n}$).

From Definition 1, it is easy to see that hermitian matrices and perhermitian matrices are special cases of κ -hermitian matrices, with $k(i) = i$ and $k(i) = n - i + 1$, respectively. Hermitian matrices and perhermitian matrices, which are one of twelve symmetry patterns of matrices [8], are applied in engineering, statistics, and so on [9, 10].

From Definition 1, it is also easy to prove the following conclusions.

- (1) $A \in KHC^{n \times n}$ if and only if $A = KA^H K$.
- (2) $A \in SKHC^{n \times n}$ if and only if $A = -KA^H K$.
- (3) If K is a fixed permutation matrix, then $KHC^{n \times n}$ and $SKHC^{n \times n}$ are the closed linear subspaces of $C^{n \times n}$ and satisfy

$$C^{n \times n} = KHC^{n \times n} \oplus SKHC^{n \times n}. \quad (2)$$

The notation $V_1 \oplus V_2$ stands for the orthogonal direct sum of linear subspace V_1 and V_2 .

- (4) $A \in KHC^{n \times n}$ if and only if there is a matrix $\tilde{A} \in HC^{n \times n}$ such that $\tilde{A} = KA$.
- (5) $A \in SKHC^{n \times n}$ if and only if there is a matrix $\tilde{A} \in SHC^{n \times n}$ such that $\tilde{A} = KA$.

Proof. (1) From Definition 1, if $A = (a_{ij}) \in KHC^{n \times n}$, then $a_{ij} = \bar{a}_{k(j)k(i)}$, this implies $A = KA^H K$, for $KA^H K = (\bar{a}_{k(j)k(i)})$.

(2) With the same method, we can prove (2). So, the proof is omitted.

- (3) (a) For any $A \in C^{n \times n}$, there exist $A_1 \in KHC^{n \times n}$, $A_2 \in SKHC^{n \times n}$ such that

$$A = A_1 + A_2, \quad (3)$$

where $A_1 = (1/2)(A + KA^H K)$, $A_2 = (1/2)(A - KA^H K)$.

- (b) If there exist another $\bar{A}_1 \in KHC^{n \times n}$, $\bar{A}_2 \in SKHC^{n \times n}$ such that

$$A = \bar{A}_1 + \bar{A}_2, \quad (4)$$

(3)-(4) yields

$$A_1 - \bar{A}_1 = -(A_2 - \bar{A}_2). \quad (5)$$

Multiplying (5) on the left and on the right by K , respectively, and according to (1) and (2), we obtain

$$A_1 - \bar{A}_1 = A_2 - \bar{A}_2. \quad (6)$$

Combining (5) and (6) gives $A_1 = \bar{A}_1$, $A_2 = \bar{A}_2$.

- (c) For any $A_1 \in KHC^{n \times n}$, $A_2 \in SKHC^{n \times n}$, we have

$$\begin{aligned} \langle A_1, A_2 \rangle &= \text{re}(\text{tr}(A_2^H A_1)) = \text{re}(\text{tr}(KA_2^H KKA_1 K)) \\ &= \text{re}(\text{tr}(-A_2^H A_1)) = -\langle A_1, A_2 \rangle. \end{aligned} \quad (7)$$

This implies $\langle A_1, A_2 \rangle = 0$. Combining (a), (b), and (c) gives (3).

(4) Let $\tilde{A} = KA$, if $A \in KHC^{n \times n}$, then $\tilde{A}^H = \tilde{A} \in HC^{n \times n}$. If $\tilde{A}^H = \tilde{A} \in HC^{n \times n}$, then $A = K\tilde{A}$ and $KA^H K = K\tilde{A}^H K = K\tilde{A} = A \in KHC^{n \times n}$.

(5) With the same method, we can prove (5). So, the proof is omitted. \square

In this paper, we suppose that K is a fixed permutation matrix and assume (λ_i, x_i) , $i = 1, \dots, h$, be right eigenpairs of A ; (μ_j, y_j) , $j = 1, \dots, l$, be left eigenpairs of A . If we let $X = (x_1, \dots, x_h) \in C^{n \times h}$, $\Lambda = \text{diag}(\lambda_1, \dots, \lambda_h) \in C^{h \times h}$; $Y = (y_1, \dots, y_l) \in C^{n \times l}$, $\Gamma = \text{diag}(\mu_1, \dots, \mu_l) \in C^{l \times l}$, then the problems studied in this paper can be described as follows.

Problem 2. Giving $X \in C^{n \times h}$, $\Lambda = \text{diag}(\lambda_1, \dots, \lambda_h) \in C^{h \times h}$; $Y \in C^{n \times l}$, $\Gamma = \text{diag}(\mu_1, \dots, \mu_l) \in C^{l \times l}$, find $A \in KHC^{n \times n}$ such that

$$\begin{aligned} AX &= X\Lambda, \\ Y^T A &= \Gamma Y^T. \end{aligned} \quad (8)$$

Problem 3. Giving $B \in C^{n \times n}$, find $\hat{A} \in S_E$ such that

$$\|B - \hat{A}\| = \min_{\hat{A} \in S_E} \|B - \hat{A}\|, \quad (9)$$

where S_E is the solution set of Problem 2.

This paper is organized as follows. In Section 2, we first obtain the equivalent problem with the properties of $KHC^{n \times n}$ and then derive the solvability conditions of Problem 2 and its general solution's expression. In Section 3, we first attest the existence and uniqueness theorem of Problem 3 then present the unique approximation solution. Finally, we provide a calculation procedure to compute the unique approximation solution and numerical experiment to illustrate the results obtained in this paper correction.

2. Solvability Conditions of Problem 2

We first discuss the properties of $KHC^{n \times n}$

Lemma 4. Denoting $M = KEKGE$, and $E \in HC^{n \times n}$, one has the following conclusions.

- (1) If $G \in KHC^{n \times n}$, then $M \in KHC^{n \times n}$.
- (2) If $G \in SKHC^{n \times n}$, then $M \in SKHC^{n \times n}$.
- (3) If $G = G_1 + G_2$, where $G_1 \in KHC^{n \times n}$, $G_2 \in SKHC^{n \times n}$, then $M \in KHC^{n \times n}$ if and only if $KEKG_2E = 0$. In addition, one has $M = KEKG_1E$.

Proof. (1) $KM^H K = KEG^H KEK = KE(KGK)KE = KEKGE = M$.

Hence, we have $M \in KHC^{n \times n}$.

(2) $KM^H K = KEG^H KEK = KE(-KGK)KE = -KEKGE = -M$.

Hence, we have $M \in SKHC^{n \times n}$.

(3) $M = KEK(G_1 + G_2)E = KEKG_1E + KEKG_2E$, we have $KEKG_1E \in KHC^{n \times n}$, $KEKG_2E \in SKHC^{n \times n}$ from (1) and (2). If $M \in KHC^{n \times n}$, then $M - KEKG_1E \in KHC^{n \times n}$, while $M - KEKG_1E = KEKG_2E \in SKHC^{n \times n}$. Therefore from the conclusion (3) of Definition 1, we have $KEKG_2E = 0$, that is, $M = KEKG_1E$. On the contrary, if $KEKG_2E = 0$, it is clear that $M = KEKG_1E \in KHC^{n \times n}$. The proof is completed. \square

Lemma 5. Let $A \in KHC^{n \times n}$, if (λ, x) is a right eigenpair of A , then $(\bar{\lambda}, K\bar{x})$ is a left eigenpair of A .

Proof. If (λ, x) is a right eigenpair of A , then we have

$$Ax = \lambda x. \quad (10)$$

From the conclusion (1) of Definition 1, it follows that

$$KA^H Kx = \lambda x. \quad (11)$$

This implies

$$(K\bar{x})^T A = \bar{\lambda}(K\bar{x})^T. \quad (12)$$

So $(\bar{\lambda}, K\bar{x})$ is a left eigenpair of A . \square

From Lemma 5, without loss of the generality, we may assume that Problem 2 is as follows.

$$\begin{aligned} X \in C^{n \times h}, \quad \Lambda = \text{diag}(\lambda_1, \dots, \lambda_h) \in C^{h \times h}, \\ Y = K\bar{X} \in C^{n \times h}, \quad \Gamma = \bar{\Lambda} \in C^{h \times h}. \end{aligned} \quad (13)$$

Combining (13) and the conclusion (4) of Definition 1, it is easy to derive the following lemma.

Lemma 6. *If X, Λ, Y, Γ are given by (13), then Problem 2 is equivalent to the following problem. If X, Λ, Y, Γ are given by (13), find $KA \in HC^{n \times n}$ such that*

$$KAX = KX\Lambda. \quad (14)$$

Lemma 7 (see [11]). *If giving $X \in C^{n \times h}, B \in C^{n \times h}$, then matrix equation $\bar{A}X = B$ has solution $\bar{A} \in HC^{n \times n}$ if and only if*

$$B = BX^+X, \quad B^H X = X^H B. \quad (15)$$

Moreover, the general solution \bar{A} can be expressed as

$$\begin{aligned} \bar{A} = BX^+ + (BX^+)^H (I_n - XX^+) \\ + (I_n - XX^+) \bar{G} (I_n - XX^+), \quad \forall \bar{G} \in HC^{n \times n}. \end{aligned} \quad (16)$$

Theorem 8. *If X, Λ, Y, Γ are given by (13), then Problem 2 has a solution in $KHC^{n \times n}$ if and only if*

$$X^H KX\Lambda = \bar{\Lambda} X^H KX, \quad X\Lambda = X\Lambda X^+ X. \quad (17)$$

Moreover, the general solution can be expressed as

$$A = A_0 + KEKG E, \quad \forall G \in KHC^{n \times n}, \quad (18)$$

where

$$A_0 = X\Lambda X^+ + K(X\Lambda X^+)^H KE, \quad E = I_n - XX^+. \quad (19)$$

Proof. Necessity: If there is a matrix $A \in KHC^{n \times n}$ such that $(AX = X\Lambda, Y^T A = \Gamma Y^T)$, then from Lemma 6, there exists a matrix $KA \in HC^{n \times n}$ such that $KAX = KX\Lambda$, and according to Lemma 7, we have

$$KX\Lambda = KX\Lambda X^+ X, \quad (KX\Lambda)^H X = X^H (KX\Lambda). \quad (20)$$

It is easy to see that (20) is equivalent to (17).

Sufficiency: If (17) holds, then (20) holds. Hence, matrix equation $KAX = KX\Lambda$ has solution $KA \in HC^{n \times n}$. Moreover, the general solution can be expressed as follows:

$$\begin{aligned} KA = KX\Lambda X^+ + (KX\Lambda X^+)^H (I_n - XX^+) \\ + (I_n - XX^+) \bar{G} (I_n - XX^+), \quad \forall \bar{G} \in HC^{n \times n}. \end{aligned} \quad (21)$$

Let

$$A_0 = X\Lambda X^+ + K(X\Lambda X^+)^H KE, \quad E = I_n - XX^+. \quad (22)$$

This implies $A = A_0 + KE\bar{G}E$. Combining the definition of K, E and the first equation of (17), we have

$$\begin{aligned} KA_0^H K &= K(X\Lambda X^+)^H K + X\Lambda X^+ - K(XX^+)^H K (X\Lambda X^+) \\ &= X\Lambda X^+ + K(X\Lambda X^+)^H K - K(X\Lambda X^+)^T KXX^+ \\ &= A_0. \end{aligned} \quad (23)$$

Hence, $A_0 \in KHC^{n \times n}$. Combining the definition of $K, E, (13)$ and (17), we have

$$\begin{aligned} A_0 X &= X\Lambda X^+ X + K(X\Lambda X^+)^H K (I_n - XX^+) X = X\Lambda, \\ Y^T A_0 &= X^H KX\Lambda X^+ + X^H KK(X\Lambda X^+)^H KE \\ &= \bar{\Lambda} X^H KXX^+ + (KX\Lambda X^+ X)^H (I_n - XX^+) \\ &= \bar{\Lambda} X^H KXX^+ + \bar{\Lambda} X^H K (I_n - XX^+) \\ &= \bar{\Lambda} X^H K = \Gamma Y^T. \end{aligned} \quad (24)$$

Therefore, A_0 is a special solution of Problem 2. Combining the conclusion (4) of Definition 1, Lemma 4, and $E = I_n - XX^+ \in HC^{n \times n}$, it is easy to prove that $A = A_0 + KEKG E \in KHC^{n \times n}$ if and only if $G \in KHC^{n \times n}$. Hence, the solution set of Problem 2 can be expressed as (18). \square

3. An Expression of the Solution of Problem 3

From (18), it is easy to prove that the solution set S_E of Problem 2 is a nonempty closed convex set if Problem 2 has a solution in $KHC^{n \times n}$. We claim that for any given $B \in R^{n \times n}$, there exists a unique optimal approximation for Problem 3.

Theorem 9. *Giving $B \in C^{n \times n}$, if the conditions of X, Y, Λ, Γ are the same as those in Theorem 8, then Problem 3 has a unique solution $\hat{A} \in S_E$. Moreover, \hat{A} can be expressed as*

$$\hat{A} = A_0 + KEKB_1 E, \quad (25)$$

where A_0, E are given by (19) and $B_1 = (1/2)(B + KB^H K)$.

Proof. Denoting $E_1 = I_n - E$, it is easy to prove that matrices E and E_1 are orthogonal projection matrices satisfying $EE_1 = 0$. It is clear that matrices KEK and $KE_1 K$ are also orthogonal

projection matrices satisfying $(KEK)(KE_1K) = 0$. According to the conclusion (3) of Definition 1, for any $B \in C^{n \times n}$, there exists unique

$$B_1 \in KHC^{n \times n}, \quad B_2 \in SKHC^{n \times n} \quad (26)$$

such that

$$B = B_1 + B_2, \quad \langle B_1, B_2 \rangle = 0, \quad (27)$$

where

$$B_1 = \frac{1}{2}(B + KB^HK), \quad B_2 = \frac{1}{2}(B - KB^HK). \quad (28)$$

Combining Theorem 8, for any $A \in S_E$, we have

$$\begin{aligned} \|B - A\|^2 &= \|B - A_0 - KEKGE\|^2 \\ &= \|B_1 + B_2 - A_0 - KEKGE\|^2 \\ &= \|B_1 - A_0 - KEKGE\|^2 + \|B_2\|^2 \\ &= \|(B_1 - A_0)(E + E_1) - KEKGE\|^2 + \|B_2\|^2 \\ &= \|(B_1 - A_0)E - KEKGE\|^2 \\ &\quad + \|(B_1 - A_0)E_1\|^2 + \|B_2\|^2 \\ &= \|K(E + E_1)K(B_1 - A_0)E - KEKGE\|^2 \end{aligned}$$

$$\begin{aligned} &+ \|(B_1 - A_0)E_1\|^2 + \|B_2\|^2 \\ &= \|KEK(B_1 - A_0)E - KEKGE\|^2 \\ &\quad + \|KE_1K(B_1 - A_0)E\|^2 \\ &\quad + \|(B_1 - A_0)E_1\|^2 + \|B_2\|^2. \end{aligned} \quad (29)$$

It is easy to prove that $KEKA_0E = 0$ according to the definitions of A_0, E . So we have

$$\begin{aligned} \|B - A\|^2 &= \|KEKB_1E - KEKGE\|^2 + \|KE_1K(B_1 - A_0)E\|^2 \\ &\quad + \|(B_1 - A_0)E_1\|^2 + \|B_2\|^2. \end{aligned} \quad (30)$$

Obviously, $\min_{A \in S_E} \|B - A\|$ is equivalent to

$$\min_{G \in KHC^{n \times n}} \|KEKB_1E - KEKGE\|. \quad (31)$$

Since $EE_1 = 0, (KEK)(KE_1K) = 0$, it is clear that $G = B_1 + KE_1K\widehat{G}E_1$, for any $\widehat{G} \in KHC^{n \times n}$, is a solution of (31). Substituting this result to (18), we can obtain (25). \square

Algorithm 10. (1) Input X, Λ, Y, Γ according to (13). (2) Compute $X^HKX\Lambda, \overline{\Lambda}X^HKX, X\Lambda X^+X, X\Lambda$, if (17) holds, then continue; otherwise stop. (3) Compute A_0 according to (19), and compute B_1 according to (28). (4) According to (25) calculate \widehat{A} .

Example 11 ($n = 8, h = l = 4$).

$$\begin{aligned} X &= \begin{pmatrix} 0.5661 & -0.2014 - 0.1422i & 0.1446 + 0.2138i & 0.524 \\ -0.2627 + 0.1875i & 0.5336 & -0.2110 - 0.4370i & -0.0897 + 0.3467i \\ -0.4132 + 0.2409i & 0.0226 - 0.0271i & -0.1095 + 0.2115i & -0.3531 - 0.0642i \\ -0.0306 + 0.2109i & -0.3887 - 0.0425i & 0.2531 + 0.2542i & 0.0094 + 0.2991i \\ 0.0842 - 0.1778i & -0.0004 - 0.3733i & 0.3228 - 0.1113i & 0.1669 + 0.1952i \\ 0.0139 - 0.3757i & -0.2363 + 0.3856i & 0.2583 + 0.0721i & 0.1841 - 0.2202i \\ 0.0460 + 0.3276i & -0.1114 + 0.0654i & -0.0521 - 0.2556i & -0.2351 + 0.3002i \\ 0.0085 - 0.1079i & 0.0974 + 0.3610i & 0.5060 & -0.2901 - 0.0268i \end{pmatrix}, \\ \Lambda &= \begin{pmatrix} -0.3967 - 0.4050i & 0 & 0 & 0 \\ 0 & -0.3967 + 0.4050i & 0 & 0 \\ 0 & 0 & 0.0001 & 0 \\ 0 & 0 & 0 & -0.0001i \end{pmatrix}, \\ K &= \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}, \\ Y &= K\overline{X}, \quad \Gamma = \overline{\Lambda} \end{aligned} \quad (32)$$

$B =$

From the first column to the fourth column

$$\begin{pmatrix} -0.5218 + 0.0406i & 0.2267 - 0.0560i & -0.1202 + 0.0820i & -0.0072 - 0.3362i \\ 0.3909 - 0.3288i & 0.2823 - 0.2064i & -0.0438 - 0.0403i & 0.2707 + 0.0547i \\ 0.2162 - 0.1144i & -0.4307 + 0.2474i & -0.0010 - 0.0412i & 0.2164 - 0.1314i \\ -0.1872 - 0.0599i & -0.0061 + 0.4698i & 0.3605 - 0.0247i & 0.4251 + 0.1869i \\ -0.1227 - 0.0194i & 0.2477 - 0.0606i & 0.3918 + 0.6340i & 0.1226 + 0.0636i \\ -0.0893 + 0.4335i & 0.0662 + 0.0199i & -0.0177 - 0.1412i & 0.4047 + 0.2288i \\ 0.1040 - 0.2015i & 0.1840 + 0.2276i & 0.2681 - 0.3526i & -0.5252 + 0.1022i \\ 0.1808 + 0.2669i & 0.2264 + 0.3860i & -0.1791 + 0.1976i & -0.0961 - 0.0117i \end{pmatrix} \quad (33)$$

From the fifth column to the eighth column

$$\begin{pmatrix} -0.2638 - 0.4952i & -0.0863 - 0.1664i & 0.2687 + 0.1958i & -0.2544 - 0.1099i \\ -0.2741 - 0.1656i & -0.0227 + 0.2684i & 0.1846 + 0.2456i & -0.0298 + 0.5163i \\ -0.1495 - 0.3205i & 0.1391 + 0.2434i & 0.1942 - 0.5211i & -0.3052 - 0.1468i \\ -0.2554 + 0.2690i & -0.4222 - 0.1080i & 0.2232 + 0.0774i & 0.0965 - 0.0421i \\ 0.3856 - 0.0619i & 0.1217 - 0.0270i & 0.1106 - 0.3090i & -0.1122 + 0.2379i \\ -0.1130 + 0.0766i & 0.7102 - 0.0901i & 0.1017 + 0.1397i & -0.0445 + 0.0038i \\ 0.1216 + 0.0076i & 0.2343 - 0.1772i & 0.5242 - 0.0089i & -0.0613 + 0.0258i \\ -0.0750 - 0.3581i & 0.0125 + 0.0964i & 0.0779 - 0.1074i & 0.6735 - 0.0266i \end{pmatrix} \quad (34)$$

It is easy to see that matrices X, Λ, Y, Γ satisfy (17). Hence, there exists the unique solution for Problem 3. Using the

software “MATLAB”, we obtain the unique solution \widehat{A} of Problem 3.

From the first column to the fourth column

$$\begin{pmatrix} -0.1983 + 0.0491i & 0.1648 + 0.0032i & 0.0002 + 0.1065i & 0.1308 + 0.2690i \\ 0.1071 - 0.2992i & 0.2106 + 0.2381i & -0.0533 - 0.3856i & -0.0946 + 0.0488i \\ 0.1935 - 0.1724i & -0.0855 - 0.0370i & 0.0200 - 0.0665i & -0.2155 - 0.0636i \\ 0.0085 - 0.2373i & -0.0843 - 0.1920i & 0.0136 + 0.0382i & -0.0328 - 0.0000i \\ -0.0529 + 0.1703i & 0.1948 - 0.0719i & 0.1266 + 0.1752i & 0.0232 - 0.2351i \\ 0.0855 + 0.1065i & 0.0325 - 0.2068i & 0.2624 + 0.0000i & 0.0136 - 0.0382i \\ 0.1283 - 0.1463i & -0.0467 + 0.0000i & 0.0325 + 0.2067i & -0.0843 + 0.1920i \\ 0.2498 + 0.0000i & 0.1283 + 0.1463i & 0.0855 - 0.1065i & 0.0086 + 0.2373i \end{pmatrix} \quad (35)$$

From the fifth column to the eighth column

$$\begin{pmatrix} 0.2399 - 0.1019i & 0.1928 - 0.1488i & -0.3480 - 0.2574i & 0.1017 - 0.0000i \\ -0.1955 + 0.0644i & 0.2925 + 0.1872i & 0.3869 + 0.0000i & -0.3481 + 0.2574i \\ 0.0074 + 0.0339i & -0.3132 - 0.0000i & 0.2926 - 0.1872i & 0.1928 + 0.1488i \\ 0.0232 + 0.2351i & -0.2154 + 0.0636i & -0.0946 - 0.0489i & 0.1309 - 0.2691i \\ -0.0545 - 0.0000i & 0.0074 - 0.0339i & -0.1955 - 0.0643i & 0.2399 + 0.1019i \\ 0.1266 - 0.1752i & 0.0200 + 0.0665i & -0.0533 + 0.3857i & 0.0002 - 0.1065i \\ 0.1949 + 0.0719i & -0.0855 + 0.0370i & 0.2106 - 0.2381i & 0.1648 - 0.0032i \\ -0.0529 - 0.1703i & 0.1935 + 0.1724i & 0.1071 + 0.2992i & -0.1983 - 0.0491i \end{pmatrix} \quad (36)$$

Conflict of Interests

There is no conflict of interests between the authors.

Acknowledgments

This research was supported by National Natural Science Foundation of China (31170532). The authors are very grateful to the referees for their valuable comments and also thank the editor for his helpful suggestions.

References

- [1] J. H. Wilkinson, *The Algebraic Eigenvalue Problem*, Oxford University Press, Oxford, UK, 1965.
- [2] A. Berman and R. J. Plemmons, *Nonnegative Matrices in the Mathematical Sciences*, vol. 9 of *Classics in Applied Mathematics*, SIAM, Philadelphia, Pa, USA, 1994.
- [3] M. Arav, D. Hershkowitz, V. Mehrmann, and H. Schneider, "The recursive inverse eigenvalue problem," *SIAM Journal on Matrix Analysis and Applications*, vol. 22, no. 2, pp. 392–412, 2000.
- [4] R. Loewy and V. Mehrmann, "A note on the symmetric recursive inverse eigenvalue problem," *SIAM Journal on Matrix Analysis and Applications*, vol. 25, no. 1, pp. 180–187, 2003.
- [5] F. L. Li, X. Y. Hu, and L. Zhang, "Left and right inverse eigenpairs problem of skew-centrosymmetric matrices," *Applied Mathematics and Computation*, vol. 177, no. 1, pp. 105–110, 2006.
- [6] F. L. Li, X. Y. Hu, and L. Zhang, "Left and right inverse eigenpairs problem of generalized centrosymmetric matrices and its optimal approximation problem," *Applied Mathematics and Computation*, vol. 212, no. 2, pp. 481–487, 2009.
- [7] R. D. Hill and S. R. Waters, "On κ -real and κ -Hermitian matrices," *Linear Algebra and its Applications*, vol. 169, pp. 17–29, 1992.
- [8] S. P. Irwin, "Matrices with multiple symmetry properties: applications of centro-Hermitian and per-Hermitian matrices," *Linear Algebra and its Applications*, vol. 284, no. 1–3, pp. 239–258, 1998.
- [9] L. C. Biedenharn and J. D. Louck, *Angular Momentum in Quantum Physics*, vol. 8 of *Encyclopedia of Mathematics and its Applications*, Addison-Wesley, Reading, Mass, USA, 1981.
- [10] W. M. Gibson and B. R. Pollard, *Symmetry Principles in Elementary Particle Physics*, Cambridge University Press, Cambridge, UK, 1976.
- [11] W. F. Trench, "Hermitian, Hermitian R -symmetric, and Hermitian R -skew symmetric Procrustes problems," *Linear Algebra and its Applications*, vol. 387, pp. 83–98, 2004.



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

