

## Research Article

# Moment Inequality for $\varphi$ -Mixing Sequences and Its Applications

Wang Xuejun, Hu Shuhe, Shen Yan, and Yang Wenzhi

School of Mathematical Science, Anhui University, Hefei, Anhui 230039, China

Correspondence should be addressed to Hu Shuhe, hushuhe@263.net

Received 11 April 2009; Accepted 21 September 2009

Recommended by Sever Silvestru Dragomir

Firstly, the maximal inequality for  $\varphi$ -mixing sequences is given. By using the maximal inequality, we study the convergence properties for  $\varphi$ -mixing sequences. The Hájek-Rényi-type inequality, strong law of large numbers, strong growth rate and the integrability of supremum for  $\varphi$ -mixing sequences are obtained.

Copyright © 2009 Wang Xuejun et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

## 1. Introduction

Let  $\{X_n, n \geq 1\}$  be a random variable sequence defined on a fixed probability space  $(\Omega, \mathcal{F}, P)$  and  $S_n = \sum_{i=1}^n X_i$  for each  $n \geq 1$ . Let  $n$  and  $m$  be positive integers. Write  $\mathcal{F}_n^m = \sigma(X_i, n \leq i \leq m)$ . Given  $\sigma$ -algebras  $\mathcal{B}, \mathcal{R}$  in  $\mathcal{F}$ , let

$$\varphi(\mathcal{B}, \mathcal{R}) = \sup_{A \in \mathcal{B}, B \in \mathcal{R}, P(A) > 0} |P(B | A) - P(B)|. \quad (1.1)$$

Define the  $\varphi$ -mixing coefficients by

$$\varphi(n) = \sup_{k \geq 1} \varphi(\mathcal{F}_1^k, \mathcal{F}_{k+n}^\infty), \quad n \geq 0. \quad (1.2)$$

*Definition 1.1.* A random variable sequence  $\{X_n, n \geq 1\}$  is said to be a  $\varphi$ -mixing random variable sequence if  $\varphi(n) \downarrow 0$  as  $n \rightarrow \infty$ .

The concept of  $\varphi$ -mixing random variables was introduced by Dobrushin [1] and many applications have been found. See, for example, Dobrushin [1], Utev [2], and Chen [3]

for central limit theorem, Herrndorf [4] and Peligrad [5] for weak invariance principle, Sen [6, 7] for weak convergence of empirical processes, Iosifescu [8] for limit theorem, Peligrad [9] for Ibragimov-Iosifescu conjecture, Shao [10] for almost sure invariance principles, Hu and Wang [11] for large deviations, and so forth. When these are compared with the corresponding results of independent random variable sequences, there still remains much to be desired. The main purpose of this paper is to study the maximal inequality for  $\varphi$ -mixing sequences, by which we can get the Hájek-Rényi-type inequality, strong law of large numbers, strong growth rate, and the integrability of supremum for  $\varphi$ -mixing sequences.

Throughout the paper,  $C$  denotes a positive constant which may be different in various places. The main results of this paper depend on the following lemmas.

**Lemma 1.2** (see Lu and Lin [12]). *Let  $\{X_n, n \geq 1\}$  be a sequence of  $\varphi$ -mixing random variables. Let  $X \in L_p(\mathcal{F}_1^k)$ ,  $Y \in L_q(\mathcal{F}_{k+n}^\infty)$ ,  $p \geq 1$ ,  $q \geq 1$ , and  $1/p + 1/q = 1$ . Then*

$$|EXY - EXEY| \leq 2(\varphi(n))^{1/p} (E|X|^p)^{1/p} (E|Y|^q)^{1/q}. \quad (1.3)$$

**Lemma 1.3** (see Shao [10]). *Let  $\{X_n, n \geq 1\}$  be a  $\varphi$ -mixing sequence. Put  $T_a(n) = \sum_{i=a+1}^{a+n} X_i$ . Suppose that there exists an array  $\{C_{a,n}\}$  of positive numbers such that*

$$ET_a^2(n) \leq C_{a,n} \quad \text{for every } a \geq 0, n \geq 1. \quad (1.4)$$

*Then for every  $q \geq 2$ , there exists a constant  $C$  depending only on  $q$  and  $\varphi(\cdot)$  such that*

$$E\left(\max_{1 \leq j \leq n} |T_a(j)|^q\right) \leq C \left[ C_{a,n}^{q/2} + E\left(\max_{a+1 \leq i \leq a+n} |X_i|^q\right) \right] \quad (1.5)$$

*for every  $a \geq 0$  and  $n \geq 1$ .*

**Lemma 1.4** (see Hu et al. [13]). *Let  $b_1, b_2, \dots$  be a nondecreasing unbounded sequence of positive numbers and let  $\alpha_1, \alpha_2, \dots$  be nonnegative numbers. Let  $r$  and  $C$  be fixed positive numbers. Assume that for each  $n \geq 1$ ,*

$$E\left(\max_{1 \leq l \leq n} |S_l|\right)^r \leq C \sum_{i=1}^n \alpha_i, \quad (1.6)$$

$$\sum_{l=1}^{\infty} \frac{\alpha_l}{b_l^r} < \infty, \quad (1.7)$$

*then*

$$\lim_{n \rightarrow \infty} \frac{S_n}{b_n} = 0 \quad \text{a.s.}, \quad (1.8)$$

*and with the growth rate*

$$\frac{S_n}{b_n} = O\left(\frac{\beta_n}{b_n}\right) \quad \text{a.s.}, \quad (1.9)$$

where

$$\beta_n = \max_{1 \leq k \leq n} b_k v_k^{\delta/r}, \quad \forall 0 < \delta < 1, \quad v_n = \sum_{k=n}^{\infty} \frac{\alpha_k}{b_k^r}, \quad \lim_{n \rightarrow \infty} \frac{\beta_n}{b_n} = 0,$$

$$E\left(\max_{1 \leq l \leq n} \left|\frac{S_l}{b_l}\right|^r\right) \leq 4C \sum_{l=1}^n \frac{\alpha_l}{b_l^r} < \infty, \tag{1.10}$$

$$E\left(\sup_{l \geq 1} \left|\frac{S_l}{b_l}\right|^r\right) \leq 4C \sum_{l=1}^{\infty} \frac{\alpha_l}{b_l^r} < \infty.$$

If further we assume that  $\alpha_n > 0$  for infinitely many  $n$ , then

$$E\left(\sup_{l \geq 1} \left|\frac{S_l}{\beta_l}\right|^r\right) \leq 4C \sum_{l=1}^{\infty} \frac{\alpha_l}{\beta_l^r} < \infty. \tag{1.11}$$

**Lemma 1.5** (see Fazekas and Klesov [14] and Hu [15]). *Let  $b_1, b_2, \dots$  be a nondecreasing unbounded sequence of positive numbers and let  $\alpha_1, \alpha_2, \dots$  be nonnegative numbers. Denote  $\Lambda_k = \alpha_1 + \alpha_2 + \dots + \alpha_k$  for  $k \geq 1$ . Let  $r$  be a fixed positive number satisfying (1.6). If*

$$\sum_{l=1}^{\infty} \Lambda_l \left(\frac{1}{b_l^r} - \frac{1}{b_{l+1}^r}\right) < \infty, \tag{1.12}$$

$$\frac{\Lambda_n}{b_n^r} \text{ is bounded}, \tag{1.13}$$

then (1.8)–(1.11) hold.

## 2. Maximal Inequality for $\varphi$ -Mixing Sequences

**Theorem 2.1.** *Let  $\{X_n, n \geq 1\}$  be a sequence of  $\varphi$ -mixing random variables satisfying  $\sum_{n=1}^{\infty} \varphi^{1/2}(n) < \infty$ . Assume that  $EX_n = 0$  and  $EX_n^2 < \infty$  for each  $n \geq 1$ . Then there exists a constant  $C$  depending only on  $\varphi(\cdot)$  such that for any  $n \geq 1$  and  $a \geq 0$*

$$E\left(\sum_{i=a+1}^{a+n} X_i\right)^2 \leq C \sum_{i=a+1}^{a+n} EX_i^2, \tag{2.1}$$

$$E\left(\max_{1 \leq j \leq n} \left|\sum_{i=a+1}^{a+j} X_i\right|^2\right) \leq C \sum_{i=a+1}^{a+n} EX_i^2. \tag{2.2}$$

In particular,

$$E\left(\sum_{i=1}^n X_i\right)^2 \leq C \sum_{i=1}^n EX_i^2, \quad (2.3)$$

$$E\left(\max_{1 \leq j \leq n} \left| \sum_{i=1}^j X_i \right|^2\right) \leq C \sum_{i=1}^n EX_i^2, \quad (2.4)$$

where  $C$  may be different in various places.

*Proof.* By Lemma 1.2 for  $p = q = 2$ , we can see that

$$\begin{aligned} E\left(\sum_{i=a+1}^{a+n} X_i\right)^2 &= \sum_{i=a+1}^{a+n} EX_i^2 + 2 \sum_{a+1 \leq i < j \leq a+n} E(X_i X_j) \\ &\leq \sum_{i=a+1}^{a+n} EX_i^2 + 4 \sum_{a+1 \leq i < j \leq a+n} \varphi^{1/2}(j-i) (EX_i^2)^{1/2} (EX_j^2)^{1/2} \\ &\leq \sum_{i=a+1}^{a+n} EX_i^2 + 2 \sum_{k=1}^{n-1} \sum_{i=a+1}^{a+n-k} \varphi^{1/2}(k) (EX_i^2 + EX_{k+i}^2) \\ &\leq \left(1 + 4 \sum_{k=1}^{\infty} \varphi^{1/2}(k)\right) \sum_{i=a+1}^{a+n} EX_i^2 \\ &\doteq C \sum_{i=a+1}^{a+n} EX_i^2, \end{aligned} \quad (2.5)$$

which implies (2.1). By (2.1) and Lemma 1.3 (take  $q = 2$ ), we can get

$$\begin{aligned} E\left(\max_{1 \leq j \leq n} \left| \sum_{i=a+1}^{a+j} X_i \right|^2\right) &\leq C \left[ \sum_{i=a+1}^{a+n} EX_i^2 + E\left(\max_{a+1 \leq i \leq a+n} X_i^2\right) \right] \\ &\leq C \sum_{i=a+1}^{a+n} EX_i^2. \end{aligned} \quad (2.6)$$

The proof is completed.  $\square$

### 3. Hájek-Rényi-Type Inequality for $\varphi$ -Mixing Sequences

In this section, we will give the Hájek-Rényi-type inequality for  $\varphi$ -mixing sequences, which can be applied to obtain the strong law of large numbers and the integrability of supremum.

**Theorem 3.1.** *Let  $\{X_n, n \geq 1\}$  be a sequence of  $\varphi$ -mixing random variables satisfying  $\sum_{n=1}^{\infty} \varphi^{1/2}(n) < \infty$  and let  $\{b_n, n \geq 1\}$  be a nondecreasing sequence of positive numbers. Then for*

any  $\varepsilon > 0$  and any integer  $n \geq 1$ ,

$$P \left\{ \max_{1 \leq k \leq n} \left| \frac{1}{b_k} \sum_{j=1}^k (X_j - EX_j) \right| \geq \varepsilon \right\} \leq \frac{4C}{\varepsilon^2} \sum_{j=1}^n \frac{\text{Var}(X_j)}{b_j^2}, \tag{3.1}$$

where  $C$  is defined in (2.4) in Theorem 2.1.

*Proof.* Without loss of generality, we assume that  $b_n \geq 1$  for all  $n \geq 1$ . Let  $\alpha = \sqrt{2}$ . For  $i \geq 0$ , define

$$A_i = \{1 \leq k \leq n : \alpha^i \leq b_k < \alpha^{i+1}\}. \tag{3.2}$$

For  $A_i \neq \emptyset$ , we let  $v(i) = \max\{k : k \in A_i\}$  and  $t_n$  be the index of the last nonempty set  $A_i$ . Obviously,  $A_i A_j = \emptyset$  if  $i \neq j$  and  $\sum_{i=0}^{t_n} A_i = \{1, 2, \dots, n\}$ . It is easily seen that  $\alpha^i \leq b_k \leq b_{v(i)} < \alpha^{i+1}$  if  $k \in A_i$  and  $\{X_n - EX_n, n \geq 1\}$  is also a sequence of  $\varphi$ -mixing random variables. By Markov's inequality and (2.4), we have

$$\begin{aligned} & P \left\{ \max_{1 \leq k \leq n} \left| \frac{1}{b_k} \sum_{j=1}^k (X_j - EX_j) \right| \geq \varepsilon \right\} \\ &= P \left\{ \max_{0 \leq i \leq t_n, A_i \neq \emptyset} \max_{k \in A_i} \left| \frac{1}{b_k} \sum_{j=1}^k (X_j - EX_j) \right| \geq \varepsilon \right\} \\ &\leq \sum_{i=0, A_i \neq \emptyset}^{t_n} P \left\{ \frac{1}{\alpha^i} \max_{1 \leq k \leq v(i)} \left| \sum_{j=1}^k (X_j - EX_j) \right| \geq \varepsilon \right\} \\ &\leq \frac{1}{\varepsilon^2} \sum_{i=0, A_i \neq \emptyset}^{t_n} \frac{1}{\alpha^{2i}} E \left\{ \max_{1 \leq k \leq v(i)} \left| \sum_{j=1}^k (X_j - EX_j) \right|^2 \right\} \\ &\leq \frac{C}{\varepsilon^2} \sum_{i=0, A_i \neq \emptyset}^{t_n} \frac{1}{\alpha^{2i}} \sum_{j=1}^{v(i)} \text{Var}(X_j) \\ &\leq \frac{C}{\varepsilon^2} \sum_{j=1}^n \text{Var}(X_j) \sum_{i=0, A_i \neq \emptyset, v(i) \geq j}^{t_n} \frac{1}{\alpha^{2i}}. \end{aligned} \tag{3.3}$$

Now we estimate  $\sum_{i=0, A_i \neq \emptyset, v(i) \geq j}^{t_n} (1/\alpha^{2i})$ . Let  $i_0 = \min\{i : A_i \neq \emptyset, v(i) \geq j\}$ . Then  $b_j \leq b_{v(i_0)} < \alpha^{i_0+1}$  follows from the definition of  $v(i)$ . Therefore,

$$\sum_{i=0, A_i \neq \emptyset, v(i) \geq j}^{t_n} \frac{1}{\alpha^{2i}} < \sum_{i=i_0}^{\infty} \frac{1}{\alpha^{2i}} = \frac{1}{1 - 1/\alpha^2} \frac{1}{\alpha^{2i_0}} < \frac{\alpha^2}{1 - 1/\alpha^2} \frac{1}{b_j^2} = \frac{4}{b_j^2}. \tag{3.4}$$

Thus, (3.1) follows from (3.3) and (3.4) immediately. □

**Theorem 3.2.** Let  $\{X_n, n \geq 1\}$  be a sequence of  $\varphi$ -mixing random variables satisfying  $\sum_{n=1}^{\infty} \varphi^{1/2}(n) < \infty$  and let  $\{b_n, n \geq 1\}$  be a nondecreasing sequence of positive numbers. Then for any  $\varepsilon > 0$  and any positive integers  $m < n$ ,

$$P\left(\max_{m \leq k \leq n} \left| \frac{1}{b_k} \sum_{j=1}^k (X_j - EX_j) \right| \geq \varepsilon\right) \leq \frac{4C}{\varepsilon^2} \left( \sum_{j=1}^m \frac{\text{Var}(X_j)}{b_m^2} + 4 \sum_{j=m+1}^n \frac{\text{Var}(X_j)}{b_j^2} \right), \quad (3.5)$$

where  $C$  is defined in (3.1).

*Proof.* Observe that

$$\max_{m \leq k \leq n} \left| \frac{1}{b_k} \sum_{j=1}^k (X_j - EX_j) \right| \leq \left| \frac{1}{b_m} \sum_{j=1}^m (X_j - EX_j) \right| + \max_{m+1 \leq k \leq n} \left| \frac{1}{b_k} \sum_{j=m+1}^k (X_j - EX_j) \right|, \quad (3.6)$$

thus

$$\begin{aligned} & P\left(\max_{m \leq k \leq n} \left| \frac{1}{b_k} \sum_{j=1}^k (X_j - EX_j) \right| \geq \varepsilon\right) \\ & \leq P\left(\left| \frac{1}{b_m} \sum_{j=1}^m (X_j - EX_j) \right| \geq \frac{\varepsilon}{2}\right) + P\left(\max_{m+1 \leq k \leq n} \left| \frac{1}{b_k} \sum_{j=m+1}^k (X_j - EX_j) \right| \geq \frac{\varepsilon}{2}\right) \doteq I + II. \end{aligned} \quad (3.7)$$

For  $I$ , by Markov's inequality and (2.3), we have

$$I \leq \frac{4}{\varepsilon^2 b_m^2} E \left( \sum_{j=1}^m (X_j - EX_j) \right)^2 \leq \frac{4C}{\varepsilon^2} \sum_{j=1}^m \frac{\text{Var}(X_j)}{b_m^2}. \quad (3.8)$$

For  $II$ , we will apply Theorem 3.1 to  $\{X_{m+i}, 1 \leq i \leq n-m\}$  and  $\{b_{m+i}, 1 \leq i \leq n-m\}$ . Noting that

$$\max_{m+1 \leq k \leq n} \left| \frac{1}{b_k} \sum_{j=m+1}^k (X_j - EX_j) \right| = \max_{1 \leq k \leq n-m} \left| \frac{1}{b_{m+k}} \sum_{j=1}^k (X_{m+j} - EX_{m+j}) \right|, \quad (3.9)$$

thus, by Theorem 3.1, we get

$$II \leq \frac{4C}{(\varepsilon/2)^2} \sum_{j=1}^{n-m} \frac{\text{Var}(X_{m+j})}{b_{m+j}^2} = \frac{16C}{\varepsilon^2} \sum_{j=m+1}^n \frac{\text{Var}(X_j)}{b_j^2}. \quad (3.10)$$

Therefore, the desired result (3.5) follows from (3.7)–(3.10) immediately.  $\square$

**Theorem 3.3.** Let  $\{X_n, n \geq 1\}$  be a sequence of  $\varphi$ -mixing random variables satisfying  $\sum_{n=1}^{\infty} \varphi^{1/2}(n) < \infty$  and let  $\{b_n, n \geq 1\}$  be a nondecreasing sequence of positive numbers. Denote  $T_n = \sum_{i=1}^n (X_i - EX_i)$  for  $n \geq 1$ . Assume that

$$\sum_{j=1}^{\infty} \frac{\text{Var}(X_j)}{b_j^2} < \infty, \quad (3.11)$$

then for any  $r \in (0, 2)$ ,

$$E\left(\sup_{n \geq 1} \left|\frac{T_n}{b_n}\right|^r\right) \leq 1 + \frac{4Cr}{2-r} \sum_{j=1}^{\infty} \frac{\text{Var}(X_j)}{b_j^2} < \infty, \quad (3.12)$$

where  $C$  is defined in (3.1). Furthermore, if  $\lim_{n \rightarrow \infty} b_n = +\infty$ , then

$$\lim_{n \rightarrow \infty} \frac{1}{b_n} \sum_{j=1}^n (X_j - EX_j) = 0 \quad \text{a.s.} \quad (3.13)$$

*Proof.* By the continuity of probability and Theorem 3.1, we get

$$\begin{aligned} E\left(\sup_{n \geq 1} \left|\frac{T_n}{b_n}\right|^r\right) &= \int_0^{\infty} P\left(\sup_{n \geq 1} \left|\frac{T_n}{b_n}\right|^r > t\right) dt \\ &= \int_0^1 P\left(\sup_{n \geq 1} \left|\frac{T_n}{b_n}\right|^r > t\right) dt + \int_1^{\infty} P\left(\sup_{n \geq 1} \left|\frac{T_n}{b_n}\right|^r > t\right) dt \\ &\leq 1 + \int_1^{\infty} P\left(\sup_{n \geq 1} \left|\frac{T_n}{b_n}\right| > t^{1/r}\right) dt \\ &\leq 1 + \int_1^{\infty} \lim_{N \rightarrow \infty} P\left(\max_{1 \leq n \leq N} \left|\frac{T_n}{b_n}\right| > t^{1/r}\right) dt \\ &\leq 1 + 4C \sum_{j=1}^{\infty} \frac{\text{Var}(X_j)}{b_j^2} \int_1^{\infty} t^{-2/r} dt \\ &= 1 + \frac{4Cr}{2-r} \sum_{j=1}^{\infty} \frac{\text{Var}(X_j)}{b_j^2} < \infty. \end{aligned} \quad (3.14)$$

Observe that

$$P\left(\bigcup_{n=m}^{\infty} \left(\left|\frac{T_n}{b_n}\right| > \varepsilon\right)\right) = P\left(\bigcup_{N=m}^{\infty} \left(\max_{m \leq n \leq N} \left|\frac{T_n}{b_n}\right| > \varepsilon\right)\right) = \lim_{N \rightarrow \infty} P\left(\max_{m \leq n \leq N} \left|\frac{T_n}{b_n}\right| > \varepsilon\right). \quad (3.15)$$

By Theorem 3.2, we have that

$$P\left(\max_{m \leq n \leq N} \left| \frac{T_n}{b_n} \right| > \varepsilon\right) \leq \frac{4C}{\varepsilon^2} \left( \sum_{j=1}^m \frac{\text{Var}(X_j)}{b_m^2} + 4 \sum_{j=m+1}^N \frac{\text{Var}(X_j)}{b_j^2} \right). \quad (3.16)$$

Hence, by (3.11) and Kronecker's Lemma, it follows that

$$\lim_{m \rightarrow \infty} P\left(\bigcup_{n=m}^{\infty} \left( \left| \frac{T_n}{b_n} \right| > \varepsilon \right)\right) = 0, \quad \forall \varepsilon > 0, \quad (3.17)$$

which is equivalent to

$$\lim_{n \rightarrow \infty} \frac{1}{b_n} \sum_{j=1}^n (X_j - EX_j) = 0 \quad \text{a.s.} \quad (3.18)$$

So the desired results are proved.  $\square$

#### 4. Strong Law of Large Numbers and Growth Rate for $\varphi$ -Mixing Sequences

**Theorem 4.1.** *Let  $\{X_n, n \geq 1\}$  be a sequence of mean zero  $\varphi$ -mixing random variables satisfying  $\sum_{n=1}^{\infty} \varphi^{1/2}(n) < \infty$  and let  $\{b_n, n \geq 1\}$  be a nondecreasing unbounded sequence of positive numbers. Assume that*

$$\sum_{n=1}^{\infty} \frac{EX_n^2}{b_n^2} < \infty, \quad (4.1)$$

then

$$\lim_{n \rightarrow \infty} \frac{S_n}{b_n} = 0 \quad \text{a.s.}, \quad (4.2)$$

and with the growth rate

$$\frac{S_n}{b_n} = O\left(\frac{\beta_n}{b_n}\right) \quad \text{a.s.}, \quad (4.3)$$



where

$$\beta_n = \max_{1 \leq k \leq n} b_k v_k^{\delta/2}, \quad \forall 0 < \delta < 1, \quad v_n = \sum_{k=n}^{\infty} \frac{\alpha_k}{b_k^2}, \quad \lim_{n \rightarrow \infty} \frac{\beta_n}{b_n} = 0, \tag{4.4}$$

$\alpha_k = CEX_k^2, \quad k \geq 1$ , where  $C$  is defined in (2.4),

$$E \left( \max_{1 \leq l \leq n} \left| \frac{S_l}{b_l} \right|^2 \right) \leq 4 \sum_{l=1}^n \frac{\alpha_l}{b_l^2} < \infty, \tag{4.5}$$

$$E \left( \sup_{l \geq 1} \left| \frac{S_l}{b_l} \right|^2 \right) \leq 4 \sum_{l=1}^{\infty} \frac{\alpha_l}{b_l^2} < \infty.$$

If further we assume that  $\alpha_n > 0$  for infinitely many  $n$ , then

$$E \left( \sup_{l \geq 1} \left| \frac{S_l}{\beta_l} \right|^2 \right) \leq 4 \sum_{l=1}^{\infty} \frac{\alpha_l}{\beta_l^2} < \infty. \tag{4.6}$$

*Proof.* By (2.4) in Theorem 2.1, we have

$$E \left( \max_{1 \leq k \leq n} |S_k|^2 \right) \leq C \sum_{i=1}^n EX_i^2 = \sum_{k=1}^n \alpha_k. \tag{4.7}$$

It follows by (4.1) that

$$\sum_{n=1}^{\infty} \frac{\alpha_n}{b_n^2} = C \sum_{n=1}^{\infty} \frac{EX_n^2}{b_n^2} < \infty. \tag{4.8}$$

Thus, (4.2)–(4.6) follow from (4.7), (4.8), and Lemma 1.4 immediately. We complete the proof of the theorem.  $\square$

**Theorem 4.2.** Let  $\{X_n, n \geq 1\}$  be a sequence of  $\varphi$ -mixing random variables with  $\sum_{n=1}^{\infty} \varphi^{1/2}(n) < \infty$ .  $1 \leq p < 2$ . Denote  $Q_n = \max_{1 \leq k \leq n} EX_k^2$  for  $n \geq 1$  and  $Q_0 = 0$ . Assume that

$$\sum_{n=1}^{\infty} \frac{Q_n}{n^{2/p}} < \infty, \tag{4.9}$$

then

$$\lim_{n \rightarrow \infty} \frac{1}{n^{1/p}} \sum_{i=1}^n (X_i - EX_i) = 0 \quad a.s., \tag{4.10}$$

and with the growth rate

$$\frac{1}{n^{1/p}} \sum_{i=1}^n (X_i - EX_i) = O\left(\frac{\beta_n}{n^{1/p}}\right) \quad a.s., \quad (4.11)$$

where

$$\beta_n = \max_{1 \leq k \leq n} k^{1/p} v_k^{\delta/2}, \quad \forall 0 < \delta < 1, \quad v_n = \sum_{k=n}^{\infty} \frac{\alpha_k}{k^{2/p}}, \quad \lim_{n \rightarrow \infty} \frac{\beta_n}{n^{1/p}} = 0, \quad (4.12)$$

$$\alpha_k = C(kQ_k - (k-1)Q_{k-1}), \quad k \geq 1, \quad \text{where } C \text{ is defined in (2.4),}$$

$$E\left(\max_{1 \leq l \leq n} \left|\frac{S_l}{l^{1/p}}\right|^2\right) \leq 4 \sum_{l=1}^n \frac{\alpha_l}{l^{2/p}} < \infty, \quad (4.13)$$

$$E\left(\sup_{l \geq 1} \left|\frac{S_l}{l^{1/p}}\right|^2\right) \leq 4 \sum_{l=1}^{\infty} \frac{\alpha_l}{l^{2/p}} < \infty. \quad (4.14)$$

If further we assume that  $\alpha_n > 0$  for infinitely many  $n$ , then

$$E\left(\sup_{l \geq 1} \left|\frac{S_l}{\beta_l}\right|^2\right) \leq 4 \sum_{l=1}^{\infty} \frac{\alpha_l}{\beta_l^2} < \infty. \quad (4.15)$$

In addition, for any  $r \in (0, 2)$ ,

$$E\left(\sup_{l \geq 1} \left|\frac{S_l}{l^{1/p}}\right|^r\right) \leq 1 + \frac{4r}{2-r} \sum_{l=1}^{\infty} \frac{\alpha_l}{l^{2/p}} < \infty. \quad (4.16)$$

*Proof.* Assume that  $EX_n = 0$ ,  $b_n = n^{1/p}$ , and  $\Lambda_n = \sum_{l=1}^n \alpha_l$ ,  $n \geq 1$ . By (2.4) in Theorem 2.1, we can see that

$$E\left(\max_{1 \leq k \leq n} \left|\sum_{i=1}^k X_i\right|^2\right) \leq C \sum_{i=1}^n EX_i^2 \leq CnQ_n = \sum_{k=1}^n \alpha_k. \quad (4.17)$$

It is a simple fact that  $\alpha_k \geq 0$  for all  $k \geq 1$ . It follows by (4.9) that

$$\begin{aligned} \sum_{l=1}^{\infty} \Lambda_l \left(\frac{1}{b_l^2} - \frac{1}{b_{l+1}^2}\right) &= C \sum_{l=1}^{\infty} lQ_l \left(\frac{1}{l^{2/p}} - \frac{1}{(l+1)^{2/p}}\right) \\ &\leq \frac{2C}{p} \sum_{l=1}^{\infty} \frac{Q_l}{l^{2/p}} < \infty. \end{aligned} \quad (4.18)$$

That is to say (1.12) holds. By Remark 2.1 in Fazekas and Klesov [14], (1.12) implies (1.13). By Lemma 1.5, we can obtain (4.10)–(4.15) immediately. By (4.14), it follows that

$$\begin{aligned}
 E\left(\sup_{l \geq 1} \left| \frac{S_l}{l^{1/p}} \right|^r\right) &= \int_0^\infty P\left(\sup_{l \geq 1} \left| \frac{S_l}{l^{1/p}} \right|^r > t\right) dt \\
 &\leq 1 + \int_1^\infty P\left(\sup_{l \geq 1} \left| \frac{S_l}{l^{1/p}} \right| > t^{1/r}\right) dt \\
 &\leq 1 + E\left(\sup_{l \geq 1} \left| \frac{S_l}{l^{1/p}} \right|^2\right) \int_1^\infty t^{-2/r} dt \\
 &\leq 1 + \frac{4r}{2-r} \sum_{l=1}^\infty \frac{\alpha_l}{l^{2/p}} < \infty.
 \end{aligned} \tag{4.19}$$

The proof is completed.  $\square$

*Remark 4.3.* By using the maximal inequality, we get the Hájek-Rényi-type inequality, the strong law of large numbers and the strong growth rate for  $\varphi$ -mixing sequences. In addition, we get some new bounds of probability inequalities for  $\varphi$ -mixing sequences, such as (3.1), (3.5), (3.12), (4.5)–(4.6), and (4.13)–(4.16).

## Acknowledgments

The authors are most grateful to the Editor Sever Silvestru Dragomir, the referee Professor Mihaly Bencze and an anonymous referee for careful reading of the manuscript and valuable suggestions and comments which helped to significantly improve an earlier version of this paper. This work was supported by the National Natural Science Foundation of China (Grant nos. 10871001, 60803059) and the Innovation Group Foundation of Anhui University.

## References

- [1] R. L. Dobrushin, "The central limit theorem for non-stationary Markov chain," *Theory of Probability and Its Applications*, vol. 1, no. 4, pp. 72–88, 1956.
- [2] S. A. Utev, "On the central limit theorem for  $\varphi$ -mixing arrays of random variables," *Theory of Probability and Its Applications*, vol. 35, no. 1, pp. 131–139, 1990.
- [3] D. C. Chen, "A uniform central limit theorem for nonuniform  $\varphi$ -mixing random fields," *The Annals of Probability*, vol. 19, no. 2, pp. 636–649, 1991.
- [4] N. Herrndorf, "The invariance principle for  $\varphi$ -mixing sequences," *Zeitschrift für Wahrscheinlichkeitstheorie und Verwandte Gebiete*, vol. 63, no. 1, pp. 97–108, 1983.
- [5] M. Peligrad, "An invariance principle for  $\varphi$ -mixing sequences," *The Annals of Probability*, vol. 13, no. 4, pp. 1304–1313, 1985.
- [6] P. K. Sen, "A note on weak convergence of empirical processes for sequences of  $\varphi$ -mixing random variables," *Annals of Mathematical Statistics*, vol. 42, no. 6, pp. 2131–2133, 1971.
- [7] P. K. Sen, "Weak convergence of multidimensional empirical processes for stationary  $\varphi$ -mixing processes," *The Annals of Probability*, vol. 2, no. 1, pp. 147–154, 1974.
- [8] J. Iosifescu, "Limit theorem for  $\varphi$ -mixing sequences," in *Proceedings of the 5th Conference on Probability Theory*, pp. 1–6, September 1977.
- [9] M. Peligrad, "On Ibragimov-Iosifescu conjecture for  $\varphi$ -mixing sequences," *Stochastic Processes and Their Applications*, vol. 35, no. 2, pp. 293–308, 1990.

- [10] Q. M. Shao, "Almost sure invariance principles for mixing sequences of random variables," *Stochastic Processes and Their Applications*, vol. 48, no. 2, pp. 319–334, 1993.
- [11] S. Hu and X. Wang, "Large deviations for some dependent sequences," *Acta Mathematica Scientia. Series B*, vol. 28, no. 2, pp. 295–300, 2008.
- [12] C. R. Lu and Z. Y. Lin, *Limit Theory for Mixing Dependent Sequences*, Science Press, Beijing, China, 1997.
- [13] S. Hu, G. Chen, and X. Wang, "On extending the Brunk-Prokhorov strong law of large numbers for martingale differences," *Statistics & Probability Letters*, vol. 78, no. 18, pp. 3187–3194, 2008.
- [14] I. Fazekas and O. Klesov, "A general approach to the strong law of large numbers," *Theory of Probability and Its Applications*, vol. 45, no. 3, pp. 436–449, 2001.
- [15] S. Hu, "Some new results for the strong law of large numbers," *Acta Mathematica Sinica*, vol. 46, no. 6, pp. 1123–1134, 2003 (Chinese).