



Permutation Groups with a Cyclic Regular Subgroup and Arc Transitive Circulants

CAI HENG LI

li@maths.uwa.edu.au

Department of Mathematics, Yunnan University, Kunming 650031, People's Republic of China;

School of Mathematics and Statistics, The University of Western Australia, Crawley, 6009 WA, Australia

Received January 29, 2003; Revised January 21, 2004; Accepted January 21, 2004

Abstract. A description is given of finite permutation groups containing a cyclic regular subgroup. It is then applied to derive a classification of arc transitive circulants, completing the work dating from 1970's. It is shown that a connected arc transitive circulant Γ of order n is one of the following: a complete graph K_n , a lexicographic product $\Sigma[K_b]$, a deleted lexicographic product $\Sigma[K_b] - b\Sigma$, where Σ is a smaller arc transitive circulant, or Γ is a normal circulant, that is, $\text{Aut}\Gamma$ has a normal cyclic regular subgroup. The description of this class of permutation groups is also used to describe the class of rotary Cayley maps in subsequent work.

Keywords: cyclic regular subgroup, arc transitive circulant

1991 MR Subject Classification: 20B15, 20B30, 05C25

1. Introduction

A finite permutation group which contains a cyclic regular subgroup is called a *c-group*, for convenience. Characterizing *c-groups* is an old topic in permutation group theory, initiated by Burnside (1900), and studied by Schur et al., see for example [17, Chapter 4], [5], [7, Theorem 1.49] and [9]. The classical result of Schur tells us that a primitive *c-group* is 2-transitive or has prime degree. Thus, based on the finite simple group classification, primitive *c-groups* are essentially known in 1980's, see [5, 7, 9]. A precise list of primitive *c-groups* was recently given in [8, 13], independently.

Proposition 1.1 (see [13, Corollary 1.2]) *A primitive permutation group X of degree n contains a cyclic regular subgroup G if and only if one of the following holds:*

- (i) $\mathbb{Z}_p \cong G \leq X \leq \text{AGL}(1, p)$, where p is a prime;
- (ii) $X = A_n$ with n odd, or S_n , where $n \geq 4$;
- (iii) $\text{PGL}(d, q) \leq X \leq \text{P}\Gamma\text{L}(d, q)$, and G is a Singer subgroup of X , and $n = \frac{q^d - 1}{q - 1}$;
- (iv) $X = \text{P}\Gamma\text{L}(2, 8)$, and $G = \langle s\sigma \rangle \cong \mathbb{Z}_9$, where $\langle s \rangle$ is a Singer subgroup of X and $\sigma \in X \setminus \text{PSL}(2, 8)$ such that $o(\sigma) = 3$;
- (v) $(X, n) = (\text{PSL}(2, 11), 11)$, $(M_{11}, 11)$, or $(M_{23}, 23)$.

The main purpose of this paper is to give a description of general *c-groups*. To state the description, we need more notation and definitions.

For a transitive permutation group X on Ω , an orbit Δ of X on $\Omega \times \Omega$ is called an *orbital*, and the digraph with vertex set Ω and arc set Δ is called an *orbital graph* of X . (An *arc* of a digraph is an ordered pair of adjacent vertices.) An arc-disjoint union of orbital graphs is called a *generalised orbital graph* of X .

As usual, denote by K_n the complete graph of order n . By \bar{K}_n we mean the complement of K_n , that is, the graph with the same vertex set as K_n and contains no edges. For a digraph Γ with vertex set V and a digraph Σ with vertex set W , the *lexicographic product* $\Gamma[\Sigma]$ of a digraph Σ by a digraph Γ is the digraph with vertex set $V \times W$ such that (v_1, w_1) is adjacent to (v_2, w_2) if and only if either v_1 is adjacent to v_2 in Γ , or $v_1 = v_2$ and w_1 is adjacent to w_2 in Σ .

Let Γ be a digraph with vertex set Ω such that $X \leq \text{Aut}\Gamma$ is transitive on Ω . For a normal subgroup $N \triangleleft X$ which is intransitive on Ω , denote by Ω_N the set of N -orbits in Ω . The *normal quotient* Γ_N of the digraph Γ induced by N is the digraph with vertex set Ω_N such that for any $B, C \in \Omega_N$, B is adjacent to C if and only if there exist $u \in B$ and $v \in C$ such that u is adjacent to v in Γ .

A c -group X is called a *normal c -group* if it contains a normal regular cyclic group. Then by Proposition 1.1, a primitive c -group is either 2-transitive, or normal. In particular, all insoluble primitive c -groups are 2-transitive.

A description of c -groups is stated in the following theorem, which shows that either the structure of X is known, or the structure of some orbital graphs of X is known.

Theorem 1.2 *Let X be a permutation group on a set Ω of degree n . Assume that X contains a cyclic regular subgroup G . Then one of the following holds:*

- (1) *X has a normal subgroup $Y = Y_1 \times Y_2 \times \cdots \times Y_r$ with $r \geq 1$ such that $G \leq Y$, and further,

 - (i) *each Y_i is a 2-transitive c -group or a normal c -group of degree n_i ,*
 - (ii) *$n = n_1 n_2 \dots n_r$, and the n_i are pairwise coprime.**
- (2) *X has a normal subgroup N such that each connected generalised orbital graph of X contains a subgraph Γ which is an orbital graph of X and is of the form $\Gamma = \Gamma_N[\bar{K}_b]$, where $b = |N \cap G|$.*

A digraph Γ is called a *Cayley graph* if there exist a group G and a subset $S \subset G$ such that the vertex set of Γ may be identified with G and u is adjacent to v if and only if $vu^{-1} \in S$. This digraph Γ is denoted by $\text{Cay}(G, S)$. A Cayley graph $\Gamma = \text{Cay}(G, S)$ has an automorphism group

$$\hat{G} = \{\hat{g} : x \mapsto xg \text{ for all } x \in G \mid g \in G\},$$

consisting all right multiplications of elements of G . Thus the subgroup $\hat{G} \leq \text{Aut}\Gamma$ acts regularly on the vertex set of Γ . In particular, Γ is *vertex transitive*. Let $\text{Aut}(G, S) = \{\sigma \in \text{Aut}(G) \mid S^\sigma = S\}$. Then each element in $\text{Aut}(G, S)$ induces an automorphism of the Cayley graph $\Gamma = \text{Cay}(G, S)$. Moreover, it is easily shown that $\mathbf{N}_{\text{Aut}\Gamma}(\hat{G}) = \hat{G}:\text{Aut}(G, S)$, see for example [6]. A Cayley graph $\Gamma = \text{Cay}(G, S)$ is called *normal* if \hat{G} is normal in $\text{Aut}\Gamma$, that is, $\text{Aut}\Gamma = \hat{G}:\text{Aut}(G, S)$.

Let G be a cyclic group, and let $\Gamma = \text{Cay}(G, S)$. Then this Cayley graph is called a *circulant*. Since $\text{Aut}\Gamma$ contains the cyclic regular subgroup \hat{G} , $\text{Aut}\Gamma$ is a c -group. If Γ is a normal Cayley graph, that is, \hat{G} is normal in $\text{Aut}\Gamma$, then Γ is called a *normal circulant*. The description of c -groups given in Theorem 1.2 can be used to study circulants. The second result of this paper is to present a classification of arc transitive circulants. (Recall that a graph Γ is *arc transitive* if $\text{Aut}\Gamma$ is transitive on the arcs of Γ .)

The problem of classifying arc transitive circulants has been investigated for a long time, dating from 1970's, and it has been solved for several special cases: A classification of 2-arc transitive circulants was obtained in [1] for the undirected case and in [16] for the directed case; while a classification of arc transitive circulants of special orders was given by a collection of articles: prime order in [2, 3], square-free order in [15], and odd prime-power order in [19].

For a positive integer b and a digraph Γ , denote by $b\Gamma$ the digraph consisting of b vertex-disjoint copies of Γ ; the digraph $\Gamma[\bar{K}_b] - b\Gamma$ is called a *deleted lexicographic product*, which is the digraph whose vertex set is the vertex set of $\Gamma[\bar{K}_b]$ and arc set equals the arc set of $\Gamma[\bar{K}_b]$ minus the arc set of $b\Gamma$.

Theorem 1.3 *Let Γ be a connected arc transitive circulant of order n which is not a complete graph. Then either*

- (1) Γ is a normal circulant, or
- (2) there exists an arc transitive circulant Σ of order m such that $n = mb$ with $b, m > 1$ and

$$\Gamma = \begin{cases} \Sigma[\bar{K}_b], & \text{or} \\ \Sigma[\bar{K}_b] - b\Sigma & \text{with } (b, m) = 1. \end{cases}$$

Remarks

- (a) After the previous version of this paper had been submitted for publication, the author was aware of that the classification of arc transitive circulants given in Theorems 1.3 was also obtained by I. Kovács [10], independently. Also the referees of this paper pointed out this to the author.
- (b) It is easily shown that a digraph is a circulant if and only if its automorphism group is a c -group. An arc transitive circulant is therefore exactly an orbital graph of a c -group. Thus the proof of Theorem 1.3 easily follows from Theorem 1.2, and will not be presented in this paper. A proof for this classification may be found in [10].
- (c) If an arc transitive circulant Γ is of the form $\Sigma[\bar{K}_b]$ or $\Sigma[\bar{K}_b] - b\Sigma$, namely Γ satisfies part (2) of Theorem 1.3, then Γ can be easily reconstructed from a smaller arc transitive circulant Σ . As the automorphism group of a cyclic group is abelian, normal circulants may be easily and explicitly constructed. Arc transitive circulants are therefore well-characterized by Theorem 1.3.
- (d) The circulants concerned in Theorem 1.3 are directed graphs. It is easily shown that an undirected edge transitive circulant Γ is arc transitive, and Theorem 1.3 may be easily changed into a version for the undirected case.

2. Proof of Theorem 1.2

This section is devoted to proving Theorem 1.2. The proof is based on a result of Evdokimov and Ponomarenko [4] regarding the 2-closures of c -groups. We here first introduce some simple properties for the 2-closures of transitive permutation groups.

Let G be a transitive permutation group on Ω . The 2-closure of X is the largest subgroup of $\text{Sym}(\Omega)$ which preserves each orbital of X ; denoted by $X^{(2)}$. We observe the following properties:

- (i) X is 2-transitive if and only if $X^{(2)} = \text{Sym}(\Omega)$;
- (ii) X is primitive if and only if $X^{(2)}$ is primitive.

Assume that X is imprimitive, and let \mathcal{B} be an *imprimitive partition* of Ω , that is, \mathcal{B} is a non-trivial X -invariant partition of Ω . For any block $B \in \mathcal{B}$, denote by X^B the restriction of X to B , that is, $X^B = \langle x \in X \mid B^x = B \rangle \leq \text{Sym}(B)$. It is easily shown that an orbital of X containing an arc (u, v) with $u, v \in B$ is also an orbital of $X^{(2)}$. It follows that B is an imprimitive block for $X^{(2)}$, and \mathcal{B} is an imprimitive partition of Ω for $X^{(2)}$. Thus we have the third observation:

- (iii) X and $X^{(2)}$ have the same imprimitive partitions of Ω .

For the action induced on a block, we further have the following conclusion.

Lemma 2.1 ([18, 5.25]) *Let X be a transitive permutation group on Ω such that Ω has a non-trivial X -invariant partition \mathcal{B} . Then for $B \in \mathcal{B}$, we have*

$$(X^{(2)})^B \leq (X^B)^{(2)}.$$

For some c -groups, the 2-closures are known.

Example 2.2 If X is a primitive c -group of degree n , then either $X^{(2)} = \text{S}_n$, or $n = p$ is a prime and $X \leq X^{(2)} \cong \mathbb{Z}_p : \mathbb{Z}_l$, where l is a proper divisor of $p - 1$.

A powerful method for studying c -groups comes from Schur ring theory, which was initiated by I. Schur and developed by H. Wielandt, see [17, Chapter 4]. A complete description of Schur rings over a cyclic group is given in [4, 11, 12]. From the description, the following result regarding c -groups may be derived.

Theorem 2.3 ([4, 11, 12]) *Let X be a c -group on Ω , and let $X^{(2)}$ be the 2-closure of X . Then one of the following statements holds:*

- (i) $X^{(2)} = X_1 \times X_2 \times \cdots \times X_r$, where $r \geq 1$, each $X_i \cong \text{S}_{n_i}$, or a normal c -group of degree n_i , such that $(n_i, n_j) = 1$ and $n = n_1 n_2 \cdots n_r$;
- (ii) X has a normal subgroup M such that each connected generalised orbital graph contains a subgraph Γ which is an orbital graph of X and is of the form $\Gamma = \Gamma_M[\bar{K}_b]$, where $b = |M \cap G|$.

The author is grateful to a referee for pointing out and formulating this theorem.

The next lemma shows that if X contains an abelian regular subgroup, then each X -invariant partition of Ω consists of orbits of a normal subgroup.

Lemma 2.4 ([13, Lemma 3.1]) *Let X be a transitive permutation group on Ω which contains an abelian regular subgroup G . Let \mathcal{B} be an X -invariant partition of Ω and $B \in \mathcal{B}$, and let K be the kernel of X acting on \mathcal{B} . Then \mathcal{B} equals the set of K -orbits in Ω , and $G \cap K$ is regular on B .*

Now we are ready to prove Theorem 1.2. In the original version of this paper, a permutation group theoretical proof of Theorem 1.2 was given, which was independent of Theorem 2.3. The following proof was suggested by one of the referees. The author is grateful to him for this suggestion.

Proof of Theorem 1.2: Let X be a c -group on Ω of degree n , and let G be a regular cyclic subgroup of X . To complete the proof of Theorem 1.2, by Theorem 2.3, we may assume that the 2-closure $X^{(2)}$ satisfies

$$X^{(2)} = X_1 \times X_2 \times \cdots \times X_r,$$

where $r \geq 1$, either $X_i \cong \mathbf{S}_{n_i}$, or X_i is a normal c -group of degree n_i , such that $(n_i, n_j) = 1$ and $n = n_1 n_2 \cdots n_r$.

If $r = 1$ and $X_1 \cong \mathbf{S}_{n_1}$, then $X^{(2)} = X_1 \cong \mathbf{S}_{n_1}$, and hence X is a 2-transitive c -group. If all X_i 's are normal c -groups, then $X^{(2)}$ is a normal c -group. Since now $G \leq X \leq X^{(2)}$, it follows that X is a normal c -group.

Thus we assume that $r \geq 2$, and that at least one of X_i is a symmetric group, say $X_1 \cong \mathbf{S}_{n_1}$, where $n_1 \geq 4$. Let $Y_i = X \cap X_i$, and let $Y = \langle Y_1, Y_2, \dots, Y_r \rangle$. Then $Y_i \triangleleft X$, and for $j \neq i$, we have $Y_i \cap Y_j = 1$. Thus $Y = Y_1 \times Y_2 \times \cdots \times Y_r$, and Y is normal in X . Let $G_i = G \cap X_i$. Then $|G_i| = n_i$, and for $j \neq i$, we have $G_i \cap G_j = 1$. Since $(n_i, n_j) = 1$, it follows that $G = G_1 \times G_2 \times \cdots \times G_r$.

Let B be an orbit of X_i in Ω , and let \mathcal{B} be the set of X_i -orbits in Ω . Then \mathcal{B} is an $X^{(2)}$ -invariant partition of Ω . It is easily shown that X_i equals the kernel of $X^{(2)}$ acting on \mathcal{B} . It follows that $(X^{(2)})^B \cong X_i$. Now G_i is cyclic and regular on B , and hence X_i is a c -group of degree n_i .

Let $Y_i = X \cap X_i$. Since $G \leq X$, we have that $G_i \leq X \cap X_i = Y_i$. Hence Y_i is a c -group on B , and \mathcal{B} is the set of Y_i -orbits in Ω . It is easily shown that $X^B \cong Y_i$. If X_i is a normal c -group, then it follows that Y_i is also a normal c -group.

Assume that $X_i \cong \mathbf{S}_{n_i}$. Then by Lemma 2.1, we have $\mathbf{S}_{n_i} \cong X_i \cong (X^{(2)})^B \leq (X^B)^{(2)}$, and thus X^B is 2-transitive. So $Y_i \cong X^B$ is a 2-transitive c -group of degree n_i .

Therefore, each Y_i is a 2-transitive c -group or a normal c -group, of degree n_i , and $G \leq Y = Y_1 \times Y_2 \times \cdots \times Y_r$, completing the proof of Theorem 1.2. \square

Acknowledgment

This work forms a part of an Australian Research Council large grant project, and the author is supported by a QEII Fellowship. The author is very grateful to one referee for his

constructive suggestions, and especially for his formulating Theorem 2.3, which leads to the very short proof for Theorem 1.2 presented in this paper.

References

1. B. Alspach, M. Conder, D. Marušič, and M.Y. Xu, "A classification of 2-arc-transitive circulants," *J. Algebraic Combin.* **5** (1996), 83–86.
2. C.Y. Chao, "On the classification of symmetric graphs with a prime number of vertices," *Trans. Amer. Math. Soc.* **158** (1971), 247–256.
3. C.Y. Chao and J.G. Wells, "A class of vertex-transitive digraphs," *J. Combin. Theory Ser. B* **14** (1973), 246–255.
4. S.A. Evdokimov and I.N. Ponomarenko, "Characterization of cyclotomic schemes and normal Schur rings over a cyclic group, (Russian)," *Algebra i Analiz* **14**(2) (2002), 11–55.
5. W. Feit, "Some consequences of the classification of finite simple groups," The Santa Cruz Conference on Finite Groups Univ. California, Santa Cruz, Calif., 1979, pp. 175–181, *Proc. Sympos. Pure Math.*, **37**, Amer. Math. Soc., Providence, R.I., 1980.
6. C.D. Godsil, "On the full automorphism group of a graph," *Combinatorica* **1** (1981), 243–256.
7. D. Gorenstein, *Finite Simple Groups*, Plenum Press, New York, 1982.
8. G. Jones, "Cyclic regular subgroups of primitive permutation groups," *J. Group Theory* **5**(4) (2002), 403–407.
9. W. Kantor, Some consequences of the classification of finite simple groups, in *Finite Groups—Coming of Age*, John McKay (Ed.), Amer. Math. Soc., 1982.
10. I. Kovács, "Classifying arc-transitive circulants," *J. Algebraic Combin.* **20** (2004), 353–358.
11. K.H. Leung and S.H. Man, "On Schur rings over cyclic groups," II, *J. Algebra* **183** (1996), 173–285.
12. K.H. Leung and S. H. Man, "On Schur rings over cyclic groups," *Israel J. Math.* **106** (1998), 251–267.
13. C.H. Li, "The finite primitive permutation groups containing an abelian regular subgroup," *Proc. London Math. Soc.* **87** (2003), 725–748.
14. C.H. Li, "Edge-transitive Cayley graphs and rotary Cayley maps," *Trans. Amer. Math. Soc.* (to appear).
15. C.H. Li, D. Marušič, and J. Morris, "A classification of circulant arc-transitive graphs of square-free order," *J. Algebraic Combin.* **14** (2001), 145–151.
16. J.X. Meng and J.Z. Wang, "A classification of 2-arc-transitive circulant digraphs," *Disc. Math.* **222** (2000), 281–284.
17. H. Wielandt, *Finite Permutation Groups*, Academic Press, New York, 1964.
18. H. Wielandt, Permutation groups through invariant relations and invariant functions, in *Mathematische Werke/Mathematical works*. Vol. 1. *Group theory*, Bertram Huppert and Hans Schneider (Eds.), Walter de Gruyter Co., Berlin, 1994, pp. 237–266.
19. M.Y. Xu, H.S. Sim, and Y.J. Baik, "Arc-transitive circulant digraphs of odd prime-power order," *Discrete Math.* **287** (2004), 113–119.