

ON INTEGRAL INEQUALITIES OF GRONWALL-BELLMAN-BIHARI TYPE IN SEVERAL VARIABLES

JAMES ADEDAYO OGUNTUASE

Department of Mathematical Sciences
University of Agriculture, Abeokuta
NIGERIA

EMail: adedayo@unaab.edu.ng



volume 1, issue 2, article 20,
2000.

*Received 23 February, 2000;
accepted 17 April 2000.*

Communicated by: G. Anastassiou

[Abstract](#)

[Contents](#)




[Home Page](#)

[Go Back](#)

[Close](#)

[Quit](#)



Abstract

We present some new results on the linear and non-linear integral inequalities of Gronwall-Bellman-Bihari type to n -dimensional integrals with a kernel of the form $k(x, t)$ where x and t are in $S \subset \mathbb{R}^n$.

These inequalities extend and compliment some existing results in the literature on Gronwall-Bellman-Bihari type inequalities.

2000 Mathematics Subject Classification: 26D15

Key words: Gronwall-Bellman-Bihari inequality, good kernel, nonincreasing function, nondecreasing function, nonnegative function.

Contents

1	Introduction	3
2	Linear Inequalities	4
3	Non-linear Inequalities	9
	References	

On Integral inequalities of Gronwall-Bellman-Bihari type in several variables

J.A. Oguntuase

Title Page

Contents



Go Back

Close

Quit

Page 2 of 15

1. Introduction

The results obtained in this paper originated from the celebrated Gronwall-Bellman-Bihari inequality which has been of vital importance in the study of existence, uniqueness, continuous dependence, comparison, perturbation, boundedness and stability of solutions of differential and integral equations (see for example [1, 2, 3, 4, 5, 6] and the references cited therein).

In the last three decades, more than one variable generalizations of these inequalities have been obtained and these results have generated a lot of research interests due to its usefulness in the theory of differential and integral equations (see for example [1, 3, 6, 7, 8, 9, 10] and the references cited therein).

The purpose of this paper is to establish some new integral inequalities in n independent variables which will compliment the existing results in the literature on Gronwall- Bellman-Bihari type inequalities in several variables.


Throughout this paper, we shall assume that S is any bounded open set in the n dimensional Euclidean space \mathfrak{R}^n and that our integrals are on $\mathfrak{R}^n (n \geq 1)$, unless otherwise specified.

For $x = (x_1, x_2, \dots, x_n)$, $x^0 = (x_1^0, x_2^0, \dots, x_n^0) \in S$, we shall denote the integral

$$\int_{x_1^0}^{x_1} \int_{x_1^0}^{x_2} \dots \int_{x_1^0}^{x_n} \dots dt_n \dots dt_1 \quad \text{by} \quad \int_{x^0}^x \dots dt$$

and $D_i = \frac{\partial}{\partial x_i}$ for $i = 1, 2, \dots, n$.

Furthermore, for $x, t \in \mathfrak{R}^n$, we shall write $t \leq x$ whenever $t_i \leq x_i, i = 1, 2, \dots, n$. Unless otherwise specified, all functions considered are functions of n -variables which are nonnegative and continuous on $[x^0, x], x \geq x^0 \geq 0$ and $x \in S$.



On Integral inequalities of Gronwall-Bellman-Bihari type in several variables

J.A. Oguntuase

Title Page

Contents



Go Back

Close

Quit

Page 3 of 15

2. Linear Inequalities

In this section, we shall obtain bounds to the linear Gronwall-Bellman-Bihari type integral inequalities for a more general kernel $k(x, t)$ and a product kernel $k(x, t) = h(x)f(t)$.

Definition 2.1. A function $k(x, t)$ of the $2n$ variables x_1, \dots, t_n is called a good kernel if

1. $k(\cdot, \cdot) \geq 0$.
2. $k(\cdot, \cdot)$ is a continuous function of its $2n$ variables.
3. $k(\cdot, \cdot)$ is monotone non-decreasing in its first n variables, i.e. $k(x, t) \geq k(y, t)$ whenever $x \geq y$.

Theorem 2.1. Let $k(x, t)$ be a good kernel, $u(x)$ is a real valued nonnegative continuous function on S and $g(x)$ be a positive, nondecreasing continuous function on S . Suppose that the following inequality


$$(2.1) \quad u(x) \leq g(x) + \int_{x^0}^x k(x, t)u(t)dt$$

holds for all $x \in S$ with $x \geq x^0$, then

$$(2.2) \quad u(x) \leq g(x) \left\{ 1 + \int_{x^0}^x k(s, s) \exp \left(\int_{x^0}^s k(t, t)dt \right) ds \right\}.$$

Proof. Since $g(x)$ is positive and nondecreasing, we can write (2.1) as

$$\frac{u(x)}{g(x)} \leq 1 + \int_{x^0}^x k(x, t) \frac{u(t)}{g(t)} dt.$$



On Integral inequalities of
Gronwall-Bellman-Bihari type in
several variables

J.A. Oguntuase

Title Page

Contents



Go Back

Close

Quit

Page 4 of 15

Setting $\frac{u(x)}{g(x)} = r(x)$, then we have

$$r(x) \leq 1 + \int_{x^0}^x k(x, t)r(t)dt.$$

Let

$$v(x) = 1 + \int_{x^0}^x k(x, t)r(t)dt.$$

Then

$$r(x) \leq v(x)$$

and $v(x^0) = 1$ or $x_i = x_i^0, i = 1, 2, \dots, n$.

Hence

$$(2.3) \quad D_1 \cdots D_n v(x) = k(x, x)r(x) \leq k(x, x)v(x).$$

From (2.3) we obtain


$$\frac{v(x)D_1 \cdots D_n v(x)}{v^2(x)} \leq k(x, x).$$

That is

$$\frac{v(x)D_1 \cdots D_n v(x)}{v^2(x)} \leq k(x, x) + \frac{(D_n v(x)) (D_1 \cdots D_{n-1} v(x))}{v^2(x)}.$$

Hence

$$D_n \left(\frac{D_1 \cdots D_{n-1} v(x)}{v(x)} \right) \leq k(x, x).$$



On Integral inequalities of Gronwall-Bellman-Bihari type in several variables

J.A. Oguntuase

Title Page

Contents



Go Back

Close

Quit

Page 5 of 15

Integrating with respect to x_n from x_n^0 to x_n , we have

$$\frac{D_1 \cdots D_{n-1} v(x)}{v(x)} \leq \int_{x_n^0}^{x_n} k(x_1, x_2, \dots, x_{n-1}, t_n, x_1, x_2, \dots, x_{n-1}, t_n) dt_n.$$

Thus

$$\begin{aligned} \frac{v(x) D_1 \cdots D_{n-1} v(x)}{v^2(x)} &\leq \int_{x_n^0}^{x_n} k(x_1, x_2, \dots, x_{n-1}, t_n, x_1, x_2, \dots, x_{n-1}, t_n) dt_n \\ &+ \frac{(D_{n-1} v(x)) (D_1 \cdots D_{n-2} v(x))}{v^2(x)}. \end{aligned}$$

That is


$$D_{n-1} \left(\frac{D_1 \cdots D_{n-2} v(x)}{v(x)} \right) \leq \int_{x_n^0}^{x_n} k(x_1, x_2, \dots, x_{n-1}, t_n, x_1, x_2, \dots, x_{n-1}, t_n) dt_n.$$

Integrating with respect to x_{n-1} from x_{n-1}^0 to x_{n-1} , we have

$$\begin{aligned} &\frac{D_1 \cdots D_{n-2} v(x)}{v(x)} \\ &\leq \int_{x_{n-1}^0}^{x_{n-1}} \int_{x_n^0}^{x_n} k(x_1, x_2, \dots, x_{n-2}, t_{n-1}, t_n, x_1, x_2, \dots, x_{n-2}, t_{n-1}, t_n) dt_n dt_{n-1}. \end{aligned}$$

Continuing this process, we obtain

$$\frac{D_1 D_2 v(x)}{v(x)} \leq \int_{x_3^0}^{x_3} \cdots \int_{x_n^0}^{x_n} k(x_1, x_2, t_3, \dots, t_n, x_1, x_2, t_3, \dots, t_n) dt_n \cdots dt_3.$$



On Integral inequalities of Gronwall-Bellman-Bihari type in several variables

J.A. Oguntuase

Title Page

Contents



Go Back

Close

Quit

Page 6 of 15

From this we obtain

$$D_2 \left(\frac{D_1 v(x)}{v(x)} \right) \leq \int_{x_3^0}^{x_3} \cdots \int_{x_n^0}^{x_n} k(x_1, x_2, t_3 \dots, t_n, x_1, x_2, t_3 \dots, t_n) dt_n \dots dt_3.$$

Integrating with respect to the x_2 component from x_2^0 to x_2 , we have

$$\frac{D_1 v(x)}{v(x)} \leq \int_{x_2^0}^{x_2} \cdots \int_{x_n^0}^{x_n} k(x_1, t_2, t_3 \dots, t_n, x_1, t_2, t_3 \dots, t_n) dt_n \dots dt_2.$$

Integrating with respect to the x_1 component from x_1^0 to x_1 , we obtain

$$\log \frac{v(x)}{v(x_1^0, x_2, \dots, x_n)} \leq \int_{x_1^0}^x k(t, t) dt.$$

That is


$$(2.4) \quad v(x) \leq \exp \left(\int_{x_1^0}^x k(t, t) dt \right).$$

Substituting (2.4) into (2.3) we have

$$D_1 \cdots D_n r(x) \leq k(x, x) v(x) \leq k(x, x) \exp \left(\int_{x_1^0}^x k(t, t) dt \right).$$

Integrating this inequality with respect to the x_n component from x_n^0 to x_n , then with respect to the x_{n-1}^0 to x_{n-1} , and continuing until finally x_1^0 to x_1 , and noting that $r(x) = 1$ at $x_i = x_i^0$, we have

$$r(x) \leq 1 + \int_{x_1^0}^x k(s, s) \exp \left(\int_{x_1^0}^s k(t, t) dt \right) ds.$$



On Integral inequalities of Gronwall-Bellman-Bihari type in several variables

J.A. Oguntuase

Title Page

Contents



Go Back

Close

Quit

Page 7 of 15

Since $\frac{u(x)}{g(x)} = r(x)$, then we obtain

$$u(x) \leq g(x) \left\{ 1 + \int_{x^0}^x k(s, s) \exp \left(\int_{x^0}^s k(t, t) dt \right) ds \right\}.$$

This completes the proof of our result. \square

Next, we shall consider the case in which $k(x, y) = h(x)f(t)$. Then we have the following result.

Theorem 2.2. *Let $h(x)$, $f(t)$, $u(x)$ be real valued nonnegative continuous functions on S and $g(x)$ be a positive, nondecreasing continuous function on S . If $h'(x) = 0$, where the prime denote $\frac{\partial^n}{\partial x_1 \dots \partial x_n}$ and the following inequality*

$$(2.5) \quad u(x) \leq g(x) + h(x) \int_{x^0}^x f(t)u(t)dt$$


holds for all $x \in S$ with $x \geq x^0$, then

$$(2.6) \quad u(x) \leq g(x) \left\{ 1 + \int_{x^0}^x h(s)f(s) \exp \left(\int_{x^0}^s h(t)f(t)dt \right) ds \right\}.$$

Proof. Similar to the proof of Theorem 2.1 and so the details are omitted. \square

Remark 2.3. *If we set $k(x, t) = f(t)$ in Theorem 2.2, then our estimate reduces to*

$$u(x) \leq g(x) \left\{ 1 + \int_{x^0}^x f(s) \exp \left(\int_{x^0}^s f(t)dt \right) ds \right\}.$$



3. Non-linear Inequalities

Definition 3.1. A function $\phi : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ is said to belong to the class \mathcal{F} if it satisfies the following conditions:

1. ϕ is nondecreasing and continuous in \mathbb{R}^+ and $\phi(u) > 0$ for $u > 0$;
2. $\frac{1}{\alpha}\phi(u) \leq \phi\left(\frac{u}{\alpha}\right)$, $u \geq 0$, $\alpha \geq 1$.

We observe from the above definition that \mathcal{F} has the following properties

1. $\phi \in \mathcal{F}$ if and only if $\frac{\phi(u)}{u}$ is nonincreasing for $u > 0$;
2. $\phi \in \mathcal{F}$ implies that ϕ is subadditive;
3. If ϕ satisfies (1) of Definition 3.1 and is concave in \mathbb{R}^+ , then $\phi \in \mathcal{F}$.

Theorem 3.1. Let $k(x, t)$ be a good kernel and $u(x)$ be a real valued nonnegative continuous function on S . If $g(x)$ be a positive, nondecreasing continuous function on S and ϕ belong to class \mathcal{F} for which the following inequality


$$(3.1) \quad u(x) \leq g(x) + \int_{x^0}^x k(x, t)\phi(u(t))dt$$

holds for all $x \in S$ with $x \geq x^0$, then for $x^0 \leq x \leq x^*$,

$$(3.2) \quad u(x) \leq g(x)G^{-1}\left(G(1) + \int_{x^0}^x k(t, t)dt\right),$$

where

$$G(z) = \int_{z^0}^z \frac{ds}{\phi(s)}, \quad z \geq z^0 > 0,$$



On Integral inequalities of Gronwall-Bellman-Bihari type in several variables

J.A. Oguntuase

Title Page

Contents



Go Back

Close

Quit

Page 9 of 15

G^{-1} is the inverse of G and x^* is chosen so that

$$G(1) + \int_{x^0}^x k(t, t) dt \in \text{Dom}(G^{-1}).$$

Proof. Since $g(x)$ is positive and nondecreasing, we can write (3.1) as

$$\frac{u(x)}{g(x)} \leq 1 + \int_{x^0}^x k(x, t) \frac{\phi(u(t))}{g(t)} dt \leq 1 + \int_{x^0}^x k(x, t) \phi\left(\frac{u(t)}{g(t)}\right) dt.$$

Setting $\frac{u(x)}{g(x)} = v(x)$, then we have

$$v(x) \leq 1 + \int_{x^0}^x k(x, t) \phi(v(t)) dt.$$

Let

$$r(x) = 1 + \int_{x^0}^x k(x, t) \phi(v(t)) dt.$$

Then


$$v(x) \leq r(x)$$

and $v(x^0) = 1$ or $x_i = x_i^0$, $i = 1, 2, \dots, n$ and

$$D_1 \cdots D_n r(x) = k(x, x) \phi(r(x)).$$

That is

$$\frac{D_1 \cdots D_n r(x)}{\phi(r(x))} \leq k(x, x).$$



On Integral inequalities of Gronwall-Bellman-Bihari type in several variables

J.A. Oguntuase

Title Page

Contents



Go Back

Close

Quit

Page 10 of 15

Since

$$D_n \left(\frac{D_1 \cdots D_{n-1} r(x)}{\phi(v(x))} \right) = \frac{D_1 \cdots D_n r(x)}{\phi(r(x))} - \frac{D_n \phi(r(x)) D_1 \cdots D_{n-1} r(x)}{\phi^2(r(x))}$$

and

$$D_n \phi(r(x)) = \phi'(r(x)) D_n r(x) \geq 0, \quad D_1 \cdots D_{n-1} r(x) \geq 0.$$

The above inequality implies

$$D_n \left(\frac{D_1 \cdots D_{n-1} r(x)}{\phi(r(x))} \right) \leq k(x, x)$$

provided $\phi'(r(x)) \geq 0$ for $r(x) \geq 0$.

Integrating with respect to x_n from x_n^0 to x_n and taking into account the fact that $D_1 \cdots D_{n-1} r(x) = 0$ for $x_n = x_n^0$, we have


$$\frac{D_1 \cdots D_{n-1} r(x)}{\phi(v(x))} \leq \int_{x_n^0}^{x_n} k(x_1, x_2, \dots, x_{n-1}, t_n, x_1, x_2, \dots, x_{n-1}, t_n) dt_n.$$

Repeating this, we find (after $n - 1$ steps) that

$$\frac{D_1 r(x)}{\phi(r(x))} \leq \int_{x_1^0}^{x_1} \cdots \left(\int_{x_n^0}^{x_n} k(x_1, \dots, x_{n-1}, t_n, x_1, \dots, x_{n-1}, t_n) dt_n \right) \cdots dt_2.$$

We note that for

$$G(s) = \int_{s^0}^s \frac{dz}{\phi(z)}, \quad s \geq s^0 > 0.$$



On Integral inequalities of Gronwall-Bellman-Bihari type in several variables

J.A. Oguntuase

Title Page

Contents



Go Back

Close

Quit

Page 11 of 15

It thus follows that

$$D_1G(r(x)) = \frac{D_1r(x)}{\phi(r(x))},$$

so that

$$D_1G(r(x)) \leq \int_{x_2^0}^{x_2} k(x_1, t_2, \dots, t_n, x_1, t_2, \dots, t_n) dt_n \dots dt_2.$$

Integrating both sides of the above inequality with respect to the component

$$G(r(x_1, \dots, x_n)) - G(r(t_1, x_2, \dots, x_n)) \leq \int_{x^0}^x k(t, t) dt.$$

Since $r(t_1, x_2, \dots, x_n) = 1$ we have

$$r(x) \leq G^{-1} \left(G(1) + \int_{x^0}^x k(t, t) dt \right).$$


From this we obtain

$$v(x) \leq r(x) \leq G^{-1} \left(G(1) + \int_{x^0}^x k(t, t) dt \right).$$

Using the fact that $\frac{u(x)}{g(x)} = v(x)$, we have

$$u(x) \leq g(x)G^{-1} \left(G(1) + \int_{x^0}^x k(t, t) dt \right)$$

which is required and the proof is complete. □



**On Integral inequalities of
Gronwall-Bellman-Bihari type in
several variables**

J.A. Oguntuase

Title Page

Contents



Go Back

Close

Quit

Page 12 of 15

If we set $k(x, t) = h(x)f(t)$, then we shall obtain the following result.

Theorem 3.2. Let $h(x)$, $f(t)$, $u(x)$ be real valued nonnegative continuous functions on S and $g(x)$ be a positive, nondecreasing continuous function on S , and ϕ belong to class \mathcal{F} . If $h'(x) = 0$ and the following inequality

$$(3.3) \quad u(x) \leq g(x) + h(x) \int_{x^0}^x f(t)\phi(u(t))dt$$

holds for all $x \in S$ with $x \geq x^0$, then for $x^0 \leq x \leq x^*$, then

$$(3.4) \quad u(x) \leq g(x)G^{-1} \left(G(1) + h(x) \int_{x^0}^x f(t)dt \right),$$

where

$$G(z) = \int_{z^0}^z \frac{ds}{\phi(s)}, \quad z \geq z^0 > 0,$$


G^{-1} is the inverse of G and x^* is chosen so that

$$G(1) + h(x) \int_{x^0}^x f(t)dt \in \text{Dom}(G^{-1}).$$

Proof. Similar to the proof of Theorem 3.1 and so the details are omitted. \square

Remark 3.3. If we set $k(x, t) = f(t)$ in Theorem 3.2, then our estimate reduces to

$$u(x) \leq g(x)G^{-1} \left(G(1) + \int_{x^0}^x f(t)dt \right).$$



On Integral inequalities of Gronwall-Bellman-Bihari type in several variables

J.A. Oguntuase

Title Page

Contents



Go Back


Close

Quit

Page 13 of 15

References

- [1] O. AKINYELE, On Gronwall-Bellman-Bihari type integral inequalities in several variables with retardation, *J. Math. Anal. Appl.*, **104** (1984), 1–24.
- [2] D. BAINOV AND P. SIMEONOV, *Integral Inequalities and Applications*, Kluwer Academic Publishers, Dordrecht, Netherlands, 1992
- [3] J. CONLAN AND C-L. WANG, Higher Dimensional Gronwall- Bellman type inequalities, *Appl. Anal.*, **18** (1984), 1–12.
- [4] F.M. DANNAN, Submultiplicative and subadditive functions and integral inequalities of Bellman-Bihari type, *J. Math. Anal. Appl.*, **120** (1986), 631–646.
- [5] U.D. DHONGADE AND S.G. DEO, Some generalization of Bellman-Bihari integral inequalities, *J. Math. Anal. Appl.*, **120** (1973), 218–226.
- [6] J.A. OGUNTUASE, Higher dimensional integral inequality of Gronwall-Bellman type, *Ann. Stiint. Univ. 'Al. I. Cuza' din Iasi*, Ser. **T.45** (1999), in press.
- [7] J.A. OGUNTUASE, Gronwall-Bellman type integral inequalities for multi-distribution, *Riv. Mat. Univ. Parma*, **6(2)** (1999), in press.
- [8] C.C. YEH, Bellman-Bihari integral inequality in n independent variables, *J. Math. Anal. Appl.*, **87** (1982), 311–321.
- [9] C.C. YEH, On some integral inequalities in n independent variables and their applications, *J. Math. Anal. Appl.*, **87** (1982), 387–410.



On Integral inequalities of
Gronwall-Bellman-Bihari type in
several variables

J.A. Oguntuase

Title Page

Contents




Go Back

Close

Quit

Page 14 of 15

[10] C.C. YEH AND M-H. SHIH, The Gronwall-Bellman inequality in several variables, *J. Math. Anal. Appl.*, **86** (1982), 157–167.



**On Integral inequalities of
Gronwall-Bellman-Bihari type in
several variables**

J.A. Oguntuase

Title Page

Contents



Go Back

Close

Quit

Page 15 of 15