


Journal of Inequalities in Pure and Applied Mathematics



ON SIMULTANEOUS APPROXIMATION FOR CERTAIN BASKAKOV DURRMEYER TYPE OPERATORS

VIJAY GUPTA, MUHAMMAD ASLAM NOOR
AND MAN SINGH BENIWAL

School of Applied Sciences
Netaji Subhas Institute of Technology
Sector 3 Dwarka
New Delhi 110075, India
EMail: vijay@nsit.ac.in

Mathematics Department
COMSATS Institute of Information Technology
Islamabad, Pakistan
EMail: noormaslam@hotmail.com

Department of Applied Science
Maharaja Surajmal Institute of Technology
C-4, Janakpuri, New Delhi - 110058, India
EMail: man_s_2005@yahoo.co.in

Department of Mathematics
Ch Charan Singh University
Meerut 250004, India
EMail: mkgupta2002@hotmail.com

volume 7, issue 4, article 125,
2006.

*Received 06 January, 2006;
accepted 16 August, 2006.*

Communicated by: N.E. Cho

[Abstract](#)

[Contents](#)



[Home Page](#)

[Go Back](#)

[Close](#)

[Quit](#)



Abstract

In the present paper, we study a certain integral modification of the well known Baskakov operators with the weight function of Beta basis function. We establish pointwise convergence, an asymptotic formula an error estimation and an inverse result in simultaneous approximation for these new operators.

2000 Mathematics Subject Classification: 41A30, 41A36.

Key words: Baskakov operators, Simultaneous approximation, Asymptotic formula, Pointwise convergence, Error estimation, Inverse theorem.

The work carried out when the second author visited Department of Mathematics and Statistics, Auburn University, USA in fall 2005.

The authors are thankful to the referee for making many valuable suggestions, leading to the better presentation of the paper.

On Simultaneous Approximation For Certain Baskakov Durrmeyer Type Operators

Vijay Gupta,
Muhammad Aslam Noor,
Man Singh Beniwal and
M. K. Gupta

Contents

1	Introduction	3
2	Basic Results	5
3	Direct Theorems	13
4	Inverse Theorem	27
References		

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

1. Introduction

For

$$f \in C_\gamma[0, \infty) \equiv \{f \in C[0, \infty) : |f(t)| \leq Mt^\gamma\}$$

for some $M > 0, \gamma > 0\}$ we consider a certain type of Baskakov-Durrmeyer operator as

$$\begin{aligned} (1.1) \quad B_n(f(t), x) &= \sum_{k=1}^{\infty} p_{n,k}(x) \int_0^{\infty} b_{n,k}(t) f(t) dt + (1+x)^{-n} f(0) \\ &= \int_0^{\infty} W_n(x, t) f(t) dt \end{aligned}$$

where


$$\begin{aligned} p_{n,k}(x) &= \binom{n+k-1}{k} \frac{x^k}{(1+x)^{n+k}}, \\ b_{n,k}(t) &= \frac{1}{B(n+1, k)} \cdot \frac{t^{k-1}}{(1+t)^{n+k+1}} \end{aligned}$$

and

$$W_n(x, t) = \sum_{k=1}^{\infty} p_{n,k}(x) b_{n,k}(t) + (1+x)^{-n} \delta(t),$$

$\delta(t)$ being the Dirac delta function. The norm- $\|\cdot\|_\gamma$ on the class $C_\gamma[0, \infty)$ is defined as $\|f\|_\gamma = \sup_{0 \leq t < \infty} |f(t)|t^{-\gamma}$.

The operators defined by (1.1) are the integral modification of the well known Baskakov operators with weight functions of some Beta basis functions. Very



On Simultaneous
Approximation For Certain
Baskakov Durrmeyer Type
Operators

Vijay Gupta,
Muhammad Aslam Noor,
Man Singh Beniwal and
M. K. Gupta

Title Page

Contents



Go Back

Close

Quit

Page 3 of 33

recently Finta [2] also studied some other approximation properties of these operators. The behavior of these operators is very similar to the operators recently introduced in [6], [9] and also studied in [8]. These operators reproduce not only the constant functions but also the linear functions, which is the interesting property of such operators. The other usual Durrmeyer type integral modification of the Baskakov operators [5] reproduce only the constant functions, so one can not apply the iterative combinations easily to improve the order of approximation for the usual Baskakov Durrmeyer operators. For recent work in this area we refer to [7]. In the present paper we study some direct results which include pointwise convergence, asymptotic formula, error estimation and inverse theorem in the simultaneous approximation for the unbounded functions of growth of order t^γ .



On Simultaneous
Approximation For Certain
Baskakov Durrmeyer Type
Operators

Vijay Gupta,
Muhammad Aslam Noor,
Man Singh Beniwal and
M. K. Gupta

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

2. Basic Results

In this section we mention certain lemmas which will be used in the sequel.

Lemma 2.1 ([3]). For $m \in N \cup \{0\}$, if the m^{th} order moment be defined as

$$U_{n,m}(x) = \sum_{k=0}^{\infty} p_{n,k}(x) \left(\frac{k}{n} - x \right)^m,$$

then $U_{n,0}(x) = 1$, $U_{n,1}(x) = 0$ and

$$nU_{n,m+1}(x) = x(1+x)(U_{n,m}^{(1)}(x) + mU_{n,m-1}(x)).$$


Consequently we have $U_{n,m}(x) = O(n^{-[(m+1)/2]})$.

Lemma 2.2. Let the function $T_{n,m}(x)$, $m \in N \cup \{0\}$, be defined as

$$\begin{aligned} T_{n,m}(x) &= B_n((t-x)^m x) \\ &= \sum_{k=1}^{\infty} p_{n,k}(x) \int_0^{\infty} b_{n,k}(t)(t-x)^m dt + (1+x)^{-n}(-x)^m. \end{aligned}$$

Then $T_{n,0}(x) = 1$, $T_{n,1} = 0$, $T_{n,2}(x) = \frac{2x(1+x)}{n-1}$ and also there holds the recurrence relation

$$(n-m)T_{n,m+1}(x) = x(1+x) [T_{n,m}^{(1)}(x) + 2mT_{n,m-1}(x)] + m(1+2x)T_{n,m}(x).$$



On Simultaneous
Approximation For Certain
Baskakov Durrmeyer Type
Operators

Vijay Gupta,
Muhammad Aslam Noor,
Man Singh Beniwal and
M. K. Gupta

Title Page

Contents



Go Back

Close

Quit

Proof. By definition, we have

$$\begin{aligned}
 T_{n,m}^{(1)}(x) &= \sum_{k=1}^{\infty} p_{n,k}^{(1)}(x) \int_0^{\infty} b_{n,k}(t)(t-x)^m dt \\
 &\quad - m \sum_{k=1}^{\infty} p_{n,k}(x) \int_0^{\infty} b_{n,k}(t)(t-x)^{m-1} dt \\
 &\quad - n(1+x)^{-n-1}(-x)^m - m(1+x)^{-n}(-x)^{m-1}.
 \end{aligned}$$

Using the identities


$$x(1+x)p_{n,k}^{(1)}(x) = (k-nx)p_{n,k}(x)$$

and

$$t(1+t)b_{n,k}^{(1)}(t) = [(k-1)-(n+2)t]b_{n,k}(t),$$

we have

$$\begin{aligned}
 &x(1+x) [T_{n,m}^{(1)}(x) + mT_{n,m-1}(x)] \\
 &= \sum_{k=1}^{\infty} p_{n,k}(x) \int_0^{\infty} (k-nx)b_{n,k}(t)(t-x)^m dt + n(1+x)^{-n}(-x)^{m+1} \\
 &= \sum_{k=1}^{\infty} p_{n,k}(x) \int_0^{\infty} [(k-1)-(n+2)t+(n+2)(t-x) \\
 &\quad + (1+2x)]b_{n,k}(t)(t-x)^m dt + n(1+x)^{-n}(-x)^{m+1}
 \end{aligned}$$



On Simultaneous Approximation For Certain Baskakov Durrmeyer Type Operators

*Vijay Gupta,
Muhammad Aslam Noor,
Man Singh Beniwal and
M. K. Gupta*

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

$$\begin{aligned}
&= \sum_{k=1}^{\infty} p_{n,k}(x) \int_0^{\infty} t(1+t)b_{n,k}^{(1)}(t)(t-x)^m dt \\
&\quad + (n+2)[T_{n,m+1}(x) - (1+x)^{-n}(-x)^{m+1}] \\
&\quad + (1+2x)[T_{n,m}(x) - (1+x)^{-n}(-x)^m] + n(1+x)^{-n}(-x)^{m+1} \\
&= -(m+1)(1+2x)[T_{n,m}(x) - (1+x)^{-n}(-x)^m] \\
&\quad - (m+2)[T_{n,m+1} - (1+x)^{-n}(-x)^{m+1}] \\
&\quad - mx(1+x)[T_{n,m-1}(x) - (1+x)^{-n}(-x)^{m-1}] \\
&\quad + (n+2)[T_{n,m+1} - (1+x)^{-n}(-x)^{m+1}] \\
&\quad + (1+2x)[T_{n,m}(x) - (1+x)^{-n}(-x)^m] + n(1+x)^{-n}(-x)^{m+1}.
\end{aligned}$$

Thus, we get

$$(n-m)T_{n,m+1}(x) = x(1+x)[T_{n,m}^{(1)}(x) + 2mT_{n,m-1}(x)] + m(1+2x)T_{n,m}(x).$$

This completes the proof of recurrence relation. From the above recurrence relation, it is easily verified for all $x \in [0, \infty)$ that

$$T_{n,m}(x) = O(n^{-[(m+1)/2]}).$$

□

Remark 1. It is easily verified from Lemma 2.1 that for each $x \in (0, \infty)$

$$B_n(t^i, x) = \frac{(n+i-1)!(n-i)!}{n!(n-1)!} x^i + i(i-1) \frac{(n+i-2)!(n-i)!}{n!(n-1)!} x^{i-1} + O(n^{-2}).$$



On Simultaneous Approximation For Certain Baskakov Durrmeyer Type Operators

*Vijay Gupta,
Muhammad Aslam Noor,
Man Singh Beniwal and
M. K. Gupta*

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

Corollary 2.3. Let δ be a positive number. Then for every $\gamma > 0, x \in (0, \infty)$, there exists a constant $M(s, x)$ independent of n and depending on s and x such that

$$\left\| \int_{|t-x|>\delta} W_n(x, t) t^\gamma dt \right\|_{C[a, b]} \leq M(s, x) n^{-s}, \quad s = 1, 2, 3, \dots$$

Lemma 2.4. There exist the polynomials $Q_{i,j,r}(x)$ independent of n and k such that


$$\{x(1+x)\}^r D^r [p_{n,k}(x)] = \sum_{\substack{2i+j \leq r \\ i, j \geq 0}} n^i (k - nx)^j Q_{i,j,r}(x) p_{n,k}(x),$$

where $D \equiv \frac{d}{dx}$.

By C_0 , we denote the class of continuous functions on the interval $(0, \infty)$ having a compact support and C_0^r is the class of r times continuously differentiable functions with $C_0^r \subset C_0$. The function f is said to belong to the generalized Zygmund class $Liz(\alpha, 1, a, b)$, if there exists a constant M such that $\omega_2(f, \delta) \leq M\delta^\alpha$, $\delta > 0$, where $\omega_2(f, \delta)$ denotes the modulus of continuity of 2nd order on the interval $[a, b]$. The class $Liz(\alpha, 1, a, b)$ is more commonly denoted by $Lip^*(\alpha, a, b)$. Suppose $G^{(r)} = \{g : g \in C_0^{r+2}, \text{ supp } g \subset [a', b'] \text{ where } [a', b'] \subset (a, b)\}$. For r times continuously differentiable functions f with $\text{supp } f \subset [a', b']$ the Peetre's K-functionals are defined as

$$K_r(\xi, f) = \inf_{g \in G^{(r)}} \left[\|f^{(r)} - g^{(r)}\|_{C[a', b']} + \xi \left\{ \|g^{(r)}\|_{C[a', b']} + \|g^{(r+2)}\|_{C[a', b']}\right\} \right],$$

$0 < \xi \leq 1.$



On Simultaneous
Approximation For Certain
Baskakov Durrmeyer Type
Operators

Vijay Gupta,
Muhammad Aslam Noor,
Man Singh Beniwal and
M. K. Gupta

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

For $0 < \alpha < 2$, $C_0^r(\alpha, 1, a, b)$ denotes the set of functions for which

$$\sup_{0 < \xi \leq 1} \xi^{-\alpha/2} K_r(\xi, f, a, b) < C.$$

Lemma 2.5. Let $0 < a' < a'' < b'' < b' < b < \infty$ and $f^{(r)} \in C_0$ with $\text{supp } f \subset [a'', b'']$ and if $f \in C_0^r(\alpha, 1, a', b')$, we have $f^{(r)} \in \text{Liz}(\alpha, 1, a', b')$ i.e. $f^{(r)} \in \text{Lip}^*(\alpha, a', b')$ where $\text{Lip}^*(\alpha, a', b')$ denotes the Zygmund class satisfying $K_r(\delta, f) \leq C\delta^{\alpha/2}$.

Proof. Let $g \in G^{(r)}$, then for $f \in C_0^r(\alpha, 1, a', b')$, we have

$$\begin{aligned} |\Delta_\delta^2 f^{(r)}(x)| &\leq |\Delta_\delta^2(f^{(r)} - g^{(r)})(x)| + |\Delta_\delta^2 g^{(r)}(x)| \\ &\leq \|\Delta_\delta^2(f^{(r)} - g^{(r)})\|_{C[a', b']} + \delta^2 \|g^{(r+2)}\|_{C[a', b']} \\ &\leq 4M_1 K_r(\delta^2, f) \leq M_2 \delta^\alpha. \end{aligned}$$

□

Lemma 2.6. If f is r times differentiable on $[0, \infty)$, such that $f^{(r-1)} = O(t^\alpha)$, $\alpha > 0$ as $t \rightarrow \infty$, then for $r = 1, 2, 3, \dots$ and $n > \alpha + r$ we have

$$B_n^{(r)}(f, x) = \frac{(n+r-1)!(n-r)!}{n!(n-1)!} \sum_{k=0}^{\infty} p_{n+r,k}(x) \int_0^\infty b_{n-r,k+r}(t) f^{(r)}(t) dt.$$

Proof. First

$$B_n^{(1)}(f, x) = \sum_{k=1}^{\infty} p_{n,k}^{(1)}(x) \int_0^\infty b_{n,k}(t) f(t) dt - n(1+x)^{-n-1} f(0).$$



**On Simultaneous
Approximation For Certain
Baskakov Durrmeyer Type
Operators**

Vijay Gupta,
Muhammad Aslam Noor,
Man Singh Beniwal and
M. K. Gupta

Title Page

Contents



Go Back

Close

Quit

Page 9 of 33

Now using the identities

$$(2.1) \quad p_{n,k}^{(1)}(x) = n[p_{n+1,k-1}(x) - p_{n+1,k}(x)],$$


$$(2.2) \quad b_{n,k}^{(1)}(t) = (n+1)[b_{n+1,k-1}(t) - b_{n+1,k}(t)].$$

for $k \geq 1$, we have

$$\begin{aligned} & B_n^{(1)}(f, x) \\ &= \sum_{k=1}^{\infty} n[p_{n+1,k-1}(x) - p_{n+1,k}(x)] \int_0^{\infty} b_{n,k}(t)f(t)dt - n(1+x)^{-n-1}f(0) \\ &= np_{n+1,0}(x) \int_0^{\infty} b_{n,1}(t)f(t)dt - n(1+x)^{-n-1}f(0) \\ &\quad + n \sum_{k=1}^{\infty} p_{n+1,k}(x) \int_0^{\infty} [b_{n,k+1}(t) - b_{n,k}(t)]f(t)dt \\ &= n(1+x)^{-n-1} \int_0^{\infty} (n+1)(1+t)^{-n-2}f(t)dt - n(1+x)^{-n-1}f(0) \\ &\quad + n \sum_{k=1}^{\infty} p_{n+1,k}(x) \int_0^{\infty} -\frac{1}{n}b_{n-1,k+1}^{(1)}(t)f(t)dt. \end{aligned}$$

Integrating by parts, we get

$$B_n^{(1)}(f, x) = n(1+x)^{-n-1}f(0) + n(1+x)^{-n-1} \int_0^{\infty} (1+t)^{-n-1}f^{(1)}(t)dt$$



On Simultaneous Approximation For Certain Baskakov Durrmeyer Type Operators

*Vijay Gupta,
Muhammad Aslam Noor,
Man Singh Beniwal and
M. K. Gupta*

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)


$$\begin{aligned}
& - n(1+x)^{-n-1}f(0) + \sum_{k=1}^{\infty} p_{n+1,k}(x) \int_0^{\infty} b_{n-1,k+1}(t)f^{(1)}(t)dt \\
& = \sum_{k=0}^{\infty} p_{n+1,k}(x) \int_0^{\infty} b_{n-1,k+1}(t)f^{(1)}(t)dt.
\end{aligned}$$

Thus the result is true for $r = 1$. We prove the result by induction method.
Suppose that the result is true for $r = i$, then

$$B_n^{(i)}(f, x) = \frac{(n+i-1)!(n-i)!}{n!(n-1)!} \sum_{k=0}^{\infty} p_{n+i,k}(x) \int_0^{\infty} b_{n-i,k+i}(t)f^{(i)}(t)dt.$$

Thus using the identities (2.1) and (2.2), we have

$$\begin{aligned}
& B_n^{(i+1)}(f, x) \\
& = \frac{(n+i-1)!(n-i)!}{n!(n-1)!} \\
& \quad \times \sum_{k=1}^{\infty} (n+i)[p_{n+i+1,k-1}(x) - p_{n+i+1,k}(x)] \int_0^{\infty} b_{n-i,k+i}(t)f^{(i)}(t)dt \\
& \quad + \frac{(n+i-1)!(n-i)!}{n!(n-1)!}(-(n+i)(1+x)^{-n-i-1}) \int_0^{\infty} b_{n-i,i}(t)f^{(i)}(t)dt \\
& = \frac{(n+i)!(n-i)!}{n!(n-1)!} p_{n+i+1,0}(x) \int_0^{\infty} b_{n-i,i+1}(t)f^{(i)}(t)dt \\
& \quad - \frac{(n+i)!(n-i)!}{n!(n-1)!} p_{n+i+1,0}(x) \int_0^{\infty} b_{n-i,i}(t)f^{(i)}(t)dt
\end{aligned}$$



On Simultaneous Approximation For Certain Baskakov Durrmeyer Type Operators

Vijay Gupta,
Muhammad Aslam Noor,
Man Singh Beniwal and
M. K. Gupta

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)


[Quit](#)

$$\begin{aligned}
& + \frac{(n+i)!(n-i)!}{n!(n-1)!} \sum_{k=1}^{\infty} p_{n+i+1,k}(x) \int_0^{\infty} [b_{n-i,k+i+1}(t) - b_{n-i,k+i}(t)] f^{(i)}(t) dt \\
& = \frac{(n+i)!(n-i)!}{n!(n-1)!} p_{n+i+1,0}(x) \int_0^{\infty} -\frac{1}{(n-i)} b_{n-i-1,i+1}^{(1)}(t) f^{(i)}(t) dt \\
& \quad + \frac{(n+i)!(n-i)!}{n!(n-1)!} \sum_{k=1}^{\infty} p_{n+i+1,k}(x) \int_0^{\infty} -\frac{1}{(n-i)} b_{n-i-1,k+i+1}^{(1)}(t) f^{(i)}(t) dt.
\end{aligned}$$

Integrating by parts, we obtain

$$B_n^{(i+1)}(f, x) = \frac{(n+i)!(n-i-1)!}{n!(n-1)!} \sum_{k=0}^{\infty} p_{n+i+1,k}(x) \int_0^{\infty} b_{n-i-1,k+i+1}(t) f^{(i+1)}(t) dt.$$

This completes the proof of the lemma. \square



On Simultaneous Approximation For Certain Baskakov Durrmeyer Type Operators

Vijay Gupta,
Muhammad Aslam Noor,
Man Singh Beniwal and
M. K. Gupta

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

3. Direct Theorems

In this section we present the following results.

Theorem 3.1. Let $f \in C_\gamma[0, \infty)$ and $f^{(r)}$ exists at a point $x \in (0, \infty)$. Then we have

$$B_n^{(r)}(f, x) = f^{(r)}(x)$$

as $n \rightarrow \infty$.


Proof. By Taylor expansion of f , we have

$$f(t) = \sum_{i=0}^r \frac{f^{(i)}(x)}{i!}(t-x)^i + \varepsilon(t, x)(t-x)^r,$$

where $\varepsilon(t, x) \rightarrow 0$ as $t \rightarrow x$. Hence

$$\begin{aligned} B_n^{(r)}(f, x) &= \int_0^\infty W_n^{(r)}(t, x)f(t)dt \\ &= \sum_{i=0}^r \frac{f^{(i)}(x)}{i!} \int_0^\infty W_n^{(r)}(t, x)(t-x)^i dt \\ &\quad + \int_0^\infty W_n^{(r)}(t, x)\varepsilon(t, x)(t-x)^r dt \\ &=: R_1 + R_2. \end{aligned}$$

First to estimate R_1 , using the binomial expansion of $(t-x)^i$ and Remark 1, we



On Simultaneous
Approximation For Certain
Baskakov Durrmeyer Type
Operators

Vijay Gupta,
Muhammad Aslam Noor,
Man Singh Beniwal and
M. K. Gupta

Title Page

Contents



Go Back

Close

Quit

Page 13 of 33

have

$$\begin{aligned}
 R_1 &= \sum_{i=0}^r \frac{f^{(i)}(x)}{i!} \sum_{v=0}^i \binom{i}{v} (-x)^{i-v} \frac{\partial^r}{\partial x^r} \int_0^\infty W_n(t, x) t^v dt \\
 &= \frac{f^{(r)}(x)}{r!} \frac{d^r}{dx^r} \left[\frac{(n+r-1)!(n-r)!}{n!(n-1)!} x^r + \text{terms containing lower powers of } x \right] \\
 &= f^{(r)}(x) \left[\frac{(n+r-1)!(n-r)!}{n!(n-1)!} \right] \rightarrow f^{(r)}(x)
 \end{aligned}$$


as $n \rightarrow \infty$. Next applying Lemma 2.4, we obtain

$$R_2 = \int_0^\infty W_n^{(r)}(t, x) \varepsilon(t, x) (t-x)^r dt,$$

$$|R_2| \leq \sum_{\substack{2i+j \leq r \\ i,j \geq 0}} n^i \frac{|Q_{i,j,r}(x)|}{\{x(1+x)\}^r} \sum_{k=1}^\infty |k-nx|^j p_{n,k}(x)$$

$$\int_0^\infty b_{n,k}(t) |\varepsilon(t, x)| |t-x|^r dt + \frac{(n+r+1)!}{(n-1)!} (1+x)^{-n-r} |\varepsilon(0, x)| x^r.$$

The second term in the above expression tends to zero as $n \rightarrow \infty$. Since $\varepsilon(t, x) \rightarrow 0$ as $t \rightarrow x$ for a given $\varepsilon > 0$ there exists a δ such that $|\varepsilon(t, x)| < \varepsilon$ whenever $0 < |t-x| < \delta$. If $\alpha \geq \max\{\gamma, r\}$, where α is any integer, then we can find a constant $M_3 > 0$, $|\varepsilon(t, x)(t-x)^\alpha| \leq M_3 |t-x|^\alpha$, for $|t-x| \geq \delta$.



On Simultaneous Approximation For Certain Baskakov Durrmeyer Type Operators

Vijay Gupta,
Muhammad Aslam Noor,
Man Singh Beniwal and
M. K. Gupta

[Title Page](#)

[Contents](#)

[◀◀](#) [▶▶](#)

[◀](#) [▶](#)

[Go Back](#)

[Close](#)

[Quit](#)

Therefore

$$\begin{aligned}
 |R_2| &\leq M_3 \sum_{\substack{2i+j \leq r \\ i,j \geq 0}} n^i \sum_{k=0}^{\infty} p_{n,k}(x) |k - nx|^j \\
 &\quad \times \left\{ \varepsilon \int_{|t-x|<\delta} b_{n,k}(t) |t-x|^r dt + \int_{|t-x|\geq\delta} b_{n,k}(t) |t-x|^\alpha dt \right\} \\
 &=: R_3 + R_4.
 \end{aligned}$$

Applying the Cauchy-Schwarz inequality for integration and summation respectively, we obtain

$$\begin{aligned}
 R_3 &\leq \varepsilon M_3 \sum_{\substack{2i+j \leq r \\ i,j \geq 0}} n^i \left\{ \sum_{k=1}^{\infty} p_{n,k}(x) (k - nx)^{2j} \right\}^{\frac{1}{2}} \\
 &\quad \times \left\{ \sum_{k=1}^{\infty} p_{n,k}(x) \int_0^{\infty} b_{n,k}(t) (t - x)^{2r} dt \right\}^{\frac{1}{2}}.
 \end{aligned}$$

Using Lemma 2.1 and Lemma 2.2, we get

$$R_3 = \varepsilon \cdot O(n^{r/2})O(n^{-r/2}) = \varepsilon \cdot o(1).$$

Again using the Cauchy-Schwarz inequality, Lemma 2.1 and Corollary 2.3, we



On Simultaneous Approximation For Certain Baskakov Durrmeyer Type Operators

Vijay Gupta,
 Muhammad Aslam Noor,
 Man Singh Beniwal and
 M. K. Gupta

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

get

$$\begin{aligned}
R_4 &\leq M_4 \sum_{\substack{2i+j \leq r \\ i,j \geq 0}} n^i \sum_{k=1}^{\infty} p_{n,k}(x) |k - nx|^j \int_{|t-x| \geq \delta} b_{n,k}(t) |t - x|^{\alpha} dt \\
&\leq M_4 \sum_{\substack{2i+j \leq r \\ i,j \geq 0}} n^i \sum_{k=1}^{\infty} p_{n,k}(x) |k - nx|^j \left\{ \int_{|t-x| \geq \delta} b_{n,k}(t) dt \right\}^{\frac{1}{2}} \\
&\quad \times \left\{ \int_{|t-x| \geq \delta} b_{n,k}(t) (t-x)^{2\alpha} dt \right\}^{\frac{1}{2}} \\
&\leq M_4 \sum_{\substack{2i+j \leq r \\ i,j \geq 0}} n^i \left\{ \sum_{k=1}^{\infty} p_{n,k}(x) (k-nx)^{2j} \right\}^{\frac{1}{2}} \\
&\quad \times \left\{ \sum_{k=1}^{\infty} p_{n,k}(x) \int_0^{\infty} b_{n,k}(t) (t-x)^{2\alpha} dt \right\}^{\frac{1}{2}} \\
&= \sum_{\substack{2i+j \leq r \\ i,j \geq 0}} n^i O(n^{j/2}) O(n^{-\alpha/2}) = O(n^{(r-\alpha)/2}) = o(1).
\end{aligned}$$

Collecting the estimates of $R_1 - R_4$, we obtain the required result. \square

Theorem 3.2. Let $f \in C_{\gamma}[0, \infty)$. If $f^{(r+2)}$ exists at a point $x \in (0, \infty)$. Then

$$\begin{aligned}
\lim_{n \rightarrow \infty} n [B_n^{(r)}(f, x) - f^{(r)}(x)] \\
= r(r-1)f^{(r)}(x) + r(1+2x)f^{(r+1)}(x) + x(1+x)f^{(r+2)}(x).
\end{aligned}$$



On Simultaneous Approximation For Certain Baskakov Durrmeyer Type Operators

Vijay Gupta,
Muhammad Aslam Noor,
Man Singh Beniwal and
M. K. Gupta

Title Page

Contents



Go Back

Close

Quit

Proof. Using Taylor's expansion of f , we have

$$f(t) = \sum_{i=0}^{r+2} \frac{f^{(i)}(x)}{i!} (t-x)^i + \varepsilon(t, x)(t-x)^{r+2},$$

where $\varepsilon(t, x) \rightarrow 0$ as $t \rightarrow x$ and $\varepsilon(t, x) = O((t-x)^\gamma)$, $t \rightarrow \infty$ for $\gamma > 0$.

Applying Lemma 2.2, we have

$$\begin{aligned} n [B_n^{(r)}(f(t), x) - f^{(r)}(x)] \\ = n \left[\sum_{i=0}^{r+2} \frac{f^{(i)}(x)}{i!} \int_0^\infty W_n^{(r)}(t, x)(t-x)^i dt - f^{(r)}(x) \right] \\ + n \int_0^\infty W_n^{(r)}(t, x)\varepsilon(t, x)(t-x)^{r+2} dt \\ =: E_1 + E_2. \end{aligned}$$

First, we have

$$\begin{aligned} E_1 &= n \sum_{i=0}^{r+2} \frac{f^{(i)}(x)}{i!} \sum_{j=0}^i \binom{i}{j} (-x)^{i-j} \int_0^\infty W_n^{(r)}(t, x)t^j dt - nf^{(r)}(x) \\ &= \frac{f^{(r)}(x)}{r!} n [B_n^{(r)}(t^r, x) - (r!)] \\ &\quad + \frac{f^{(r+1)}(x)}{(r+1)!} n [(r+1)(-x)B_n^{(r)}(t^r, x) + B_n^{(r)}(t^{r+1}, x)] \\ &\quad + \frac{f^{(r+2)}(x)}{(r+2)!} n \left[\frac{(r+2)(r+1)}{2} x^2 B_n^{(r)}(t^r, x) \right. \end{aligned}$$



On Simultaneous Approximation For Certain Baskakov Durrmeyer Type Operators

*Vijay Gupta,
Muhammad Aslam Noor,
Man Singh Beniwal and
M. K. Gupta*

[Title Page](#)


[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)



Therefore, by Remark 1, we have

$$\begin{aligned}
 E_1 &= nf^{(r)}(x) \left[\frac{(n+r-1)!(n-r)!}{n!(n-1)!} - 1 \right] \\
 &\quad + \frac{nf^{(r+1)}(x)}{(r+1)!} \left[(x-1)(-x) \left\{ \frac{(n+r-1)!(n-r)!}{n!(n-1)!} \right\} \right. \\
 &\quad + \left\{ \frac{(n+r)!(n-r-1)!}{n!(n-1)!} (r+1)!x + (r+1)r \frac{(n+r-1)!(n-r-1)!}{n!(n-1)!} r! \right\} \\
 &\quad + \frac{nf^{(r+2)}(x)}{(r+2)!} \left[\frac{(r+2)(r+1)}{2} x^2 \frac{(n+r-1)!(n-r)!}{n!(n-1)!} r! \right] \\
 &\quad + (r+2)(-x) \left\{ \frac{(n+r)!(n-r-1)!}{2} x(r+1)! \right. \\
 &\quad \left. + (r+1)r \frac{(n-r-1)!(n-r-1)!}{n!(n-1)!} r! \right\} \\
 &\quad + \left\{ \frac{(n+r+1)!(n-r-2)!}{n!(n-1)!} \frac{(r+2)!}{2} x^2 \right. \\
 &\quad \left. + (r+2)(r+1) \frac{(n+r)!(n-r-2)!}{n!(n-1)!} (r+1)!x \right\} \Big] + O(n^{-2}).
 \end{aligned}$$

In order to complete the proof of the theorem it is sufficient to show that $E_2 \rightarrow 0$ as $n \rightarrow \infty$ which easily follows proceeding along the lines of the proof of Theorem 3.1 and by using Lemma 2.1, Lemma 2.2 and Lemma 2.4. \square

On Simultaneous Approximation For Certain Baskakov Durrmeyer Type Operators

Vijay Gupta,
 Muhammad Aslam Noor,
 Man Singh Beniwal and
 M. K. Gupta

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

Lemma 3.3. Let $0 < \alpha < 2$, $0 < a < a' < a'' < b'' < b' < b < \infty$. If $f \in C_0$ with $\text{supp } f \subset [a'', b'']$ and $\left\| B_n^{(r)}(f, \cdot) - f^{(r)} \right\|_{C[a,b]} = O(n^{-\alpha/2})$, then

$$K_r(\xi, f) = M_5 \left\{ n^{-\alpha/2} + n\xi K_r(n^{-1}, f) \right\}.$$

Consequently $K_r(\xi, f) \leq M_6 \xi^{\alpha/2}$, $M_6 > 0$.

Proof. It is sufficient to prove

$$K_r(\xi, f) = M_7 \left\{ n^{-\alpha/2} + n\xi K_r(n^{-1}, f) \right\},$$

for sufficiently large n . Because $\text{supp } f \subset [a'', b'']$ therefore by Theorem 3.2 there exists a function $h^{(i)} \in G^{(r)}$, $i = r, r+2$, such that


$$\left\| B_n^{(r)}(f, \bullet) - h^{(i)} \right\|_{C[a', b']} \leq M_8 n^{-1}.$$

Therefore,

$$\begin{aligned} K_r(\xi, f) &\leq 3M_9 n^{-1} + \left\| B_n^{(r)}(f, \bullet) - f^{(r)} \right\|_{C[a', b']} \\ &\quad + \xi \left\{ \left\| B_n^{(r)}(f, \bullet) \right\|_{C[a', b']} + \left\| B_n^{(r+2)}(f, \bullet) \right\|_{C[a', b']} \right\}. \end{aligned}$$

Next, it is sufficient to show that there exists a constant M_{10} such that for each $g \in G^{(r)}$

$$(3.1) \quad \left\| B_n^{(r+2)}(f, \bullet) \right\|_{C[a', b']} \leq M_{10} n \left\{ \left\| f^{(r)} - g^{(r)} \right\|_{C[a', b']} + n^{-1} \left\| g^{(r+2)} \right\|_{C[a', b']} \right\}.$$



**On Simultaneous
Approximation For Certain
Baskakov Durrmeyer Type
Operators**

Vijay Gupta,
Muhammad Aslam Noor,
Man Singh Beniwal and
M. K. Gupta

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

Also using the linearity property, we have

$$(3.2) \quad \begin{aligned} \|B_n^{(r+2)}(f, \bullet)\|_{C[a', b']} &\leq \|B_n^{(r+2)}(f - g, \bullet)\|_{C[a', b']} + \|B_n^{(r+2)}(g, \bullet)\|_{C[a', b']} . \end{aligned}$$

Applying Lemma 2.4, we get

$$\begin{aligned} \int_0^\infty \left| \frac{\partial^{r+2}}{\partial x^{r+2}} W_n(x, t) \right| dt &\leq \sum_{\substack{2i+j \leq r+2 \\ i, j \geq 0}} \sum_{k=1}^{\infty} n^i |k - nx|^j \frac{|Q_{i,j,r+2}(x)|}{\{x(1+x)\}^{r+2}} \\ &\quad \times p_{n,k}(x) \int_0^\infty b_{n,k}(t) dt + \frac{d^{r+2}}{dx^{r+2}} [(1+x)^{-n}] . \end{aligned}$$

Therefore by the Cauchy-Schwarz inequality and Lemma 2.1, we obtain

$$(3.3) \quad \|B_n^{(r)}(f - g, \bullet)\|_{C[a', b']} \leq M_{11} n \|f^{(r)} - g^{(r)}\|_{C[a', b']} ,$$

where the constant M_{11} is independent of f and g . Next by Taylor's expansion, we have

$$g(t) = \sum_{i=0}^{r+1} \frac{g^{(i)}(x)}{i!} (t-x)^i + \frac{g^{(r+2)}(\xi)}{(r+2)!} (t-x)^{r+2},$$

where ξ lies between t and x . Using the above expansion and the fact that $\int_0^\infty \frac{\partial^m}{\partial x^m} W_n(x, t) (t-x)^i dt = 0$ for $m > i$, we get

$$(3.4) \quad \begin{aligned} \|B_n^{(r+2)}(g, \bullet)\|_{C[a', b']} &\leq M_{12} \|g^{(r+2)}\|_{C[a', b']} \cdot \left\| \int_0^\infty \frac{\partial^{r+2}}{\partial x^{r+2}} W_n(x, t) (t-x)^{r+2} dt \right\|_{C[a', b']} . \end{aligned}$$



On Simultaneous Approximation For Certain Baskakov Durrmeyer Type Operators

*Vijay Gupta,
Muhammad Aslam Noor,
Man Singh Beniwal and
M. K. Gupta*

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

Also by Lemma 2.4 and the Cauchy-Schwarz inequality, we have

$$\begin{aligned}
E &\equiv \int_0^\infty \left| \frac{\partial^{r+2}}{\partial x^{r+2}} W_n(x, t) \right| (t-x)^{r+2} dt \\
&\leq \sum_{\substack{2i+j \leq r+2 \\ i,j \geq 0}} \sum_{k=1}^{\infty} n^i p_{n,k}(x) |k-nx|^j \frac{|Q_{i,j,r+2}(x)|}{\{x(1+x)\}^{r+2}} \int_0^\infty b_{n,k}(t)(t-x)^{r+2} dt \\
&\quad + \frac{d^{r+2}}{dx^{r+2}} [(-x)^{r+2}(1+x)^{-n}] \\
&\leq \sum_{\substack{2i+j \leq r+2 \\ i,j \geq 0}} \frac{|Q_{i,j,r+2}(x)|}{\{x(1+x)\}^{r+2}} \left(\sum_{k=1}^{\infty} p_{n,k}(x)(k-nx)^{2j} \right)^{\frac{1}{2}} \\
&\quad \times \left(\sum_{k=1}^{\infty} p_{n,k}(x) \int_0^\infty b_{n,k}(t)(t-x)^{2r+4} dt \right)^{\frac{1}{2}} \left(\int_0^\infty b_{n,k}(t) dt \right)^{\frac{1}{2}} \\
&\quad + \frac{d^{r+2}}{dx^{r+2}} [(-x)^{r+2}(1+x)^{-n}] \\
&= \sum_{\substack{2i+j \leq r+2 \\ i,j \geq 0}} n^i \frac{|Q_{i,j,r+2}(x)|}{\{x(1+x)\}^{r+2}} O(n^{j/2}) O\left(n^{-\left(1+\frac{r}{2}\right)}\right).
\end{aligned}$$

Hence

$$(3.5) \quad \|B_n^{(r+2)}(g, \bullet)\|_{C[a', b']} \leq M_{13} \|g^{(r+2)}\|_{C[a', b']}.$$

Combining the estimates of (3.2)-(3.5), we get (3.1). The other consequence follows from [1]. This completes the proof of the lemma. \square



On Simultaneous Approximation For Certain Baskakov Durrmeyer Type Operators

*Vijay Gupta,
Muhammad Aslam Noor,
Man Singh Beniwal and
M. K. Gupta*

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

Theorem 3.4. Let $f \in C_\gamma[0, \infty)$ and suppose $0 < a < a_1 < b_1 < b < \infty$. Then for all n sufficiently large, we have

$$\|B_n^{(r)}(f, \bullet) - f^{(r)}\|_{C[a_1, b_1]} \leq \max \left\{ M_{14} \omega_2 \left(f^{(r)}, n^{-\frac{1}{2}}, a, b \right) + M_{15} n^{-1} \|f\|_\gamma \right\},$$

where $M_{14} = M_{14}(r)$, $M_{15} = M_{15}(r, f)$.

Proof. For sufficiently small $\delta > 0$, we define a function $f_{2,\delta}(t)$ corresponding to $f \in C_\gamma[0, \infty)$ by

$$f_{2,\delta}(t) = \delta^{-2} \int_{-\frac{\delta}{2}}^{\frac{\delta}{2}} \int_{-\frac{\delta}{2}}^{\frac{\delta}{2}} (f(t) - \Delta_\eta^2 f(t)) dt_1 dt_2,$$

where $\eta = \frac{t_1 + t_2}{2}$, $t \in [a, b]$ and $\Delta_\eta^2 f(t)$ is the second forward difference of f with step length η . Following [4] it is easily checked that:

- (i) $f_{2,\delta}$ has continuous derivatives up to order $2k$ on $[a, b]$,
- (ii) $\|f_{2,\delta}^{(r)}\|_{C[a_1, b_1]} \leq \widehat{M}_1 \delta^{-r} \omega_2(f, \delta, a, b)$,
- (iii) $\|f - f_{2,\delta}\|_{C[a_1, b_1]} \leq \widehat{M}_2 \omega_2(f, \delta, a, b)$,
- (iv) $\|f_{2,\delta}\|_{C[a_1, b_1]} \leq \widehat{M}_3 \|f\|_\gamma$,

where \widehat{M}_i , $i = 1, 2, 3$ are certain constants that depend on $[a, b]$ but are independent of f and n [4].



**On Simultaneous
Approximation For Certain
Baskakov Durrmeyer Type
Operators**

Vijay Gupta,
Muhammad Aslam Noor,
Man Singh Beniwal and
M. K. Gupta

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

We can write

$$\begin{aligned} & \|B_n^{(r)}(f, \bullet) - f^{(r)}\|_{C[a_1, b_1]} \\ & \leq \|B_n^{(r)}(f - f_{2,\delta}, \bullet)\|_{C[a_1, b_1]} + \|B_n^{(r)}(f_{2,\delta}, \bullet) - f_{2,\delta}^{(r)}\|_{C[a_1, b_1]} + \|f^{(r)} - f_{2,\delta}^{(r)}\|_{C[a_1, b_1]} \\ & =: H_1 + H_2 + H_3. \end{aligned}$$

Since $f_{2,\delta}^{(r)} = (f^{(r)})_{2,\delta}(t)$, by property (iii) of the function $f_{2,\delta}$, we get

$$H_3 \leq \widehat{M}_4 \omega_2(f^{(r)}, \delta, a, b).$$

Next on an application of Theorem 3.2, it follows that


$$H_2 \leq \widehat{M}_5 n^{-1} \sum_{j=r}^{r+2} \|f_{2,\delta}^{(j)}\|_{C[a,b]}.$$

Using the interpolation property due to Goldberg and Meir [4], for each $j = r, r+1, r+2$, it follows that

$$\|f_{2,\delta}^{(j)}\|_{C[a_1, b_1]} \leq \widehat{M}_6 \left\{ \|f_{2,\delta}\|_{C[a,b]} + \|f_{2,\delta}^{(r+2)}\|_{C[a,b]} \right\}.$$

Therefore by applying properties (iii) and (iv) of the function $f_{2,\delta}$, we obtain

$$H_2 \leq \widehat{M}_7 4 \cdot n^{-1} \{ \|f\|_\gamma + \delta^{-2} \omega_2(f^{(r)}, \delta) \}.$$



On Simultaneous Approximation For Certain Baskakov Durrmeyer Type Operators

Vijay Gupta,
Muhammad Aslam Noor,
Man Singh Beniwal and
M. K. Gupta

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

Finally we shall estimate H_1 , choosing a^*, b^* satisfying the conditions $0 < a < a^* < a_1 < b_1 < b^* < b < \infty$. Suppose $\psi(t)$ denotes the characteristic function of the interval $[a^*, b^*]$, then

$$\begin{aligned} H_1 &\leq \|B_n^{(r)}(\psi(t)(f(t) - f_{2,\delta}(t)), \bullet)\|_{C[a_1, b_1]} \\ &\quad + \|B_n^{(r)}((1 - \psi(t))(f(t) - f_{2,\delta}(t)), \bullet)\|_{C[a_1, b_1]} \\ &=: H_4 + H_5. \end{aligned}$$

Using Lemma 2.6, it is clear that

$$\begin{aligned} &B_n^{(r)}(\psi(t)(f(t) - f_{2,\delta}(t)), x) \\ &= \frac{(n+r-1)!(n-r)!}{n!(n-1)!} \sum_{k=0}^{\infty} p_{n+r,k}(x) \int_0^{\infty} b_{n-r,k+r}(t) \psi(t) (f^{(r)}(t) - f_{2,\delta}^{(r)}(t)) dt. \end{aligned}$$

Hence

$$\|B_n^{(r)}(\psi(t)(f(t) - f_{2,\delta}(t)), \bullet)\|_{C[a_1, b_1]} \leq \widehat{M}_8 \left\| f^{(r)} - f_{2,\delta}^{(r)} \right\|_{C[a^*, b^*]}.$$

Next for $x \in [a_1, b_1]$ and $t \in [0, \infty) \setminus [a^*, b^*]$, we choose a $\delta_1 > 0$ satisfying $|t - x| \geq \delta_1$.

Therefore by Lemma 2.4 and the Cauchy-Schwarz inequality, we have

$$I \equiv B_n^{(r)}((1 - \psi(t))(f(t) - f_{2,\delta}(t), x)$$



**On Simultaneous
Approximation For Certain
Baskakov Durrmeyer Type
Operators**

Vijay Gupta,
Muhammad Aslam Noor,
Man Singh Beniwal and
M. K. Gupta

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

and

$$\begin{aligned}
 |I| &\leq \sum_{\substack{2i+j \leq r \\ i,j \geq 0}} n^i \frac{|Q_{i,j,r}(x)|}{\{x(1+x)\}^r} \sum_{k=1}^{\infty} p_{n,k}(x) |k - nx|^j \\
 &\quad \times \int_0^{\infty} b_{n,k}(t) (1 - \psi(t)) |f(t) - f_{2,\delta}(t)| dt \\
 &\quad + \frac{(n+r-1)!}{(n-1)!} (1+x)^{-n-r} (1 - \psi(0)) |f(0) - f_{2,\delta}(0)|.
 \end{aligned}$$

For sufficiently large n , the second term tends to zero. Thus

$$\begin{aligned}
 |I| &\leq \widehat{M}_9 \|f\|_{\gamma} \sum_{\substack{2i+j \leq r \\ i,j \geq 0}} n^i \sum_{k=1}^{\infty} p_{n,k}(x) |k - nx|^j \int_{|t-x| \geq \delta_1} b_{n,k}(t) dt \\
 &\leq \widehat{M}_9 \|f\|_{\gamma} \delta_1^{-2m} \sum_{\substack{2i+j \leq r \\ i,j \geq 0}} n^i \sum_{k=1}^{\infty} p_{n,k}(x) |k - nx|^j \left(\int_0^{\infty} b_{n,k}(t) dt \right)^{\frac{1}{2}} \\
 &\quad \times \left(\int_0^{\infty} b_{n,k}(t) (t-x)^{4m} dt \right)^{\frac{1}{2}} \\
 &\leq \widehat{M}_9 \|f\|_{\gamma} \delta_1^{-2m} \sum_{\substack{2i+j \leq r \\ i,j \geq 0}} n^i \left\{ \sum_{k=1}^{\infty} p_{n,k}(x) (k - nx)^{2j} \right\}^{\frac{1}{2}} \\
 &\quad \times \left\{ \sum_{k=1}^{\infty} p_{n,k}(x) \int_0^{\infty} b_{n,k}(t) (t-x)^{4m} dt \right\}^{\frac{1}{2}}.
 \end{aligned}$$



On Simultaneous Approximation For Certain Baskakov Durrmeyer Type Operators

Vijay Gupta,
 Muhammad Aslam Noor,
 Man Singh Beniwal and
 M. K. Gupta

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

Hence by using Lemma 2.1 and Lemma 2.2, we have

$$I \leq \widehat{M}_{10} \|f\|_{\gamma} \delta_1^{-2m} O\left(n^{(i+\frac{j}{2}-m)}\right) \leq \widehat{M}_{11} n^{-q} \|f\|_{\gamma},$$

where $q = m - \frac{r}{2}$. Now choosing $m > 0$ satisfying $q \geq 1$, we obtain $I \leq \widehat{M}_{11} n^{-1} \|f\|_{\gamma}$. Therefore by property (iii) of the function $f_{2,\delta}(t)$, we get

$$\begin{aligned} H_1 &\leq \widehat{M}_8 \left\| f^{(r)} - f_{2,\delta}^{(r)} \right\|_{C[a^*, b^*]} + \widehat{M}_{11} n^{-1} \|f\|_{\gamma} \\ &\leq \widehat{M}_{12} \omega_2(f^{(r)}, \delta, a, b) + \widehat{M}_{11} n^{-1} \|f\|_{\gamma}. \end{aligned}$$

Choosing $\delta = n^{-\frac{1}{2}}$, the theorem follows. □



**On Simultaneous
Approximation For Certain
Baskakov Durrmeyer Type
Operators**

*Vijay Gupta,
Muhammad Aslam Noor,
Man Singh Beniwal and
M. K. Gupta*

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

4. Inverse Theorem

This section is devoted to the following inverse theorem in simultaneous approximation:

Theorem 4.1. *Let $0 < \alpha < 2$, $0 < a_1 < a_2 < b_2 < b_1 < \infty$ and suppose $f \in C_\gamma[0, \infty)$. Then in the following statements $(i) \Rightarrow (ii)$*

$$(i) \|B_n^{(r)}(f, \bullet)\|_{C[a_1, b_1]} = O(n^{-\alpha/2}),$$

$$(ii) f^{(r)} \in Lip^*(\alpha, a_2, b_2),$$


where $Lip^*(\alpha, a_2, b_2)$ denotes the Zygmund class satisfying $\omega_2(f, \delta, a_2, b_2) \leq M\delta^\alpha$.

Proof. Let us choose a', a'', b', b'' in such a way that $a_1 < a' < a'' < a_2 < b_2 < b'' < b' < b_1$. Also suppose $g \in C_0^\infty$ with $\text{supp } g \in [a'', b'']$ and $g(x) = 1$ on the interval $[a_2, b_2]$. For $x \in [a', b']$ with $D \equiv \frac{d}{dx}$, we have

$$\begin{aligned} B_n^{(r)}(fg, x) - (fg)^{(r)}(x) &= D^r(B_n((fg)(t) - (fg)(x)), x) \\ &= D^r(B_n(f(t)(g(t) - g(x)), x)) + D^r(B_n(g(x)(f(t) - f(x)), x)) \\ &=: J_1 + J_2. \end{aligned}$$

Using the Leibniz formula, we have

$$J_1 = \frac{\partial^r}{\partial x^r} \int_0^\infty W_n(x, t) f(t) [g(t) - g(x)] dt$$



On Simultaneous Approximation For Certain Baskakov Durrmeyer Type Operators

Vijay Gupta,
Muhammad Aslam Noor,
Man Singh Beniwal and
M. K. Gupta

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

$$\begin{aligned}
&= \sum_{i=0}^r \binom{r}{i} \int_0^\infty W_n^{(i)}(x, t) \frac{\partial^{r-i}}{\partial x^{r-i}} [f(t)(g(t) - g(x))] dt \\
&= - \sum_{i=0}^{r-1} \binom{r}{i} g^{(r-i)}(x) B_n^{(i)}(f, x) + \int_0^\infty W_n^{(r)}(x, t) f(t)(g(t) - g(x)) dt \\
&=: J_3 + J_4.
\end{aligned}$$

Applying Theorem 3.4, we have


$$J_3 = - \sum_{i=0}^{r-1} \binom{r}{i} g^{(r-i)}(x) f^{(i)}(x) + O(n^{-\frac{\alpha}{2}}),$$

uniformly in $x \in [a', b']$. Applying Theorem 3.2, the Cauchy-Schwarz inequality, Taylor's expansions of f and g and Lemma 2.2, we are led to

$$\begin{aligned}
J_4 &= \sum_{i=0}^r \frac{g^{(i)}(x) f^{(r-i)}(x)}{i!(r-i)!} r! + o(n^{-\frac{1}{2}}) \\
&= \sum_{i=0}^r \binom{r}{i} g^{(i)}(x) f^{(r-i)}(x) + o(n^{-\frac{\alpha}{2}}),
\end{aligned}$$

uniformly in $x \in [a', b']$. Again using the Leibniz formula, we have

$$\begin{aligned}
J_2 &= \sum_{i=0}^r \binom{r}{i} \int_0^\infty W_n^{(i)}(x, t) \frac{\partial^{r-i}}{\partial x^{r-i}} [g(t)(f(t) - f(x))] dt \\
&= \sum_{i=0}^r \binom{r}{i} g^{(r-i)}(x) B_n^{(i)}(f, x) - (fg)^{(r)}(x)
\end{aligned}$$



On Simultaneous Approximation For Certain Baskakov Durrmeyer Type Operators

*Vijay Gupta,
Muhammad Aslam Noor,
Man Singh Beniwal and
M. K. Gupta*

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)


$$\begin{aligned}
&= \sum_{i=0}^r \binom{r}{i} g^{(r-i)}(x) f^{(i)}(x) - (fg)^{(r)}(x) + o(n^{-\alpha/2}) \\
&= O\left(n^{-\frac{\alpha}{2}}\right),
\end{aligned}$$

uniformly in $x \in [a', b']$. Combining the above estimates, we get

$$\|B_n^{(r)}(fg, \bullet) - (fg)^{(r)}\|_{C[a', b']} = O\left(n^{-\frac{\alpha}{2}}\right).$$

Thus by Lemma 2.4 and Lemma 2.5, we have $(fg)^{(r)} \in Lip^*(\alpha, a', b')$ also $g(x) = 1$ on the interval $[a_2, b_2]$, it proves that $f^{(r)} \in Lip^*(\alpha, a_2, b_2)$. This completes the validity of the implication $(i) \Rightarrow (ii)$ for the case $0 < \alpha \leq 1$. To prove the result for $1 < \alpha < 2$ for any interval $[a^*, b^*] \subset (a_1, b_1)$, let a_2^*, b_2^* be such that $(a_2, b_2) \subset (a_2^*, b_2^*)$ and $(a_2^*, b_2^*) \subset (a_1^*, b_1^*)$. Letting $\delta > 0$ we shall prove the assertion $\alpha < 2$. From the previous case it implies that $f^{(r)}$ exists and belongs to $Lip(1 - \delta, a_1^*, b_1^*)$. Let $g \in C_0^\infty$ be such that $g(x) = 1$ on $[a_2, b_2]$ and $\text{supp } g \subset (a_2^*, b_2^*)$. Then with $\chi_2(t)$ denoting the characteristic function of the interval $[a_1^*, b_1^*]$, we have

$$\begin{aligned}
&\|B_n^{(r)}(fg, \bullet) - (fg)^{(r)}\|_{C[a_2^*, b_2^*]} \\
&\leq \|D^r[B_n(g(\cdot)(f(t) - f(\cdot)), \bullet)]\|_{C[a_2^*, b_2^*]} \\
&\quad + \|D^r[B_n(f(t)(g(t) - g(\cdot)), \bullet)]\|_{C[a_2^*, b_2^*]} \\
&=: P_1 + P_2.
\end{aligned}$$



On Simultaneous Approximation For Certain Baskakov Durrmeyer Type Operators

Vijay Gupta,
Muhammad Aslam Noor,
Man Singh Beniwal and
M. K. Gupta

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

To estimate P_1 , by Theorem 3.4, we have

$$\begin{aligned}
 P_1 &= \|D^r[B_n(g(\cdot)(f(t), \bullet)] - (fg)^{(r)}\|_{C[a_2^*, b_2^*]} \\
 &= \left\| \sum_{i=0}^r \binom{r}{i} g^{(r-i)}(\cdot) B_n^{(i)}(f, \bullet) - (fg)^{(r)} \right\|_{C[a_2^*, b_2^*]} \\
 &= \left\| \sum_{i=0}^r \binom{r}{i} g^{(r-i)}(\cdot) f^{(i)} - (fg)^{(r)} \right\|_{C[a_2^*, b_2^*]} + O(n^{-\alpha/2}) \\
 &= O\left(n^{-\frac{\alpha}{2}}\right).
 \end{aligned}$$

Also by the Leibniz formula and Theorem 3.2, have

$$\begin{aligned}
 P_2 &\leq \left\| \sum_{i=0}^r \binom{r}{i} g^{(r-i)}(\cdot) B_n(f, \bullet) + B_n^{(r)}(f(t)(g(t) - g(\cdot))\chi_2(t), \bullet) \right\|_{C[a_2^*, b_2^*]} \\
 &\quad + O(n^{-1}) \\
 &=: \|P_3 + P_4\|_{C[a_2^*, b_2^*]} + O(n^{-1}).
 \end{aligned}$$

Then by Theorem 3.4, we have

$$P_3 = \sum_{i=0}^{r-1} \binom{r}{i} g^{(r-i)}(x) f^{(i)}(x) + O\left(n^{-\frac{\alpha}{2}}\right),$$



On Simultaneous Approximation For Certain Baskakov Durrmeyer Type Operators

Vijay Gupta,
 Muhammad Aslam Noor,
 Man Singh Beniwal and
 M. K. Gupta

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

uniformly in $x \in [a_2^*, b_2^*]$. Applying Taylor's expansion of f , we have


$$\begin{aligned} P_4 &= \int_{i=0}^{\infty} W_n^{(r)}(x, t) [f(t)(g(t) - g(x))\chi_2(t)] dt \\ &= \sum_{i=0}^r \frac{f^{(i)}(x)}{i!} \int_0^{\infty} W_n^{(r)}(x, t) (t-x)^i (g(t) - g(x)) dt \\ &\quad + \int_0^{\infty} W_n^{(r)}(x, t) \frac{(f^{(r)}(\xi) - f^{(r)}(x))}{r!} (t-x)^r (g(t) - g(x)) \chi_2(t) dt, \end{aligned}$$

where ξ lying between t and x . Next by Theorem 3.4, the first term in the above expression is given by

$$\sum_{m=0}^r \binom{r}{m} g^{(m)} f^{(r-m)}(x) + O(n^{-\frac{\alpha}{2}}),$$

uniformly in $x \in [a_2^*, b_2^*]$. Also by mean value theorem and using Lemma 2.4, we can obtain the second term as follows:

$$\begin{aligned} &\left\| \int_0^{\infty} W_n^{(r)}(x, t) \frac{(f^{(r)}(\xi) - f^{(r)}(x))}{r!} (t-x)^r (g(t) - g(x)) \chi_2(t) dt \right\|_{C[a_2^*, b_2^*]} \\ &\leq \sum_{\substack{2m+s \leq r \\ m, s \geq 0}} n^{m+s} \left\| \frac{|Q_{m,s,r}(x)|^r}{x(1+x)} \int_0^{\infty} W_n(x, t) |t-x|^{\delta+r+1} \right. \\ &\quad \times \left. \frac{|f^{(r)}(\xi) - f^{(r)}(x)|}{r!} |g'(\eta)| \chi_2(t) dt \right\|_{C[a_2^*, b_2^*]} \\ &= O\left(n^{-\frac{\delta}{2}}\right), \end{aligned}$$



On Simultaneous Approximation For Certain Baskakov Durrmeyer Type Operators

Vijay Gupta,
Muhammad Aslam Noor,
Man Singh Beniwal and
M. K. Gupta

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)


[Quit](#)

choosing δ such that $0 \leq \delta \leq 2 - \alpha$. Combining the above estimates we get

$$\|B_n^{(r)}(fg, \bullet) - (fg)^{(r)}\|_{C[a_2^*, b_2^*]} = O\left(n^{-\frac{\alpha}{2}}\right).$$

Since $\text{supp } fg \subset (a_2^*, b_2^*)$, it follows from Lemma 2.4 and Lemma 2.5 that $(fg)^{(r)} \in \text{Liz}(\alpha, 1, a_2^*, b_2^*)$. Since $g(x) = 1$ on $[a_2, b_2]$, we have $f^{(r)} \in \text{Liz}(\alpha, 1, a_2^*, b_2^*)$. This completes the proof of the theorem. \square

Remark 2. As noted in the first section, these operators also reproduce the linear functions so we can easily apply the iterative combinations to the operators (1.1) to improve the order of approximation.



On Simultaneous
Approximation For Certain
Baskakov Durrmeyer Type
Operators

Vijay Gupta,
Muhammad Aslam Noor,
Man Singh Beniwal and
M. K. Gupta

[Title Page](#)

[Contents](#)



[Go Back](#)


[Close](#)

[Quit](#)

Page 32 of 33

References

- [1] H. BERENS AND G.G. LORENTZ, Inverse theorem for Bernstein polynomials, *Indiana Univ. Math. J.*, **21** (1972), 693–708.
- [2] Z. FINTA, On converse approximation theorems, *J. Math. Anal. Appl.*, **312**(1) (2005), 159–180.
- [3] G. FREUD AND V. POPOV, On approximation by Spline functions, *Proceeding Conference on Constructive Theory Functions*, Budapest (1969), 163–172.
- [4] S. GOLDBERG AND V. MEIR, Minimum moduli of ordinary differential operators, *Proc. London Math. Soc.*, **23** (1971), 1–15.
- [5] V. GUPTA, A note on modified Baskakov type operators, *Approximation Theory Appl.*, **10**(3) (1994), 74–78.
- [6] V. GUPTA, M.K. GUPTA AND V. VASISHTHA, Simultaneous approximation by summation-integral type operators, *Nonlinear Funct. Anal. Appl.*, **8**(3), (2003), 399–412.
- [7] V. GUPTA AND M.A. NOOR, Convergence of derivatives for certain mixed Szasz Beta operators, *J. Math. Anal. Appl.*, **321**(1) (2006), 1–9.
- [8] N. ISPIR AND I. YUKSEL, On the Bezier variant of Srivastava-Gupta operators, *Applied Math. E Notes*, 5 (2005), 129–137.
- [9] H.M. SRIVASTAVA AND V. GUPTA, A certain family of summation integral type operators, *Math. Comput. Modelling*, **37** (2003), 1307–1315.



On Simultaneous
Approximation For Certain
Baskakov Durrmeyer Type
Operators

Vijay Gupta,
Muhammad Aslam Noor,
Man Singh Beniwal and
M. K. Gupta

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)