

Journal of Inequalities in Pure and Applied Mathematics

HISTORY, GENERALIZATIONS AND UNIFIED TREATMENT OF TWO OSTROWSKI'S INEQUALITIES

SANJA VAROŠANEC

Department of Mathematics
University of Zagreb,
Zagreb, Croatia.

EMail: varosans@math.hr

©2000 Victoria University
ISSN (electronic): 1443-5756
022-04



volume 5, issue 2, article 23,
2004.

*Received 28 January, 2004;
accepted 28 January, 2004.*

Communicated by: J.E. Pečarić

Abstract

Contents



Home Page

Go Back

Close

Quit

Abstract

In this paper we present a historical review of the investigation of two Ostrowski inequalities and describe several distinct streams for their generalizations. Also we point out some new methods to obtain known results and give a number of new results related to Ostrowski's inequalities.

2000 Mathematics Subject Classification: Primary 26D15, Secondary 46C05.

Key words: Ostrowski's inequalities, Cauchy-Buniakowski-Schwarz inequality, Gram's determinant, Unitary vector space, Superadditive function, Interpolation.

Contents

1	History and Generalizations	3
2	Further Generalizations of Ostrowski's Inequalities	13
3	Interpolation	16
References		



History, Generalizations And Unified Treatment Of Two Ostrowski's Inequalities

Sanja Varošanec

Title Page

Contents



Go Back

Close

Quit

Page 2 of 20

1. History and Generalizations

In his book *Vorlesungen über Differential und Integralrechnung II*, A. Ostrowski presented the following interesting inequalities.

Theorem 1.1. [12, p. 289, problem 61], [10, pp. 92–93]. *The minimum of the sum $x_1^2 + \cdots + x_n^2$ under the conditions*

$$(1.1) \quad \sum_{i=1}^n a_i x_i = 0 \quad \text{and} \quad \sum_{i=1}^n b_i x_i = 1$$

is

$$(1.2) \quad \frac{\sum_{i=1}^n a_i^2}{\sum_{i < j} (a_i b_j - a_j b_i)^2}, \quad \left(\sum_{i=1}^n a_i^2 + \sum_{i=1}^n b_i^2 > 0 \right).$$


Theorem 1.2. [12, p. 290, problem 63], [10, p. 94]. *The maximum of the sum $(\sum_{i=1}^n b_i x_i)^2$ under the conditions*

$$(1.3) \quad \sum_{i=1}^n a_i x_i = 0 \quad \text{and} \quad \sum_{i=1}^n x_i^2 = 1$$

is

$$(1.4) \quad \frac{\sum_{i < j} (a_i b_j - a_j b_i)^2}{\sum_{i=1}^n a_i^2}, \quad \left(\sum_{i=1}^n a_i^2 > 0 \right).$$

According to the Lagrange identity [10, p. 84], Theorem 1.1 can be rewritten in the following form.



History, Generalizations And
Unified Treatment Of Two
Ostrowski's Inequalities

Sanja Varošanec

Title Page

Contents



Go Back

Close

Quit

Page 3 of 20

Theorem 1.3. Let $a = (a_1, \dots, a_n)$ and $b = (b_1, \dots, b_n)$ be two nonproportional sequences of real numbers and let $x = (x_1, \dots, x_n)$ be any real sequence which satisfies

$$(1.5) \quad \sum_{i=1}^n a_i x_i = 0 \quad \text{and} \quad \sum_{i=1}^n b_i x_i = 1.$$

Then

$$(1.6) \quad \sum_{i=1}^n x_i^2 \geq \frac{\sum_{i=1}^n a_i^2}{(\sum_{i=1}^n a_i^2)(\sum_{i=1}^n b_i^2) - (\sum_{i=1}^n a_i b_i)^2}.$$


The second Ostrowski problem can also be written in the analogue form. In the literature those forms are used more frequently than the original and have been extended, improved and generalized in different ways.

The aim of this paper is to give a brief historical review and to carry those ideas somewhat further.

K. Fan and J. Todd, [8], using Theorem 1.1, i.e. Theorem 1.3, they established the following theorem.

Theorem 1.4. Let $a = (a_1, \dots, a_n)$ and $b = (b_1, \dots, b_n)$ ($n \geq 2$) be two sequences of real numbers such that $a_i b_j \neq a_j b_i$ for $i \neq j$. Then

$$(1.7) \quad \frac{\sum_{i=1}^n a_i^2}{(\sum_{i=1}^n a_i^2)(\sum_{i=1}^n b_i^2) - (\sum_{i=1}^n a_i b_i)^2} \leq \left(\frac{2}{n(n-1)} \right)^2 \sum_{i=1}^n \left(\sum_{j=1, j \neq i}^n \frac{a_j}{a_j b_i - a_i b_j} \right)^2.$$



Title Page

Contents



Go Back

Close

Quit

Page 4 of 20

They also generalized Theorem 1.4 using more than two vectors.

Another direction of generalization has arisen from the fact that the map $(x_1, \dots, x_n) \mapsto \sqrt{\sum_{i=1}^n x_i^2}$ is a Euclidean norm in \mathbb{R}^n generated by the inner product $\langle x, y \rangle = \sum_{i=1}^n x_i y_i$. It is natural to consider an arbitrary inner product instead of the Euclidean inner product. The first generalization of that kind was done by Ž. Mitrović, [11] and after that some similar results were given in [6], [7] and [15]. Here we quote Mitrović's result.

Theorem 1.5. *Let a and b be linearly independent vectors of a unitary complex vector space V and let x be a vector in V such that*

$$(1.8) \quad \langle x, a \rangle = \alpha \text{ and } \langle x, b \rangle = \beta.$$

Then

$$(1.9) \quad G(a, b) \|x\|^2 \geq \|\bar{\alpha}b - \bar{\beta}a\|^2,$$

where $G(a, b)$ is the Gram determinant of vectors a and b . Equality holds if and only if

$$(1.10) \quad x = \frac{1}{G(a, b)} (\langle a, \bar{\beta}a - \bar{\alpha}b \rangle b - \langle b, \bar{\beta}a - \bar{\alpha}b \rangle a).$$

Here we present a rough outline of Mitrović's proof. Let y be a vector in V given by

$$y = \frac{1}{G(a, b)} (\langle a, \bar{\beta}a - \bar{\alpha}b \rangle b - \langle b, \bar{\beta}a - \bar{\alpha}b \rangle a).$$



If vector x satisfies conditions (1.8), then $\langle y, y \rangle = \langle x, y \rangle = \frac{1}{G(a,b)} \|\bar{\alpha}b - \bar{\beta}a\|^2$ and $\|x - y\|^2 = \|x\|^2 - \|y\|^2$. Since $\|x - y\|^2 \geq 0$, we obtain

$$\|x\|^2 \geq \|y\|^2 = \frac{1}{G(a,b)} \|\bar{\alpha}b - \bar{\beta}a\|^2$$

and inequality (1.9) holds.

Remark 1.1. Now, we point out another proof of Theorem 1.5. It is well known that Gram's determinant of the vectors x_1, x_2, x_3 is nonnegative, i.e. inequality

$$G(x_1, x_2, x_3) \geq 0$$

holds with equality iff the vectors x_1, x_2, x_3 are linearly dependent. Putting $x_1 = x, x_2 = a, x_3 = b$ and using notations $\langle x, a \rangle = \alpha$ and $\langle x, b \rangle = \beta$ we have the following

$$\begin{aligned} 0 &\leq G(x, a, b) \\ &= \begin{vmatrix} \langle x, x \rangle & \langle x, a \rangle & \langle x, b \rangle \\ \langle a, x \rangle & \langle a, a \rangle & \langle a, b \rangle \\ \langle b, x \rangle & \langle b, a \rangle & \langle b, b \rangle \end{vmatrix} \\ &= G(a, b)\|x\|^2 - \langle x, a \rangle \begin{vmatrix} \langle a, x \rangle & \langle a, b \rangle \\ \langle b, x \rangle & \langle b, b \rangle \end{vmatrix} + \langle x, b \rangle \begin{vmatrix} \langle a, x \rangle & \langle a, a \rangle \\ \langle b, x \rangle & \langle b, a \rangle \end{vmatrix} \\ &= G(a, b)\|x\|^2 - \alpha(\bar{\alpha}\langle b, b \rangle - \bar{\beta}\langle a, b \rangle) + \beta(\bar{\alpha}\langle b, a \rangle - \bar{\beta}\langle a, a \rangle), \end{aligned}$$

$$G(a, b)\|x\|^2 \geq |\alpha|^2\langle b, b \rangle - \alpha\bar{\beta}\langle a, b \rangle - \beta\bar{\alpha}\langle b, a \rangle + |\beta|^2\langle a, a \rangle = \|\bar{\alpha}b - \bar{\beta}a\|^2.$$



History, Generalizations And
Unified Treatment Of Two
Ostrowski's Inequalities

Sanja Varošanec

Title Page

Contents



Go Back

Close

Quit

Page 6 of 20

Equality holds iff vectors x, a and b are linearly dependent, i.e. there exist scalars λ and μ such that

$$x = \lambda a + \mu b.$$

Multiplying that identity by a and b respectively, we obtain $\alpha = \lambda\langle a, a \rangle + \mu\langle b, a \rangle$ and $\beta = \lambda\langle a, b \rangle + \mu\langle b, b \rangle$ from where we easily find that

$$\lambda = \frac{1}{G(a, b)}(\alpha\langle b, b \rangle - \beta\langle b, a \rangle), \quad \mu = \frac{1}{G(a, b)}(\beta\langle a, a \rangle - \alpha\langle a, b \rangle).$$

So, x is the vector given in (1.10).

In the same paper [11] a generalization of Fan-Todd's result is given. Furthermore, in the paper [2] P.R. Beesack noticed that inequality (1.9) and a fortiori also Ostrowski's inequality (1.6) can be regarded as a special case of the Bessel inequality for non-orthonormal vectors.


Theorem 1.6. [2] *Let a_1, \dots, a_k , ($k \geq 1$) be linearly independent vectors of a Hilbert space H and let $\alpha_1, \dots, \alpha_k$ be given scalars. If $x \in H$ satisfies*

$$(1.11) \quad \langle x, a_i \rangle = \alpha_i \quad 1 \leq i \leq k,$$

then

$$(1.12) \quad G(a_1, \dots, a_k)^2 \|x\|^2 \geq \left\| \sum_{i=1}^k \gamma_i^{(k)} a_i \right\|^2,$$

where $G(a_1, \dots, a_k)$ is the Gram determinant of a_1, \dots, a_k and $\gamma_i^{(k)}$ is the determinant obtained from G by replacing the elements of the i^{th} row of G by



History, Generalizations And
Unified Treatment Of Two
Ostrowski's Inequalities

Sanja Varošanec

Title Page

Contents



Go Back

Close

Quit

Page 7 of 20

$(\alpha_1, \dots, \alpha_k)$. Moreover, equality holds if and only if

$$(1.13) \quad G(a_1, \dots, a_k)x = \sum_{i=1}^k \gamma_i^{(k)} a_i.$$

Finally, an analogue of Theorem 1.5 and related generalizations in 2-inner and n -inner spaces are given in [4] and [5].

The second stream of generalization of Ostrowski's inequality (1.6) was started by Madevski's paper [9]. He used Theorem 1.3 to obtain inequalities between certain statistical central moments. Also, he gave the following p -version of Ostrowski's inequality.


Theorem 1.7. [9] *Let $a = (a_1, \dots, a_n)$ and $b = (b_1, \dots, b_n)$ be two nonproportional sequences of real numbers and let $x = (x_1, \dots, x_n)$ be any real sequence which satisfies*

$$(1.14) \quad \sum_{i=1}^n a_i x_i = 0 \quad \text{and} \quad \sum_{i=1}^n b_i x_i = 1.$$

If p is an integer, then

$$(1.15) \quad \left(\sum_{i=1}^n x_i^2 \right)^p \geq \frac{(\sum_{i=1}^n a_i^2)^p}{(\sum_{i=1}^n a_i^2)^p (\sum_{i=1}^n b_i^2)^p - (\sum_{i=1}^n a_i b_i)^{2p}}.$$

In [1] M. Alić and J. Pečarić proved that the integer p can be substituted by an arbitrary real number $p \geq 1$. In the same paper a sequence of results involving moments of discrete distribution function has been given. An integral version



History, Generalizations And
Unified Treatment Of Two
Ostrowski's Inequalities

Sanja Varošanec

Title Page

Contents



Go Back

Close

Quit

Page 8 of 20

of those results and some generalizations of known statistical inequalities given in [9], [14] and [16] are obtained in [13].

Recently, Theorems 1.1 and 1.2 have been the focus of investigation. In the papers [6] and [7] the authors have used elementary arguments and the Cauchy-Buniakowski-Schwarz inequality to obtain Ostrowski type inequalities in unitary space. Indeed, the following theorems are obtained.

Theorem 1.8. [7] *Let a and b be linearly independent vectors of a real or complex unitary vector space V and let x be a vector in V such that*

$$(1.16) \quad \langle x, a \rangle = 0 \quad \text{and} \quad |\langle x, b \rangle| = 1.$$

Then

$$(1.17) \quad \|x\|^2 \geq \frac{\|a\|^2}{\|a\|^2\|b\|^2 - |\langle a, b \rangle|^2}.$$

Equality holds if and only if


$$(1.18) \quad x = \mu \left(b - \frac{\langle a, b \rangle}{\|a\|^2} a \right),$$

where $\mu \in \mathbb{K}$ ($\mathbb{K} = \mathbb{R}, \mathbb{C}$) is such that

$$|\mu| = \frac{\|a\|^2}{\|a\|^2\|b\|^2 - |\langle a, b \rangle|^2}.$$

Theorem 1.9. [6] *Let a and b be linearly independent vectors of a real or complex unitary vector space V and let x be a vector in V such that*

$$(1.19) \quad \langle x, a \rangle = 0 \quad \text{and} \quad \|x\| = 1.$$



**History, Generalizations And
Unified Treatment Of Two
Ostrowski's Inequalities**

Sanja Varošanec

Title Page

Contents



Go Back

Close

Quit

Page 9 of 20

Then

$$(1.20) \quad |\langle x, b \rangle|^2 \leq \frac{\|a\|^2 \|b\|^2 - |\langle a, b \rangle|^2}{\|a\|^2}.$$

Equality holds if and only if

$$(1.21) \quad x = \nu \left(b - \frac{\langle b, a \rangle}{\|a\|^2} a \right),$$

where $\nu \in \mathbb{K} (\mathbb{K} = \mathbb{R}, \mathbb{C})$ is so that

$$|\nu| = \frac{\|a\|}{(\|a\|^2 \|b\|^2 - |\langle a, b \rangle|^2)^{\frac{1}{2}}}.$$


It is obvious that these results are special cases of Theorem 1.5 but we mentioned it because the method of proving is different from Mitrović's method and leads to another generalization which will be given in the next section. Proofs of the previous two theorems are based on the Cauchy-Buniakowski-Schwarz inequality:

$$\|u\|^2 \|v\|^2 \geq |\langle u, v \rangle|^2, \quad u, v \in V.$$

Applying it on vectors $u = z - \frac{\langle z, c \rangle}{\|c\|^2} c$ and $v = d - \frac{\langle d, c \rangle}{\|c\|^2} c$, where $c \neq 0$ and taking into account that

$$(1.22) \quad \left\| z - \frac{\langle z, c \rangle}{\|c\|^2} c \right\|^2 = \frac{\|z\|^2 \|c\|^2 - |\langle z, c \rangle|^2}{\|c\|^2},$$

$$(1.23) \quad \left\| d - \frac{\langle d, c \rangle}{\|c\|^2} c \right\|^2 = \frac{\|d\|^2 \|c\|^2 - |\langle d, c \rangle|^2}{\|c\|^2},$$



History, Generalizations And
Unified Treatment Of Two
Ostrowski's Inequalities

Sanja Varošanec

Title Page

Contents



Go Back

Close

Quit

Page 10 of 20

and

$$(1.24) \quad \left\langle z - \frac{\langle z, c \rangle}{\|c\|^2} c, d - \frac{\langle d, c \rangle}{\|c\|^2} c \right\rangle = \frac{\langle z, d \rangle \|c\|^2 - \langle z, c \rangle \langle c, d \rangle}{\|c\|^2}$$

we have the following inequality

$$(1.25) \quad (\|z\|^2 \|c\|^2 - |\langle z, c \rangle|^2) (\|d\|^2 \|c\|^2 - |\langle d, c \rangle|^2) \geq |\langle z, d \rangle \|c\|^2 - \langle z, c \rangle \langle c, d \rangle|^2.$$

Putting in inequality (1.25) $z = x$, $c = a$ and $d = b$ where a and x satisfy $\langle x, a \rangle = 0$ and $\|x\| = 1$ we get inequality (1.20), while if a and x satisfy $\langle x, a \rangle = 0$ and $|\langle x, b \rangle| = 1$ inequality (1.17) is obtained.


Remark 1.2. Let us mention that inequality (1.9) also can be obtained by the above-mentioned method. In fact, putting in inequality (1.25) $z = x$, $c = a$ and $d = b$, $\langle x, a \rangle = \alpha$ and $\langle x, b \rangle = \beta$ we get

$$(\|x\|^2 \|a\|^2 - |\langle x, a \rangle|^2) (\|b\|^2 \|a\|^2 - |\langle b, a \rangle|^2) \geq |\beta \|a\|^2 - \alpha \langle a, b \rangle|^2,$$

$$\|x\|^2 \|a\|^2 G(a, b) \geq |\beta \|a\|^2 - \alpha \langle a, b \rangle|^2 + |\alpha|^2 G(a, b) = \|a\|^2 \|\bar{\alpha} b - \bar{\beta} a\|^2$$

from where inequality (1.9) occurs. Using the fact that in the Cauchy-Buniakowski-Schwarz inequality, equality holds iff vectors are proportional, we get (1.10).

Remark 1.3. Inequality (1.25) is a special case of the more general result related to Gram's determinant given in [10, p. 599]. That result is as follows.



History, Generalizations And
Unified Treatment Of Two
Ostrowski's Inequalities

Sanja Varošanec

Title Page

Contents



Go Back

Close


Quit

Page 11 of 20

Theorem 1.10. For vectors x_1, \dots, x_n and y_1, \dots, y_n from unitary space V the following inequality holds

$$\left| \det \begin{bmatrix} \langle x_1, y_1 \rangle & \dots & \langle x_1, y_n \rangle \\ \vdots & & \vdots \\ \langle x_n, y_1 \rangle & \dots & \langle x_n, y_n \rangle \end{bmatrix} \right|^2 \leq G(x_1, \dots, x_n)G(y_1, \dots, y_n),$$

with equality iff the vectors x_1, \dots, x_n span the same subspace as the vectors y_1, \dots, y_n .



**History, Generalizations And
Unified Treatment Of Two
Ostrowski's Inequalities**

Sanja Varošanec

Title Page

Contents



Go Back

Close

Quit

Page 12 of 20

2. Further Generalizations of Ostrowski's Inequalities

In this section we extend results from papers [1], [6], [7], [13] introducing super(sub)additive function.

Theorem 2.1. *Let a and b be linearly independent vectors of a unitary complex vector space V and let x be a vector in V such that*

$$(2.1) \quad \langle x, a \rangle = \alpha \quad \text{and} \quad \langle x, b \rangle = \beta.$$

If $\phi : [0, \infty) \rightarrow \mathbb{R}$ is a nondecreasing, superadditive function, then

$$(2.2) \quad \phi(\|a\|^2\|b\|^2) - \phi(|\langle a, b \rangle|^2) \geq \phi\left(\frac{\|\bar{\alpha}b - \bar{\beta}a\|^2}{\|x\|^2}\right).$$

If ϕ is a nonincreasing, subadditive function then a reverse in (2.2) holds.

Proof. Let us suppose that ϕ is a superadditive nondecreasing function. Then we have

$$(2.3) \quad \phi(u) = \phi((u-v) + v) \geq \phi(u-v) + \phi(v), \quad \text{i.e.} \quad \phi(u) - \phi(v) \geq \phi(u-v).$$

Taking into account the nondecreasing property of ϕ , results of Theorem 1.5 and inequality (2.3) we conclude

$$\begin{aligned} \phi(\|a\|^2\|b\|^2) - \phi(|\langle a, b \rangle|^2) &\geq \phi(\|a\|^2\|b\|^2) - (|\langle a, b \rangle|^2) \\ &\geq \phi\left(\frac{\|\bar{\alpha}b - \bar{\beta}a\|^2}{\|x\|^2}\right). \end{aligned}$$



History, Generalizations And
Unified Treatment Of Two
Ostrowski's Inequalities

Sanja Varošanec

Title Page

Contents



Go Back

Close

Quit

Page 13 of 20

The case when ϕ is a nonincreasing and subadditive function has been done similarly. \square

In particular, inequality (2.2) holds for any nondecreasing convex function ϕ , while its reverse holds for any nonincreasing concave function. The result of Theorem 2.1 can be improved if function ϕ is a power function. In that case we have the following result.

Theorem 2.2. *Suppose that a, b and x are as in Theorem 2.1. If $p \geq 1$, then*

$$(2.4) \quad \|x\|^{2p} \|a\|^{2p} (\|a\|^{2p} \|b\|^{2p} - |\langle a, b \rangle|^{2p}) \\ \geq \max \left\{ \|a\|^{2p} \|\bar{\alpha}b - \bar{\beta}a\|^{2p}, \right. \\ \left. |\beta\|a\|^2 - \alpha\langle a, b \rangle|^{2p} + |\alpha|^{2p} (\|a\|^{2p} \|b\|^{2p} - |\langle a, b \rangle|^{2p}) \right\}.$$

Proof. The function $\phi(x) = x^p, p \geq 1$ is a nondecreasing superadditive function so, a direct consequence of the previous theorem is that for a, b and x which satisfy assumptions of Theorem 2.1 we have the following inequality

$$(2.5) \quad \|a\|^{2p} \|b\|^{2p} - |\langle a, b \rangle|^{2p} \geq \frac{\|\bar{\alpha}b - \bar{\beta}a\|^{2p}}{\|x\|^{2p}}.$$

Applying the method of proving in Theorem 2.1 on inequality (1.25) we get

$$(2.6) \quad (\|z\|^{2p} \|c\|^{2p} - |\langle z, c \rangle|^{2p}) (\|d\|^{2p} \|c\|^{2p} - |\langle d, c \rangle|^{2p}) \\ \geq |\langle z, d \rangle \|c\|^2 - \langle z, c \rangle \langle c, d \rangle|^{2p}$$



History, Generalizations And
Unified Treatment Of Two
Ostrowski's Inequalities

Sanja Varošanec

Title Page

Contents



Go Back

Close

Quit


Page 14 of 20

i.e. putting $z = x$, $c = a$, $d = b$ and taking into account that $\langle x, a \rangle = \alpha$ and $\langle x, b \rangle = \beta$ we have

$$(2.7) \quad (\|x\|^{2p}\|a\|^{2p} - |\alpha|^{2p}) (\|b\|^{2p}\|a\|^{2p} - |\langle b, a \rangle|^{2p}) \geq |\beta\|a\|^2 - \alpha\langle a, b \rangle|^{2p}.$$

After simple calculations, inequalities (2.6) and (2.7) give inequality (2.4). \square

Remark 2.1. *If $p = 1$, then the two terms on the righthand side of inequality (2.4) are equal, but if $p > 1$ terms $\|a\|^{2p}\|\bar{\alpha}b - \bar{\beta}a\|^{2p}$ and $|\beta\|a\|^2 - \alpha\langle a, b \rangle|^{2p} + |\alpha|^{2p}(\|a\|^{2p}\|b\|^{2p} - |\langle a, b \rangle|^{2p})$ are not comparable. For example, if $p = 2$, $\|a\| = 1$, $\|b\| = 1$, $\langle a, b \rangle = \frac{1}{2}$ and $\alpha = 1$, $\beta \in \mathbb{R}$ the first term is equal to $(\beta^2 - \beta + 1)^2$, while the second term is equal to $(\beta^2 - \beta + \frac{1}{4})^2 + \frac{15}{16}$. If $\beta \in (0, 1)$ the first term is less than the second term and if $\beta > 1$ the opposite inequality holds.*



History, Generalizations And
Unified Treatment Of Two
Ostrowski's Inequalities

Sanja Varošanec

Title Page

Contents



Go Back

Close

Quit

Page 15 of 20

3. Interpolation

Some results about refinements of the original first Ostrowski's inequality are given in [3]. Here we give more general results in which we consider refinements of Ostrowski's inequalities in arbitrary unitary complex vector spaces.

Theorem 3.1. *Let a and b be linearly independent vectors in a unitary complex vector space V and let x be a vector in V such that*

$$(3.1) \quad \langle x, a \rangle = \alpha \quad \text{and} \quad \langle x, b \rangle = \beta.$$

Let y be a vector defined by

$$(3.2) \quad y = \frac{1}{G(a, b)} (\langle a, \bar{\beta}a - \bar{\alpha}b \rangle b - \langle b, \bar{\beta}a - \bar{\alpha}b \rangle a).$$

Then the vector $F(x) = \theta x + (1 - \theta)y$, $\theta \in [0, 1]$, satisfies

$$(3.3) \quad \|x\|^2 \geq \|F(x)\|^2$$

and


$$(3.4) \quad G(a, b) \|F(x)\|^2 \geq \|\bar{\alpha}b - \bar{\beta}a\|^2.$$

Proof. Let us note that y is a vector for which equality in (1.9) holds, i.e.

$$(3.5) \quad G(a, b) \|y\|^2 = \|\bar{\alpha}b - \bar{\beta}a\|^2.$$

So, without any calculation we conclude that $\langle y, a \rangle = \alpha$ and $\langle y, b \rangle = \beta$. Now,

$$(3.6) \quad \langle F(x), a \rangle = \langle \theta x + (1 - \theta)y, a \rangle = \theta\alpha + (1 - \theta)\alpha = \alpha.$$



History, Generalizations And
Unified Treatment Of Two
Ostrowski's Inequalities

Sanja Varošanec

Title Page

Contents



Go Back

Close

Quit

Page 16 of 20

Similarly, we obtain

$$(3.7) \quad \langle F(x), b \rangle = \beta.$$

According to Theorem 1.5 and in view of (3.6) and (3.7) we get

$$G(a, b) \|F(x)\|^2 \geq \|\bar{\alpha}b - \bar{\beta}a\|^2.$$

Let us calculate the product $\langle y, x \rangle$.


$$\begin{aligned} G(a, b) \langle y, x \rangle &= \langle \langle a, \bar{\beta}a - \bar{\alpha}b \rangle b - \langle b, \bar{\beta}a - \bar{\alpha}b \rangle a, x \rangle \\ &= \langle a, \bar{\beta}a - \bar{\alpha}b \rangle \langle b, x \rangle - \langle b, \bar{\beta}a - \bar{\alpha}b \rangle \langle a, x \rangle \\ &= \bar{\beta}(\beta \langle a, a \rangle - \alpha \langle a, b \rangle) - \bar{\alpha}(\beta \langle b, a \rangle - \alpha \langle b, b \rangle) \\ &= |\beta|^2 \|a\|^2 - \alpha \bar{\beta} \langle a, b \rangle - \bar{\alpha} \beta \langle b, a \rangle + |\alpha|^2 \|b\|^2 \\ &= \|\bar{\alpha}b - \bar{\beta}a\|^2. \end{aligned}$$

Comparing this result with (3.5) we have $\langle y, x \rangle = \langle y, y \rangle = \langle x, y \rangle$. Using these equalities we obtain

$$\begin{aligned} \|F(x)\|^2 &= \langle F(x), F(x) \rangle \\ &= \theta^2 \|x\|^2 + \theta(1 - \theta) \langle x, y \rangle + (1 - \theta)\theta \langle y, x \rangle + (1 - \theta)^2 \langle y, y \rangle \\ &= \theta^2 \|x\|^2 + (1 - \theta^2) \|y\|^2. \end{aligned}$$

$$\|x\|^2 - \|F(x)\|^2 = (1 - \theta^2)(\|x\|^2 - \|y\|^2) = (1 - \theta^2)(\|x - y\|^2) \geq 0$$

and inequality (3.3) has been established. \square



History, Generalizations And
Unified Treatment Of Two
Ostrowski's Inequalities

Sanja Varošanec

Title Page

Contents



Go Back

Close

Quit

Page 17 of 20


Thus we obtain a sequence of successive approximations

$$x, F(x), F^2(x), \dots, F^n(x), \dots$$

converging to y for $\theta < 1$ which interpolate inequality (1.9)

$$\|x\|^2 \geq \|F(x)\|^2 \geq \|F^2(x)\|^2 \geq \dots \geq \|F^n(x)\|^2 \geq \dots \geq \|y\|^2 = \frac{\|\bar{\alpha}b - \bar{\beta}a\|^2}{G(a, b)}.$$

If $\alpha = 0$, $\beta = 1$, $\theta = \frac{1}{2}$ and $\|x\|^2 = \sum_{i=1}^n x_i^2$, then we get a result of M. Bjelica, [3].



History, Generalizations And
Unified Treatment Of Two
Ostrowski's Inequalities

Sanja Varošanec

Title Page

Contents



Go Back

Close

Quit

Page 18 of 20

References

- [1] M. ALIĆ AND J. PEČARIĆ, On some inequalities of Ž. Madevski and A.M. Ostrowski, *Rad HAZU, Matematičke znanosti*, **472** (1997), 77–82.
- [2] P.R. BEESACK, On Bessel's inequality and Ostrowski's, *Univ. Beograd Publ.Elektrotehn. Fak. Ser. Mat. Fiz.*, No **498-541** (1975), 69–71.
- [3] M. BJELICA, Refinements of Ostrowski's and Fan-Todd's inequalities, in *Recent Progress in Inequalities*, G.V. Milovanović (Ed.), Kluwer Academic Publishers, Dordrecht, 1998, 445–448.
- [4] Y.J. CHO, M. MATIĆ AND J. PEČARIĆ, On Gram's determinant in 2-inner product spaces, *J. Korean Math. Soc.*, **38**(6) (2001), 1125–1156.
- [5] Y.J. CHO, M. MATIĆ AND J. PEČARIĆ, On Gram's determinant in n-inner product spaces, *Bull. Korean Math. Soc.*, to appear.
- [6] S.S. DRAGOMIR AND A.C. GOŠA, A generalization of an Ostrowski inequality in inner product spaces, *RGMI Research Report Collection*, **6**(2) (2003), Article 20. [ONLINE] <http://rgmia.vu.edu.au/v6n2.html>.
- [7] S.S. DRAGOMIR, Ostrowski's inequality in complex inner product spaces, *RGMI Research Report Collection* **6**(Supp.) (2003), Article 6. [ONLINE] [http://rgmia.vu.edu.au/v6\(E\).html](http://rgmia.vu.edu.au/v6(E).html).
- [8] K. FAN AND J. TODD, A determinantal inequality, *J. London Math. Soc.*, **30** (1955), 58–64.



History, Generalizations And
Unified Treatment Of Two
Ostrowski's Inequalities

Sanja Varošanec

Title Page

Contents




Go Back

Close

Quit

Page 19 of 20

- [9] Ž. MADEVSKI, Quelques conséquences d'une inégalité d'Ostrowski, *Univ. Beograd Publ. Elektrotehn. Fak. Ser. Mat. Fiz.*, No **412-460** (1973), 168–170.
- [10] D.S. MITRINOVIĆ, J.E. PEČARIĆ AND A.M. FINK, *Classical and New Inequalities*, Kluwer Academic Publishers, Dordrecht, 1993.
- [11] Ž. MITROVIĆ, On a generalization of Fan-Todd's inequality, *Univ. Beograd Publ. Elektrotehn. Fak. Ser. Mat. Fiz.* No **412-460** (1973), 151–154.
- [12] A. OSTROWSKI, *Vorlesungen über Differential und Integralrechnung II*, Verlag Birkhäuser Basel, 1951.
- [13] C.E.M. PEARCE, J. PEČARIĆ AND S. VAROŠANEC, An integral analogue of the Ostrowski inequality, *J. Ineq. Appl.*, **2** (1998), 275–283.
- [14] K. PEARSON, Mathematical contributions to the theory of evolution. XIX: second supplement to a memoir on skew variation, *Phil.Trans.Roy. Soc. A*, **216** (1916), 432.
- [15] H. ŠIKIĆ AND T. ŠIKIĆ, A note on Ostrowski's inequality, *Math. Ineq. Appl.*, **4**(2) (2001), 297–299.
- [16] J.E. WILKINS, A note on skewness and kurtosis, *Ann. Math. Statist.*, **15** (1944), 333–335.



History, Generalizations And
Unified Treatment Of Two
Ostrowski's Inequalities

Sanja Varošanec

Title Page

Contents



Go Back

Close

Quit

Page 20 of 20