

IDRISSA LY

Laboratoire de Mathématiques et Applications de Metz  
Ile du Saulcy, 57045 Metz Cedex 01, France.

EMail: [idrissa@math.univ-metz.fr](mailto:idrissa@math.univ-metz.fr)



---

volume 6, issue 3, article 91,  
2005.

*Received 12 February, 2005;*  
*accepted 17 June, 2005.*

Communicated by: *S.S. Dragomir*

---

[Abstract](#)

[Contents](#)



[Home Page](#)

[Go Back](#)

[Close](#)

[Quit](#)

## Abstract

In this paper, using the Hausdorff topology in the space of open sets under some capacity constraints on geometrical domains we prove the strong continuity with respect to the moving domain of the solutions of a  $p$ -Laplacian Dirichlet problem. We are also interested in the minimization of the first eigenvalue of the  $p$ -Laplacian with Dirichlet boundary conditions among open sets and quasi open sets of given measure.

*2000 Mathematics Subject Classification:* 35J70, 35P30, 35R35.

*Key words:*  $p$ -Laplacian, Nonlinear eigenvalue problems, Shape optimization.

## Contents

1	Introduction .....	3
2	Definition of the First and Second Eigenvalues .....	6
3	Properties of the Geometric Variations .....	7
4	Shape Optimization Result .....	12
5	Domain in Box .....	21
	References	



### The First Eigenvalue for the $p$ -Laplacian Operator

Idrissa Ly

Title Page

Contents



Go Back

Close

Quit

Page 2 of 28

# 1. Introduction

Let  $\Omega$  be an open subset of a fixed ball  $D$  in  $\mathbb{R}^N$ ,  $N \geq 2$  and  $1 < p < +\infty$ . Consider the Sobolev space  $W_0^{1,p}(\Omega)$  which is the closure of  $C^\infty$  functions compactly supported in  $\Omega$  for the norm

$$\|u\|_{1,p}^p = \int_{\Omega} |u(x)|^p dx + \int_{\Omega} |\nabla u(x)|^p dx.$$

The  $p$ -Laplacian is the operator defined by

$$\begin{aligned} \Delta_p : W_0^{1,p}(\Omega) &\longrightarrow W^{-1,q}(\Omega) \\ u &\longmapsto \Delta_p u = \operatorname{div}(|\nabla u|^{p-2} \nabla u), \end{aligned}$$

where  $W^{-1,q}(\Omega)$  is the dual space of  $W_0^{1,p}(\Omega)$  and we have  $1 < p, q < \infty$ ,  $\frac{1}{p} + \frac{1}{q} = 1$ .


We are interested in the nonlinear eigenvalue problem

$$(1.1) \quad \begin{cases} -\Delta_p u - \lambda |u|^{p-2} u &= 0 \text{ in } \Omega, \\ u &= 0 \text{ on } \partial\Omega. \end{cases}$$

Let  $u$  be a function of  $W_0^{1,p}(\Omega)$ , not identically 0. The function  $u$  is called an eigenfunction if

$$\int_{\Omega} |\nabla u(x)|^{p-2} \nabla u \nabla \phi dx = \lambda \int_{\Omega} |u(x)|^{p-2} u \phi dx$$

for all  $\phi \in C_0^\infty(\Omega)$ . The corresponding real number  $\lambda$  is called an eigenvalue.



The First Eigenvalue for the  $p$ -Laplacian Operator

Idrissa Ly

Title Page

Contents



Go Back

Close

Quit

Page 3 of 28

Contrary to the Laplace operator, the  $p$ -Laplacian spectrum has not been proved to be discrete. In [15], the first eigenvalue and the second eigenvalue are described.


Let  $D$  be a bounded domain in  $\mathbb{R}^N$  and  $c > 0$ . Let us denote  $\lambda_1^p(\Omega)$  as the first eigenvalue for the  $p$ -Laplacian operator. The aim of this paper is to study the isoperimetric inequality

$$\min\{\lambda_1^p(\Omega), \Omega \subseteq D \text{ and } |\Omega| = c\}$$

and its continuous dependence with respect to the domain. We extend the Rayleigh-Faber-Khran inequality to the  $p$ -Laplacian operator and study the minimization of the first eigenvalue in two dimensions when  $D$  is a box. By considering a class of simply connected domains, we study the stability of the minimizer  $\Omega_p$  of the first eigenvalue with respect to  $p$  that is if  $\Omega_p$  is a minimizer of the first eigenvalue for the  $p$ -Laplacian Dirichlet, when  $p$  goes to 2,  $\Omega_2$  is also a minimizer of the first eigenvalue of the Laplacian Dirichlet. Thus we will give a formal justification of the following conjecture: " $\Omega$  is a minimizer of given volume  $c$ , contained in a fixed box  $D$  and if  $D$  is too small to contain a ball of the same volume as  $\Omega$ . Are the free parts of the boundary of  $\Omega$  pieces of circle?"

Henrot and Oudet solved this question and proved by using the Hölmgren uniqueness theorem, that the free part of the boundary of  $\Omega$  cannot be pieces of circle, see [10].

The structure of this paper is as follows: The first section is devoted to the definition of two eigenvalues. In the second section, we study the properties of geometric variations for the first eigenvalue. The third section is devoted to the minimization of the first eigenvalue among open (or, if specified, quasi open)



## The First Eigenvalue for the $p$ -Laplacian Operator

Idrissa Ly

Title Page

Contents



Go Back


Close

Quit

Page 4 of 28

sets of given volume. In the fourth part we discuss the minimization of the first eigenvalue in a box in two dimensions.

Let  $D$  be a bounded open set in  $\mathbb{R}^N$  which contains all the open (or, if specified, quasi open) subsets used.



---

## The First Eigenvalue for the $p$ -Laplacian Operator

Idrissa Ly

---

Title Page

Contents



Go Back

Close

Quit

Page 5 of 28

## 2. Definition of the First and Second Eigenvalues

The first eigenvalue is defined by the nonlinear Rayleigh quotient

$$\lambda_1(\Omega) = \min_{\phi \in W_0^{1,p}(\Omega), \phi \neq 0} \frac{\int_{\Omega} |\nabla \phi(x)|^p dx}{\int_{\Omega} |\phi(x)|^p} = \frac{\int_{\Omega} |\nabla u_1(x)|^p dx}{\int_{\Omega} |u_1(x)|^p dx},$$

where the minimum is achieved by  $u_1$  which is a weak solution of the Euler-Lagrange equation

$$(2.1) \quad \begin{cases} -\Delta_p u - \lambda |u|^{p-2} u = 0 & \text{in } \Omega \\ u = 0 & \text{on } \partial\Omega. \end{cases}$$

The first eigenvalue has many special properties, it is strictly positive, simple in any bounded connected domain see [15]. And  $u_1$  is the only positive eigenfunction for the  $p$ -Laplacian Dirichlet see also [15].


In [15], the second eigenvalue is defined by

$$\lambda_2(\Omega) = \inf_{C \in \mathcal{C}_2} \max_C \frac{\int_{\Omega} |\nabla \phi(x)|^p dx}{\int_{\Omega} |\phi(x)|^p},$$

where

$$\mathcal{C}_2 := \{C \in W_0^{1,p}(\Omega) : C = -C \text{ such that } \text{genus}(C) \geq 2\}.$$

In [1], Anane and Tsouli proved that there does not exist any eigenvalue between the first and the second ones.



The First Eigenvalue for the  
 $p$ -Laplacian Operator

Idrissa Ly

Title Page

Contents



Go Back

Close

Quit

Page 6 of 28

### 3. Properties of the Geometric Variations

In this section we are interested in the continuity of the map

$$\Omega \longmapsto \lambda_1(\Omega).$$

Then, we have to fix topology on the space of the open subsets of  $D$ . On the family of the open subsets of  $D$ , we define the Hausdorff complementary topology, denoted  $H^c$  given by the metric

$$d_{H^c}(\Omega_1^c, \Omega_2^c) = \sup_{x \in \mathbb{R}^N} |d(x, \Omega_1^c) - d(x, \Omega_2^c)|.$$


The  $H^c$ -topology has some good properties for example the space of the open subsets of  $D$  is compact. Moreover if  $\Omega_n \xrightarrow{H^c} \Omega$ , then for any compact  $K \subset \subset \Omega$  we have  $K \subset \subset \Omega_n$  for  $n$  large enough.

However, perturbations in this topology may be very irregular and in general situations the continuity of the mapping  $\Omega \longmapsto \lambda_1(\Omega)$  fails, see [4].

In order to obtain a compactness result we impose some additional constraints on the space of the open subsets of  $D$  which are expressed in terms of the Sobolev capacity. There are many ways to define the Sobolev capacity, we use the local capacity defined in the following way.

**Definition 3.1.** For a compact set  $K$  contained in a ball  $B$ ,

$$\text{cap}(K, B) := \inf \left\{ \int_B |\nabla \phi|^p, \phi \in C_0^\infty(B), \phi \geq 1 \text{ on } K \right\}.$$



The First Eigenvalue for the  
 $p$ -Laplacian Operator

Idrissa Ly

Title Page

Contents



Go Back

Close

Quit

Page 7 of 28

### Definition 3.2.

1. It is said that a property holds  $p$ -quasi everywhere (abbreviated as  $p$ -q.e) if it holds outside a set of  $p$ -capacity zero.
2. A set  $\Omega \subset \mathbb{R}^N$  is said to be quasi open if for every  $\epsilon > 0$  there exists an open set  $\Omega_\epsilon$  such that  $\Omega \subseteq \Omega_\epsilon$ , and  $\text{cap}(\Omega_\epsilon \setminus \Omega) < \epsilon$ .
3. A function  $u : \mathbb{R}^N \longrightarrow \mathbb{R}$  is said  $p$ -quasi continuous if for every  $\epsilon > 0$  there exists an open set  $\Omega_\epsilon$  such that  $\text{cap}(\Omega_\epsilon) < \epsilon$  and  $u|_{\mathbb{R} \setminus \Omega_\epsilon}$  is continuous in  $\mathbb{R} \setminus \Omega_\epsilon$ .

It is well known that every Sobolev function  $u \in W^{1,p}(\mathbb{R}^N)$  has a  $p$ -quasi continuous representative which we still denote  $u$ . Therefore, level sets of Sobolev functions are  $p$ -quasi open sets; in particular  $\Omega_v = \{x \in D; |v(x)| > 0\}$  is quasi open subsets of  $D$ .


**Definition 3.3.** We say that an open set  $\Omega$  has the  $p - (r, c)$  capacity density condition if

$$\forall x \in \partial\Omega, \quad \forall 0 < \delta < r, \quad \frac{\text{cap}(\Omega^c \cap \bar{B}(x, \delta), B(x, 2\delta))}{\text{cap}(\bar{B}(x, \delta), B(x, 2\delta))} \geq c$$

where  $B(x, \delta)$  denotes the ball of radius  $\delta$ , centred at  $x$ .

**Definition 3.4.** We say that the sequence of the spaces  $W_0^{1,p}(\Omega_n)$  converges in the sense of Mosco to the space  $W_0^{1,p}(\Omega)$  if the following conditions hold

1. The first Mosco condition: For all  $\phi \in W_0^{1,p}(\Omega)$ , there exists a sequence  $\phi_n \in W_0^{1,p}(\Omega_n)$  such that  $\phi_n$  converges strongly in  $W_0^{1,p}(D)$  to  $\phi$ .



The First Eigenvalue for the  
 $p$ -Laplacian Operator

Idrissa Ly

Title Page

Contents



Go Back

Close

Quit

Page 8 of 28



2. *The second Mosco condition: For every sequence  $\phi_{n_k} \in W_0^{1,p}(\Omega_{n_k})$  weakly convergent in  $W_0^{1,p}(D)$  to a function  $\phi$ , we have  $\phi \in W_0^{1,p}(\Omega)$ .*

**Definition 3.5.** *We say a sequence  $(\Omega_n)$  of open subsets of a fixed ball  $D$   $\gamma_p$ -converges to  $\Omega$  if for any  $f \in W^{-1,q}(\Omega)$  the solutions of the Dirichlet problem*

$$-\Delta_p u_n = f \quad \text{in } \Omega_n, u_n \in W_0^{1,p}(\Omega_n)$$

*converge strongly in  $W_0^{1,p}(D)$ , as  $n \rightarrow +\infty$ , to the solution of the corresponding problem in  $\Omega$ , see [7], [8].*


By  $\mathcal{O}_{p-(r,c)}(D)$ , we denote the family of all open subsets of  $D$  which satisfy the  $p - (r, c)$  capacity density condition. This family is compact in the  $H^c$  topology see [4]. In [2], D. Bucur and P. Trebeschi, using capacity constraints analogous to those introduced in [3] and [4] for the linear case, prove the  $\gamma_p$ -compactness result for the  $p$ -Laplacian. In the same way, they extend the continuity result of Šverák [19] to the  $p$ -Laplacian for  $p \in (N - 1, N]$ ,  $N \geq 2$ . The reason of the choice of  $p$  is that in  $\mathbb{R}^N$  the curves have  $p$  positive capacity if  $p > N - 1$ . The case  $p > N$  is trivial since all functions in  $W^{1,p}(\mathbb{R}^N)$  are continuous.

Let us denote by

$$\mathcal{O}_l(D) = \{\Omega \subseteq D, \#\Omega^c \leq l\}$$

where  $\#$  denotes the number of the connected components. We have the following theorem.

**Theorem 3.1 (Bucur-Trebeschi).** *Let  $N \geq p > N - 1$ . Consider the sequence  $(\Omega_n) \subseteq \mathcal{O}_l(D)$  and assume that  $\Omega_n$  converges in Hausdorff complementary topology to  $\Omega$ . Then  $\Omega \subseteq \mathcal{O}_l(D)$  and  $\Omega_n$   $\gamma_p$ -converges to  $\Omega$ .*



The First Eigenvalue for the  $p$ -Laplacian Operator

Idrissa Ly

Title Page

Contents



Go Back

Close

Quit

Page 9 of 28

*Proof of Theorem 3.1.* See [2]. □

For  $N = 2$  and  $p = 2$ , Theorem 3.1 becomes the continuity result of Šverák [19].

Back to the continuity result, we use the above results to prove the following theorem.

**Theorem 3.2.** *Consider the sequence  $(\Omega_n) \subseteq \mathcal{O}_l(D)$ . Assume that  $\Omega_n$  converges in Hausdorff complementary topology to  $\Omega$ . Then  $\lambda_1(\Omega_n)$  converges to  $\lambda_1(\Omega)$ .*

*Proof of Theorem 3.2.* Let us take

$$\lambda_1(\Omega_n) = \min_{\phi_n \in W_0^{1,p}(\Omega_n), \phi_n \neq 0} \frac{\int_{\Omega_n} |\nabla \phi_n(x)|^p dx}{\int_{\Omega_n} |\phi_n(x)|^p} = \frac{\int_{\Omega_n} |\nabla u_n(x)|^p dx}{\int_{\Omega_n} |u_n(x)|^p},$$


where the minimum is attained by  $u_n$ , and

$$\lambda_1(\Omega) = \min_{\phi \in W_0^{1,p}(\Omega), \phi \neq 0} \frac{\int_{\Omega} |\nabla \phi(x)|^p dx}{\int_{\Omega} |\phi(x)|^p} = \frac{\int_{\Omega} |\nabla u_1(x)|^p dx}{\int_{\Omega} |u_1(x)|^p dx},$$

where the minimum is achieved by  $u_1$ .

By the Bucur and Trebeschi theorem,  $\Omega_n \xrightarrow{\gamma_p} \Omega$ . This implies  $W_0^{1,p}(\Omega_n)$  converges in the sense of Mosco to  $W_0^{1,p}(\Omega)$ .

If the sequence  $(u_n)$  is bounded in  $W_0^{1,p}(D)$ , then there exists a subsequence still denoted  $u_n$  such that  $u_n$  converges weakly in  $W_0^{1,p}(D)$  to a function  $u$ . The second condition of Mosco implies that  $u \in W_0^{1,p}(\Omega)$ .



The First Eigenvalue for the  $p$ -Laplacian Operator

Idrissa Ly

Title Page

Contents



Go Back

Close

Quit

Page 10 of 28

Using the weak lower semicontinuity of the  $L^p$ -norm, we have the inequality

$$\liminf_{n \rightarrow +\infty} \frac{\int_D |\nabla u_n(x)|^p dx}{\int_D |u_n(x)|^p} \geq \frac{\int_\Omega |\nabla u(x)|^p dx}{\int_\Omega |u(x)|^p} \geq \frac{\int_\Omega |\nabla u_1(x)|^p dx}{\int_\Omega |u_1(x)|^p},$$

then

$$(3.1) \quad \liminf_{n \rightarrow +\infty} \lambda_1(\Omega_n) \geq \lambda_1(\Omega).$$

Using the first condition of Mosco, there exists a sequence  $(v_n) \in W_0^{1,p}(\Omega_n)$  such that  $v_n$  converges strongly in  $W_0^{1,p}(D)$  to  $u_1$ .

We have

$$\lambda_1(\Omega_n) \leq \frac{\int_D |\nabla v_n(x)|^p dx}{\int_D |v_n(x)|^p}$$


this implies that

$$\begin{aligned} \limsup_{n \rightarrow +\infty} \lambda_1(\Omega_n) &\leq \limsup_{n \rightarrow +\infty} \frac{\int_D |\nabla v_n(x)|^p dx}{\int_D |v_n(x)|^p} \\ &= \lim_{n \rightarrow +\infty} \frac{\int_D |\nabla v_n(x)|^p dx}{\int_D |v_n(x)|^p} = \frac{\int_\Omega |\nabla u_1(x)|^p dx}{\int_\Omega |u_1(x)|^p} \end{aligned}$$

then

$$(3.2) \quad \limsup_{n \rightarrow +\infty} \lambda_1(\Omega_n) \leq \lambda_1(\Omega).$$

By the relations (3.1) and (3.2) we conclude that  $\lambda_1(\Omega_n)$  converges to  $\lambda_1(\Omega)$ . □



### The First Eigenvalue for the $p$ -Laplacian Operator

Idrissa Ly

Title Page

Contents



Go Back

Close

Quit

Page 11 of 28

## 4. Shape Optimization Result

We extend the classical inequality of Faber-Krahn for the first eigenvalue of the Dirichlet Laplacian to the Dirichlet  $p$ -Laplacian. We study this inequality when  $\Omega$  is a quasi open subset of  $D$ .

**Definition 4.1.** Let  $\Omega$  be an open subset and bounded in  $\mathbb{R}^N$ . We denote by  $B$  the ball centred at the origin with the same volume as  $\Omega$ . Let  $u$  be a non negative function in  $\Omega$ , which vanishes on  $\partial\Omega$ . For all  $c > 0$ , the set  $\{x \in \Omega, u(x) > c\}$  is called the level set of  $u$ .

The function  $u^*$  which has the following level set

$$\forall c > 0, \quad \{x \in B, u^*(x) > c\} = \{x \in \Omega, u(x) > c\}^*$$


is called the Schwarz rearrangement of  $u$ . The level sets of  $u^*$  are the balls that we obtain by rearranging the sets of the same volume of  $u$ .

We have the following lemma.

**Lemma 4.1.** Let  $\Omega$  be an open subset in  $\mathbb{R}^N$ . Let  $\psi$  be any continuous function on  $\mathbb{R}_+^*$ , we have

- $\int_{\Omega} \psi(u(x))dx = \int_{\Omega^*} \psi(u^*(x))dx$   $u^*$  is equi-mesurable with  $u$ .
- $\int_{\Omega} u(x)v(x)dx \leq \int_{\Omega^*} u^*(x)v^*(x)dx$ .
- If  $u \in W_0^{1,p}(\Omega), p > 1$  then  $u^* \in W_0^{1,p}(\Omega^*)$  and

$$\int_{\Omega} |\nabla u(x)|^p dx \geq \int_{\Omega^*} |\nabla u^*(x)|^p dx \quad \text{P\`olya inequality.}$$



The First Eigenvalue for the  $p$ -Laplacian Operator

Idrissa Ly

Title Page

Contents



Go Back

Close

Quit

Page 12 of 28

*Proof of Lemma 4.1.* See [12]. □

The basic result for the minimization of eigenvalues is the conjecture of Lord Rayleigh: “The disk should minimize the first eigenvalue of the Laplacian Dirichlet among every open set of given measure”. We extend the Rayleigh-Faber-Krahn inequality to the  $p$ -Laplacian operator.

Let  $\Omega$  be any open set in  $\mathbb{R}^N$  with finite measure. We denote by  $\lambda_1(\Omega)$  the first eigenvalue for the  $p$ -Laplacian operator with Dirichlet boundary conditions. We have the following theorem.

**Theorem 4.2.** *Let  $B$  be the ball of the same volume as  $\Omega$ , then*

$$\lambda_1(B) = \min\{\lambda_1(\Omega), \Omega \text{ open set of } \mathbb{R}^N, |\Omega| = |B|\}.$$

*Proof of Theorem 4.2.* Let  $u_1$  be the first eigenfunction of  $\lambda_1(\Omega)$ , it is strictly positive see [15]. By Lemma 4.1, equi-mesurability of the function  $u_1$  and its Schwarz rearrangement  $u_1^*$  gives


$$\int_{\Omega} |u_1(x)|^p dx = \int_B |u_1^*(x)|^p dx.$$

The Pòlya inequality implies that

$$\int_{\Omega} |\nabla u_1(x)|^p dx \geq \int_B |\nabla u_1^*(x)|^p dx.$$

By the two conditions, it becomes

$$\frac{\int_B |\nabla u_1^*(x)|^p dx}{\int_B |u_1^*(x)|^p dx} \leq \frac{\int_{\Omega} |\nabla u_1(x)|^p dx}{\int_{\Omega} |u_1(x)|^p dx} = \lambda_1(\Omega).$$




---

**The First Eigenvalue for the  
 $p$ -Laplacian Operator**

Idrissa Ly

---

Title Page

Contents



Go Back

Close

Quit

Page 13 of 28

This implies that

$$\lambda_1(B) = \min_{v \in W_0^{1,p}(B), v \neq 0} \frac{\int_B |\nabla v(x)|^p dx}{\int_B |v(x)|^p dx} \leq \frac{\int_B |\nabla u_1^*(x)|^p dx}{\int_B |u_1^*(x)|^p dx} \leq \lambda_1(\Omega).$$

□

**Remark 1.** *The solution  $\Omega$  must satisfy an optimality condition. We suppose that  $\Omega \in C^2$  – regular to compute the shape derivative. We deform the domain  $\Omega$  with respect to an admissible vector field  $V$  to compute the shape derivative*

$$dJ(\Omega; V) = \lim_{t \rightarrow 0} \frac{J(\text{Id} + t\Omega) - J(\Omega)}{t}.$$


We have the variation calculation

$$\begin{aligned} -\text{div}(|\nabla u|^{p-2} \nabla u) &= \lambda |u|^{p-2} u \\ -\int_{\Omega} \text{div}(|\nabla u|^{p-2} \nabla u) \phi dx &= \int_{\Omega} \lambda |u|^{p-2} u \phi dx, \quad \text{for all } \phi \in \mathcal{D}(\Omega) \\ \int_{\Omega} |\nabla u|^{p-2} \nabla u \nabla \phi dx &= \int_{\Omega} \lambda |u|^{p-2} u \phi dx, \quad \text{for all } \phi \in \mathcal{D}(\Omega) \end{aligned}$$

Let us take  $J(\Omega) = \int_{\Omega} |\nabla u|^{p-2} \nabla u \nabla \phi dx$  and  $J_1(\Omega) = \int_{\Omega} \lambda |u|^{p-2} u \phi dx$ . We have  $dJ(\Omega; V) = dJ_1(\Omega; V)$ .

We use the classical Hadamard formula to compute the Eulerian derivative of the functional  $J$  at the point  $\Omega$  in the direction  $V$ .

$$dJ(\Omega; V) = \int_{\Omega} (|\nabla u|^{p-2} \nabla u \nabla \phi)' dx + \int_{\Omega} \text{div}(|\nabla u|^{p-2} \nabla u \nabla \phi \cdot V(0)) dx.$$



The First Eigenvalue for the  
 $p$ -Laplacian Operator

Idrissa Ly

Title Page

Contents



Go Back

Close

Quit

Page 14 of 28

We have

$$\int_{\Omega} (|\nabla u|^{p-2} \nabla u \nabla \phi)' dx = \int_{\Omega} (|\nabla u|^{p-2})' \nabla u \nabla \phi dx + \int_{\Omega} |\nabla u|^{p-2} \nabla u (\nabla \phi)' dx + \int_{\Omega} |\nabla u|^{p-2} (\nabla u)' \nabla \phi dx.$$


We have the expression

$$\begin{aligned} (|\nabla u|^{p-2})' &= \left( (|\nabla u|^2)^{\frac{p-2}{2}} \right)' \\ &= \frac{p-2}{2} (|\nabla u|^2)' (|\nabla u|^2)^{\frac{p-4}{2}} \\ (|\nabla u|^{p-2})' &= (p-2) \nabla u \nabla u' |\nabla u|^{p-4}. \end{aligned}$$

Then

$$\begin{aligned} dJ(\Omega; V) &= (p-2) \int_{\Omega} |\nabla u|^{p-4} |\nabla u|^2 \nabla u' \nabla \phi dx \\ &\quad - \int_{\Omega} \operatorname{div}(|\nabla u|^{p-2} \nabla u) \phi' dx - \int_{\Omega} \operatorname{div}(|\nabla u|^{p-2} (\nabla u)') \phi dx \end{aligned}$$

$$\begin{aligned} dJ(\Omega, V) &= (p-2) \int_{\Omega} |\nabla u|^{p-2} \nabla u' \nabla \phi dx \\ &\quad - \int_{\Omega} \operatorname{div}(|\nabla u|^{p-2} \nabla u) \phi' dx - \int_{\Omega} \operatorname{div}(|\nabla u|^{p-2} (\nabla u)') \phi dx \end{aligned}$$



The First Eigenvalue for the  
 $p$ -Laplacian Operator

Idrissa Ly

Title Page

Contents



Go Back

Close

Quit

Page 15 of 28

because

$$\int_{\Omega} \operatorname{div}(|\nabla u|^{p-2} \nabla u \nabla \phi \cdot V(0)) dx = \int_{\partial\Omega} |\nabla u|^{p-2} \nabla u \nabla \phi \cdot V(0) \cdot \nu ds = 0.$$

We obtain

$$dJ(\Omega; V) = -(p-1) \int_{\Omega} \operatorname{div}(|\nabla u|^{p-2} \nabla u') \phi dx - \int_{\Omega} \operatorname{div}(|\nabla u|^{p-2} \nabla u) \phi' dx$$

We have also

$$\begin{aligned} dJ_1(\Omega; V) &= \int_{\Omega} \lambda' |u|^{p-2} u \phi dx + \int_{\Omega} \lambda |u|^{p-2} u' \phi dx \\ &\quad + \int_{\Omega} \lambda |u|^{p-2} u \phi' dx + (p-2) \int_{\Omega} \lambda |u|^{p-2} u' \phi dx, \end{aligned}$$


$$dJ_1(\Omega; V) = \int_{\Omega} \lambda' |u|^{p-2} u \phi dx + \int_{\Omega} \lambda |u|^{p-2} u \phi' dx + (p-1) \int_{\Omega} \lambda |u|^{p-2} u' \phi dx$$

$dJ(\Omega; V) = dJ_1(\Omega, V)$  implies

$$\begin{aligned} &-(p-1) \int_{\Omega} \operatorname{div}(|\nabla u|^{p-2} \nabla u') \phi dx - \int_{\Omega} \operatorname{div}(|\nabla u|^{p-2} \nabla u) \phi' dx \\ &= \int_{\Omega} \lambda' |u|^{p-2} u \phi dx + \int_{\Omega} \lambda |u|^{p-2} u \phi' dx + (p-1) \int_{\Omega} \lambda |u|^{p-2} u' \phi dx. \end{aligned}$$

By simplification we get

$$\begin{aligned} &-(p-1) \int_{\Omega} \operatorname{div}(|\nabla u|^{p-2} \nabla u') \phi dx - \int_{\Omega} \operatorname{div}(|\nabla u|^{p-2} \nabla u) \phi' dx \\ &= \int_{\Omega} \lambda' |u|^{p-2} u \phi dx + (p-1) \int_{\Omega} \lambda |u|^{p-2} u' \phi dx, \quad \text{for all } \phi \in \mathcal{D}(\Omega). \end{aligned}$$



## The First Eigenvalue for the $p$ -Laplacian Operator

Idrissa Ly

Title Page

Contents



Go Back

Close

Quit

Page 16 of 28



This implies that

$$(4.1) \quad \left\{ -(p-1) \operatorname{div}(|\nabla u|^{p-2} \nabla u') = \lambda' |u|^{p-2} u + (p-1) \lambda |u|^{p-2} u' \text{ in } \mathcal{D}'(\Omega) \right.$$

We multiply the equation (4.1) by  $u$  and by Green's formula we get

$$\begin{aligned} -(p-1) \left[ \int_{\Omega} \operatorname{div}(|\nabla u|^{p-2} \nabla u) u' dx + \int_{\partial\Omega} |\nabla u|^{p-2} \nabla u \cdot \nu u' ds \right] \\ = \lambda' + (p-1) \int_{\Omega} \lambda |u|^{p-2} u u' dx. \end{aligned}$$

Finally we obtain the expression of

$$\lambda'(\Omega; V) = -(p-1) \int_{\partial\Omega} |\nabla u|^p \nabla u \cdot \nu u' ds$$


where  $u'$  satisfies  $u' = -\frac{\partial u}{\partial \nu} V(0) \cdot \nu$  on  $\partial\Omega$ . Then

$$\lambda'(\Omega, V) = -(p-1) \int_{\partial\Omega} |\nabla u|^p V \cdot \nu ds.$$

We have a similar formula for the variation of the volume  $dJ_2(\Omega, V) = \int_{\partial\Omega} V \cdot \nu ds$ , where  $J_2(\Omega) = \int_{\Omega} dx - c$ .

If  $\Omega$  is an optimal domain then there exists a Lagrange multiplier  $a < 0$  such that

$$-(p-1) \int_{\partial\Omega} |\nabla u|^p V \cdot \nu ds = a \int_{\partial\Omega} V \cdot \nu ds.$$



The First Eigenvalue for the  
 $p$ -Laplacian Operator

Idrissa Ly

Title Page

Contents



Go Back

Close

Quit

Page 17 of 28

Then we obtain

$$|\nabla u| = \left( \frac{-a}{p-1} \right)^{\frac{1}{p}} \quad \text{on } \partial\Omega.$$

Since  $\Omega$  is  $C^2$ -regular and  $u = 0$  on  $\partial\Omega$ , then we get

$$-\frac{\partial u}{\partial \nu} = \left( \frac{-a}{p-1} \right)^{\frac{1}{p}} \quad \text{on } \partial\Omega.$$

We are also interested the existence of a minimizer for the following problem

$$\min\{\lambda_1(\Omega), \Omega \in \mathcal{A}, |\Omega| \leq c\},$$

where  $\mathcal{A}$  is a family of admissible domain defined by

$$\mathcal{A} = \{\Omega \subseteq D, \Omega \text{ is quasi open}\}$$


and  $\lambda_1(\Omega)$  is defined by

$$\lambda_1(\Omega) = \min_{\phi \in W_0^{1,p}(\Omega), \phi \neq 0} \frac{\int_{\Omega} |\nabla \phi(x)|^p dx}{\int_{\Omega} |\phi(x)|^p dx} = \frac{\int_{\Omega} |\nabla u_1(x)|^p dx}{\int_{\Omega} |u_1(x)|^p dx}.$$

The Sobolev space  $W_0^{1,p}(\Omega)$  is seen as a closed subspace of  $W_0^{1,p}(D)$  defined by

$$W_0^{1,p}(\Omega) = \{u \in W_0^{1,p}(D) : u = 0 \text{ p. q.e. on } D \setminus \Omega\}.$$

The problem is to look for weak topology constraints which would make the class  $\mathcal{A}$  sequentially compact. This convergence is called weak  $\gamma_p$ -convergence for quasi open sets.



## The First Eigenvalue for the $p$ -Laplacian Operator

Idrissa Ly

Title Page

Contents



Go Back

Close

Quit

Page 18 of 28

**Definition 4.2.** We say that a sequence  $(\Omega_n)$  of  $\mathcal{A}$  weakly  $\gamma_p$ -converges to  $\Omega \in \mathcal{A}$  if the sequence  $u_n$  converges weakly in  $W_0^{1,p}(D)$  to a function  $u \in W_0^{1,p}(D)$  (that we may take as quasi-continuous) such that  $\Omega = \{u > 0\}$ .

We have the following theorem.

**Theorem 4.3.** *The problem*

$$(4.2) \quad \min\{\lambda_1(\Omega), \Omega \in \mathcal{A}, |\Omega| \leq c\}$$

*admits at least one solution.*

*Proof of Theorem 4.3.* Let us take

$$\lambda_1(\Omega_n) = \min_{\phi_n \in W_0^{1,p}(\Omega_n), \phi_n \neq 0} \frac{\int_{\Omega_n} |\nabla \phi_n(x)|^p dx}{\int_{\Omega_n} |\phi_n(x)|^p dx} = \frac{\int_{\Omega_n} |\nabla u_n(x)|^p dx}{\int_{\Omega_n} |u_n(x)|^p dx}.$$


Suppose that  $(\Omega_n)_{(n \in \mathbb{N})}$  is a minimizing sequence of domain for the problem (4.2). We denote by  $u_n$  a first eigenfunction on  $\Omega_n$ , such that  $\int_{\Omega_n} |u_n(x)|^p dx = 1$ .

Since  $u_n$  is the first eigenfunction of  $\lambda_1(\Omega_n)$ ,  $u_n$  is strictly positive, cf [15], then the sequence  $(\Omega_n)$  is defined by  $\Omega_n = \{u_n > 0\}$ .

If the sequence  $(u_n)$  is bounded in  $W_0^{1,p}(D)$ , then there exists a subsequence still denoted by  $u_n$  such that  $u_n$  converges weakly in  $W_0^{1,p}(D)$  to a function  $u$ . By compact injection, we have that  $\int_{\Omega} |u(x)|^p dx = 1$ .

Let  $\Omega$  be quasi open and defined by  $\Omega = \{u > 0\}$ , this implies that  $u \in W_0^{1,p}(\Omega)$ . As the sequence  $(u_n)$  is bounded in  $W_0^{1,p}(D)$ , then

$$\liminf_{n \rightarrow +\infty} \frac{\int_{\Omega_n} |\nabla u_n(x)|^p dx}{\int_{\Omega_n} |u_n(x)|^p dx} \geq \frac{\int_{\Omega} |\nabla u(x)|^p dx}{\int_{\Omega} |u(x)|^p dx} \geq \frac{\int_{\Omega} |\nabla u_1(x)|^p dx}{\int_{\Omega} |u_1(x)|^p dx} = \lambda_1(\Omega).$$



The First Eigenvalue for the  $p$ -Laplacian Operator

Idrissa Ly

Title Page

Contents



Go Back


Close

Quit

Page 19 of 28

Now we show that  $|\Omega| \leq c$ .

We know that if the sequence  $\Omega_n$  weakly  $\gamma_p$ -converges to  $\Omega$  and the Lebesgue measure is weakly  $\gamma_p$ -lower semicontinuous on the class  $\mathcal{A}$  (see [5]), then we obtain  $|\{u > 0\}| \leq \liminf_{n \rightarrow +\infty} |\{u_n > 0\}| \leq c$  this implies that  $|\Omega| \leq c$ .  $\square$



---

### The First Eigenvalue for the $p$ -Laplacian Operator

Idrissa Ly

---

Title Page

Contents



Go Back

Close

Quit

Page 20 of 28

## 5. Domain in Box

Now let us take  $N = 2$ . We consider the class of admissible domains defined by

$$\mathcal{C} = \{\Omega, \Omega \text{ open subsets of } D \text{ and simply connected, } |\Omega| = c\}.$$

- For  $p > 2$ , the  $p$ -capacity of a point is strictly positive and every  $W_0^{1,p}$  function has a continuous representative. For this reason, a property which holds  $p - q.e$  with  $p > 2$  holds in fact everywhere. For  $p > 2$ , the domain  $\Omega_p$  is a minimizer of the problem  $\min\{\lambda_1^p(\Omega_p), \Omega_p \in \mathcal{C}\}$ .


Consider the sequence  $(\Omega_{p_n}) \subseteq \mathcal{C}$  and assume that  $\Omega_{p_n}$  converges in Hausdorff complementary topology to  $\Omega_2$ , when  $p_n$  goes to 2 and  $p_n > 2$ . Then  $\Omega_2 \subseteq \mathcal{C}$  and  $\Omega_{p_n}$   $\gamma_2$ -converges to  $\Omega_2$ .

By the Sobolev embedding theorem, we have  $W_0^{1,p_n}(\Omega_{p_n}) \hookrightarrow H_0^1(\Omega_{p_n})$ . The  $\gamma_2$ -convergence implies that  $H_0^1(\Omega_{p_n})$  converges in the sense of Mosco to  $H_0^1(\Omega_2)$ . For  $p_n > 2$ , by the Hölder inequality we have

$$\left( \int |\nabla u_{p_n}|^2 dx \right)^{\frac{1}{2}} \leq |\Omega_{p_n}|^{\frac{1}{2} - \frac{1}{p_n}} \left( \int |\nabla u_{p_n}|^{p_n} dx \right)^{\frac{1}{p_n}}$$

$$\left( \int |\nabla u_{p_n}|^2 dx \right)^{\frac{1}{2}} dx \leq c^{\frac{1}{2} - \frac{1}{p_n}} \lambda_1^{p_n}(\Omega_{p_n}).$$

Then the sequence  $(u_{p_n})$  is uniformly bounded in  $H_0^1(\Omega_{p_n})$ . There exists a subsequence still denoted  $u_{p_n}$  such that  $u_{p_n}$  converges weakly in  $H_0^1(D)$  to a function  $u$ . The second condition of Mosco implies that  $u \in H_0^1(\Omega_2)$ .



The First Eigenvalue for the  
 $p$ -Laplacian Operator

Idrissa Ly

Title Page

Contents




Go Back

Close

Quit

Page 21 of 28



**The First Eigenvalue for the  
 $p$ -Laplacian Operator**

Idrissa Ly

Title Page

Contents



Go Back

Close

Quit

Page 22 of 28

For  $p > 2$ , we have the Sobolev embedding theorem  $W_0^{1,p}(D) \hookrightarrow C^{0,\alpha}(\bar{D})$ .

Ascoli's theorem implies that  $u_{p_n} \rightarrow u$  and  $\nabla u_{p_n} \rightarrow \nabla u$  locally uniformly in  $\Omega_2$ , when  $p_n$  goes to 2 and  $p_n > 2$ .

Now show that

$$\lim_{p_n \rightarrow 2} \int |u_{p_n}|^2 dx = 1 \quad \text{i.e.} \quad \int |u|^2 dx = 1.$$


For  $\epsilon > 0$  small, we have  $p_n > 2 - \epsilon$ . Noting that

$$\begin{aligned} \left( \int |\nabla u_{p_n}|^{2-\epsilon} dx \right)^{\frac{1}{2-\epsilon}} &\leq |\Omega_{p_n}|^{\frac{1}{2-\epsilon} - \frac{1}{p_n}} \left( \int |\nabla u_{p_n}|^{p_n} dx \right)^{\frac{1}{p_n}} \\ &= c^{\frac{1}{2-\epsilon} - \frac{1}{p_n}} \lambda_1^{p_n}(\Omega_{p_n}), \end{aligned}$$

this implies that the sequence  $u_{p_n}$  is uniformly bounded in  $W_0^{1,2-\epsilon}(\Omega_{p_n})$ . Then there exists a subsequence still denoted  $u_{p_n}$  such that  $u_{p_n}$  is weakly convergent in  $H_0^1(D)$  to  $u$ . By the second condition of Mosco we get  $u \in W_0^{1,2-\epsilon}(\Omega_2)$ . It follows that

$$\begin{aligned} \int |u|^{2-\epsilon} dx &= \lim_{p_n \rightarrow 2} \int |u_{p_n}|^{2-\epsilon} dx \\ &\leq \lim_{p_n \rightarrow 2} |\Omega_{p_n}|^{1 - \frac{2-\epsilon}{p_n}} \left( \int |u_{p_n}|^{p_n} dx \right)^{\frac{2-\epsilon}{p_n}} = c^\epsilon. \end{aligned}$$

Letting  $\epsilon \rightarrow 0$ , we obtain  $\int |u|^2 dx \leq 1$ .



**The First Eigenvalue for the  $p$ -Laplacian Operator**

Idrissa Ly

Title Page

Contents



Go Back

Close

Quit

Page 23 of 28

On the other hand, Lemma 4.2 of [14] implies that

$$\int |u|^{p_n} dx \geq \int |u_{p_n}|^{p_n} dx + p_n \int |u|^{p_n-2} dx u_{p_n} (u - u_{p_n}).$$

The second integral on the right-hand side approaches 0 as  $p_n \rightarrow 2$ . Thus we get  $\int |u|^2 dx \geq 1$ , and we conclude that  $\int |u|^2 dx = 1$ .

In [11, Theorem 2.1 p. 3350],  $\lambda_k^p$  is continuous in  $p$  for  $k = 1, 2$ , where  $\lambda_k^p$  is the  $k$ -th eigenvalue for the  $p$ -Laplacian operator.

We have

$$(5.1) \quad \int |\nabla u_{p_n}|^{p_n-2} \nabla u_{p_n} \nabla \phi dx = \int \lambda_1^{p_n} |u_{p_n}|^{p_n-2} u_{p_n} \phi dx, \quad \text{for all } \phi \in \mathcal{D}(\Omega_2).$$

Letting  $p_n$  go to 2,  $p_n > 2$  in (5.1), and noting that  $u_{p_n}$  converges uniformly to  $u$  on the support of  $\phi$ , we obtain

$$\int \nabla u \nabla \phi dx = \int \lambda_1^2 u \phi dx, \quad \text{for all } \phi \in \mathcal{D}(\Omega_2),$$

whence we have

$$\begin{cases} -\Delta u = \lambda_1^2 u & \text{in } \mathcal{D}'(\Omega_2) \\ u = 0 & \text{on } \partial\Omega_2. \end{cases}$$

We conclude that when  $p \rightarrow 2$  and  $p > 2$  the free parts of the boundary of  $\Omega_p$  cannot be pieces of circle.

- For  $p \leq 2$ , we consider the sequence  $(\Omega_{p_n}) \subseteq \mathcal{C}$  and assume that  $\Omega_{p_n}$  converges in Hausdorff complementary topology to  $\Omega_2$ , when  $p_n$  goes to 2 and  $p_n \leq 2$ . Then by Theorem 3.1, we get  $\Omega_2 \subseteq \mathcal{C}$  and  $\Omega_{p_n} \gamma_{2-\epsilon}$ -converges to  $\Omega_2$ .

In [16], the sequence  $(u_{p_n})$  is bounded in  $W^{1,2-\epsilon}(D)$ ,  $0 < \epsilon < 1$  that is  $\nabla u_{p_n}$  converges weakly in  $L^{2-\epsilon}(D)$  to  $\nabla u$  and  $u_{p_n}$  converges strongly in  $L^{2-\epsilon}(D)$  to  $u$ . In [16], we get also  $\int |\nabla u|^2 dx \leq \beta$  and  $\int |u|^2 dx < \infty$ .

By Lemma 4.2 of [14], we have

$$\int |u|^{p_n} dx \geq \int |u_{p_n}|^{p_n} dx + p_n \int |u|^{p_n-2} dx u_{p_n} (u - u_{p_n}).$$


The second integral on the right-hand side approaches 0 as  $p_n \rightarrow 2$ . Thus we get  $\int |u|^2 dx \geq 1$ . This implies that

$$\lim_{p_n \rightarrow 2} \int |u_{p_n}|^{2-\epsilon} dx = \int |u|^{2-\epsilon} dx = 1.$$

Letting  $\epsilon \rightarrow 0$ , we obtain  $\int |u|^2 dx = 1$ .

The  $\gamma_{2-\epsilon}$ -convergence implies that  $u_{p_n}$  converges strongly in  $W_0^{1,2-\epsilon}(\Omega_2)$  to  $u$ . According to P. Lindqvist see [16], we have  $u \in H^1(D)$ , and we can deduce that  $u \in H_0^1(\Omega_2)$ . As the first eigenvalue for the  $p$ -Laplacian operator is continuous in  $p$  cf [11], we have

$$(5.2) \quad \int |\nabla u_{p_n}|^{p_n-2} \nabla u_{p_n} \nabla \phi dx = \int \lambda_1^{p_n} |u_{p_n}|^{p_n-2} u_{p_n} \phi dx, \quad \text{for all } \phi \in \mathcal{D}(\Omega_2).$$



## The First Eigenvalue for the $p$ -Laplacian Operator

Idrissa Ly

Title Page

Contents



Go Back

Close

Quit

Page 24 of 28




Letting  $p_n$  go to 2,  $p_n \leq 2$  in (5.2), and noting that  $u_{p_n}$  converges uniformly to  $u$  on the support of  $\phi$ , we obtain

$$\int \nabla u \nabla \phi dx = \int \lambda_1^2 u \phi dx, \quad \text{for all } \phi \in \mathcal{D}(\Omega_2),$$

whence we have

$$\begin{cases} -\Delta u = \lambda_1^2 u & \text{in } \mathcal{D}'(\Omega_2) \\ u = 0 & \text{on } \partial\Omega_2. \end{cases}$$

We conclude that when  $p \rightarrow 2$  and  $p \leq 2$  the free parts of the boundary of  $\Omega_p$  cannot be pieces of circle.




---

The First Eigenvalue for the  
 $p$ -Laplacian Operator

Idrissa Ly

---

Title Page

Contents



Go Back


Close

Quit

Page 25 of 28

## References

- [1] A. ANANE AND N. TSOULI, The second eigenvalue of the  $p$ -Laplacian, *Nonlinear Partial Differential Equations* (Fès, 1994), Pitman Research Notes Math. Ser., 343, Longman, Harlow, (1996), 1–9.
- [2] D. BUCUR AND P. TREBESCHI, Shape optimization problem governed by nonlinear sate equation, *Proc. Roy. Edinburgh*, **128 A**(1998), 945–963.
- [3] D. BUCUR AND J.P. ZOLÉSIO, Wiener’s criterion and shape continuity for the Dirichlet problem, *Boll. Un. Mat. Ital.*, **B 11** (1997), 757–771.
- [4] D. BUCUR AND J.P. ZOLÉSIO, N-Dimensional shape Optimization under Capacity Constraints, *J. Differential Equations*, **123**(2) (1995), 504–522.
- [5] D. BUCUR AND G. BUTTAZZO, *Variational Methods in Shape Optimization Problems*, SNS Pisa 2002.
- [6] G. BUTTAZZO AND G. DAL MASO, An existence result for a class of shape optimization problems, *Arch. Rational. Mech. Anal.*, **122** (1993), 183–195.
- [7] G. DAL MASO,  $\Gamma$ -convergence and  $\mu$  capacities, *Ann. Scuola Norm. Sup. Pisa*, **14** (1988), 423–464.
- [8] G. DAL MASO AND A. DEFRANCESCHI, Limits of nonlinear Dirichlet problems in varying domains, *Manuscripta Math.*, **61** (1988), 251–278.



---

The First Eigenvalue for the  
 $p$ -Laplacian Operator

Idrissa Ly

---

Title Page

Contents




Go Back

Close

Quit

Page 26 of 28

- [9] G. FABER, Bewiw, dass unter allen homogenen Membranen von gleicher Fläche und gleicher Spannung die kreisförmige den tiefsten Grundton gibt, *Sitz. Ber. Bayer. Akad. Wiss.*, (1923), 169–172.
- [10] A. HENROT AND E. OUDET, Minimizing the second eigenvalue of the Laplace operator with Dirichlet boundary conditions, *Arch. Ration. Mech. Anal.*, **169**(1) (2003), 73–87.
- [11] Y.X. HUANG, On the eigenvalue of the  $p$ -Laplacian with varying  $p$ , *Proc. Amer. Math. Soc.*, **125**(11) (1997), 3347–3354.
- [12] B. KAWOHL, *Rearrangements and Convexity of Level Sets in PDE*, (Lecture Notes In Mathematics 1150), Springer-Verlag, Heidelberg.
- [13] E. KRAHN, Über eine von Rayleigh formulierte Minimaleigenschaft des Kreises, *Math. Ann.*, **94** (1924), 97–100.
- [14] P. LINDQVIST, On the equation  $\operatorname{div}(|\nabla u|^{p-2}\nabla u) + \lambda|u|^{p-2}u = 0$ , *Proc. Amer. Math. Soc.*, **109**(1) (1990), 157–164. Addendum, *ibidem*, **116**(2) (1992), 583–584.
- [15] P. LINDQVIST, On a nonlinear eigenvalue problem, *Berichte Univ. Jyväskylä Math. Inst.*, **68** (1995), 33–54.
- [16] P. LINDQVIST, Stability for the solutions of  $\operatorname{div}(|\nabla u|^{p-2}\nabla u) = f$  with varying  $p$ , *J. Math. Anal. Appl.*, **127**(1) (1987), 93–102.
- [17] J. SIMON, Differential with respect to the domain in boundary value problems, *Numer. Funct. Anal. and Optimiz.*, **2**(7-8) (1980), 649–687.




---

The First Eigenvalue for the  
 $p$ -Laplacian Operator

Idrissa Ly

---

Title Page

Contents




Go Back

Close

Quit

Page 27 of 28

- [18] J. SOKOŁOWSKI AND J.P. ZOLESIO, *Introduction to Shape Optimization. Shape Sensitivity Analysis*. Springer series in Computational Mathematics, 16. Springer-Verlag, Berlin (1992).
- [19] V. ŠVERÀK, On optimal shape design, *J. Math. Pures Appl.*, **72** (1993), 537–551.



---

**The First Eigenvalue for the  
 $p$ -Laplacian Operator**

Idrissa Ly

---

Title Page

Contents



Go Back

Close

Quit

Page 28 of 28