

Journal of Inequalities in Pure and Applied Mathematics

BOUNDS FOR LINEAR RECURRENCES WITH RESTRICTED COEFFICIENTS

KENNETH S. BERENHAUT AND ROBERT LUND

Department of Mathematics
Wake Forest University
Winston-Salem, NC 27109.
EMail: berenhks@wfu.edu

Department of Statistics
The University of Georgia
Athens, GA 30602-1952.
EMail: lund@stat.uga.edu

©2000 Victoria University
ISSN (electronic): 1443-5756
054-02



volume 4, issue 2, article 26,
2003.

*Received 16 May, 2002;
accepted 14 February, 2003.*

Communicated by: I. Pressman

[Abstract](#)

[Contents](#)



[Home Page](#)

[Go Back](#)

[Close](#)

[Quit](#)

Abstract

This paper derives inequalities for general linear recurrences. Optimal bounds for solutions to the recurrence are obtained when the coefficients of the recursion lie in intervals that include zero. An important aspect of the derived bounds is that they are easily computable. The results bound solutions of triangular matrix equations and coefficients of ratios of power series.

2000 Mathematics Subject Classification: 39A10, 30B10, 15A45, 15A24, 11B37.

Key words: Recurrence, Restricted Coefficients, Power Series, Triangular Matrices.

We are grateful to Andrew Granville for many helpful suggestions. The comments made by the referee greatly improved readability of discourse. The authors acknowledge financial support from NSF Grant DMS 0071383.

Contents

1	Introduction	3
2	Results	7
3	Triangular Linear Systems with Restricted Entries	12
4	Ratios of Power Series	14
5	Proofs	18
	References	



Bounds for Linear Recurrences with Restricted Coefficients

Kenneth S. Berenhaut and
Robert Lund

Title Page

Contents



Go Back

Close

Quit

Page 2 of 34

1. Introduction

This paper derives bounds for solutions to the linear recurrence

$$(1.1) \quad b_n = \sum_{k=1}^{n-1} \alpha_{n,k} b_k, \quad n \geq 2.$$


Throughout, we assume that $b_1 \neq 0$ as $b_1 = 0$ implies that $b_n = 0$ for all $n \geq 2$. Our results bound $\{b_n\}_{n=1}^{\infty}$ in a term-by-term manner with a second order time-homogeneous linear recursion that is readily analyzable.

Our motivation for studying (1.1) lies in applied probability. There it is useful to have a bound for coefficients of a ratio of power series when limited information is available on the constituent series (cf. Kijima [14], Kendall [13], Heathcote [11], Feller [6]). The series comprising the ratio are often probability generating functions. Linear algebra is another setting where (1.1) arises.

Example 1.1. *What is the largest $|b_5|$ possible in (1.1) when $b_1 = -1$ and $\alpha_{n,k} \in [-3, 0]$ for all n and k ? In Section 2, we show that $|b_5| \leq 99$ for such situations, and that this value is produced by $\alpha_{n,k}$ having the alternating form*

$$(1.2) \quad \begin{array}{cccc} & \alpha_{n,1} & \alpha_{n,2} & \alpha_{n,3} & \alpha_{n,4} \\ n = 2 & -3 & & & \\ n = 3 & 0 & -3 & & \\ n = 4 & -3 & 0 & -3 & \\ n = 5 & 0 & -3 & 0 & -3 \end{array} .$$

Specifically, these $\alpha_{n,k}$ give $b_2 = 3, b_3 = -9, b_4 = 30, \text{ and } b_5 = -99$. We return to this example in Section 2.



**Bounds for Linear Recurrences
with Restricted Coefficients**

Kenneth S. Berenhaut and
Robert Lund

Title Page

Contents



Go Back

Close

Quit

Page 3 of 34

Example 1.2. For a fixed $I \subset \mathfrak{R}$, let \mathcal{F}_I be the set of I -power series defined by

$$(1.3) \quad \mathcal{F}_I = \left\{ f : f(z) = 1 + \sum_{k=1}^{\infty} a_k z^k \text{ and } a_k \in I \text{ for each } k \geq 1 \right\}.$$

Flatto, Lagarias, and Poonen [7] and Solomyak [22] proved independently that if z is a root of a series in $\mathcal{F}_{[0,1]}$, then $|z| \geq 2/(1 + \sqrt{5})$. As $z = -2/(1 + \sqrt{5})$ is a root of $1 + z + z^3 + z^5 + \dots$, this bound is tight over $\mathcal{F}_{[0,1]}$. The coefficients of the multiplicative inverse of a series in $\mathcal{F}_{[0,1]}$ cannot increase at a rate larger than the golden ratio.


We will show later that the coefficients of the multiplicative inverse of a power series in $\mathcal{F}_{[0,1]}$ are bounded by the ubiquitous Fibonacci numbers. This gives a “first constant” for the aforementioned rate. Observe that

$$(1.4) \quad \left(1 + \sum_{n=1}^{\infty} z^{2n-1} \right)^{-1} = 1 - z + z^2 - 2z^3 + 3z^4 - 5z^4 + \dots,$$

the coefficients on the right hand side of (1.4) having the magnitude of the Fibonacci numbers. Hence, the first constant is also good. We return to this setting in Section 4.

Example 1.3. Consider the lower triangular linear system $L\vec{x} = \vec{b}$ where L is the 10×10 matrix with (i, j) th entry

$$(1.5) \quad L_{i,j} = \begin{cases} 1, & \text{if } i = j \\ 10, & \text{if } i > j \\ 0, & \text{if } i < j \end{cases},$$



**Bounds for Linear Recurrences
with Restricted Coefficients**

Kenneth S. Berenhaut and
Robert Lund

Title Page

Contents



Go Back

Close

Quit

Page 4 of 34


and the i th component of \vec{b} is $b_i = i^2$ for $1 \leq i \leq 10$. The exact solution is

$$(1.6) \quad \vec{x} = \begin{bmatrix} 1 \\ -6 \\ 59 \\ -524 \\ 4725 \\ -42514 \\ 382639 \\ -3443736 \\ 30993641 \\ -278942750 \end{bmatrix} = L^{-1}\vec{b}.$$

The condition number of L is 26633841560.0; this essentially drives the rate of growth of x_i in i (cf. Trefethen and Bau [23] for general discussion). Our results will imply that all matrix equations $L\vec{x} = \vec{b}$, with L an $n \times n$ unit lower triangular matrix with $L_{i,j} \in [0, 10]$ for $1 \leq i < j \leq n$ and $|b_i| \leq i^2$, have solutions whose i th component x_i is bounded by (coefficients rounded to three decimal places)

$$(1.7) \quad |x_i| \leq (0.142) 10.099^i + 3.538 (-0.099)^i - 0.400 i + 0.320, \quad 1 \leq i \leq n.$$

The first four values of the right hand side of (1.7) are 1, 14, 145, and 1472. These show essentially the same order of magnitude as the x_i 's; hence the bound is performing reasonably. We return to this example in Section 3.



**Bounds for Linear Recurrences
with Restricted Coefficients**

Kenneth S. Berenhaut and
Robert Lund

Title Page

Contents



Go Back

Close

Quit


Page 5 of 34

Recurrences with varying or random coefficients have been studied by many previous authors. A partial survey of such literature contains Viswanath [24] and [25], Viswanath and Trefethen [26], Embree and Trefethen [5], Wright and Trefethen [28], Mallik [16], Popenda [20], Kittapa [15], and Odlyzko [19].

Our methods of proof are based on a careful analysis of sign changes in solutions to (1.1). This differs considerably from past authors, who typically take a more analytic approach. An advantage of our discourse is that it is entirely elementary, discrete, and self-contained. A disadvantage of our arguments lie with laborious bookkeeping.

Study of (1.1) could alternatively be based on linear algebraic or analytic techniques. Some of the applications considered here, namely solutions of linear matrix equations and coefficients of ratios of power series, are indeed classical problems. However, linear algebraic and analytic techniques have yielded disappointing explicit bounds to date. Hence, this paper explores alternative methods.

The rest of this paper proceeds as follows. Section 2 presents the main theorem, some variants of this result, and discussion of the hypotheses and optimality. Sections 3 and 4 consider application of the results to lower triangular linear systems and coefficients of ratios of power series, respectively. Proofs are deferred to Section 5. There, a simple case of our main result is first proven to convey the logic of our sign change analyses.



Bounds for Linear Recurrences with Restricted Coefficients

Kenneth S. Berenhaut and
Robert Lund

Title Page

Contents



Go Back

Close

Quit

Page 6 of 34

2. Results

The general form of our main result is the following.


Theorem 2.1. *Suppose that $A \geq 1$ and $0 \leq B \leq A$ are constants and that $\{D_n\}_{n=2}^\infty$ is a nondecreasing sequence of nonnegative real numbers. Suppose that the coefficients in (1.1) are restricted to intervals: $\alpha_{n,1} \in [-D_n, D_n]$ for $n \geq 2$ and $\alpha_{n,k} \in [-A, B]$ for $n \geq 2$ and $2 \leq k \leq n - 1$. Then solutions to (1.1) satisfy $|b_n|/|b_1| \leq U_n$ for all $n \geq 1$, where*

$$(2.1) \quad U_n = \begin{cases} 1, & \text{if } n = 1 \\ D_2, & \text{if } n = 2 \\ AD_2 + D_3, & \text{if } n = 3 \\ AU_{n-1} + (1 + B)U_{n-2} + D_n - D_{n-2}, & \text{if } n > 3 \end{cases}.$$

Neglecting the bookkeeping complications induced by a general $\{D_n\}$, the difference equation in (2.1) is second-order, time-homogeneous, and linear. In many cases, one can solve (2.1) explicitly for U_n . As such, we view U_n as being “easy to compute”. The generality added by a non-decreasing $\{D_n\}$ is relevant in probabilistic settings where generalized renewal equations are common (cf. Feller [6] and Heathcote [11]).

For cases where asymmetric bounds on $\alpha_{n,1}$ are available, we offer the following.

Theorem 2.2. *Suppose that $A \geq 1$ and that $C \geq 0$ and $D \geq 0$. If $\alpha_{n,1} \in [-C, D]$ and $\alpha_{n,k} \in [-A, 0]$ for all $n \geq 2$ and $2 \leq k \leq n - 1$, then $|b_n|/|b_1| \leq$*



**Bounds for Linear Recurrences
with Restricted Coefficients**

Kenneth S. Berenhaut and
Robert Lund

Title Page

Contents



Go Back

Close

Quit

Page 7 of 34

U_n for all $n \geq 1$, where

$$(2.2) \quad U_n = \begin{cases} 1, & \text{if } n = 1 \\ \max(C, D), & \text{if } n = 2 \\ A \max(C, D) + \min(C, D), & \text{if } n = 3 \\ AU_{n-1} + U_{n-2}, & \text{if } n > 3 \end{cases}.$$

Theorems 2.1 and 2.2 are proven in Section 5. There, we first prove the results in the simple setting where $A = C = \Delta > 1$, $D = 0$, and $b_1 = -1$ to convey the basic ideas of a sign change analysis. In particular, we prove the following Corollary.

Corollary 2.3. *Suppose that $b_1 = -1$ and that $\alpha_{n,k} \in [-\Delta, 0]$ for all n, k where $\Delta \geq 1$. Then $|b_n| \leq U_n$ for all $n \geq 1$, where $\{U_n\}$ satisfies*


$$(2.3) \quad U_n = \begin{cases} \Delta^{n-1}, & \text{if } n \leq 2 \\ \Delta U_{n-1} + U_{n-2}, & \text{if } n \geq 3 \end{cases}.$$

Solving (2.3) explicitly for U_n gives

$$(2.4) \quad U_n = \frac{\Delta}{\sqrt{\Delta^2 + 4}} \left(r_1^{n-1} - \left(-\frac{1}{r_1} \right)^{n-1} \right),$$

for $n \geq 2$, where r_1 is the root

$$(2.5) \quad r_1 = \frac{\Delta + \sqrt{\Delta^2 + 4}}{2}$$



**Bounds for Linear Recurrences
with Restricted Coefficients**

Kenneth S. Berenhaut and
Robert Lund

Title Page

Contents



Go Back

Close

Quit

Page 8 of 34

of the characteristic polynomial associated with (2.3). The other root of the characteristic polynomial in (2.3) is $r_2 = 2^{-1}(\Delta - \sqrt{\Delta^2 + 4})$. Observe that $|r_1| > |r_2|$ and $r_1 r_2 = -1$.

The flexibility allowed in bounds for $\alpha_{n,1}$ in Theorems 2.1 and 2.2 comes at a bookkeeping price during the proof in Section 5. The benefits of such generality will become apparent in Sections 3 and 4 where we bound solutions of nonhomogeneous (rather than merely homogeneous) matrix equations and the coefficients of power series ratios (rather than merely reciprocals).

This section concludes with some comments on the assumptions and optimality of Theorems 2.1 and 2.2.

Remark 2.1. (Optimality of Theorems 2.1 and 2.2). For a given b_1 , $\{D_n\}_{n=2}^\infty$, A , and B , the bound in (2.1) cannot be improved upon. To see this, set

$$(2.6) \quad \alpha_{n,1} = \begin{cases} -D_n & \text{if } n \text{ is odd} \\ D_n & \text{if } n \text{ is even} \end{cases}$$


and

$$(2.7) \quad \alpha_{n,k} = \begin{cases} -A & \text{if } n+k \text{ is odd} \\ B & \text{if } n+k \text{ is even} \end{cases}$$

for $n \geq 2$ and $1 < k \leq n-1$. It is easy to verify from (1.1) that $b_n = (-1)^n U_n b_1$ for $n \geq 2$, implying that the bound in Theorem 2.1 is achieved. A similar construction shows that the bound in Theorem 2.2 is also optimal.

For completeness, we also consider situations where $0 \leq A \leq B$. In this case, a straightforward analysis will yield the following bound for solutions to (1.1).

Remark 2.2.



**Bounds for Linear Recurrences
with Restricted Coefficients**

Kenneth S. Berenhaut and
Robert Lund

Title Page

Contents



Go Back

Close

Quit

Page 9 of 34

Consider the setup in Theorem 2.1 except that $0 \leq A \leq B$. Then $\{U_n^*\}_{n=1}^\infty$ defined by

$$(2.8) \quad U_n^* = \begin{cases} 1, & \text{if } n = 1 \\ D_2, & \text{if } n = 2 \\ BD_2 + D_3, & \text{if } n = 3 \\ (B + 1)U_{n-1}^* + D_n - D_{n-1}, & \text{if } n > 3 \end{cases}$$


is a bound satisfying $|b_n|/|b_1| \leq U_n^*$ for all $n \geq 1$. This bound is achieved in the case where $\alpha_{n,1} = D_n$ and $\alpha_{n,k} = B$ for $n \geq 2$ and $2 \leq k \leq n - 1$.

The above results provide optimal bounds for $|b_n|$ when $\alpha_{n,k} \in [-A, B]$ except when $0 \leq B < A < 1$. As our next remark shows, the condition $A \geq 1$ is essential for optimality.

Remark 2.3. *Optimality of Theorem 2.1 may not occur when $A < 1$. To see this, suppose that $B < A < 1$ and consider $\{b_n\}_{n=1}^\infty$ satisfying (1.1) with $b_1 = -1$, $\alpha_{2,1} = D_2$, $\alpha_{3,1} = D_3$, $\alpha_{3,2} = B$, $\alpha_{4,1} = -D_4$, $\alpha_{4,2} = -A$, and $\alpha_{4,3} = -A$. Then (1.1) gives $b_2 = -D_2$, $b_3 = -(BD_2 + D_3)$, and*

$$(2.9) \quad \begin{aligned} b_4 &= D_4 + A(BD_2 + D_3) + AD_2 \\ &= (A + AB)D_2 + AD_3 + D_4 \\ &> (A^2 + B)D_2 + AD_3 + D_4, \end{aligned}$$

where the strict inequality above follows from $A + AB > A^2 + B$ (which follows



Bounds for Linear Recurrences with Restricted Coefficients

Kenneth S. Berenhaut and
Robert Lund

Title Page

Contents



Go Back

Close

Quit


Page 10 of 34

from $B < A < 1$). Applying (2.1) now gives

$$\begin{aligned}
 b_4 &> A(AD_2 + D_3) + (B + 1)D_2 + D_4 - D_2 \\
 &= AU_3 + (1 + B)U_2 + D_4 - D_2 \\
 (2.10) \quad &= U_4.
 \end{aligned}$$

Hence, U_n may not bound $|b_n|$ in this setting.

Example 2.1. In the setting of Example 1.1, the $\{\alpha_{n,k}\}$ producing the maximal $\{|b_n|\}$ are obtained via the argument in Remark 2.1. When $\alpha_{n,k} \in [-3, 0]$ for all n and k , the maximal $|b_n|$'s are produced with $\alpha_{n,k}$ either -3 or 0 in the alternating fashion depicted in the table in Example 1.1.



**Bounds for Linear Recurrences
with Restricted Coefficients**

Kenneth S. Berenhaut and
Robert Lund

Title Page

Contents



Go Back

Close

Quit

Page 11 of 34

3. Triangular Linear Systems with Restricted Entries

Theorems 2.1 and 2.2 have applications to systems of linear equations. Consider the lower triangular linear system

$$(3.1) \quad \begin{bmatrix} l_{1,1} & 0 & \cdots & 0 \\ l_{2,1} & l_{2,2} & \ddots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ l_{n,1} & l_{n,2} & \cdots & l_{n,n} \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} = \begin{bmatrix} c_1 \\ c_2 \\ \vdots \\ c_n \end{bmatrix},$$

with $l_{i,i} \neq 0$ for $1 \leq i \leq n$. Solving this for $\{x_j\}$ gives


$$(3.2) \quad x_m = \frac{c_m}{l_{m,m}} x_0 - \sum_{k=1}^{m-1} \frac{l_{m,k}}{l_{m,m}} x_k, \quad 1 \leq m \leq n,$$

with $x_0 = 1$. Letting $b_{m+1} = x_m$ for $0 \leq m \leq n$ produces

$$(3.3) \quad b_{m+1} = \frac{c_m}{l_{m,m}} b_1 - \sum_{k=2}^m \frac{l_{m,k-1}}{l_{m,m}} b_k$$

which is (1.1) with $\alpha_{m,1} = c_{m-1}/l_{m-1,m-1}$ and $\alpha_{m,k} = -l_{m-1,k-1}/l_{m-1,m-1}$ for $2 \leq k \leq m-1$. Hence, Theorems 2.1 and 2.2 become the following.

Corollary 3.1. *Consider the linear system in (3.1). Suppose that $0 \leq B \leq A$ and that D_k is nondecreasing in k . Then*



Bounds for Linear Recurrences
with Restricted Coefficients

Kenneth S. Berenhaut and
Robert Lund

Title Page

Contents



Go Back

Close

Quit

Page 12 of 34

(i) If $c_i/l_{i,i} \in [-D_{i+1}, D_{i+1}]$ for $1 \leq i \leq n$ and $l_{i,j}/l_{i,i} \in [-B, A]$ for $2 \leq i \leq n$ and $1 \leq j \leq i$, then $|x_i| \leq U_{i+1}$ for $2 \leq i \leq n$ where $\{U_k\}$ is as in (2.1).

(ii) If $c_i/l_{i,i} \in [-C, D]$ for $1 \leq i \leq n$ and $l_{i,j}/l_{i,i} \in [0, A]$ for $2 \leq i \leq n$ and $1 \leq j \leq i$, then $|x_i| \leq U_{i+1}$ for $1 \leq i \leq n$ where $\{U_k\}$ is as in (2.2).

Example 3.1. Returning to Example 1.3, the bound in (1.7) follows from Part (i) of Corollary 3.1 with $D_i = (i - 1)^2$, $A = 10$, and $B = 0$. The difference equation in (2.1) simplifies to


$$(3.4) \quad U_n = 10U_{n-1} + U_{n-2} + 4n - 8.$$

Corollary 3.1 compares favorably to the bounds for matrix equation solutions with coefficients that are restricted to more general intervals in Neumaier [17], Hansen [9] and [8], Hansen and Smith [10], and Kearfott [12]. Here, optimal bounds are obtained regardless of interval widths and dimension; moreover, the computational burden is limited to solving the second-order linear recurrences in (2.1) or (2.2).

If $c_i = 0$ for $i \geq 2$ in (3.1) (this situation is discussed further in Viswanath and Trefethen [26]), then (3.2) is

$$(3.5) \quad x_m = - \sum_{k=1}^{m-1} \frac{l_{m,k}}{l_{m,m}} x_k, \quad 1 \leq m \leq n,$$

with $x_1 = c_1/l_{1,1}$. One can now bound $|x_n|$ via Theorem 2.1 or 2.2.



**Bounds for Linear Recurrences
with Restricted Coefficients**

Kenneth S. Berenhaut and
Robert Lund

Title Page

Contents



Go Back

Close

Quit

Page 13 of 34

4. Ratios of Power Series

The recurrence equation (1.1) arises when computing coefficients of ratios of formal power series. Equating coefficients in the expansion

$$(4.1) \quad h_0 + h_1z + h_2z^2 + \cdots = \frac{g_0 + g_1z + g_2z^2 + \cdots}{f_0 + f_1z + f_2z^2 + \cdots}$$

(take $f_0 = 1$ and $g_0 = 1$ for simplicity) gives $h_0 = 1$ and


$$(4.2) \quad h_n = (g_n - f_n)h_0 - \sum_{j=1}^{n-1} f_{n-j}h_j, \quad n \geq 1.$$

The theorems in Section 2 translate to the following.

Corollary 4.1. *Suppose that $0 \leq B \leq A$, that $\{D_n\}_{n=2}^{\infty}$ is a nondecreasing sequence of nonnegative real numbers, and that $\{f_n\}_{n=0}^{\infty}$, $\{g_n\}_{n=0}^{\infty}$, and $\{h_n\}_{n=0}^{\infty}$ satisfy (4.1) with $f_0 = g_0 = 1$.*

- (i) *If $g_n - f_n \in [-D_{n+1}, D_{n+1}]$ for all $n \geq 1$ and $f_n \in [-B, A]$ for all $n \geq 0$, then $|h_n| \leq U_{n+1}$ for all $n \geq 0$ where $\{U_n\}_{n=1}^{\infty}$ is as in (2.1).*
- (ii) *If $g_n - f_n \in [-C, D]$ for $n \geq 1$ and $f_n \in [0, A]$ for $n \geq 0$, then $|h_n| \leq U_{n+1}$ for $n \geq 0$ where $\{U_n\}_{n=1}^{\infty}$ is as in (2.2).*

Merely inverting a power series simplifies the statements in Corollary 4.1. Here, $g_k = 0$ for all $k \geq 1$ and $g_0 = 1$. Using this in (4.2), applying Part (i) of Corollary 3.1 (with $D_n \equiv A$) and Part (ii) of Corollary 3.1 (with $C = A$ and $D = 0$), and solving (2.1) and (2.2) for $\{U_n\}$ gives the following results.



**Bounds for Linear Recurrences
with Restricted Coefficients**

Kenneth S. Berenhaut and
Robert Lund

Title Page

Contents



Go Back

Close

Quit

Page 14 of 34

Corollary 4.2. Suppose that $0 \leq B \leq A$ and that $g_k = 0$ for $k \geq 1$ and $g_0 = 1$.
Let

$$(4.3) \quad r_1 = \frac{A + \sqrt{A^2 + 4(1+B)}}{2}$$

be a root of the characteristic polynomial in (2.1).

(i) If $f_n \in [-B, A]$ for all $n \geq 0$, then

$$(4.4) \quad |h_n| \leq \kappa_1 r_1^n + \kappa_2 \left[\frac{-(B+1)}{r_1} \right]^n$$

for all $n \geq 1$ where

$$(4.5) \quad \kappa_1 = \frac{2(B+1) - A + \sqrt{A^2 + 4(1+B)}}{2(B+1)\sqrt{A^2 + 4(1+B)}},$$


and

$$(4.6) \quad \kappa_2 = -\frac{2(B+1) - A - \sqrt{A^2 + 4(1+B)}}{2(B+1)\sqrt{A^2 + 4(1+B)}}.$$

(ii) If $f_n \in [0, A]$ for all $n \geq 0$ ($B = 0$), then

$$(4.7) \quad |h_n| \leq \frac{A}{\sqrt{A^2 + 4}} r_1^n - \frac{A}{\sqrt{A^2 + 4}} \left[\frac{-1}{r_1} \right]^n$$

for all $n \geq 1$.



**Bounds for Linear Recurrences
with Restricted Coefficients**

Kenneth S. Berenhaut and
Robert Lund

Title Page

Contents



Go Back

Close

Quit

Page 15 of 34

Remark 4.1. Corollary 4.2 (ii) is optimal as the bound is attained for $f(z) = 1 + Az + Az^3 + Az^5 + \dots$. Regarding the sharpness of Corollary 4.2 (i), set $f(z) = 1 + Az - Bz^2 + Az^3 + \dots$. If u_n is taken to be the bound on the right hand side of (4.4) then it is not difficult to show that u_n and h_n are similar in magnitude: $u_n/|h_n| \leq 1 + 2(A - B)/A^2$ and

$$(4.8) \quad \lim_{n \rightarrow \infty} \frac{u_n}{|h_n|} = 1 + \frac{(A - B)(\sqrt{A^2 + 4(1 + B)} - A)}{AB + 2A + B\sqrt{A^2 + 4(1 + B)}}.$$


Hence, the rate is again sharp.

Corollaries 4.1 and 4.2 are useful when generating functions or formal power series are utilized such as in enumerative combinatorics and stochastic processes (cf. Wilf [27], Feller [6], Kijima [14]).

The above results provide bounds for the location of the smallest root of a complex valued power series. Power series with restricted coefficients have been studied in the context of determining distributions of zeroes (cf. Flatto *et al.* [7], Solomyak [22], Beaucoup *et al.* [1], [2], and Pinner [21]). Related problems for polynomials have been considered by Odlyzko and Poonen [18], Yamamoto [29], Borwein and Pinner [4], and Borwein and Erdelyi [3]. As mentioned above, Flatto *et al.* [7] and Solomyak [22] independently proved that if z is a root of a series in $\mathcal{F}_{[0,1]}$, then $|z| \geq 2/(1 + \sqrt{5})$. The following extension of this result is a consequence of Corollary 4.2.

Corollary 4.3. If z is a root of a power series in $\mathcal{F}_{[-B,A]}$ with $0 \leq B \leq A$, then

$$(4.9) \quad |z| \geq \frac{2}{A + \sqrt{A^2 + 4(1 + B)}}.$$



**Bounds for Linear Recurrences
with Restricted Coefficients**

Kenneth S. Berenhaut and
Robert Lund

Title Page

Contents



Go Back


Close

Quit

Page 16 of 34

Proof. Suppose that $f \in \mathcal{F}_{[-B,A]}$. Apply Part (i) of Corollary 4.2 and note from (4.4) that $f(z)^{-1}$ is finite for $|z| < r_1^{-1}$ (Observe that r_1 is the root of the characteristic polynomial with largest magnitude). If f had a root in $\{z : |z| < r_1^{-1}\}$, say at $z = z_0$, then we would have the contradiction $|f(z_0)|^{-1} = \infty$. \square

The result in Corollary 4.3 is again optimal: for given $0 \leq B \leq A$, $f(z) = 1 + Az - Bz^2 + Az^3 - Bz^4 + \dots$ has a root at $z = -r_1^{-1}$.



**Bounds for Linear Recurrences
with Restricted Coefficients**

Kenneth S. Berenhaut and
Robert Lund

Title Page

Contents



Go Back

Close

Quit


Page 17 of 34

5. Proofs

This section proves Theorem 2.1. As the arguments for Theorem 2.2 are similar, we concentrate on Theorem 2.1 only. While the proof of Theorem 2.1 is self-contained and elementary, it does employ a “sign change analysis” of $\{b_n\}_{n=1}^{\infty}$ which is case-by-case intensive and delicate. Attempts to find a direct analytic argument, by other authors as well as ourselves, have been unsuccessful to date. In particular, standard manipulations with classical inequalities do not yield the sharpness or generality of Theorem 2.1. The rudimentary structure of the problem emerges with the sign change arguments. Moreover, the arguments provide both a convergence rate and explicit “first constant” bound for the rate. Obtaining an explicit first constant, a practical matter needed to apply the bounds, takes considerably more effort in general.

The sign-change arguments below first bound all solutions to (1.1) that have a particular sign configuration; in the notation below, this is $|b_n| \leq |B_n|$ for all $n \geq 1$. A subsequent analysis is needed to bound $|B_n|$ by an accessible quantity; in the notation below, this is $|B_n| \leq U_n$ where U_n is defined in (2.1). We first consider the arguments for Corollary 2.3 as these are reasonably brief and convey the essence of the general analysis.

Arguments for Corollary 2.3. Suppose that $b_1 = -1$ and let $P = \{n \geq 1 : b_n \geq 0\}$ and $N = \{n \geq 1 : b_n < 0\}$ partition the sign configuration of $\{b_n\}_{n=1}^{\infty}$. Now define B_n recursively in n from N and P via $B_1 = -1$ and



Bounds for Linear Recurrences
with Restricted Coefficients

Kenneth S. Berenhaut and
Robert Lund

Title Page

Contents



Go Back

Close

Quit

Page 18 of 34

$$(5.1) \quad B_n = \begin{cases} \Delta - \Delta \sum_{\substack{2 \leq r \leq n-1 \\ r \in N}} B_r, & n \in P \\ -\Delta \sum_{\substack{2 \leq r \leq n-1 \\ r \in P}} B_r, & n \in N \end{cases}$$

for $n \geq 2$. A simple induction with (5.1) will show that B_n and b_n have the same sign for $n \geq 1$.

We now prove by induction that $|b_n| \leq |B_n|$ for all $n > 1$. First, assume that $n > 1$ and that $n \in P$. Returning to (1.1) and collecting positive and negative terms gives


$$(5.2) \quad b_n = \alpha_{n,1}b_1 + \sum_{\substack{2 \leq r \leq n-1 \\ r \in P}} \alpha_{n,r}b_r + \sum_{\substack{2 \leq r \leq n-1 \\ r \in N}} \alpha_{n,r}b_r.$$

Using $b_1 = -1$, the bound $\alpha_{n,k} \in [-\Delta, 0]$ for all n, k , and neglecting the first summation in (5.2) gives

$$(5.3) \quad \begin{aligned} b_n &\leq \Delta + \sum_{\substack{2 \leq r \leq n-1 \\ r \in N}} -\Delta b_r \\ &= \Delta + \Delta \sum_{\substack{2 \leq r \leq n-1 \\ r \in N}} |b_r|. \end{aligned}$$

Using the inductive hypothesis and the fact that $|b_n| = b_n$ in (5.3) produces

$$|b_n| \leq \Delta + \Delta \sum_{\substack{2 \leq r \leq n-1 \\ r \in N}} |B_r|$$



Title Page

Contents




Go Back

Close

Quit

Page 19 of 34



Title Page

Contents



Go Back

Close

Quit

Page 20 of 34

$$= \Delta - \Delta \sum_{\substack{2 \leq r \leq n-1 \\ r \in N}} B_r$$

$$(5.4) \quad = B_n$$

after (5.1) is applied. An analogous argument works when $n \in N$.

We now finish the arguments for Corollary 2.3 by inductively showing that $|B_n| \leq U_n$ from (5.1). First, it is easy to verify that $|B_i| \leq U_i$, for $1 \leq i \leq 3$ for all possible sign configurations of $\{B_1, B_2, B_3\}$. Now assume that $n \in P$ ($B_n \geq 0$) where $n > 3$. If $n - 1 \in P$ ($B_{n-1} \geq 0$), then $B_n = B_{n-1}$ by (5.1) and $|B_n| = |B_{n-1}| \leq U_{n-1} \leq U_n$ since U_n is nondecreasing in n (this follows from $\Delta \geq 1$). So we need only consider the case where $n - 1 \in N$ ($B_{n-1} < 0$). If $r \in N$ for all $r \leq n - 1$ ($B_r < 0$ for $1 \leq r \leq n - 1$), then $B_2 = B_3 = \dots = B_{n-1} = 0$ by (5.1) and we have $B_n = \Delta = U_3 \leq U_n$.

Finally, consider the case where a non-negative element in $\{B_1, \dots, B_{n-2}\}$ exists; that is, $r \in P$ for some $2 \leq r \leq n - 2$. Let r^* be the largest such integer and set $k = n - r^* - 1$. For signs of $\{B_n\}$, we have $B_{n-k-1} \geq 0$ ($B_{n-k-1} \in P$) and $B_j < 0$ for $n - k \leq j \leq n - 1$. Using these in (5.1) gives $B_{n-1} = \dots = B_{n-k}$. Applying (5.1) yet again produces

$$\begin{aligned} B_n &= \Delta - \Delta \sum_{\substack{2 \leq r \leq n-1 \\ r \in N}} B_r \\ &= \Delta - \Delta \sum_{r=n-k}^{n-1} B_r - \Delta \sum_{\substack{2 \leq r \leq n-k-2 \\ r \in N}} B_r \end{aligned}$$

$$(5.5) \quad = B_{n-k-1} - \Delta k B_{n-k}.$$

Applying the induction hypothesis and the triangle inequality in (5.5) produces

$$(5.6) \quad |B_n| \leq U_{n-k-1} + \Delta k U_{n-k},$$

and the difference equation in (2.3) can be used to increase the smallest subscript appearing on the right hand side of (5.6) to $n - k$:

$$(5.7) \quad |B_n| \leq U_{n-k+1} + \Delta(k-1)U_{n-k}.$$

Since U_n is nondecreasing in n and $\Delta(k-1) \geq 1$, we may swap the coefficients on U_{n-k+1} and U_{n-k} in (5.7) to obtain

$$(5.8) \quad |B_n| \leq U_{n-k} + \Delta(k-1)U_{n-k+1}.$$

Note that (5.8) is (5.6) with k replaced by $k-1$. As the discourse from (5.6) – (5.8) is merely algebraic, we iterate the above arguments to obtain

$$(5.9) \quad |B_n| \leq U_{n-(k-j)-1} + \Delta(k-j)U_{n-(k-j)}$$


for each $0 \leq j \leq k-1$. In particular, taking $j = k-1$ in (5.9) now gives

$$(5.10) \quad |B_n| \leq U_{n-2} + \Delta U_{n-1}.$$

Applying (2.3) in (5.10) immediately gives the required bound $|B_n| \leq U_n$ and finishes our work. The arguments for the case where $n \in N$ are similar.

□

Following the logic of the above arguments, we now present the proof of Theorem 2.1 in its generality.



**Bounds for Linear Recurrences
with Restricted Coefficients**

Kenneth S. Berenhaut and
Robert Lund

Title Page

Contents



Go Back

Close

Quit

Page 21 of 34

Proof of Theorem 2.1. We first reduce to the case where $b_1 = -1$ by examining b_n/b_1 . Again let $P = \{n \geq 1 : b_n \geq 0\}$ and $N = \{n \geq 1 : b_n < 0\}$ be the sign partition for $\{b_n\}_{n=1}^\infty$. This time, define a bounding sequence $\{B_n\}_{n=1}^\infty$ for this sign configuration recursively in n via $B_1 = -1$, and for $n \geq 2$ by

$$(5.11) \quad B_n = \begin{cases} D_n - A \sum_{\substack{2 \leq r \leq n-1 \\ r \in N}} B_r + B \sum_{\substack{2 \leq r \leq n-1 \\ r \in P}} B_r, & n \in P \\ -D_n - A \sum_{\substack{2 \leq r \leq n-1 \\ r \in P}} B_r + B \sum_{\substack{2 \leq r \leq n-1 \\ r \in N}} B_r, & n \in N \end{cases}.$$


As before, an induction will show that B_n and b_n have the same sign for each $n \geq 1$. This fact will be used repeatedly in the discourse below.

We now justify the majorizing properties of $\{B_n\}$ by inductively showing that $|b_n| \leq |B_n|$ for all $n \geq 1$. First, consider the case where $n \in P$. Now partition positive and negative terms in (1.1) and apply the bounds assumed on the $\alpha_{n,k}$'s in Theorem 2.1 to get

$$(5.12) \quad b_n \leq -D_n b_1 + B \sum_{\substack{2 \leq r \leq n-1 \\ r \in P}} b_r - A \sum_{\substack{2 \leq r \leq n-1 \\ r \in N}} b_r.$$

Applying $b_1 = -1$ and the induction hypothesis, and then (5.11) gives

$$b_n \leq D_n + B \sum_{\substack{2 \leq r \leq n-1 \\ r \in P}} |B_r| + A \sum_{\substack{2 \leq r \leq n-1 \\ r \in N}} |B_r|$$



Bounds for Linear Recurrences with Restricted Coefficients

Kenneth S. Berenhaut and
Robert Lund

Title Page

Contents




Go Back

Close

Quit

Page 22 of 34



Title Page

Contents



Go Back

Close

Quit

Page 23 of 34

$$= D_n + B \sum_{\substack{2 \leq r \leq n-1 \\ r \in P}} B_r - A \sum_{\substack{2 \leq r \leq n-1 \\ r \in N}} B_r$$

$$(5.13) \quad = B_n.$$

Similar arguments tackle the case where $n \in N$. Equation (5.13) represents the core of our arguments. The remainder of our work lies with devising a useful bound for the B_n 's in (5.11).

To complete the proof of Theorem 2.1, it remains to show that $|B_n| \leq U_n$ for all $n \geq 1$. For this it will be convenient to have the following technical lemma which we prove after the arguments for Theorem 2.1 (one can verify non-circularity of discourse).

Lemma 5.1. *Consider the setup in Theorem 2.1 and define $\{E_n\}_{n=1}^\infty$ via $E_0 = 1$, $E_1 = A$, $E_2 = A^2 + B$, and $E_j = AE_{j-1} + (1 + B)E_{j-2}$ for $j \geq 3$. Then U_n can be expressed as*

$$(5.14) \quad U_n = D_n + \sum_{j=2}^{n-1} E_{n-j} D_j,$$

for $n \geq 2$, with the inequality

$$(5.15) \quad U_n - (1 + B)U_{n-1} \geq D_n - D_{n-1}$$

holding for $n \geq 3$. Finally, in the case where $n \geq 2$ and $B_j < 0$ for $1 \leq j \leq n - 1$ ($j \in N$ for $1 \leq j \leq n - 1$) and $B_n \geq 0$ ($n \in P$), we have

$$(5.16) \quad B_n = D_n + \sum_{j=2}^{n-1} A(1 + B)^{n-j-1} D_j.$$

We now return to the proof of Theorem 2.1. Assume first that $n \in P$ ($B_n > 0$). We start inductive verification that $|B_j| \leq U_j$ for all $j \geq 1$ by noting that $|B_1| = U_1 = 1$ and $|B_2| = D_2 = U_2$. For B_3 , first note that if $B_2 \geq 0$ and $B_3 \geq 0$ ($\{2, 3\} \subset P$), then

$$(5.17) \quad \begin{aligned} |B_3| &= D_3 + BD_2 \\ &\leq U_3, \end{aligned}$$

where the inequality in (5.17) follows from (2.1), $D_j \geq 0$ for all j , and $B \leq A$. In the case where $B_2 < 0$ and $B_3 < 0$ ($\{2, 3\} \subset N$), then (5.17) again holds. In the cases where there is one negative and one positive sign amongst $\{B_2, B_3\}$, one can verify that

$$(5.18) \quad \begin{aligned} |B_3| &= D_3 + AD_2 \\ &\leq U_3 \end{aligned}$$


by direct application of (2.1).

Now assume that $|B_k| \leq U_k$ for $1 \leq k \leq n-1$. When $n-1 \in P$ ($B_{n-1} \geq 0$), use (5.11) to get

$$(5.19) \quad B_n = (1 + B)B_{n-1} + D_n - D_{n-1}.$$

Applying the induction hypothesis that $B_{n-1} \leq U_{n-1}$ and (5.15) in (5.19) produces

$$(5.20) \quad \begin{aligned} B_n &\leq (1 + B)U_{n-1} + D_n - D_{n-1} \\ &\leq U_n \end{aligned}$$



**Bounds for Linear Recurrences
with Restricted Coefficients**

Kenneth S. Berenhaut and
Robert Lund

Title Page

Contents



Go Back

Close

Quit

Page 24 of 34

as claimed.

It remains to consider the case where $n - 1 \in N$. First suppose that $r \in N$ for all $r \leq n - 1$. From Lemma 5.1, $E_1 = A$ and $E_2 = A^2 + B \geq A(1 + B)$ since $A \geq 1$ and $A \geq B$. Using $A \geq B$ and Lemma 5.1 in an induction argument will easily verify the inequality $E_j \geq A(1 + B)^{j-1}$ for all $j \geq 1$. Comparing coefficients in (5.16) and (5.14) now yields $|B_n| \leq U_n$ as claimed.


Having dealt with the case where the B_j are negative for all $1 \leq j \leq n - 1$, now suppose that there exists a non-negative B_j amongst the first $n - 1$ indices. In particular, suppose that $r \in P$ for some $2 \leq r \leq n - 2$ and let r^* denote the largest such integer. Set $k = n - r^* - 1$. For signs of $\{B_n\}$, we have $B_{n-k-1} \geq 0$ ($n - k - 1 \in P$), $B_j < 0$ ($j \in N$ for $n - k \leq j \leq n - 1$), and our standing assumption that $B_n \geq 0$ ($n \in P$). Using these facts in (5.11) produces

$$(5.21) \quad B_n = D_n - A \sum_{n-k \leq r \leq n-1} B_r + A \sum_{\substack{2 \leq r \leq n-k-2 \\ r \in N}} B_r + B \sum_{\substack{2 \leq r \leq n-k-2 \\ r \in P}} B_r + B|B_{n-k-1}|.$$

Now combine the definition of B_{n-k-1} in (5.11) with (5.21) to get

$$(5.22) \quad B_n = D_n + (1 + B)|B_{n-k-1}| - D_{n-k-1} - A \sum_{n-k \leq r \leq n-1} B_r.$$

Returning to (5.11) with the fact that $B_j < 0$ for $n - k \leq j \leq n - 1$ identifies



**Bounds for Linear Recurrences
with Restricted Coefficients**

Kenneth S. Berenhaut and
Robert Lund

Title Page

Contents



Go Back

Close

Quit

Page 25 of 34

the rightmost summation in (5.22):


$$(5.23) \quad \sum_{n-k \leq r \leq n-1} B_r = -|B_{n-k}| \sum_{i=0}^{k-1} (1+B)^i + D_{n-k} \sum_{i=0}^{k-2} (1+B)^i - \sum_{i=1}^{k-1} (1+B)^{i-1} D_{n-i}.$$

Combining (5.22) and (5.23) expresses B_n explicitly in terms of B_{n-k} and B_{n-k-1} :

$$(5.24) \quad B_n = D_n + (1+B)|B_{n-k-1}| - D_{n-k-1} + A|B_{n-k}| \sum_{i=0}^{k-1} (1+B)^i - AD_{n-k} \sum_{i=0}^{k-2} (1+B)^i + A \sum_{i=1}^{k-1} (1+B)^{i-1} D_{n-i}.$$

The induction hypothesis gives $|B_{n-k-1}| \leq U_{n-k-1}$ and $|B_{n-k}| \leq U_{n-k}$; using these in (5.24) along with $B_n = |B_n|$ gives the bound

$$(5.25) \quad |B_n| \leq D_n + (1+B)U_{n-k-1} - D_{n-k-1} + AU_{n-k} \sum_{i=0}^{k-1} (1+B)^i - AD_{n-k} \sum_{i=0}^{k-2} (1+B)^i + A \sum_{i=1}^{k-1} (1+B)^{i-1} D_{n-i}.$$



Bounds for Linear Recurrences with Restricted Coefficients

Kenneth S. Berenhaut and Robert Lund

Title Page

Contents



Go Back

Close

Quit

Page 26 of 34

Making the substitution $J_i = A \sum_{m=0}^i (1+B)^m$ into (5.25) now yields

$$(5.26) \quad |B_n| \leq D_n + (1+B)U_{n-k-1} - D_{n-k-1} + U_{n-k}J_{k-1} \\ - D_{n-k}J_{k-2} + A \sum_{i=1}^{k-1} (1+B)^{i-1} D_{n-i}.$$

The difference equation (2.1) gives

$$U_{n-k+1} = AU_{n-k} + (1+B)U_{n-k-1} + D_{n-k+1} - D_{n-k-1}.$$

Using this in (5.26) and algebraically simplifying produces

$$(5.27) \quad |B_n| \leq U_{n-k+1} - D_{n-k+1} + (1+B)U_{n-k}J_{k-2} \\ + D_n - D_{n-k}J_{k-2} + A \sum_{i=1}^{k-1} (1+B)^{i-1} D_{n-i},$$

where the fact that $J_{k-1} - A = (1+B)J_{k-2}$ has been applied. An algebraic rearrangement of the right hand side of (5.27) now produces

$$(5.28) \quad |B_n| \leq (1 - J_{k-2})[U_{n-k+1} - (1+B)U_{n-k}] \\ + J_{k-2}U_{n-k+1} + (1+B)U_{n-k} - D_{n-k+1} \\ + D_n - D_{n-k}J_{k-2} + A \sum_{i=1}^{k-1} (1+B)^{i-1} D_{n-i}.$$



**Bounds for Linear Recurrences
with Restricted Coefficients**

Kenneth S. Berenhaut and
Robert Lund

Title Page

Contents



Go Back

Close

Quit

Page 27 of 34

Noting that $J_{k-2} \geq 1$ for all k and applying (5.15) to the bracketed term in the right hand side of (5.28) now produces

$$|B_n| \leq (1 - J_{k-2})[D_{n-k+1} - D_{n-k}] + J_{k-2}U_{n-k+1} + (1 + B)U_{n-k} - D_{n-k+1} + D_n - D_{n-k}J_{k-2} + A \sum_{i=1}^{k-1} (1 + B)^{i-1} D_{n-i}.$$

Invoking the difference equation in (2.1) again will give

$$(5.29) \quad |B_n| \leq U_{n-k+2} - D_{n-k+2} + (1 + B)U_{n-k+1}J_{k-3} + D_n - D_{n-k+1}J_{k-3} + A \sum_{i=1}^{k-2} (1 + B)^{i-1} D_{n-i}.$$

The discourse between (5.27) – (5.29) is purely algebraic, justified via the difference equation in (2.1). Observe that the bounds for $|B_n|$ in (5.27) and (5.29) are similar in form, except that k is replaced by $k - 1$. As such, one can continue iterating the arguments in (5.27) – (5.29) until $k = 3$. This will give

$$(5.30) \quad |B_n| \leq U_{n-1} - D_{n-1} + (1 + B)U_{n-2}J_0 + D_n - D_{n-2}J_0 + AD_{n-1}.$$

Now use $J_0 = A$ in (5.30), employ (2.1) and regroup terms to get

$$(5.31) \quad |B_n| \leq U_n + D_{n-2} + (1 - A)[U_{n-1} - (1 + B)U_{n-2}] - D_{n-1} - D_{n-2}A + AD_{n-1}.$$



**Bounds for Linear Recurrences
with Restricted Coefficients**

Kenneth S. Berenhaut and
Robert Lund

Title Page

Contents



Go Back

Close

Quit

Page 28 of 34

Applying (5.20) once more to the bracketed terms in (5.31) and $A \geq 1$ to get

$$\begin{aligned}
 |B_n| &\leq U_n + D_{n-2} + (1 - A)(D_{n-1} - D_{n-2}) \\
 &\quad - D_{n-1} - D_{n-2}A + AD_{n-1} \\
 (5.32) \quad &= U_n.
 \end{aligned}$$

This completes the arguments for Theorem 2.1 in the case where $n \in P$. The discourse for the case where $n \in N$ is similar and is hence omitted. \square


Proof of Lemma 5.1. The convolution identity (5.14) is easy to verify directly from (2.1). To prove (5.16), return to (5.11) with the facts that $j \in N$ for $1 \leq j \leq n - 1$ to get $|B_2| = D_2$, $B_n = A \sum_{j=2}^{n-1} |B_j| + D_n$, and $|B_j| = (1 + B)|B_{j-1}| - D_{j-1} + D_j$ for $3 \leq j \leq n - 1$.

To prove (5.15), we get an induction started by applying (2.1) with $n = 2$ and $n = 3$:

$$\begin{aligned}
 U_3 - (1 + B)U_2 &= AD_2 + D_3 - (1 + B)D_2 \\
 &= (A - B)D_2 + D_3 - D_2 \\
 (5.33) \quad &\geq 0,
 \end{aligned}$$

where the last inequality follows from $A \geq B$, $D_2 \geq 0$ and $D_3 \geq D_2$. Equation (5.15) with $i = 4$ follows from the inequalities $A \geq 1$ and $A \geq B$:

$$\begin{aligned}
 U_4 - (1 + B)U_3 &= [AU_3 + (1 + B)U_2 + D_4 - D_2] - (1 + B)[AD_2 + D_3] \\
 &= (A - 1)(A - B)D_2 + (A - B)D_3 + D_4 - D_3 \\
 (5.34) \quad &\geq D_4 - D_3,
 \end{aligned}$$



**Bounds for Linear Recurrences
with Restricted Coefficients**

Kenneth S. Berenhaut and
Robert Lund

Title Page

Contents



Go Back

Close

Quit

Page 29 of 34

where the last inequality follows from $A \geq 1$, $A \geq B$, $D_3 \geq 0$ and $D_4 \geq D_3$.

For the general inductive step, take an $n > 4$ and suppose that $U_i - (1 + B)U_{i-1} \geq D_i - D_{i-1}$ for $3 \leq i \leq n - 1$. Then (2.1) gives

$$\begin{aligned}
 U_n - (1 + B)U_{n-1} &= [AU_{n-1} + (1 + B)U_{n-2} + D_n - D_{n-2}] \\
 &\quad - (1 + B)[AU_{n-2} + (1 + B)U_{n-3} + D_{n-1} - D_{n-3}] \\
 &= A[U_{n-1} - (1 + B)U_{n-2}] + (1 + B)[U_{n-2} - (1 + B)U_{n-3}] \\
 (5.35) \quad &\quad + D_n - D_{n-2} - (1 + B)D_{n-1} + (1 + B)D_{n-3}.
 \end{aligned}$$

Applying the inductive hypothesis to the bracketed terms in (5.35) and collecting terms gives the inequality

$$\begin{aligned}
 U_n - (1 + B)U_{n-1} &\geq A(D_{n-1} - D_{n-2}) + (1 + B)(D_{n-2} - D_{n-3}) + D_n - D_{n-2} \\
 &\quad - (1 + B)D_{n-1} + (1 + B)D_{n-3} \\
 (5.36) \quad &= D_n - D_{n-1} + (A - B)[D_{n-1} - D_{n-2}].
 \end{aligned}$$

The assumed monotonicity of D_k in k and $A \geq B$ give

$$(5.37) \quad U_n - (1 + B)U_{n-1} \geq D_n - D_{n-1}$$

and the proof is complete. \square



Bounds for Linear Recurrences with Restricted Coefficients

Kenneth S. Berenhaut and
Robert Lund

Title Page

Contents



Go Back


Close

Quit

Page 30 of 34

References

- [1] F. BEAUCOUP, P. BORWEIN, D.W. BOYD, AND C. PINNER, Power series with restricted coefficients and a root on a given ray, *Math. Comp.*, **222** (1998), 715–736.
- [2] F. BEAUCOUP, P. BORWEIN, D.W. BOYD, AND C. PINNER, Multiple roots of $[-1,1]$ power series, *J. London Math. Soc.*, **57** (1998), 135–147.
- [3] P. BORWEIN, AND T. ERDLYI, On the zeros of polynomials with restricted coefficients, *Illinois J. Math.*, **46** (1997), 667–675.
- [4] P. BORWEIN, AND C.G. PINNER, Polynomials with 0,+1,-1 coefficients and a root close to a given point, *Canad. J. Math.*, **49** (1997), 887–915.
- [5] M. EMBREE, AND L.N. TREFETHEN, Growth and decay of random Fibonacci sequences, *The Royal Society of London Proceedings, Series A, Mathematical, Physical and Engineering Sciences*, **455** (1999), 2471–2485.
- [6] W. FELLER, *An Introduction to Probability Theory and its Applications, Volume I*, 3rd Edition, John Wiley and Sons, New York (1968).
- [7] L. FLATTO, J.C. LAGARIAS, AND B. POONEN, Lap numbers and periodic points of the beta transformation, *Ergodic Theory Dynam. Systems*, (2) **14** (1994), 237–266.
- [8] E. HANSEN, Interval Arithmetic in Matrix Computations, Part II, *SIAM J. Numer. Anal.* , **2**(2), (1965), 308–320.



**Bounds for Linear Recurrences
with Restricted Coefficients**

Kenneth S. Berenhaut and
Robert Lund

Title Page

Contents




Go Back

Close

Quit

Page 31 of 34

- [9] E.R. HANSEN, *Global Optimization Using Interval Analysis*, Marcel Dekker, New York (1992).
- [10] E. HANSEN AND R. SMITH, Interval Arithmetic in Matrix Computations, Part I, *SIAM J. Numer. Anal.*, **4**(1) (1967), 1–9.
- [11] C.R. HEATHCOTE, Complete exponential convergence and related topics, *J. Appl. Probab.*, **4** (1967), 1–40.
- [12] R.B. KEARFOTT, *Rigorous Global Search: Continuous Problems*, Kluwer Academic Publishers, Dordrecht, The Netherlands (1996).
- [13] D.G. KENDALL, Unitary dilations of Markov transition operators and the corresponding integral representations for transition probability matrices, In: *Probability and Statistics*, Edited by U. Grenander, 139-161, Wiley, New York, (1959).
- [14] M. KIJIMA, *Markov Processes for Stochastic Modeling*, Chapman and Hall, London (1997).
- [15] R.K. KITTAPPA, A representation of the solution of the n th order linear difference equations with variable coefficients, *Lin. Algebra Applics.*, **193** (1993), 211–222.
- [16] R.K. MALLIK, On the solution of a linear homogeneous difference equation with varying coefficients, *SIAM J. Math. Anal.*, **31** (2000), 375–385.
- [17] A. NEUMAIER, *Interval methods for systems of equations*, Encyclopedia of Mathematics and its Applications, **37**. Cambridge University Press, Cambridge, (1990).



**Bounds for Linear Recurrences
with Restricted Coefficients**

Kenneth S. Berenhaut and
Robert Lund

Title Page

Contents




Go Back

Close

Quit

Page 32 of 34

- [18] A. ODLYZKO AND B. POONEN, Zeros of polynomials with 0,1 coefficients, *L'Enseignement Mathematique*, **39** (1993), 317–348.
- [19] A.M. ODLYZKO, *Asymptotic enumeration methods*, In: Handbook of Combinatorics (R. Graham, M. Groetschel, and L. Lovasz, Editors), Volume II, Elsevier, Amsterdam, (1995), 1063–1229.
- [20] J. POPENDA, One expression for the solutions of second order difference equations, *Proc. Amer. Math. Soc.*, **100** (1987), 87–93.
- [21] C.G. PINNER, Double roots of $[-1,1]$ power series and related matters, *Math. Comp.*, **68** (1999), 1149–1178.
- [22] B. SOLOMYAK, Conjugates of beta-numbers and the zero-free domain for a class of analytic functions, *Proc. London Math. Soc.*, **68** (1994), 477–498.
- [23] L.N. TREFETHEN AND D. BAU, *Numerical Linear Algebra*, SIAM, Philadelphia, PA, (1997).
- [24] D. VISWANATH, *Lyapunov exponents from random Fibonacci sequences to the Lorenz equations*, Ph.D. Thesis, Department of Computer Science, Cornell University, (1998).
- [25] D. VISWANATH, Random Fibonacci sequences and the number 1.13198824..., *Math. Comp.*, **69** (2000), 1131–1155.
- [26] D. VISWANATH, AND L.N. TREFETHEN, Condition numbers of random triangular matrices, *SIAM Journal on Matrix Analysis and Applications*, **19** (1998), 564–581.



**Bounds for Linear Recurrences
with Restricted Coefficients**

Kenneth S. Berenhaut and
Robert Lund

Title Page

Contents




Go Back

Close

Quit

Page 33 of 34

- [27] H.S. WILF, *Generatingfunctionology*, Second Edition, Academic Press, Boston, (1994).
- [28] T.G. WRIGHT AND L.N. TREFETHEN, Computing Lyapunov constants for random recurrences with smooth coefficients, *J. Comput. Appl. Math.*, **132**(2) (2001), 331–340.
- [29] O. YAMAMOTO, On some bounds for zeros of norm-bounded polynomials, *J. Symbolic Comput.*, **18**(5) (1994), 403–427.



**Bounds for Linear Recurrences
with Restricted Coefficients**

Kenneth S. Berenhaut and
Robert Lund

Title Page

Contents



Go Back

Close

Quit

Page 34 of 34