


Journal of Inequalities in Pure and Applied Mathematics

SOME INEQUALITIES FOR A CLASS OF GENERALIZED MEANS

KAI-ZHONG GUAN

Department of Mathematics and Physics
Nanhua University, Hengyang, Hunan 421001
The Peoples' Republic of China
EMail: kaizhongguan@yahoo.com.cn

©2000 Victoria University
ISSN (electronic): 1443-5756
054-04



volume 5, issue 3, article 69,
2004.

*Received 03 March, 2004;
accepted 30 March, 2004.*

Communicated by: D. Stefanescu

[Abstract](#)

[Contents](#)



[Home Page](#)

[Go Back](#)

[Close](#)

[Quit](#)

Abstract

In this paper, we define a symmetric function, show its properties, and establish several analytic inequalities, some of which are "Ky Fan" type inequalities. The harmonic-geometric mean inequality is refined.

2000 Mathematics Subject Classification: 05E05, 26D20.

Key words: Symmetric function ; Ky Fan inequality ; Harmonic-geometric mean inequality.

Contents

1	Introduction	3
2	Lemmas	5
3	Refinement of the Harmonic-Geometric Mean Inequality ...	8
4	Schur-convexity of the Function $H_n^r(x)$	11
5	Some "Ky Fan" Type Inequalities	16
	References	



Some Inequalities for a Class of Generalized Means

Kai-Zhong Guan

Title Page

Contents



Go Back

Close

Quit

Page 2 of 20

1. Introduction

Let $x = (x_1, x_2, \dots, x_n)$ be an n -tuple of positive numbers. The un-weighted arithmetic, geometric and harmonic means of x , denoted by $A_n(x)$, $G_n(x)$, $H_n(x)$, respectively, are defined as follows

$$A_n(x) = \frac{1}{n} \sum_{i=1}^n x_i, \quad G_n(x) = \left(\prod_{i=1}^n x_i \right)^{\frac{1}{n}}, \quad H_n(x) = \frac{n}{\sum_{i=1}^n \frac{1}{x_i}}.$$

Assume that $0 \leq x_i < 1$, $1 \leq i \leq n$ and define $1 - x = (1 - x_1, 1 - x_2, \dots, 1 - x_n)$. Throughout the sequel the symbols $A_n(1 - x)$, $G_n(1 - x)$, $H_n(1 - x)$ will stand for the un-weighted arithmetic, geometric, harmonic means of $1 - x$.

A remarkable new counterpart of the inequality $G_n \leq A_n$ has been published in [1].

Theorem 1.1. *If $0 < x_i \leq \frac{1}{2}$, for all $i = 1, 2, \dots, n$, then*

$$(1.1) \quad \frac{G_n(x)}{G_n(1 - x)} \leq \frac{A_n(x)}{A_n(1 - x)}$$

with equality if and only if all the x_i are equal.

This result, commonly referred to as the Ky Fan inequality, has stimulated the interest of many researchers. New proofs, improvements and generalizations of the inequality (1.1) have been found. For more details, interested readers can see [2], [3] and [4].

W.-L. Wang and P.-F. Wang [5] have established a counterpart of the classical inequality $H_n \leq G_n \leq A_n$. Their result reads as follows.



Some Inequalities for a Class of Generalized Means

Kai-Zhong Guan

Title Page

Contents




Go Back

Close

Quit

Page 3 of 20



Title Page

Contents



Go Back

Close

Quit

Page 4 of 20

Theorem 1.2. If $0 < x_i \leq \frac{1}{2}$, for all $i = 1, 2, \dots, n$, then

$$(1.2) \quad \frac{H_n(x)}{H_n(1-x)} \leq \frac{G_n(x)}{G_n(1-x)} \leq \frac{A_n(x)}{A_n(1-x)}.$$

All kinds of means about numbers and their inequalities have stimulated the interest of many researchers. Here we define a new mean, that is:

Definition 1.1. Let $x \in \mathbb{R}_+^n = \{x | x = (x_1, x_2, \dots, x_n) | x_i > 0, i = 1, 2, \dots, n\}$, we define the symmetric function as follows

$$H_n^r(x) = H_n^r(x_1, x_2, \dots, x_n) = \left[\prod_{1 \leq i_1 < \dots < i_r \leq n} \left(\frac{r}{\sum_{i=1}^r x_{i_j}^{-1}} \right) \right]^{\frac{1}{\binom{n}{r}}}.$$

Clearly $H_n^n(x) = H_n(x)$, $H_n^1(x) = G_n(x)$, where $\binom{n}{r} = \frac{n!}{r!(n-r)!}$.

The Schur-convex function was introduced by I. Schur in 1923 [7]. Its definition is as follows:

Definition 1.2. $f : I^n \rightarrow \mathbb{R} (n > 1)$ is called Schur-convex if $x \prec y$, then

$$(1.3) \quad f(x) \leq f(y)$$

for all $x, y \in I^n = I \times I \times \dots \times I$ (n copies). It is called strictly Schur-convex if the inequality is strict; f is called Schur-concave (resp. strictly Schur-concave) if the inequality (1.3) is reversed. For more details, interested readers can see [6], [7] and [8].

The paper is organized as follows. A refinement of harmonic-geometric mean inequality is obtained in Section 3. In Section 4, we investigate the Schur-convexity of the symmetric function. Several ‘‘Ky Fan’’ type inequalities are obtained in Section 5.

2. Lemmas

In this section, we give the following lemmas for the proofs of our main results.

Lemma 2.1. ([5]) If $0 < x_i \leq \frac{1}{2}$, for all $i = 1, 2, \dots, n$, then

$$(2.1) \quad \frac{\sum_{i=1}^n \frac{1}{1-x_i}}{\sum_{i=1}^n \frac{1}{x_i}} \leq \left[\frac{\prod_{i=1}^n \frac{1}{1-x_i}}{\prod_{i=1}^n \frac{1}{x_i}} \right]^{\frac{1}{n}} \quad \text{or} \quad \frac{H_n(x)}{H_n(1-x)} \leq \frac{G_n(x)}{G_n(1-x)}.$$

Lemma 2.2. If $0 < x_i \leq \frac{1}{2}$, for all $i = 1, 2, \dots, n+1$, and $S_{n+1} = \sum_{i=1}^{n+1} \frac{1}{x_i}$, $\bar{S}_{n+1} = \sum_{i=1}^{n+1} \frac{1}{1-x_i}$, then

$$(2.2) \quad \left[\frac{\sum_{i=1}^{n+1} \left(\bar{S}_{n+1} - \frac{1}{1-x_i} \right)}{\sum_{i=1}^{n+1} \left(S_{n+1} - \frac{1}{x_i} \right)} \right]^n \leq \left[\frac{\prod_{i=1}^{n+1} \left(\bar{S}_{n+1} - \frac{1}{1-x_i} \right)}{\prod_{i=1}^{n+1} \left(S_{n+1} - \frac{1}{x_i} \right)} \right]^{\frac{1}{n+1}}.$$

Proof. Inequality (2.2) is equivalent to the following

$$n \ln \frac{\bar{S}_{n+1}}{S_{n+1}} \leq \frac{1}{n+1} \ln \left[\frac{\prod_{i=1}^{n+1} \left(\bar{S}_{n+1} - \frac{1}{1-x_i} \right)}{\prod_{i=1}^{n+1} \left(S_{n+1} - \frac{1}{x_i} \right)} \right]$$

Since $0 < x_i \leq \frac{1}{2}$, and $1 - x_i \geq x_i$, it follows that

$$\begin{aligned} \frac{\bar{S}_{n+1} - \frac{1}{1-x_j}}{S_{n+1} - \frac{1}{x_j}} &= \frac{\frac{1}{1-x_1} + \dots + \frac{1}{1-x_{j-1}} + \frac{1}{1-x_{j+1}} + \dots + \frac{1}{1-x_{n+1}}}{\frac{1}{x_1} + \dots + \frac{1}{x_{j-1}} + \frac{1}{x_{j+1}} + \dots + \frac{1}{x_{n+1}}} \\ &\geq \frac{\frac{1}{1-x_1} \dots \frac{1}{1-x_{j-1}} \frac{1}{1-x_{j+1}} \dots \frac{1}{1-x_{n+1}}}{\frac{1}{x_1} \dots \frac{1}{x_{j-1}} \frac{1}{x_{j+1}} \dots \frac{1}{x_{n+1}}}. \end{aligned}$$



Some Inequalities for a Class of Generalized Means

Kai-Zhong Guan

Title Page

Contents



Go Back

Close

Quit

Page 5 of 20

By the above inequality and Lemma 2.1, we have

$$\begin{aligned} \frac{1}{n+1} \ln \prod_{i=1}^{n+1} \frac{\bar{S}_{n+1} - \frac{1}{1-x_i}}{S_{n+1} - \frac{1}{x_i}} &\geq \frac{1}{n+1} \ln \prod_{i=1}^{n+1} \left[\left(\frac{1}{1-x_i} \right) / \left(\frac{1}{x_i} \right) \right]^n \\ &= n \ln \prod_{i=1}^{n+1} \left[\left(\frac{1}{1-x_i} \right) / \left(\frac{1}{x_i} \right) \right]^{n+1} \\ &\geq n \ln \frac{\frac{1}{1-x_1} + \dots + \frac{1}{1-x_{n+1}}}{\frac{1}{x_1} + \dots + \frac{1}{x_{n+1}}}, \end{aligned}$$

or

$$n \ln \frac{\bar{S}_{n+1}}{S_{n+1}} \leq \frac{1}{n+1} \ln \left[\frac{\prod_{i=1}^{n+1} \left(\bar{S}_{n+1} - \frac{1}{1-x_i} \right)}{\prod_{i=1}^{n+1} \left(S_{n+1} - \frac{1}{x_i} \right)} \right].$$

□

Lemma 2.3. [6, p. 259]. Let $f(x) = f(x_1, x_2, \dots, x_n)$ be symmetric and have continuous partial derivatives on I^n , where I is an open interval. Then $f : I^n \rightarrow \mathbb{R}$ is Schur-convex if and only if

$$(2.3) \quad (x_i - x_j) \left(\frac{\partial f}{\partial x_i} - \frac{\partial f}{\partial x_j} \right) \geq 0$$

on I^n . It is strictly Schur-convex if (2.3) is a strict inequality for $x_i \neq x_j$, $1 \leq i, j \leq n$.



Some Inequalities for a Class of Generalized Means

Kai-Zhong Guan

Title Page

Contents



Go Back

Close

Quit

Page 6 of 20


Since $f(x)$ is symmetric, Schur's condition can be reduced as [7, p. 57]

$$(2.4) \quad (x_1 - x_2) \left(\frac{\partial f}{\partial x_1} - \frac{\partial f}{\partial x_2} \right) \geq 0,$$

and f is strictly Schur-convex if (2.4) is a strict inequality for $x_1 \neq x_2$. The Schur condition that guarantees a symmetric function being Schur-concave is the same as (2.3) or (2.4) except the direction of the inequality.

In Schur's condition, the domain of $f(x)$ does not have to be a Cartesian product I^n . Lemma 2.3 remains true if we replace I^n by a set $A \subseteq \mathbb{R}^n$ with the following properties ([7, p. 57]):

- (i) A is convex and has a nonempty interior;
- (ii) A is symmetric in the sense that $x \in A$ implies $Px \in A$ for any $n \times n$ permutation matrix P .



Some Inequalities for a Class of Generalized Means

Kai-Zhong Guan

Title Page

Contents



Go Back

Close

Quit

Page 7 of 20

3. Refinement of the Harmonic-Geometric Mean Inequality


The goal of this section is to obtain the basic inequality of $H_n^r(x)$, and give a refinement of the Harmonic-Geometric mean inequality.

Theorem 3.1. Let $x \in \mathbb{R}_+^n = \{x | x = (x_1, x_2, \dots, x_n) | x_i > 0, i = 1, 2, \dots, n\}$, then

$$(3.1) \quad H_n^{r+1}(x) \leq H_n^r(x), \quad r = 1, 2, \dots, n - 1.$$

Proof. By the arithmetic-geometric mean inequality and the monotonicity of the function $y = \ln x$, we have

$$\begin{aligned} & \binom{n}{r+1} \ln H_n^{r+1}(x) \\ &= \sum_{1 \leq i_1 < \dots < i_{r+1} \leq n} \ln \frac{r+1}{\sum_{j=1}^{r+1} x_{i_j}^{-1}} \\ &= \sum_{1 \leq i_1 < \dots < i_{r+1} \leq n} \ln \left[\frac{(r+1)r}{(r+1) \sum_{k=1}^{r+1} x_{i_k}^{-1} - \sum_{j=1}^{r+1} x_{i_j}^{-1}} \right] \\ &= \sum_{1 \leq i_1 < \dots < i_{r+1} \leq n} \ln \left\{ \frac{r}{\left[\sum_{j=1}^{r+1} \left(\sum_{k=1}^{r+1} x_{i_k}^{-1} - x_{i_j}^{-1} \right) \right] / (r+1)} \right\} \end{aligned}$$



Some Inequalities for a Class of Generalized Means

Kai-Zhong Guan

Title Page

Contents




Go Back

Close

Quit

Page 8 of 20



Title Page

Contents



Go Back

Close

Quit

Page 9 of 20

$$\begin{aligned}
 &\leq \sum_{1 \leq i_1 < \dots < i_{r+1} \leq n} \ln \left[\frac{r}{\left(\prod_{j=1}^{r+1} \left(\sum_{k=1}^{r+1} x_{i_k}^{-1} - x_{i_j}^{-1} \right) \right)^{\frac{1}{r+1}}} \right] \\
 &= \sum_{1 \leq i_1 < \dots < i_{r+1} \leq n} \ln \left[\prod_{j=1}^{r+1} \frac{r}{\sum_{k=1}^{r+1} x_{i_k}^{-1} - x_{i_j}^{-1}} \right]^{\frac{1}{r+1}} \\
 &= \frac{1}{r+1} \sum_{1 \leq i_1 < \dots < i_{r+1} \leq n} \left[\sum_{j=1}^{r+1} \ln \frac{r}{\sum_{k=1}^{r+1} x_{i_k}^{-1} - x_{i_j}^{-1}} \right] \\
 &= \frac{1}{r+1} \sum_{j=1}^n \sum_{1 \leq i_1 < \dots < i_r \leq n, i_1, \dots, i_r \neq j} \ln \frac{r}{\sum_{k=1}^r x_{i_k}^{-1}}.
 \end{aligned}$$

Let

$$S_j = \sum_{1 \leq i_1 < \dots < i_r \leq n, i_1, \dots, i_r \neq j} \ln \frac{r}{\sum_{k=1}^r x_{i_k}^{-1}}, \quad j = 1, 2, \dots, n.$$

We can easily get

$$\sum_{j=1}^n S_j = (n-r) \sum_{1 \leq i_1 < \dots < i_r \leq n} \ln \frac{r}{\sum_{k=1}^r x_{i_k}^{-1}} = (n-r) \binom{n}{r} \ln H_n^r(x).$$

Thus

$$\binom{n}{r+1} \ln H_n^{r+1}(x) \leq \frac{n-r}{r+1} \binom{n}{r} \ln H_n^r(x) = \binom{n}{r+1} \ln H_n^r(x),$$

or


$$H_n^{r+1}(x) \leq H_n^r(x), \quad r = 1, 2, \dots, n-1.$$

□

Corollary 3.2. Let $x \in \mathbb{R}_+^n = \{x | x = (x_1, x_2, \dots, x_n) | x_i > 0, i = 1, 2, \dots, n\}$, then

$$(3.2) \quad H_n(x) \leq H_n^{n-1}(x) \leq \dots \leq H_n^2(x) \leq H_n^1(x) = G_n(x).$$

Remark 3.1. The corollary refines the harmonic-geometric mean inequality.



Some Inequalities for a Class of
Generalized Means

Kai-Zhong Guan

Title Page

Contents



Go Back

Close

Quit

Page 10 of 20

4. Schur-convexity of the Function $H_n^r(x)$

In this section, we investigate the Schur-convexity of the function $H_n^r(x)$, and establish several analytic inequalities by use of the theory of majorization.

Theorem 4.1. Let $\mathbb{R}_+^n = \{x | x = (x_1, x_2, \dots, x_n) | x_i > 0, i = 1, 2, \dots, n\}$, then the function $H_n^r(x)$ is Schur-concave in \mathbb{R}_+^n .

Proof. It is clear that $H_n^r(x)$ is symmetric and has continuous partial derivatives on \mathbb{R}_+^n . By Lemma 2.3, we only need to prove


$$(x_1 - x_2) \left(\frac{\partial H_n^r(x)}{\partial x_1} - \frac{\partial H_n^r(x)}{\partial x_2} \right) \leq 0.$$

As matter of fact, we can easily derive

$$\ln H_n^r(x) = \frac{1}{\binom{n}{r}} \sum_{2 \leq i_1 < \dots < i_r \leq n} \ln \frac{r}{\sum_{j=1}^r x_{i_j}^{-1}} + \sum_{2 \leq i_1 < \dots < i_{r-1} \leq n} \ln \frac{r}{x_1^{-1} + \sum_{j=1}^{r-1} x_{i_j}^{-1}}.$$

Differentiating $\ln H_n^r(x)$ with respect to x_1 , we have

$$\begin{aligned} \frac{\partial H_n^r(x)}{\partial x_1} &= \frac{H_n^r(x)}{\binom{n}{r}} \left(\sum_{2 \leq i_1 < \dots < i_{r-1} \leq n} \frac{1}{x_1^{-1} + \sum_{j=1}^{r-1} x_{i_j}^{-1}} \right) \cdot \frac{1}{x_1^2} \\ &= \frac{H_n^r(x)}{\binom{n}{r}} \cdot \frac{1}{x_1^2} \left[\left(\sum_{3 \leq i_1 < \dots < i_{r-1} \leq n} \frac{1}{x_1^{-1} + \sum_{j=1}^{r-1} x_{i_j}^{-1}} \right) \right. \\ &\quad \left. + \sum_{3 \leq i_1 < \dots < i_{r-2} \leq n} \frac{1}{\left(x_1^{-1} + x_2^{-1} + \sum_{j=1}^{r-2} x_{i_j}^{-1} \right)} \right]. \end{aligned}$$



Some Inequalities for a Class of Generalized Means

Kai-Zhong Guan

Title Page

Contents



Go Back

Close

Quit

Page 11 of 20

Similar to the above, we can also obtain

$$\begin{aligned} \frac{\partial H_n^r(x)}{\partial x_2} &= \frac{H_n^r(x)}{\binom{n}{r}} \left(\sum_{2 \leq i_1 < \dots < i_{r-1} \leq n} \frac{1}{x_2^{-1} + \sum_{j=1}^{r-1} x_{i_j}^{-1}} \right) \cdot \frac{1}{x_2^2} \\ &= \frac{H_n^r(x)}{\binom{n}{r}} \cdot \frac{1}{x_2^2} \left[\left(\sum_{3 \leq i_1 < \dots < i_{r-1} \leq n} \frac{1}{x_2^{-1} + \sum_{j=1}^{r-1} x_{i_j}^{-1}} \right) \right. \\ &\quad \left. + \sum_{3 \leq i_1 < \dots < i_{r-2} \leq n} \frac{1}{\left(x_1^{-1} + x_2^{-1} + \sum_{j=1}^{r-2} x_{i_j}^{-1} \right)} \right]. \end{aligned}$$

Thus

$$\begin{aligned} &(x_1 - x_2) \left(\frac{\partial H_n^r(x)}{\partial x_1} - \frac{\partial H_n^r(x)}{\partial x_2} \right) \\ &= (x_1 - x_2) \frac{H_n^r(x)}{\binom{n}{r}} \left[\left(\sum_{2 \leq i_1 < \dots < i_{r-1} \leq n} \frac{1}{x_1^{-1} + \sum_{j=1}^{r-1} x_{i_j}^{-1}} \right) \cdot \frac{1}{x_1^2} \right. \\ &\quad - \left(\sum_{2 \leq i_1 < \dots < i_{r-1} \leq n} \frac{1}{x_2^{-1} + \sum_{j=1}^{r-1} x_{i_j}^{-1}} \right) \cdot \frac{1}{x_2^2} \\ &\quad \left. + \sum_{3 \leq i_1 < \dots < i_{r-2} \leq n} \frac{1}{\left(x_1^{-1} + x_2^{-1} + \sum_{j=1}^{r-2} x_{i_j}^{-1} \right)} \left(\frac{1}{x_1^2} - \frac{1}{x_2^2} \right) \right] \end{aligned}$$



Some Inequalities for a Class of Generalized Means

Kai-Zhong Guan

Title Page

Contents



Go Back

Close

Quit

Page 12 of 20

$$\begin{aligned}
&= -(x_1 - x_2)^2 \left[\frac{(x_1 + x_2)}{x_1^2 x_2^2} \sum_{3 \leq i_1 < \dots < i_{r-2} \leq n} \frac{1}{x_1^{-1} + x_2^{-1} + \sum_{j=1}^{r-2} x_{i_j}^{-1}} \right. \\
&\quad \left. + \sum_{2 \leq i_1 < \dots < i_{r-1} \leq n} \frac{1 + (x_1 + x_2) \sum_{j=1}^{r-1} x_{i_j}^{-1}}{x_1^2 x_2^2 \left(x_1^{-1} + \sum_{j=1}^{r-1} x_{i_j}^{-1} \right) \left(x_2^{-1} + \sum_{j=1}^{r-1} x_{i_j}^{-1} \right)} \right] \\
&\leq 0.
\end{aligned}$$

□

Corollary 4.2. Let $x_i > 0, i = 1, 2, \dots, n, n \geq 2$, and $\sum_{i=1}^n x_i = s, c > 0$, then

$$(4.1) \quad \frac{H_n^r(c+x)}{H_n^r(x)} \geq \left(\frac{nc}{s} + 1 \right)^{\frac{1}{\binom{n}{r}}}, \quad r = 1, 2, \dots, n,$$


where $c+x = (c+x_1, c+x_2, \dots, c+x_n)$.

Proof. By [9], we have

$$\frac{c+x}{nc+s} = \left(\frac{c+x_1}{nc+s}, \dots, \frac{c+x_n}{nc+s} \right) \prec \left(\frac{x_1}{s}, \dots, \frac{x_n}{s} \right) = \frac{x}{s}.$$

Using Theorem 4.1, we obtain

$$H_n^r \left(\frac{c+x}{nc+s} \right) \geq H_n^r \left(\frac{x}{s} \right).$$



Some Inequalities for a Class of Generalized Means

Kai-Zhong Guan

Title Page

Contents



Go Back

Close

Quit

Page 13 of 20

Or

$$\frac{H_n^r(c+x)}{H_n^r(x)} \geq \left(\frac{nc}{s} + 1\right)^{\left(\frac{1}{r}\right)}.$$

□

Corollary 4.3. Let $x_i > 0, i = 1, 2, \dots, n, n \geq 2$, and $\sum_{i=1}^n x_i = s, c \geq s$, then

$$(4.2) \quad \frac{H_n^r(c-x)}{H_n^r(x)} \geq \left(\frac{nc}{s} - 1\right)^{\left(\frac{1}{r}\right)}, \quad r = 1, 2, \dots, n,$$

where $c - x = (c - x_1, c - x_2, \dots, c - x_n)$.

Proof. By [9], we have

$$\frac{c-x}{nc-s} = \left(\frac{c-x_1}{nc-s}, \dots, \frac{c-x_n}{nc-s}\right) \prec \left(\frac{x_1}{s}, \dots, \frac{x_n}{s}\right) = \frac{x}{s}.$$


Using Theorem 4.1, we obtain

$$H_n^r\left(\frac{c-x}{nc-s}\right) \geq H_n^r\left(\frac{x}{s}\right),$$

or

$$\frac{H_n^r(c-x)}{H_n^r(x)} \geq \left(\frac{nc}{s} - 1\right)^{\left(\frac{1}{r}\right)}.$$

□



Some Inequalities for a Class of Generalized Means

Kai-Zhong Guan

Title Page

Contents



Go Back

Close

Quit


Page 14 of 20

Remark 4.1. Let $c = s = 1$, we can obtain

$$\frac{H_n^r(1-x)}{H_n^r(x)} \geq (n-1)^{\frac{1}{\binom{n}{r}}}, \quad r = 1, 2, \dots, n.$$

In particular,

$$\frac{H_n(1-x)}{H_n(x)} \geq (n-1), \quad \frac{G_n(1-x)}{G_n(x)} \geq \sqrt[n]{n-1}.$$



Some Inequalities for a Class of
Generalized Means

Kai-Zhong Guan

Title Page

Contents



Go Back

Close

Quit

Page 15 of 20

5. Some “Ky Fan” Type Inequalities

In this section, some “Ky Fan” type inequalities are established, the Ky Fan inequality is generalized.

Theorem 5.1. Assume that $0 < x_i \leq \frac{1}{2}$, $i = 1, 2, \dots, n$, then


$$(5.1) \quad \frac{H_n^{r+1}(x)}{H_n^{r+1}(1-x)} \leq \left[\frac{H_n^r(x)}{H_n^r(1-x)} \right]^{\frac{1}{r}}, \quad r = 1, 2, \dots, n-1.$$

Proof. Set

$$\varphi_r = \frac{H_n^r(x)}{H_n^r(1-x)} = \prod_{1 \leq i_1 < \dots < i_r \leq n} \left[\frac{\sum_{j=1}^r \frac{1}{1-x_{i_j}}}{\sum_{j=1}^r \frac{1}{x_{i_j}}} \right]^{\frac{1}{r}}.$$

By Lemma 2.2 and the monotonicity of the function $y = \ln x$, we have

$$\begin{aligned} \binom{n}{r+1} \ln \phi_{r+1} &= \sum_{1 \leq i_1 < \dots < i_{r+1} \leq n} \ln \frac{\sum_{j=1}^{r+1} \frac{1}{1-x_{i_j}}}{\sum_{j=1}^{r+1} \frac{1}{x_{i_j}}} \\ &= \sum_{1 \leq i_1 < \dots < i_{r+1} \leq n} \ln \frac{\sum_{j=1}^{r+1} \left(\sum_{k=1}^{r+1} \frac{1}{1-x_{i_k}} - \frac{1}{1-x_{i_j}} \right)}{\sum_{j=1}^{r+1} \left(\sum_{k=1}^{r+1} \frac{1}{x_{i_k}} - \frac{1}{x_{i_j}} \right)} \\ &\leq \sum_{1 \leq i_1 < \dots < i_{r+1} \leq n} \ln \left[\prod_{j=1}^{r+1} \frac{\sum_{k=1}^{r+1} \frac{1}{1-x_{i_k}} - \frac{1}{1-x_{i_j}}}{\sum_{k=1}^{r+1} \frac{1}{x_{i_k}} - \frac{1}{x_{i_j}}} \right]^{\frac{1}{r(r+1)}} \end{aligned}$$



Some Inequalities for a Class of Generalized Means

Kai-Zhong Guan

Title Page

Contents



Go Back

Close

Quit

Page 16 of 20

$$= \frac{1}{r(r+1)} \sum_{j=1}^n \sum_{1 \leq i_1 < \dots < i_r \leq n, i_1, \dots, i_r \neq j} \ln \frac{\sum_{k=1}^r \frac{1}{1-x_{i_k}}}{\sum_{k=1}^r \frac{1}{x_{i_k}}}.$$

Similar to Theorem 3.1, we can derive

$$\binom{n}{r+1} \ln \phi_{r+1} \leq \frac{1}{r(r+1)} (n-r) \binom{n}{r} \ln \phi_r = \frac{1}{r} \binom{n}{r+1} \ln \phi_r.$$

Thus

$$(\phi_r)^{\frac{1}{r}} \geq \phi_{r+1},$$

or

$$\frac{H_n^{r+1}(x)}{H_n^{r+1}(1-x)} \leq \left[\frac{H_n^r(x)}{H_n^r(1-x)} \right]^{\frac{1}{r}}, \quad r = 1, 2, \dots, n-1.$$

□

Remark 5.1. By Theorem 5.1, we can obtain


$$(5.2) \quad \frac{H_n^2(x)}{H_n^2(1-x)} \leq \frac{H_n^1(x)}{H_n^1(1-x)} = \frac{G_n(x)}{G_n(1-x)} \leq \frac{A_n(x)}{A_n(1-x)}.$$

This is a generalization of the “Ky Fan” inequality.

By Lemma 2.1 and the proof of Theorem 3.1, we have the following

Theorem 5.2. If $0 < x_i \leq \frac{1}{2}$, $i = 1, 2, \dots, n$, then

$$(5.3) \quad \frac{\prod_{i=1}^n (x_i)}{\prod_{i=1}^n (1-x_i)} \leq \frac{H_n(x)}{H_n(1-x)} \leq \frac{G_n(x)}{G_n(1-x)} \leq \frac{A_n(x)}{A_n(1-x)}.$$



Some Inequalities for a Class of Generalized Means

Kai-Zhong Guan

Title Page

Contents



Go Back

Close

Quit

Page 17 of 20

The inequality (5.3) generalizes the inequality (1.2).

Theorem 5.3. *If $0 < x_i \leq \frac{1}{2}$, $i = 1, 2, \dots, n$, then*


$$(5.4) \quad \frac{H_n^r(x)}{H_n^r(1-x)} \leq \frac{H_n^1(x)}{H_n^1(1-x)} \\ = \frac{G_n(x)}{G_n(1-x)} \leq \frac{A_n(x)}{A_n(1-x)}, \quad r = 2, \dots, n.$$

Proof. Set

$$\varphi_r = \frac{H_n^r(x)}{H_n^r(1-x)} = \prod_{1 \leq i_1 < \dots < i_r \leq n} \left[\frac{\sum_{j=1}^r \frac{1}{1-x_{i_j}}}{\sum_{j=1}^r \frac{1}{x_{i_j}}} \right]^{\frac{1}{r}}.$$

By Lemma 2.1 and the monotonicity of the function $y = \ln x$, we have

$$\begin{aligned} \binom{n}{r} \ln \varphi_r &= \sum_{1 \leq i_1 < \dots < i_{r+1} \leq n} \ln \frac{\sum_{j=1}^r \frac{1}{1-x_{i_j}}}{\sum_{j=1}^r \frac{1}{x_{i_j}}} \\ &\leq \sum_{1 \leq i_1 < \dots < i_{r+1} \leq n} \ln \left[\frac{\prod_{j=1}^r \frac{1}{1-x_{i_j}}}{\prod_{j=1}^r \frac{1}{x_{i_j}}} \right]^{\frac{1}{r}} \\ &= \frac{1}{r} \sum_{1 \leq i_1 < \dots < i_r \leq n} \sum_{j=1}^r \ln \frac{1}{\frac{1-x_{i_j}}{x_{i_j}}}. \end{aligned}$$



Some Inequalities for a Class of Generalized Means

Kai-Zhong Guan

Title Page

Contents



Go Back

Close

Quit

Page 18 of 20

By knowledge of combinations, we can easily find

$$\begin{aligned} \binom{n}{r} \ln \phi_r &\leq \frac{1}{r} \ln \left[\prod_{i=1}^n \frac{1-x_i}{x_i} \right]^{\binom{n-1}{r-1}} \\ &= \frac{1}{r} \binom{n-1}{r-1} \ln \left[\prod_{i=1}^n \frac{1-x_i}{x_i} \right] \\ &= \frac{1}{r} \binom{n-1}{r-1} \ln \phi_1 = \binom{n}{r} \ln \phi_1. \end{aligned}$$

Thus


$$\phi_r \leq \phi_1, \quad r = 2, \dots, n,$$

or

$$(5.5) \quad \frac{H_n^r(x)}{H_n^r(1-x)} \leq \frac{G_n(x)}{G_n(1-x)}, \quad r = 2, \dots, n.$$

□

The inequality (5.5) generalizes the “Ky Fan” inequality.



Some Inequalities for a Class of Generalized Means

Kai-Zhong Guan

Title Page

Contents



Go Back


Close

Quit

Page 19 of 20

References

- [1] E.F. BECKENBACH AND R. BELLMAN, *Inequalities*, Springer-Verlag, New York, 1965.
- [2] H. ALZER, The inequality of Ky Fan and related results, *Acta. Appl. Math.*, **38** (1995), 305–354.
- [3] J. SÁNDOR, On an inequality of Ky Fan, Babeş-Bolyai Univ., *Fac. Math. Phys., Res. Semin.*, **7** (1990), 29–34.
- [4] J. SÁNDOR AND T. TRIF, A new refinement of the Ky Fan inequality, *Math. Inequal. Appl.*, **2** (1999), 529–533.
- [5] W.-L. WANG AND P.-F. WANG, A class of inequalities for symmetric functions (in chinese), *Acta. Math. Sinica*, **27** (1984), 485–497.
- [6] A.W. ROBERTS AND D.E. VARBERG, *Convex Function*, Academic Press, New York, San Francisco, London, 1973.
- [7] A.W. MARSHALL AND I. OLKIN, *Inequalities: Theory of Majorization and Its Applications*, Academic Press, 1979.
- [8] D.S. MITRINOVIĆ, *Analytic Inequalities*, Springer-Verlag, New York, 1970.
- [9] HUANNAN SHI, Refinement and generalization of a class of inequalities for symmetric functions (in Chinese), *Mathematics in Practices and Theory*, **4** (1999), 81–84.



Some Inequalities for a Class of
Generalized Means

Kai-Zhong Guan

Title Page

Contents



Go Back

Close

Quit

Page 20 of 20