

AN INEQUALITY IMPROVING THE FIRST HERMITE-HADAMARD INEQUALITY FOR CONVEX FUNCTIONS DEFINED ON LINEAR SPACES AND APPLICATIONS FOR SEMI-INNER PRODUCTS

S.S. DRAGOMIR

School of Communications and Informatics
Victoria University of Technology
PO Box 14428
Melbourne City MC
8001 Victoria, Australia

*E*Mail: sever@matilda.vu.edu.au

*U*RL: <http://rgmia.vu.edu.au/SSDragomirWeb.html>



volume 3, issue 2, article 31,
2002.

Received 16 November, 2001;
accepted 15 February, 2002.

Communicated by: C.P. Niculescu

Abstract

Contents



Home Page

Go Back

Close

Quit

Abstract


An integral inequality for convex functions defined on linear spaces is obtained which contains in a particular case a refinement for the first part of the celebrated Hermite-Hadamard inequality. Applications for semi-inner products on normed linear spaces are also provided.

2000 Mathematics Subject Classification: Primary 26D15, 26D10; Secondary 46B10.

Key words: Hermite-Hadamard integral inequality, Convex functions, Semi-Inner Products.

Contents

1	Introduction	3
2	The Results	6
3	Applications for Semi-Inner Products	15
	References	



An Inequality Improving the First Hermite-Hadamard Inequality for Convex Functions Defined on Linear Spaces and Applications for Semi-Inner Products

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 2 of 19

1. Introduction

Let X be a real linear space, $a, b \in X$, $a \neq b$ and let $[a, b] := \{(1 - \lambda)a + \lambda b, \lambda \in [0, 1]\}$ be the *segment* generated by a and b . We consider the function $f : [a, b] \rightarrow \mathbb{R}$ and the attached function $g(a, b) : [0, 1] \rightarrow \mathbb{R}$, $g(a, b)(t) := f[(1 - t)a + tb]$, $t \in [0, 1]$.

It is well known that f is convex on $[a, b]$ iff $g(a, b)$ is convex on $[0, 1]$, and the following lateral derivatives exist and satisfy

- (i) $g'_{\pm}(a, b)(s) = (\nabla_{\pm} f[(1 - s)a + sb])(b - a)$, $s \in (0, 1)$
- (ii) $g'_+(a, b)(0) = (\nabla_+ f(a))(b - a)$
- (iii) $g'_-(a, b)(1) = (\nabla_- f(b))(b - a)$

where $(\nabla_{\pm} f(x))(y)$ are the *Gâteaux lateral derivatives*, we recall that

$$\begin{aligned}(\nabla_+ f(x))(y) &:= \lim_{h \rightarrow 0^+} \left[\frac{f(x + hy) - f(x)}{h} \right], \\(\nabla_- f(x))(y) &:= \lim_{k \rightarrow 0^-} \left[\frac{f(x + ky) - f(x)}{k} \right], \quad x, y \in X.\end{aligned}$$

The following inequality is the well-known Hermite-Hadamard integral inequality for convex functions defined on a segment $[a, b] \subset X$:

$$(HH) \quad f\left(\frac{a+b}{2}\right) \leq \int_0^1 f[(1-t)a + tb] dt \leq \frac{f(a) + f(b)}{2},$$



An Inequality Improving the First Hermite-Hadamard Inequality for Convex Functions Defined on Linear Spaces and Applications for Semi-Inner Products

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 3 of 19

which easily follows by the classical Hermite-Hadamard inequality for the convex function $g(a, b) : [0, 1] \rightarrow \mathbb{R}$

$$g(a, b) \left(\frac{1}{2} \right) \leq \int_0^1 g(a, b)(t) dt \leq \frac{g(a, b)(0) + g(a, b)(1)}{2}.$$

For other related results see the monograph on line [1].

Now, assume that $(X, \|\cdot\|)$ is a normed linear space. The function $f_0(s) = \frac{1}{2} \|x\|^2$, $x \in X$ is convex and thus the following limits exist

$$(iv) \langle x, y \rangle_s := (\nabla_+ f_0(y))(x) = \lim_{t \rightarrow 0^+} \left[\frac{\|y+tx\|^2 - \|y\|^2}{2t} \right];$$

$$(v) \langle x, y \rangle_i := (\nabla_- f_0(y))(x) = \lim_{s \rightarrow 0^-} \left[\frac{\|y+sx\|^2 - \|y\|^2}{2s} \right];$$

for any $x, y \in X$. They are called the *lower* and *upper semi-inner* products associated to the norm $\|\cdot\|$.


For the sake of completeness we list here some of the main properties of these mappings that will be used in the sequel (see for example [2]), assuming that $p, q \in \{s, i\}$ and $p \neq q$:

$$(a) \langle x, x \rangle_p = \|x\|^2 \text{ for all } x \in X;$$

$$(aa) \langle \alpha x, \beta y \rangle_p = \alpha\beta \langle x, y \rangle_p \text{ if } \alpha, \beta \geq 0 \text{ and } x, y \in X;$$

$$(aaa) \left| \langle x, y \rangle_p \right| \leq \|x\| \|y\| \text{ for all } x, y \in X;$$

$$(av) \langle \alpha x + y, x \rangle_p = \alpha \langle x, x \rangle_p + \langle y, x \rangle_p \text{ if } x, y \in X \text{ and } \alpha \in \mathbb{R};$$



An Inequality Improving the First Hermite-Hadamard Inequality for Convex Functions Defined on Linear Spaces and Applications for Semi-Inner Products

S.S. Dragomir

Title Page

Contents




Go Back

Close

Quit

Page 4 of 19



An Inequality Improving the
First Hermite-Hadamard
Inequality for Convex Functions
Defined on Linear Spaces and
Applications for Semi-Inner
Products

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 5 of 19

(v) $\langle -x, y \rangle_p = -\langle x, y \rangle_q$ for all $x, y \in X$;

(va) $\langle x + y, z \rangle_p \leq \|x\| \|z\| + \langle y, z \rangle_p$ for all $x, y, z \in X$;

(vaa) The mapping $\langle \cdot, \cdot \rangle_p$ is continuous and subadditive (superadditive) in the first variable for $p = s$ (or $p = i$);

(vaaa) The normed linear space $(X, \|\cdot\|)$ is smooth at the point $x_0 \in X \setminus \{0\}$ if and only if $\langle y, x_0 \rangle_s = \langle y, x_0 \rangle_i$ for all $y \in X$; in general $\langle y, x \rangle_i \leq \langle y, x \rangle_s$ for all $x, y \in X$;

(ax) If the norm $\|\cdot\|$ is induced by an inner product $\langle \cdot, \cdot \rangle$, then $\langle y, x \rangle_i = \langle y, x \rangle = \langle y, x \rangle_s$ for all $x, y \in X$.

Applying inequality (HH) for the convex function $f_0(x) = \frac{1}{2} \|x\|^2$, one may deduce the inequality

$$(1.1) \quad \left\| \frac{x+y}{2} \right\|^2 \leq \int_0^1 \|(1-t)x + ty\|^2 dt \leq \frac{\|x\|^2 + \|y\|^2}{2}$$

for any $x, y \in X$. The same (HH) inequality applied for $f_1(x) = \|x\|$, will give the following refinement of the triangle inequality:

$$(1.2) \quad \left\| \frac{x+y}{2} \right\| \leq \int_0^1 \|(1-t)x + ty\| dt \leq \frac{\|x\| + \|y\|}{2}, \quad x, y \in X.$$

In this paper we point out an integral inequality for convex functions which is related to the first Hermite-Hadamard inequality in (HH) and investigate its applications for semi-inner products in normed linear spaces.

2. The Results

We start with the following lemma which is also of interest in itself.

Lemma 2.1. *Let $h : [\alpha, \beta] \subset \mathbb{R} \rightarrow \mathbb{R}$ be a convex function on $[\alpha, \beta]$. Then for any $\gamma \in [\alpha, \beta]$ one has the inequality*

$$\begin{aligned}
 (2.1) \quad & \frac{1}{2} [(\beta - \gamma)^2 h'_+(\gamma) - (\gamma - \alpha)^2 h'_-(\gamma)] \\
 & \leq \int_{\alpha}^{\beta} h(t) dt - (\beta - \alpha) h(\gamma) \\
 & \leq \frac{1}{2} [(\beta - \gamma)^2 h'_-(\beta) - (\gamma - \alpha)^2 h'_+(\alpha)].
 \end{aligned}$$

The constant $\frac{1}{2}$ is sharp in both inequalities.

The second inequality also holds for $\gamma = \alpha$ or $\gamma = \beta$.


Proof. It is easy to see that for any locally absolutely continuous function $h : (\alpha, \beta) \rightarrow \mathbb{R}$, we have the identity

$$(2.2) \quad \int_{\alpha}^{\gamma} (t - \alpha) h'(t) dt + \int_{\gamma}^{\beta} (t - \beta) h'(t) dt = h(\gamma) - \int_{\alpha}^{\beta} h(t) dt$$

for any $\gamma \in (\alpha, \beta)$, where h' is the derivative of h which exists a.e. on (α, β) .

Since h is convex, then it is locally Lipschitzian and thus (2.2) holds. Moreover, for any $\gamma \in (\alpha, \beta)$, we have the inequalities

$$(2.3) \quad h'(t) \leq h'_-(\gamma) \text{ for a.e. } t \in [\alpha, \gamma]$$



An Inequality Improving the
First Hermite-Hadamard
Inequality for Convex Functions
Defined on Linear Spaces and
Applications for Semi-Inner
Products

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 6 of 19

and

$$(2.4) \quad h'(t) \geq h'_+(\gamma) \text{ for a.e. } t \in [\gamma, \beta].$$

If we multiply (2.3) by $t - \alpha \geq 0$, $t \in [\alpha, \gamma]$ and integrate on $[\alpha, \gamma]$, we get

$$(2.5) \quad \int_{\alpha}^{\gamma} (t - \alpha) h'(t) dt \leq \frac{1}{2} (\gamma - \alpha)^2 h'_-(\gamma)$$

and if we multiply (2.4) by $\beta - t \geq 0$, $t \in [\gamma, \beta]$, and integrate on $[\gamma, \beta]$, we also have

$$(2.6) \quad \int_{\gamma}^{\beta} (\beta - t) h'(t) dt \geq \frac{1}{2} (\beta - \gamma)^2 h'_+(\gamma).$$


If we subtract (2.6) from (2.5) and use the representation (2.2), we deduce the first inequality in (2.1).

Now, assume that the first inequality (2.1) holds with $C > 0$ instead of $\frac{1}{2}$, i.e.,

$$(2.7) \quad C [(\beta - \gamma)^2 h'_+(\gamma) - (\gamma - \alpha)^2 h'_-(\gamma)] \leq \int_{\alpha}^{\beta} h(t) dt - (\beta - \alpha) h(\gamma).$$

Consider the convex function $h_0(t) := k \left| t - \frac{\alpha + \beta}{2} \right|$, $k > 0$, $t \in [\alpha, \beta]$. Then

$$h'_{0+} \left(\frac{\alpha + \beta}{2} \right) = k, \quad h'_{0-} \left(\frac{\alpha + \beta}{2} \right) = -k, \quad h_0 \left(\frac{\alpha + \beta}{2} \right) = 0$$



An Inequality Improving the First Hermite-Hadamard Inequality for Convex Functions Defined on Linear Spaces and Applications for Semi-Inner Products

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 7 of 19

and

$$\int_{\alpha}^{\beta} h_0(t) dt = \frac{1}{4}k(\beta - \alpha)^2.$$

If in (2.7) we choose $h = h_0$, $\gamma = \frac{\alpha+\beta}{2}$, then we get

$$C \left[\frac{1}{4}(\beta - \alpha)^2 k + \frac{1}{4}(\beta - \alpha)^2 k \right] \leq \frac{1}{4}k(\beta - \alpha)^2$$

which gives $C \leq \frac{1}{2}$ and the sharpness of the constant in the first part of (2.1) is proved.

If either $h'_+(\alpha) = -\infty$ or $h'_-(\beta) = -\infty$, then the second inequality in (2.1) holds true.

Assume that $h'_+(\alpha)$ and $h'_-(\beta)$ are finite. Since h is convex on $[\alpha, \beta]$, we have


$$(2.8) \quad h'(t) \geq h'_+(\alpha) \text{ for a.e. } t \in [\alpha, \gamma] \quad (\gamma \text{ may be equal to } \beta)$$

and

$$(2.9) \quad h'(t) \leq h'_-(\beta) \text{ for a.e. } t \in [\gamma, \beta] \quad (\gamma \text{ may be equal to } \alpha).$$

If we multiply (2.8) by $t - \alpha \geq 0$, $t \in [\alpha, \gamma]$ and integrate on $[\alpha, \gamma]$, then we deduce

$$(2.10) \quad \int_{\alpha}^{\gamma} (t - \alpha) h'(t) dt \geq \frac{1}{2}(\gamma - \alpha)^2 h'_+(\alpha)$$



**An Inequality Improving the
First Hermite-Hadamard
Inequality for Convex Functions
Defined on Linear Spaces and
Applications for Semi-Inner
Products**

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 8 of 19

and if we multiply (2.9) by $\beta - t \geq 0$, $t \in [\gamma, \beta]$, and integrate on $[\gamma, \beta]$, then we also have

$$(2.11) \quad \int_{\gamma}^{\beta} (\beta - t) h'(t) dt \leq \frac{1}{2} (\beta - \gamma)^2 h'_-(\beta).$$

Finally, if we subtract (2.10) from (2.11) and use the representation (2.2), we deduce the second inequality in (2.1). Now, assume that the second inequality in (2.1) holds with a constant $D > 0$ instead of $\frac{1}{2}$, i.e.,

$$(2.12) \quad \int_{\alpha}^{\beta} h(t) dt - (\beta - \alpha) h(\gamma) \leq D [(\beta - \gamma)^2 h'_-(\beta) - (\gamma - \alpha)^2 h'_+(\alpha)].$$


If we consider the convex function $h_0(t) = k |t - \frac{\alpha+\beta}{2}|$, $k > 0$, $t \in [\alpha, \beta]$, then we have $h'_{0-}(\beta) = k$, $h'_{0+}(\alpha) = -k$ and by (2.12) applied for h_0 in $\gamma = \frac{\alpha+\beta}{2}$ we get

$$\frac{1}{4} k (\beta - \alpha)^2 \leq D \left[\frac{1}{4} k (\beta - \alpha)^2 + \frac{1}{4} k (\beta - \alpha)^2 \right],$$

giving $D \geq \frac{1}{2}$ which proves the sharpness of the constant $\frac{1}{2}$ in the second inequality in (2.1). \square

Corollary 2.2. *With the assumptions of Lemma 2.1 and if $\gamma \in (\alpha, \beta)$ is a point of differentiability for h , then*

$$(2.13) \quad \left(\frac{\alpha + \beta}{2} - \gamma \right) h'(\gamma) \leq \frac{1}{\beta - \alpha} \int_{\alpha}^{\beta} h(t) dt - h(\gamma).$$



An Inequality Improving the
First Hermite-Hadamard
Inequality for Convex Functions
Defined on Linear Spaces and
Applications for Semi-Inner
Products

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 9 of 19

Now, recall that the following inequality, which is well known in the literature as the Hermite-Hadamard inequality for convex functions, holds

$$(2.14) \quad h\left(\frac{\alpha + \beta}{2}\right) \leq \frac{1}{\beta - \alpha} \int_{\alpha}^{\beta} h(t) dt \leq \frac{h(\alpha) + h(\beta)}{2}.$$

The following corollary provides both a sharper lower bound for the difference,

$$\frac{1}{\beta - \alpha} \int_{\alpha}^{\beta} h(t) dt - h\left(\frac{\alpha + \beta}{2}\right),$$

which we know is nonnegative, and an upper bound.

Corollary 2.3. *Let $h : [\alpha, \beta] \rightarrow \mathbb{R}$ be a convex function on $[\alpha, \beta]$. Then we have the inequality*

$$(2.15) \quad \begin{aligned} 0 &\leq \frac{1}{8} \left[h'_+ \left(\frac{\alpha + \beta}{2} \right) - h'_- \left(\frac{\alpha + \beta}{2} \right) \right] (\beta - \alpha) \\ &\leq \frac{1}{\beta - \alpha} \int_{\alpha}^{\beta} h(t) dt - h\left(\frac{\alpha + \beta}{2}\right) \\ &\leq \frac{1}{8} [h'_-(\beta) - h'_+(\alpha)] (\beta - \alpha). \end{aligned}$$

The constant $\frac{1}{8}$ is sharp in both inequalities.

Example 2.1. *Assume that $-\infty < \alpha < 0 < \beta < \infty$ and consider the convex function $h : [\alpha, \beta] \rightarrow \mathbb{R}$, $h(x) = \exp|x|$. We have*

$$h'(x) = \begin{cases} -e^{-x} & \text{if } x < 0, \\ e^x & \text{if } x > 0; \end{cases}$$



An Inequality Improving the
First Hermite-Hadamard
Inequality for Convex Functions
Defined on Linear Spaces and
Applications for Semi-Inner
Products

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 10 of 19

and $h'_-(0) = -1$, $h'_+(0) = 1$. Also,

$$\int_{\alpha}^{\beta} h(t) dt = \int_{\alpha}^0 e^{-x} dx + \int_0^{\beta} e^x dx = \exp(\beta) + \exp(-\alpha) - 2.$$

Now, if $\frac{\alpha+\beta}{2} \neq 0$, then by (2.15) we deduce the elementary inequality

$$(2.16) \quad 0 \leq \frac{\exp(\beta) + \exp(-\alpha) - 2}{\beta - \alpha} - \exp\left|\frac{\alpha + \beta}{2}\right| \\ \leq \frac{1}{8} [\exp(\beta) + \exp(-\alpha)] (\beta - \alpha).$$

If $\frac{\alpha+\beta}{2} = 0$ and if we denote $\beta = a$, $a > 0$, thus $\alpha = -a$ and by (2.15) we also have

$$(2.17) \quad \frac{1}{2}a \leq \frac{\exp(a) - 1}{a} - 1 \leq \frac{1}{2}a \exp(a).$$

The reader may produce other elementary inequalities by choosing in an appropriate way the convex function h . We omit the details.

We are now able to state the corresponding result for convex functions defined on linear spaces.

Theorem 2.4. *Let X be a linear space, $a, b \in X$, $a \neq b$ and $f : [a, b] \subset X \rightarrow \mathbb{R}$ be a convex function on the segment $[a, b]$. Then for any $s \in (0, 1)$ one has the inequality*

$$(2.18) \quad \frac{1}{2} [(1-s)^2 (\nabla_+ f [(1-s)a + sb]) (b-a) \\ - s^2 (\nabla_- f [(1-s)a + sb]) (b-a)]$$



An Inequality Improving the
First Hermite-Hadamard
Inequality for Convex Functions
Defined on Linear Spaces and
Applications for Semi-Inner
Products

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 11 of 19

$$\begin{aligned} &\leq \int_0^1 f [(1-t)a + tb] dt - f [(1-s)a + sb] \\ &\leq \frac{1}{2} [(1-s)^2 (\nabla_- f(b)) (b-a) - s^2 (\nabla_+ f(a)) (b-a)]. \end{aligned}$$

The constant $\frac{1}{2}$ is sharp in both inequalities.

The second inequality also holds for $s = 0$ or $s = 1$.

Proof. Follows by Lemma 2.1 applied for the convex function $h(t) = g(a, b)(t) = f[(1-t)a + tb]$, $t \in [0, 1]$, and the choices $\alpha = 0$, $\beta = 1$, and $\gamma = s$. \square


Corollary 2.5. *If $f : [a, b] \rightarrow \mathbb{R}$ is as in Theorem 2.4 and Gâteaux differentiable in $c := (1-\lambda)a + \lambda b$, $\lambda \in (0, 1)$ along the direction $(b-a)$, then we have the inequality:*

$$(2.19) \quad \left(\frac{1}{2} - \lambda\right) (\nabla f(c)) (b-a) \leq \int_0^1 f [(1-t)a + tb] dt - f(c).$$

The following result related to the first Hermite-Hadamard inequality for functions defined on linear spaces also holds.

Corollary 2.6. *If f is as in Theorem 2.4, then*

$$(2.20) \quad \begin{aligned} 0 &\leq \frac{1}{8} \left[\nabla_+ f \left(\frac{a+b}{2} \right) (b-a) - \nabla_- f \left(\frac{a+b}{2} \right) (b-a) \right] \\ &\leq \int_0^1 f [(1-t)a + tb] dt - f \left(\frac{a+b}{2} \right) \\ &\leq \frac{1}{8} [(\nabla_- f(b)) (b-a) - (\nabla_+ f(a)) (b-a)]. \end{aligned}$$



An Inequality Improving the
First Hermite-Hadamard
Inequality for Convex Functions
Defined on Linear Spaces and
Applications for Semi-Inner
Products

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 12 of 19

The constant $\frac{1}{8}$ is sharp in both inequalities.

Now, let $\Omega \subset \mathbb{R}^n$ be an open and convex set in \mathbb{R}^n .

If $F : \Omega \rightarrow \mathbb{R}$ is a differentiable convex function on Ω , then, obviously, for any $\bar{c} \in \Omega$ we have

$$\nabla F(\bar{c})(\bar{y}) = \sum_{i=1}^n \frac{\partial F(\bar{c})}{\partial x_i} \cdot y_i, \quad \bar{y} \in \mathbb{R}^n,$$

where $\frac{\partial F}{\partial x_i}$ are the partial derivatives of F with respect to the variable x_i ($i = 1, \dots, n$).

Using (2.18), we may state that

$$\begin{aligned} (2.21) \quad & \left(\frac{1}{2} - \lambda\right) \sum_{i=1}^n \frac{\partial F(\lambda\bar{a} + (1-\lambda)\bar{b})}{\partial x_i} \cdot (b_i - a_i) \\ & \leq \int_0^1 F[(1-t)\bar{a} + t\bar{b}] dt - F((1-\lambda)\bar{a} + \lambda\bar{b}) \\ & \leq (1-\lambda)^2 \sum_{i=1}^n \frac{\partial F(\bar{b})}{\partial x_i} \cdot (b_i - a_i) - \lambda^2 \sum_{i=1}^n \frac{\partial F(\bar{a})}{\partial x_i} \cdot (b_i - a_i) \end{aligned}$$

for any $\bar{a}, \bar{b} \in \Omega$ and $\lambda \in (0, 1)$.

In particular, for $\lambda = \frac{1}{2}$, we get

$$\begin{aligned} (2.22) \quad 0 & \leq \int_0^1 F[(1-t)\bar{a} + t\bar{b}] dt - F\left(\frac{\bar{a} + \bar{b}}{2}\right) \\ & \leq \frac{1}{8} \sum_{i=1}^n \left(\frac{\partial F(\bar{b})}{\partial x_i} - \frac{\partial F(\bar{a})}{\partial x_i} \right) \cdot (b_i - a_i). \end{aligned}$$



An Inequality Improving the
First Hermite-Hadamard
Inequality for Convex Functions
Defined on Linear Spaces and
Applications for Semi-Inner
Products

S.S. Dragomir

Title Page

Contents




Go Back

Close

Quit

Page 13 of 19

In (2.22) the constant $\frac{1}{8}$ is sharp.



**An Inequality Improving the
First Hermite-Hadamard
Inequality for Convex Functions
Defined on Linear Spaces and
Applications for Semi-Inner
Products**

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 14 of 19

3. Applications for Semi-Inner Products

Let $(X, \|\cdot\|)$ be a real normed linear space. We may state the following results for the semi-inner products $\langle \cdot, \cdot \rangle_i$ and $\langle \cdot, \cdot \rangle_s$.

Proposition 3.1. For any $x, y \in X$ and $\sigma \in (0, 1)$ we have the inequalities:

$$(3.1) \quad (1 - \sigma)^2 \langle y - x, (1 - \sigma)x + \sigma y \rangle_s - \sigma^2 \langle y - x, (1 - \sigma)x + \sigma y \rangle_i \\ \leq \int_0^1 \|(1 - t)x + ty\|^2 dt - \|(1 - \sigma)x + \sigma y\|^2 \\ \leq (1 - \sigma)^2 \langle y - x, y \rangle_i - \sigma^2 \langle y - x, y \rangle_s.$$

The second inequality in (3.1) also holds for $\sigma = 0$ or $\sigma = 1$.


The proof is obvious by Theorem 2.4 applied for the convex function $f(x) = \frac{1}{2} \|x\|^2$, $x \in X$.

If the space is *smooth*, then we may put $[x, y] = \langle x, y \rangle_i = \langle x, y \rangle_s$ for each $x, y \in X$ and the first inequality in (3.1) becomes

$$(3.2) \quad (1 - 2\sigma) [y - x, (1 - \sigma)x + \sigma y] \\ \leq \int_0^1 \|(1 - t)x + ty\|^2 dt - \|(1 - \sigma)x + \sigma y\|^2.$$

An interesting particular case one can get from (3.1) is the one for $\sigma = \frac{1}{2}$,

$$(3.3) \quad 0 \leq \frac{1}{8} [\langle y - x, y + x \rangle_s - \langle y - x, y + x \rangle_i] \\ \leq \int_0^1 \|(1 - t)x + ty\|^2 dt - \left\| \frac{x + y}{2} \right\|^2$$



An Inequality Improving the
First Hermite-Hadamard
Inequality for Convex Functions
Defined on Linear Spaces and
Applications for Semi-Inner
Products

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 15 of 19

$$\leq \frac{1}{4} [\langle y - x, y \rangle_i - \langle y - x, x \rangle_s].$$

The inequality (3.3) provides a refinement and a counterpart for the first inequality (1.1).

If we consider now two linearly independent vectors $x, y \in X$ and apply Theorem 2.4 for $f(x) = \|x\|$, $x \in X$, then we get

Proposition 3.2. *For any linearly independent vectors $x, y \in X$ and $\sigma \in (0, 1)$, one has the inequalities:*

$$(3.4) \quad \frac{1}{2} \left[(1 - \sigma)^2 \frac{\langle y - x, (1 - \sigma)x + \sigma y \rangle_\sigma}{\|(1 - \sigma)x + \sigma y\|} - \sigma^2 \frac{\langle y - x, (1 - \sigma)x + \sigma y \rangle_i}{\|(1 - \sigma)x + \sigma y\|} \right]$$

$$\leq \int_0^1 \|(1 - t)x + ty\| dt - \|(1 - \sigma)x + \sigma y\|$$


$$\leq \frac{1}{2} \left[(1 - \sigma)^2 \frac{\langle y - x, y \rangle_i}{\|y\|} - \sigma^2 \frac{\langle y - x, x \rangle_s}{\|x\|} \right].$$

The second inequality also holds for $\sigma = 0$ or $\sigma = 1$.

We note that if the space is smooth, then we have

$$(3.5) \quad \left(\frac{1}{2} - \sigma \right) \cdot \frac{[y - x, (1 - \sigma)x + \sigma y]}{\|(1 - \sigma)x + \sigma y\|}$$

$$\leq \int_0^1 \|(1 - t)x + ty\| dt - \|(1 - \sigma)x + \sigma y\|,$$



An Inequality Improving the
First Hermite-Hadamard
Inequality for Convex Functions
Defined on Linear Spaces and
Applications for Semi-Inner
Products

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 16 of 19

and for $\sigma = \frac{1}{2}$, (3.4) will give the simple inequality

$$\begin{aligned}
 (3.6) \quad 0 &\leq \frac{1}{8} \left[\left\langle y - x, \frac{\frac{x+y}{2}}{\left\| \frac{x+y}{2} \right\|} \right\rangle_s - \left\langle y - x, \frac{\frac{x+y}{2}}{\left\| \frac{x+y}{2} \right\|} \right\rangle_i \right] \\
 &\leq \int_0^1 \|(1-t)x + ty\| dt - \left\| \frac{x+y}{2} \right\| \\
 &\leq \frac{1}{8} \left[\left\langle y - x, \frac{y}{\|y\|} \right\rangle_i - \left\langle y - x, \frac{x}{\|x\|} \right\rangle_s \right].
 \end{aligned}$$

The inequality (3.6) provides a refinement and a counterpart for the first inequality in (1.2).

Moreover, if we assume that $(H, \langle \cdot, \cdot \rangle)$ is an inner product space, then by (3.6) we get for any $x, y \in H$ with $\|x\| = \|y\| = 1$ that


$$(3.7) \quad 0 \leq \int_0^1 \|(1-t)x + ty\| dt - \left\| \frac{x+y}{2} \right\| \leq \frac{1}{8} \|y - x\|^2.$$

The constant $\frac{1}{8}$ is sharp.

Indeed, if $H = \mathbb{R}$, $\langle a, b \rangle = a \cdot b$, then taking $x = -1$, $y = 1$, we obtain equality in (3.7).

We give now some examples.

1. Let $\ell^2(\mathbb{K})$, $\mathbb{K} = \mathbb{C}, \mathbb{R}$; be the Hilbert space of sequences $x = (x_i)_{i \in \mathbb{N}}$ with



**An Inequality Improving the
First Hermite-Hadamard
Inequality for Convex Functions
Defined on Linear Spaces and
Applications for Semi-Inner
Products**

S.S. Dragomir

Title Page

Contents




Go Back

Close

Quit

Page 17 of 19



An Inequality Improving the First Hermite-Hadamard Inequality for Convex Functions Defined on Linear Spaces and Applications for Semi-Inner Products

S.S. Dragomir

Title Page

Contents

⏪

⏩

◀

▶

Go Back

Close

Quit

Page 18 of 19

$\sum_{i=0}^{\infty} |x_i|^2 < \infty$. Then, by (3.7), we have the inequalities

$$(3.8) \quad 0 \leq \int_0^1 \left(\sum_{i=0}^{\infty} |(1-t)x_i + ty_i|^2 \right)^{\frac{1}{2}} dt - \left(\sum_{i=0}^{\infty} \left| \frac{x_i + y_i}{2} \right|^2 \right)^{\frac{1}{2}} \leq \frac{1}{8} \cdot \sum_{i=0}^{\infty} |y_i - x_i|^2,$$

for any $x, y \in \ell^2(\mathbb{K})$ provided $\sum_{i=0}^{\infty} |x_i|^2 = \sum_{i=0}^{\infty} |y_i|^2 = 1$.

2. Let μ be a positive measure, $L_2(\Omega)$ the Hilbert space of μ -measurable functions on Ω with complex values that are 2-integrable on Ω , i.e., $f \in L_2(\Omega)$ iff $\int_{\Omega} |f(t)|^2 d\mu(t) < \infty$. Then, by (3.7), we have the inequalities

$$(3.9) \quad 0 \leq \int_0^1 \left(\int_{\Omega} |(1-\lambda)f(t) + \lambda g(t)|^2 d\mu(t) \right)^{\frac{1}{2}} d\lambda - \left(\int_{\Omega} \left| \frac{f(t) + g(t)}{2} \right|^2 d\mu(t) \right)^{\frac{1}{2}} \leq \frac{1}{8} \cdot \int_{\Omega} |f(t) - g(t)|^2 d\mu(t)$$

for any $f, g \in L_2(\Omega)$ provided $\int_{\Omega} |f(t)|^2 d\mu(t) = \int_{\Omega} |g(t)|^2 d\mu(t) = 1$.

References

- [1] I. CIORANESCU, *Geometry of Banach Spaces, Duality Mappings and Nonlinear Problems*, Kluwer Academic Publishers, Dordrecht, 1990.
- [2] S.S. DRAGOMIR AND C.E.M. PEARCE, *Selected Topics on Hermite-Hadamard Inequalities and Applications*, RGMIA Monographs, Victoria University, 2000. (ONLINE: <http://rgmia.vu.edu.au/monographs>)



**An Inequality Improving the
First Hermite-Hadamard
Inequality for Convex Functions
Defined on Linear Spaces and
Applications for Semi-Inner
Products**

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 19 of 19
