


Journal of Inequalities in Pure and Applied Mathematics



NEW INEQUALITIES OF HADAMARD'S TYPE FOR LIPSCHITZIAN MAPPINGS

LIANG-CHENG WANG

School of Mathematics Scientia
Chongqing Institute of Technology
Xingsheng Lu 4, Yangjia Ping 400050
Chongqing City, China.

EMail: wlc@cqit.edu.cn

volume 6, issue 2, article 37,
2005.

*Received 16 September, 03;
accepted 06 February, 2005.*

Communicated by: F. Qi

[Abstract](#)

[Contents](#)



[Home Page](#)

[Go Back](#)

[Close](#)

[Quit](#)



Abstract

In this paper, we study several new inequalities of Hadamard's type for Lipschitzian mappings.

2000 Mathematics Subject Classification: Primary 26D07; Secondary 26B25, 26D15.

Key words: Lipschitzian mappings, Hadamard inequality, Convex function.

This author is partially supported by the Key Research Foundation of the Chongqing Institute of Technology under Grant 2004ZD94.

New Inequalities of Hadamard's Type for Lipschitzian Mappings

Liang-Cheng Wang

Contents

1	Introduction	3
2	Main Results	5
References		

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

Page 2 of 18

1. Introduction

Let us begin by defining some mappings that we shall need. If f is a continuous function on a closed interval $[a, b]$ ($a < b$), and for any $t \in (0, 1)$, let $u = ta + (1 - t)b$, then we can define

$$A(t) \triangleq \frac{1}{t(1-t)(b-a)^2} \int_a^u \left[\int_u^b f(tx + (1-t)y) dy \right] dx,$$

$$\begin{aligned} B(t) &\triangleq \frac{1}{(1-t)(b-a)^2} \int_a^u \left[\int_u^b f\left(\frac{(b-y)x + (y-u)u}{t(b-a)}\right) dy \right] dx \\ &\quad + \frac{1}{t(b-a)^2} \int_a^u \left[\int_u^b f\left(\frac{(u-x)u + (x-a)y}{(1-t)(b-a)}\right) dy \right] dx, \end{aligned}$$


$$C(t) \triangleq \frac{t}{(1-t)(b-a)} \int_a^u f(x) dx + \frac{1-t}{t(b-a)} \int_u^b f(y) dy.$$

Let f be a continuous convex function on $[a, b]$, then

$$(1.1) \quad f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b f(x) dx \leq \frac{f(a) + f(b)}{2}.$$

The inequalities in (1.1) are called Hadamard inequalities [1] – [7]. In [2], the author of this paper gave extensions and refinements of the inequalities in (1.1). He obtained the following:

$$(1.2) \quad f(ta + (1-t)b) \leq A(t) \leq B(t) \leq C(t) \leq tf(a) + (1-t)f(b).$$



New Inequalities of Hadamard's Type for Lipschitzian Mappings

Liang-Cheng Wang

Title Page

Contents



Go Back

Close

Quit

Page 3 of 18

Recently, Dragomir *et al.* [3], Yang and Tseng [4] and Matic and Pečarić [5] proved some results for Lipschitzian mappings related to (1.1). In this paper, we will prove some new inequalities for Lipschitzian mappings related to the mappings A , B and C (or to (1.2)).



New Inequalities of Hadamard's Type for Lipschitzian Mappings

Liang-Cheng Wang

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

Page 4 of 18

2. Main Results

For the mapping $A(t)$, we have the following theorem:

Theorem 2.1. Let $f : [a, b] \rightarrow \mathbb{R}$ be a M -Lipschitzian mapping. For any $t \in (0, 1)$, then

$$(2.1) \quad |A(t) - f(ta + (1-t)b)| \leq \frac{M}{3}t(1-t)(b-a).$$

Proof. Obviously, we have

$$(2.2) \quad b - u = t(b - a), u - a = (1 - t)(b - a).$$

From integral properties and (2.2), we have

$$(2.3) \quad f(ta + (1-t)b) = \frac{1}{t(1-t)(b-a)^2} \int_a^u \left[\int_u^b f(ta + (1-t)y) dy \right] dx.$$

For $x \in [a, u]$, when $y \in [u, -\frac{t}{1-t}x + \frac{u}{1-t}]$, then we have


$$(2.4) \quad ta + (1-t)b = u \geq tx + (1-t)y,$$

if $y \in [-\frac{t}{1-t}x + \frac{u}{1-t}, b]$, then the inequality (2.4) reverses.

Using (2.2) – (2.4) and integral properties, we obtain

$$|A(t) - f(ta + (1-t)b)|$$

$$\leq \frac{1}{t(1-t)(b-a)^2} \int_a^u \left[\int_u^b |f(tx + (1-t)y) - f(u)| dy \right] dx$$



New Inequalities of Hadamard's Type for Lipschitzian Mappings

Liang-Cheng Wang

Title Page

Contents



Go Back

Close

Quit

Page 5 of 18

$$\begin{aligned}
&\leq \frac{M}{t(1-t)(b-a)^2} \int_a^u \left[\int_u^b |tx + (1-t)y - u| dy \right] dx \\
&= \frac{M}{t(1-t)(b-a)^2} \int_a^u \left[\int_{-\frac{t}{1-t}x+\frac{u}{1-t}}^b (u - (tx + (1-t)y)) dy \right] dx \\
&\quad + \frac{M}{t(1-t)(b-a)^2} \int_a^u \left[\int_{-\frac{t}{1-t}x+\frac{u}{1-t}}^b (tx + (1-t)y - u) dy \right] dx \\
&= \frac{M}{t(1-t)(b-a)^2} \int_a^u \left[\frac{t^2}{1-t}x^2 + t \left(b - \frac{1+t}{1-t}u \right) x \right. \\
&\quad \left. + \frac{1+t^2}{2(1-t)}u^2 + \frac{1-t}{2}b^2 - bu \right] dx \\
&= \frac{M}{t(b-a)} \left[\frac{t^2}{3(1-t)}(u^2 + ua + a^2) + \frac{t}{2} \left(b - \frac{1+t}{1-t}u \right) (u + a) \right. \\
&\quad \left. + \frac{1+t^2}{2(1-t)}u^2 + \frac{1-t}{2}b^2 - bu \right] \\
&= \frac{M}{t(b-a)} \left[\frac{2t^2 - 3t + 3}{6(1-t)} (ta + (1-t)b)^2 \right. \\
&\quad \left. + \frac{-t(t+3)a - 3(t^2 - 3t + 2)b}{6(1-t)} (ta + (1-t)b) \right. \\
&\quad \left. + \frac{t^2}{3(1-t)}a^2 + \frac{t}{2}ab + \frac{1-t}{2}b^2 \right] = \frac{M}{3}t(1-t)(b-a).
\end{aligned}$$

This completes the proof of Theorem 2.1. \square



New Inequalities of Hadamard's Type for Lipschitzian Mappings

Liang-Cheng Wang

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

Page 6 of 18

For the mapping $B(t)$, we have the following theorem:

Theorem 2.2. Let f be defined as in Theorem 2.1. For any $t \in (0, 1)$, then

$$(2.5) \quad |B(t) - A(t)| \leq \frac{M}{2}t(1-t)(b-a),$$

and

$$(2.6) \quad |B(t) - f(ta + (1-t)b)| \leq \frac{M}{2}t(1-t)(b-a).$$

Proof. From $\frac{1}{t(1-t)} = \frac{1}{t} + \frac{1}{1-t}$, we have


$$(2.7) \quad A(t) = \left(\frac{1}{t} + \frac{1}{1-t} \right) \frac{1}{(b-a)^2} \int_a^u \left[\int_u^b f(tx + (1-t)y) dy \right] dx.$$

Let F be defined by

$$F(y) = tx + (1-t)y - \frac{(b-y)x + (y-u)u}{t(b-a)}.$$

Obviously, if $F(y)$ is the function of first degree for y , then $F(y)$ is monotone with y on $[u, b]$. And for any $x \in [a, u]$, $F(u) = tx + (1-t)u - x \geq 0$, $F(b) = tx + (1-t)b - u \geq 0$. Hence for $x \in [a, u]$ and $y \in [u, b]$, we get $F(y) \geq 0$, i.e.,

$$(2.8) \quad tx + (1-t)y \geq \frac{(b-y)x + (y-u)u}{t(b-a)}.$$



New Inequalities of Hadamard's Type for Lipschitzian Mappings

Liang-Cheng Wang

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

Page 7 of 18

Let G be defined by

$$G(x) = \frac{(u-x)u + (x-a)y}{(1-t)(b-a)} - (tx + (1-t)y).$$


Using $G(x)$ and the same method as in the proof of (2.8), we obtain

$$(2.9) \quad \frac{(u-x)u + (x-a)y}{(1-t)(b-a)} \geq tx + (1-t)y,$$

where, $x \in [a, u]$ and $y \in [u, b]$.

Using (2.7) – (2.9) and (2.2), we obtain

$$\begin{aligned} & |B(t) - A(t)| \\ & \leq \frac{1}{(1-t)(b-a)^2} \int_a^u \left[\int_u^b \left| f\left(\frac{(b-y)x + (y-u)u}{t(b-a)}\right) - f(tx + (1-t)y) \right| dy \right] dx \\ & \quad + \frac{1}{t(b-a)^2} \int_a^u \left[\int_u^b \left| f\left(\frac{(u-x)u + (x-a)y}{(1-t)(b-a)}\right) - f(tx + (1-t)y) \right| dy \right] dx \\ & \leq \frac{M}{(1-t)(b-a)^2} \int_a^u \left[\int_u^b \left| \frac{(b-y)x + (y-u)u}{t(b-a)} - (tx + (1-t)y) \right| dy \right] dx \\ & \quad + \frac{M}{t(b-a)^2} \int_a^u \left[\int_u^b \left| \frac{(u-x)u + (x-a)y}{(1-t)(b-a)} - (tx + (1-t)y) \right| dy \right] dx \\ & = \frac{M}{(1-t)(b-a)^2} \int_a^u \left[\int_u^b \left(tx + (1-t)y - \frac{(b-y)x + (y-u)u}{t(b-a)} \right) dy \right] dx \\ & \quad + \frac{M}{t(b-a)^2} \int_a^u \left[\int_u^b \left(\frac{(u-x)u + (x-a)y}{(1-t)(b-a)} - (tx + (1-t)y) \right) dy \right] dx \end{aligned}$$



New Inequalities of Hadamard's Type for Lipschitzian Mappings

Liang-Cheng Wang

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

Page 8 of 18

$$\begin{aligned}
&= \frac{M}{t(1-t)(b-a)^3} \int_a^u \left[\int_u^b ((b-a)(2t-1)(tx+(1-t)y) \right. \\
&\quad \left. + 2xy - (b+u)x - (a+u)y + 2u^2) dy \right] dx \\
&= \frac{M}{(1-t)(b-a)^2} \int_a^u \left[\frac{1}{2} ((b-a)(2t-1)(1-t) - (a+u))(b+u) \right. \\
&\quad \left. + (b-a)(2t-1)tx + 2u^2 \right] dx \\
&= \frac{M}{2(b-a)} \left[((b-a)(2t-1)(1-t) - (a+u))(b+u) \right. \\
&\quad \left. + (b-a)(2t-1)t(u+a) + 4u^2 \right] \\
&= \frac{M}{2} t(1-t)(b-a).
\end{aligned}$$


This completes the proof of inequality (2.5).

From (2.3) and $\frac{1}{t(1-t)} = \frac{1}{t} + \frac{1}{1-t}$, we have

$$\begin{aligned}
(2.10) \quad &f(ta + (1-t)b) \\
&= \left(\frac{1}{t} + \frac{1}{1-t} \right) \frac{1}{(b-a)^2} \int_a^u \left[\int_u^b f(ta + (1-t)b) dy \right] dx.
\end{aligned}$$

If $x \in [a, u]$ and $y \in [u, b]$, then we have

$$(2.11) \quad ta + (1-t)b = u \geq \frac{(b-y)x + (y-u)u}{t(b-a)},$$



New Inequalities of Hadamard's Type for Lipschitzian Mappings

Liang-Cheng Wang

Title Page	
Contents	
	
	
Go Back	
Close	
Quit	
Page 9 of 18	

$$\frac{(u-x)u + (x-a)y}{(1-t)(b-a)} \geq ta + (1-t)b = u.$$

Using (2.10)-(2.11) and (2.2), we obtain

$$\begin{aligned}
& |B(t) - f(ta + (1-t)b)| \\
& \leq \frac{1}{(1-t)(b-a)^2} \int_a^u \left[\int_u^b \left| f\left(\frac{(b-y)x + (y-u)u}{t(b-a)}\right) - f(u) \right| dy \right] dx \\
& \quad + \frac{1}{t(b-a)^2} \int_a^u \left[\int_u^b \left| f\left(\frac{(u-x)u + (x-a)y}{(1-t)(b-a)}\right) - f(u) \right| dy \right] dx \\
& \leq \frac{M}{(1-t)(b-a)^2} \int_a^u \left[\int_u^b \left| \frac{(b-y)x + (y-u)u}{t(b-a)} - u \right| dy \right] dx \\
& \quad + \frac{M}{t(b-a)^2} \int_a^u \left[\int_u^b \left| \frac{(u-x)u + (x-a)y}{(1-t)(b-a)} - u \right| dy \right] dx \\
& = \frac{M}{(1-t)(b-a)^2} \int_a^u \left[\int_u^b \left(u - \frac{(b-y)x + (y-u)u}{t(b-a)} \right) dy \right] dx \\
& \quad + \frac{M}{t(b-a)^2} \int_a^u \left[\int_u^b \left(\frac{(u-x)u + (x-a)y}{(1-t)(b-a)} - u \right) dy \right] dx \\
& = \frac{M}{t(1-t)(b-a)^3} \int_a^u \left[\int_u^b \left((b-a)(2t-1)u \right. \right. \\
& \quad \left. \left. + (2x-u-a)y - (b+u)x + 2u^2 \right) dy \right] dx \\
& = \frac{M}{(1-t)(b-a)^2} \int_a^u \left[-\frac{1}{2}(u+a)(b+u) + (b-a)(2t-1)u + 2u^2 \right] dx
\end{aligned}$$



New Inequalities of Hadamard's Type for Lipschitzian Mappings

Liang-Cheng Wang

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

Page 10 of 18

$$\begin{aligned}
&= \frac{M}{b-a} \left[-\frac{1}{2}(u+a)(b+u) + (b-a)(2t-1)u + 2u^2 \right] \\
&= \frac{M}{2}t(1-t)(b-a).
\end{aligned}$$

This completes the proof of inequality (2.6).

This completes the proof of Theorem (2.2). \square

For the mapping $C(t)$, we have the following theorem:

Theorem 2.3. Let f be defined as in Theorem 2.1. For any $t \in (0, 1)$, then

$$(2.12) \quad |C(t) - B(t)| \leq \frac{M}{2}t(1-t)(b-a),$$

and


$$(2.13) \quad |C(t) - (tf(a) + (1-t)f(b))| \leq Mt(1-t)(b-a).$$

Proof. From the integral property and (2.2), we have

$$\begin{aligned}
(2.14) \quad C(t) &= \frac{1}{(1-t)(b-a)^2} \int_a^u \left[\int_u^b f(x)dy \right] dx \\
&\quad + \frac{1}{t(b-a)^2} \int_a^u \left[\int_u^b f(y)dy \right] dx.
\end{aligned}$$

If $x \in [a, u]$ and $y \in [u, b]$, then we have

$$(2.15) \quad x \leq \frac{(b-y)x + (y-u)u}{t(b-a)}, \quad \frac{(u-x)u + (x-a)y}{(1-t)(b-a)} \leq y.$$



New Inequalities of Hadamard's Type for Lipschitzian Mappings

Liang-Cheng Wang

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

Page 11 of 18

Using (2.14) – (2.15) and (2.2), we obtain

$$\begin{aligned}
 & |C(t) - B(t)| \\
 & \leq \frac{1}{(1-t)(b-a)^2} \int_a^u \left[\int_u^b \left| f\left(\frac{(b-y)x + (y-u)u}{t(b-a)}\right) - f(x) \right| dy \right] dx \\
 & \quad + \frac{1}{t(b-a)^2} \int_a^u \left[\int_u^b \left| f\left(\frac{(u-x)u + (x-a)y}{(1-t)(b-a)}\right) - f(y) \right| dy \right] dx \\
 & \leq \frac{M}{(1-t)(b-a)^2} \int_a^u \left[\int_u^b \left| \frac{(b-y)x + (y-u)u}{t(b-a)} - x \right| dy \right] dx \\
 & \quad + \frac{1}{t(b-a)^2} \int_a^u \left[\int_u^b \left| \frac{(u-x)u + (x-a)y}{(1-t)(b-a)} - y \right| dy \right] dx \\
 & = \frac{M}{(1-t)(b-a)^2} \int_a^u \left[\int_u^b \left(\frac{(b-y)x + (y-u)u}{t(b-a)} - x \right) dy \right] dx \\
 & \quad + \frac{1}{t(b-a)^2} \int_a^u \left[\int_u^b \left(y - \frac{(u-x)u + (x-a)y}{(1-t)(b-a)} \right) dy \right] dx \\
 & = \frac{M}{t(1-t)(b-a)^3} \int_a^u \left[\int_u^b \left((b-a)((1-t)y - tx) \right. \right. \\
 & \quad \left. \left. - 2xy + (b+u)x + (a+u)y - 2u^2 \right) dy \right] dx \\
 & = \frac{M}{(1-t)(b-a)^2} \int_a^u \left[\frac{1}{2} ((b-a)(1-t) + (a+u))(b+u) \right. \\
 & \quad \left. - (b-a)tx - 2u^2 \right] dx
 \end{aligned}$$



New Inequalities of Hadamard's Type for Lipschitzian Mappings

Liang-Cheng Wang

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

[Page 12 of 18](#)

$$\begin{aligned}
&= \frac{M}{b-a} \left[\frac{1}{2} ((b-a)(1-t) + (a+u)) (b+u) - (b-a)t(u+a) - 2u^2 \right] \\
&= \frac{M}{2} t(1-t)(b-a).
\end{aligned}$$

This completes the proof of inequality (2.12).

From integral property and (2.2), we have


$$(2.16) \quad tf(a) + (1-t)f(b) = \frac{t}{(1-t)(b-a)} \int_a^u f(a)dx + \frac{1-t}{t(b-a)} \int_u^b f(b)dy.$$

Using (2.16) and (2.2), we obtain

$$\begin{aligned}
&|C(t) - (tf(a) + (1-t)f(b))| \\
&\leq \frac{1}{b-a} \left[\frac{t}{1-t} \int_a^u |f(x) - f(a)| dx + \frac{1-t}{t} \int_u^b |f(y) - f(b)| dy \right] \\
&\leq \frac{M}{b-a} \left[\frac{t}{1-t} \int_a^u |x-a| dx + \frac{1-t}{t} \int_u^b |y-b| dy \right] \\
&= \frac{M}{b-a} \left[\frac{t}{1-t} \int_a^u (x-a) dx + \frac{1-t}{t} \int_u^b (b-y) dy \right] \\
&= \frac{M}{b-a} \left[\frac{t}{1-t} \left(\frac{1}{2}(u^2 - a^2) - a(u-a) \right) + \frac{1-t}{t} \left(b(b-u) - \frac{1}{2}(b^2 - u^2) \right) \right] \\
&= Mt(1-t)(b-a).
\end{aligned}$$

This completes the proof of inequality (2.13).

This completes the proof of Theorem 2.3. □



New Inequalities of Hadamard's Type for Lipschitzian Mappings

Liang-Cheng Wang

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

[Page 13 of 18](#)

Corollary 2.4. Let f be defined as in Theorem 2.1. For any $t \in (0, 1)$, then

$$(2.17) \quad |C(t) - f(ta + (1-t)b)| \leq Mt(1-t)(b-a),$$

$$(2.18) \quad |A(t) - C(t)| \leq Mt(1-t)(b-a),$$

$$(2.19) \quad |A(t) - (tf(a) + (1-t)f(b))| \leq 2Mt(1-t)(b-a),$$

$$(2.20) \quad |B(t) - (tf(a) + (1-t)f(b))| \leq \frac{3M}{2}t(1-t)(b-a),$$

and

$$(2.21) \quad |f(ta + (1-t)b) - (tf(a) + (1-t)f(b))| \leq 2Mt(1-t)(b-a).$$

Proof. Using (2.5) and (2.12), we get inequality (2.18):

$$|A(t) - C(t)| \leq |A(t) - B(t)| + |B(t) - C(t)| \leq Mt(1-t)(b-a).$$

Using the same method as the proof of (2.18), from (2.6) and (2.12), (2.13) and (2.18), (2.12) and (2.13), and (2.13) and (2.17), we get (2.17), (2.19), (2.20) and (2.21), respectively.

This completes the proof of Corollary 2.4. \square



New Inequalities of Hadamard's Type for Lipschitzian Mappings

Liang-Cheng Wang

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

[Page 14 of 18](#)

Remark 1. If we let $t = \frac{1}{2}$, then (2.13), (2.17) and (2.21) reduce to

$$(2.22) \quad \left| \frac{1}{b-a} \int_a^b f(x)dx - \frac{f(a) + f(b)}{2} \right| \leq \frac{M}{4}(b-a),$$

$$(2.23) \quad \left| \frac{1}{b-a} \int_a^b f(x)dx - f\left(\frac{a+b}{2}\right) \right| \leq \frac{M}{4}(b-a),$$


$$(2.24) \quad \left| f\left(\frac{a+b}{2}\right) - \frac{f(a) + f(b)}{2} \right| \leq \frac{M}{2}(b-a).$$

(2.22) is better than (2.2) in [3]. (2.23) and (2.24) are (2.1) and (2.4) in [3], respectively.

Corollary 2.5. Let f be a convex mapping on $[a, b]$, with $f'_+(a)$ and $f'_-(b)$ existing. For any $t \in (0, 1)$, then we have

$$(2.25) \quad \begin{aligned} 0 &\leq A(t) - f(ta + (1-t)b) \\ &\leq \frac{\max\{|f'_+(a)|, |f'_-(b)|\}}{3} t(1-t)(b-a), \end{aligned}$$

$$(2.26) \quad \begin{aligned} 0 &\leq B(t) - A(t) \\ &\leq \frac{\max\{|f'_+(a)|, |f'_-(b)|\}}{2} t(1-t)(b-a), \end{aligned}$$



New Inequalities of Hadamard's Type for Lipschitzian Mappings

Liang-Cheng Wang

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

Page 15 of 18

$$(2.27) \quad \begin{aligned} 0 &\leq B(t) - f(ta + (1-t)b) \\ &\leq \frac{\max\{|f'_+(a)|, |f'_-(b)|\}}{2} t(1-t)(b-a), \end{aligned}$$


$$(2.28) \quad \begin{aligned} 0 &\leq C(t) - B(t) \\ &\leq \frac{\max\{|f'_+(a)|, |f'_-(b)|\}}{2} t(1-t)(b-a), \end{aligned}$$

$$(2.29) \quad \begin{aligned} 0 &\leq tf(a) + (1-t)f(b) - C(t) \\ &\leq \max\{|f'_+(a)|, |f'_-(b)|\} t(1-t)(b-a), \end{aligned}$$

$$(2.30) \quad \begin{aligned} 0 &\leq C(t) - f(ta + (1-t)b) \\ &\leq \max\{|f'_+(a)|, |f'_-(b)|\} t(1-t)(b-a), \end{aligned}$$

$$(2.31) \quad \begin{aligned} 0 &\leq C(t) - A(t) \\ &\leq \max\{|f'_+(a)|, |f'_-(b)|\} t(1-t)(b-a), \end{aligned}$$

$$(2.32) \quad \begin{aligned} 0 &\leq tf(a) + (1-t)f(b) - A(t) \\ &\leq 2 \max\{|f'_+(a)|, |f'_-(b)|\} t(1-t)(b-a), \end{aligned}$$



New Inequalities of Hadamard's Type for Lipschitzian Mappings

Liang-Cheng Wang

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

[Page 16 of 18](#)

$$(2.33) \quad \begin{aligned} 0 &\leq tf(a) + (1-t)f(b) - B(t) \\ &\leq \frac{3}{2} \max\{|f'_+(a)|, |f'_-(b)|\} t(1-t)(b-a), \end{aligned}$$

and

$$(2.34) \quad \begin{aligned} 0 &\leq tf(a) + (1-t)f(b) - f(ta + (1-t)b) \\ &\leq 2 \max\{|f'_+(a)|, |f'_-(b)|\} t(1-t)(b-a). \end{aligned}$$

Proof. For any $x, y \in [a, b]$, from the properties of convex functions, we have the following $\max\{|f'_+(a)|, |f'_-(b)|\}$ -Lipschitzian inequality (see [7]):

$$(2.35) \quad |f(x) - f(y)| \leq \max\{|f'_+(a)|, |f'_-(b)|\} |x - y|.$$

Using (1.2), (2.35), Theorem 2.1 – 2.2 and Corollary 2.4, we obtain (2.25) – (2.34).

This completes the proof of Corollary 2.5. \square

Remark 2. The condition in Corollary 2.5 is better than that in Corollary 2.2, Corollary 4.2 and Theorem 3.3 of [3]. Actually, it is that f is a differentiable convex mapping on $[a, b]$ with $M = \sup_{t \in [a, b]} |f'(t)| < \infty$.



New Inequalities of Hadamard's Type for Lipschitzian Mappings

Liang-Cheng Wang

Title Page

Contents



Go Back


Close

Quit

Page 17 of 18

References

- [1] J. HADAMARD, Etude sur les propriétés des fonctions entières et en particulier d'une fonction considérée par Riemann, *J. Math. Pures Appl.*, **58** (1893), 171–215.
- [2] L.-C. WANG, On extensions and refinements of Hermite-Hadamard inequalities for convex functions, *Math. Ineq. Appl.*, **6**(4) (2003), 659–666.
- [3] S.S. DRAGOMIR, Y.J. CHO AND S.S. KIM, Inequalities of Hadamard's type for Lipschitzian mappings and their applications, *J. Math. Anal. Appl.*, **245** (2000), 489–501.
- [4] G.-S. YANG AND K.-L. TSENG, Inequalities of Hadamard's type for Lipschitzian mappings, *J. Math. Anal. Appl.*, **260** (2001), 230–238.
- [5] M. MATIC AND J. PEĆARIĆ, Note on inequalities of Hadamard's type for Lipschitzian mappings, *Tamkang J. Math.*, **32**(2) (2001), 127–130.
- [6] L.-C. WANG, On some extention of Hadamard inequalities of convex function, *Math. in Practice and Theory*, **32** (2002), 1027–1030. (In Chinese).
- [7] L.-C. WANG, *Convex Functions and Their Inequalities*, Sichuan University Press, Chengdu, China, 2001, pp.55. (In Chinese).



New Inequalities of Hadamard's Type for Lipschitzian Mappings

Liang-Cheng Wang

Title Page

Contents



Go Back

Close

Quit

Page 18 of 18