

Journal of Inequalities in Pure and Applied Mathematics

GENERALIZATIONS OF THE TRIANGLE INEQUALITY

SABUROU SAITOH

Department of Mathematics,
Faculty of Engineering
Gunma University,
Kiryu 376-8515, Japan.

EMail: ssaitoh@math.sci.gunma-u.ac.jp



volume 4, issue 3, article 62,
2003.

*Received 09 December, 2002;
accepted 15 March, 2003.*

Communicated by: T. Ando

[Abstract](#)

[Contents](#)




[Home Page](#)

[Go Back](#)

[Close](#)

[Quit](#)





Abstract

The triangle inequality $\|x + y\| \leq (\|x\| + \|y\|)$ is well-known and fundamental. Since the 8th General Inequalities meeting in Hungary (September 15-21, 2002), the author has been considering an idea that as triangle inequality, the inequality $\|x + y\|^2 \leq 2(\|x\|^2 + \|y\|^2)$ may be more suitable. The triangle inequality $\|x + y\|^2 \leq 2(\|x\|^2 + \|y\|^2)$ will be naturally generalized for some natural sum of any two members f_j of any two Hilbert spaces \mathcal{H}_j . We shall introduce a natural sum Hilbert space for two arbitrary Hilbert spaces.

2000 Mathematics Subject Classification: 30C40, 46E32, 44A10, 35A22.

Key words: Triangle inequality, Hilbert space, Sum of two Hilbert spaces, various operators among Hilbert spaces, Reproducing kernel, Linear mapping, Norm inequality.

Contents

1	A General Concept	3
2	Sum	6
3	Example	10
4	Discussions	12
	References	

Generalizations of the Triangle Inequality

Saburou Saitoh

Title Page

Contents



Go Back

Close

Quit

Page 2 of 13

1. A General Concept

Following [1], we shall introduce a general theory for linear mappings in the framework of Hilbert spaces.

Let \mathcal{H} be a Hilbert (possibly finite-dimensional) space. Let E be an abstract set and \mathbf{h} be a Hilbert \mathcal{H} -valued function on E . Then we shall consider the linear mapping

$$(1.1) \quad f(p) = (\mathbf{f}, \mathbf{h}(p))_{\mathcal{H}}, \quad \mathbf{f} \in \mathcal{H}$$

from \mathcal{H} into the linear space $\mathcal{F}(E)$ comprising all the complex-valued functions on E . In order to investigate the linear mapping (1.1), we form a positive matrix $K(p, q)$ on E defined by

$$(1.2) \quad K(p, q) = (\mathbf{h}(q), \mathbf{h}(p))_{\mathcal{H}} \text{ on } E \times E.$$


Then, we obtain the following:

- (I) The range of the linear mapping (1.1) by \mathcal{H} is characterized as the reproducing kernel Hilbert space $H_K(E)$ admitting the reproducing kernel $K(p, q)$.
- (II) In general, we have the inequality

$$\|f\|_{H_K(E)} \leq \|\mathbf{f}\|_{\mathcal{H}}.$$

Here, for a member f of $H_K(E)$ there exists a uniquely determined $\mathbf{f}^* \in \mathcal{H}$ satisfying

$$f(p) = (\mathbf{f}^*, \mathbf{h}(p))_{\mathcal{H}} \text{ on } E$$



Generalizations of the Triangle
Inequality

Saburou Saitoh

Title Page

Contents



Go Back

Close

Quit

Page 3 of 13

and

$$\|f\|_{H_K(E)} = \|\mathbf{f}^*\|_{\mathcal{H}}.$$

(III) In general, we have the inversion formula in (1.1) in the form

$$(1.3) \quad f \rightarrow \mathbf{f}^*$$

in (II) by using the reproducing kernel Hilbert space $H_K(E)$. However, this formula is, in general, involved and intricate. Indeed, we need specific arguments. Here, we assume that the inversion formula (1.3) may be, in general, established.

Now we shall consider two systems

$$(1.4) \quad f_j(p) = (\mathbf{f}_j, \mathbf{h}_j(p))_{\mathcal{H}_j}, \quad \mathbf{f}_j \in \mathcal{H}_j$$

in the above way by using $\{\mathcal{H}_j, E, \mathbf{h}_j\}_{j=1}^2$. Here, we assume that E is the same set for both systems in order to have the output functions $f_1(p)$ and $f_2(p)$ on the same set E .


For example, we can consider the usual operator

$$f_1(p) + f_2(p)$$

in $\mathcal{F}(E)$. Then, we can consider the following problem:

How do we represent the sum $f_1(p) + f_2(p)$ on E in terms of their inputs \mathbf{f}_1 and \mathbf{f}_2 through one system?

We shall see that by using the theory of reproducing kernels we can give a natural answer to this problem. Following similar ideas, we can consider



Generalizations of the Triangle Inequality

Saburou Saitoh

Title Page

Contents




Go Back

Close

Quit

Page 4 of 13

various operators among Hilbert space (see [2] for the details). In particular, for the product of two Hilbert spaces, the idea gives generalizations of convolutions and the related natural convolution norm inequalities. These norm inequalities gave various generalizations and applications to forward and inverse problems for linear mappings in the framework of Hilbert spaces, see for example, [3, 4, 5]. Furthermore, surprisingly enough, for some very general nonlinear systems, we can consider similar problems. For its importance in general inequalities, we shall refer to natural generalizations of the triangle inequality $\|\mathbf{x} + \mathbf{y}\|^2 \leq 2(\|\mathbf{x}\|^2 + \|\mathbf{y}\|^2)$ for some natural sum of two arbitrary Hilbert spaces. And then, we may consider this inequality to be more suitable as a triangle inequality than the usual triangle inequality.



Generalizations of the Triangle Inequality

Saburou Saitoh

Title Page

Contents



Go Back

Close

Quit

Page 5 of 13

2. Sum

By (I), $f_1 \in H_{K_1}(E)$ and $f_2 \in H_{K_2}(E)$, and we note that for the reproducing kernel Hilbert space $H_{K_1+K_2}(E)$ admitting the reproducing kernel

$$K_1(p, q) + K_2(p, q) \text{ on } E,$$

$H_{K_1+K_2}(E)$ is composed of all functions

$$(2.1) \quad f(p) = f_1(p) + f_2(p); \quad f_j \in H_{K_j}(E)$$

and its norm $\|f\|_{H_{K_1+K_2}(E)}$ is given by

$$(2.2) \quad \|f\|_{H_{K_1+K_2}(E)}^2 = \min \left\{ \|f_1\|_{H_{K_1}(E)}^2 + \|f_2\|_{H_{K_2}(E)}^2 \right\}$$

where the minimum is taken over $f_j \in H_{K_j}(E)$ satisfying (2.1) for f . Hence, in general, we have the inequality

$$(2.3) \quad \|f_1 + f_2\|_{H_{K_1+K_2}(E)}^2 \leq \|f_1\|_{H_{K_1}(E)}^2 + \|f_2\|_{H_{K_2}(E)}^2.$$


In particular, note that for the same K_1 and K_2 , we have, for $K_1 = K_2 = K$

$$\|f_1 + f_2\|_{H_K(E)}^2 \leq 2(\|f_1\|_{H_K(E)}^2 + \|f_2\|_{H_K(E)}^2).$$

Furthermore, the particular inequality (2.3) may be considered as a natural triangle inequality for the sum of reproducing kernel Hilbert spaces $H_{K_j}(E)$.

For the positive matrix $K_1 + K_2$ on E , we assume the expression in the form

$$(2.4) \quad K_1(p, q) + K_2(p, q) = (\mathbf{h}_S(q), \mathbf{h}_S(p))_{\mathcal{H}_S} \text{ on } E \times E$$



Generalizations of the Triangle Inequality

Sabrou Saitoh

Title Page

Contents



Go Back

Close

Quit

Page 6 of 13

with a Hilbert space \mathcal{H}_S -valued function on E and further we assume that

$$(2.5) \quad \{\mathbf{h}_S(p); p \in E\} \text{ is complete in } \mathcal{H}_S.$$

Such a representation is, in general, possible ([1, page 36 and see Chapter 1, §5]). Then, we can consider the linear mapping from \mathcal{H}_S onto $H_{K_1+K_2}(E)$

$$(2.6) \quad f_S(p) = (\mathbf{f}_S, \mathbf{h}_S(p))_{\mathcal{H}_S}, \mathbf{f}_S \in \mathcal{H}_S$$

and we obtain the isometric identity

$$(2.7) \quad \|f_S\|_{H_{K_1+K_2}(E)} = \|\mathbf{f}_S\|_{\mathcal{H}_S}.$$

Hence, for such representations (2.4) with (2.5), we obtain the isometric relations among the Hilbert spaces \mathcal{H}_S .


Now, for the sum $f_1(p) + f_2(p)$ there exists a uniquely determined $\mathbf{f}_S \in \mathcal{H}_S$ satisfying

$$(2.8) \quad f_1(p) + f_2(p) = (\mathbf{f}_S, \mathbf{h}_S(p))_{\mathcal{H}_S} \text{ on } E.$$

Then, \mathbf{f}_S will be considered as a sum of \mathbf{f}_1 and \mathbf{f}_2 through these mappings and so, we shall introduce the notation

$$(2.9) \quad \mathbf{f}_S = \mathbf{f}_1[+]\mathbf{f}_2.$$

This sum for the members $\mathbf{f}_1 \in \mathcal{H}_1$ and $\mathbf{f}_2 \in \mathcal{H}_2$ is introduced through the three mappings induced by $\{\mathcal{H}_j, E, \mathbf{h}_j\}$ ($j = 1, 2$) and $\{\mathcal{H}_S, E, \mathbf{h}_S\}$.



Generalizations of the Triangle Inequality

Sabourou Saitoh

Title Page

Contents



Go Back

Close

Quit

Page 7 of 13

The operator $\mathbf{f}_1[+]\mathbf{f}_2$ is expressible in terms of \mathbf{f}_1 and \mathbf{f}_2 by the inversion formula

$$(2.10) \quad (\mathbf{f}_1, \mathbf{h}_1(p))_{\mathcal{H}_1} + (\mathbf{f}_2, \mathbf{h}_2(p))_{\mathcal{H}_2} \longrightarrow \mathbf{f}_1[+]\mathbf{f}_2$$

in the sense (II) from $H_{K_1+K_2}(E)$ onto \mathcal{H}_S . Then, from (II) and (2.5) we obtain the beautiful triangle inequality

Theorem 2.1. *We have the triangle inequality*

$$(2.11) \quad \|\mathbf{f}_1[+]\mathbf{f}_2\|_{\mathcal{H}_S}^2 \leq \|\mathbf{f}_1\|_{\mathcal{H}_1}^2 + \|\mathbf{f}_2\|_{\mathcal{H}_2}^2.$$

If $\{\mathbf{h}_j(p); p \in E\}$ are complete in \mathcal{H}_j ($j = 1, 2$), then \mathcal{H}_j and H_{K_j} are isometrical, respectively. By using the isometric mappings induced by Hilbert space valued functions \mathbf{h}_j ($j = 1, 2$) and \mathbf{h}_S , we can introduce the sum space of \mathcal{H}_1 and \mathcal{H}_2 in the form

$$(2.12) \quad \mathcal{H}_1[+]\mathcal{H}_2$$


through the mappings. Of course, the sum is a Hilbert space. Furthermore, such spaces are determined in the framework of isometric relations.

For example, if for some positive number γ

$$(2.13) \quad K_1 \ll \gamma^2 K_2 \text{ on } E$$

that is, if $\gamma^2 K_2 - K_1$ is a positive matrix on E , then we have

$$(2.14) \quad H_{K_1}(E) \subset H_{K_2}(E)$$



Generalizations of the Triangle
Inequality

Sabrou Saitoh

Title Page

Contents



Go Back

Close

Quit

Page 8 of 13


and

$$(2.15) \quad \|f_1\|_{H_{K_2}(E)} \leq \gamma \|f_1\|_{H_{K_1}(E)} \text{ for } f_1 \in H_{K_1}(E)$$

([1, page 37]). Hence, in this case, we need not to introduce a new Hilbert space \mathcal{H}_S and the linear mapping (2.6) in Theorem 2.1 and we can use the linear mapping

$$(\mathbf{f}_2, \mathbf{h}_2(p))_{\mathcal{H}_2}, \quad \mathbf{f}_2 \in \mathcal{H}_2$$

instead of (2.6) in Theorem 2.1.



Generalizations of the Triangle Inequality

Saburou Saitoh

Title Page

Contents



Go Back

Close

Quit

Page 9 of 13

3. Example

We shall consider two linear transforms

$$(3.1) \quad f_j(p) = \int_T F_j(t) \overline{h(t, p)} \rho_j(t) dm(t), \quad p \in E$$

where ρ_j are positive continuous functions on T ,

$$(3.2) \quad \int_T |h(t, p)|^2 \rho_j(t) dm(t) < \infty \text{ on } p \in E$$

and

$$(3.3) \quad \int_T |F_j(t)|^2 \rho_j(t) dm(t) < \infty.$$

We assume that $\{h(t, p); p \in E\}$ is complete in the spaces satisfying (3.3).

Then, we consider the associated reproducing kernels on E


$$K_j(p, q) = \int_T h(t, p) \overline{h(t, q)} \rho_j(t) dm(t)$$

and, for example we consider the expression

$$(3.4) \quad K_1(p, q) + K_2(p, q) = \int_T h(t, p) \overline{h(t, q)} (\rho_1(t) + \rho_2(t)) dm(t).$$

So, we can consider the linear transform

$$(3.5) \quad f(p) = \int_T F(t) \overline{h(t, p)} (\rho_1(t) + \rho_2(t)) dm(t)$$



Generalizations of the Triangle Inequality

Sabrou Saitoh

Title Page

Contents



Go Back

Close

Quit


Page 10 of 13

for functions F satisfying

$$\int_T |F(t)|^2 (\rho_1(t) + \rho_2(t)) dm(t) < \infty.$$

Hence, through the three transforms (3.1) and (3.5) we have the sum

$$(3.6) \quad (F_1[+]F_2)(t) = \frac{F_1(t)\rho_1(t) + F_2(t)\rho_2(t)}{\rho_1(t) + \rho_2(t)}.$$



Generalizations of the Triangle Inequality

Saburou Saitoh

Title Page

Contents



Go Back


Close

Quit

Page 11 of 13

4. Discussions

- Which is more general the classical one and the second triangle inequality (2.11)? We stated: for two arbitrary Hilbert spaces, we can introduce the natural sum and the second triangle inequality (2.11) is valid in a natural way.
- Which is more beautiful the classical one and the second one (2.11)? The author believes the second one is more beautiful. (Gods love two; and 2 is better than 1/2).
- Which is more widely applicable?
- The author thinks the second triangle inequality (2.11) is superior to the classical one as a triangle inequality. However, for the established long tradition, how will be:
 - for the classical one, the triangle inequality of the first kind and
 - for the second inequality (2.11), the triangle inequality of the second kind?



Generalizations of the Triangle Inequality

Saburou Saitoh

Title Page

Contents



Go Back


Close

Quit

Page 12 of 13

References

- [1] S. SAITOH, *Integral Transforms, Reproducing Kernels and their Applications*, Pitman Research Notes in Mathematics Series, **369**, Addison Wesley Longman, UK (1998). Introduction to the theory of reproducing kernels, Makino-shoten (2002) (in Japanese).
- [2] S. SAITOH, Various operators in Hilbert space introduced by transforms, *International. J. Appl. Math.*, **1** (1999), 111–126.
- [3] S. SAITOH, Weighted L_p -norm inequalities in convolutions, *Survey on Classical Inequalities*, Kluwer Academic Publishers, The Netherlands, (2000), 225–234.
- [4] S. SAITOH, V.K. TUAN AND M. YAMAMOTO, Reverse weighted L_p -norm inequalities in convolutions and stability in inverse problems, *J. Ineq. Pure and Appl. Math.*, **1**(1) (2000), Art. 7. [ONLINE: http://jipam.vu.edu.au/v1n1/018_99.html].
- [5] S. SAITOH, V.K. TUAN AND M. YAMAMOTO, Reverse convolution inequalities and applications to inverse heat source problems, *J. Ineq. Pure and Appl. Math.*, **3**(5) (2003), Art. 80. [ONLINE: http://jipam.vu.edu.au/v3n5/029_02.html].



Generalizations of the Triangle Inequality

Saburou Saitoh

Title Page

Contents



Go Back

Close

Quit

Page 13 of 13