

Journal of Inequalities in Pure and Applied Mathematics

CONVOLUTION CONDITIONS FOR SPIRALLIKENESS AND CONVEX SPIRALLIKENESS OF CERTAIN MEROMORPHIC p -VALENT FUNCTIONS



V. RAVICHANDRAN, S. SIVAPRASAD KUMAR AND
K.G. SUBRAMANIAN

Department of Computer Applications
Sri Venkateswara College of Engineering
Pennalur, Sriperumbudur 602 105, India.
EMail: vravi@svce.ac.in

Department of Mathematics
Sindhi College
123 P. H. Road, Numbal
Chennai 600 077, India.
EMail: sivpk71@yahoo.com

Department of Mathematics
Madras Christian College
Tambaram, Chennai 600 059, India.
EMail: kgsmani@vsnl.net

volume 5, issue 1, article 11,
2004.

Received 27 October, 2003;
accepted 11 December, 2003.

Communicated by: *H. Silverman*

Abstract

Contents



Home Page

Go Back

Close

Quit



Abstract

In the present investigation, the authors derive necessary and sufficient conditions for spirallikeness and convex spirallikeness of a suitably normalized meromorphic p -valent function in the punctured unit disk, using convolution. Also we give an application of our result to obtain a convolution condition for a class of meromorphic functions defined by a linear operator.

2000 Mathematics Subject Classification: Primary 30C45, Secondary 30C80.

Key words: Meromorphic p -valent functions, Analytic functions, Starlike functions, Convex functions, Spirallike functions, Convex Spirallike functions, Hadamard product (or Convolution), Subordination, Linear operator.

Contents

1	Introduction	3
2	Convolution condition for the class $S_p^\lambda(\varphi)$	7
3	Convolution condition for the class $C_p^\lambda(\varphi)$	10
4	Convolution conditions for a class of function defined by linear operator	11

References

Convolution Conditions for Spirallikeness and Convex Spirallikeness of Certain Meromorphic p -valent Functions

V. Ravichandran, S. Sivaprasad Kumar
and K.G. Subramanian

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

Page 2 of 16

1. Introduction

Let Σ_p be the class of *meromorphic* functions

$$(1.1) \quad f(z) = \frac{1}{z^p} + \sum_{n=1-p}^{\infty} a_n z^n \quad (p \in \mathbb{N} := \{1, 2, 3, \dots\}),$$

which are *analytic* and p -valent in the punctured unit disk

$$\mathbb{E}^* := \{z : z \in \mathbb{C} \text{ and } 0 < |z| < 1\} = \mathbb{E} \setminus \{0\},$$

where $\mathbb{E} := \{z : z \in \mathbb{C} \text{ and } |z| < 1\}$.

For two functions f and g analytic in \mathbb{E} , we say that the function $f(z)$ is subordinate to $g(z)$ in \mathbb{E} and write

$$f \prec g \quad \text{or} \quad f(z) \prec g(z) \quad (z \in \mathbb{E}),$$

if there exists a Schwarz function $w(z)$, analytic in \mathbb{E} with

$$w(0) = 0 \quad \text{and} \quad |w(z)| < 1 \quad (z \in \mathbb{E}),$$


such that

$$(1.2) \quad f(z) = g(w(z)) \quad (z \in \mathbb{E}).$$

In particular, if the function g is univalent in \mathbb{E} , the above subordination is equivalent to

$$f(0) = g(0) \quad \text{and} \quad f(\mathbb{E}) \subset g(\mathbb{E}).$$

We define two subclasses of meromorphic p -valent functions in the following:



Convolution Conditions for
Spirallikeness and Convex
Spirallikeness of Certain
Meromorphic p -valent
Functions

V. Ravichandran, S. Sivaprasad Kumar
and K.G. Subramanian

Title Page

Contents



Go Back

Close

Quit

Page 3 of 16

Definition 1.1. Let $|\lambda| < \frac{\pi}{2}$ and $p \in \mathbb{N}$. Let φ be an analytic function in the unit disk \mathbb{E} . We define the classes $S_p^\lambda(\varphi)$ and $C_p^\lambda(\varphi)$ by

$$(1.3) \quad S_p^\lambda(\varphi) := \left\{ f \in \Sigma_p : \frac{zf'(z)}{f(z)} \prec -pe^{-i\lambda} [\cos \lambda \varphi(z) + i \sin \lambda] \right\},$$

$$(1.4) \quad C_p^\lambda(\varphi) := \left\{ f \in \Sigma_p : 1 + \frac{zf''(z)}{f'(z)} \prec -pe^{-i\lambda} [\cos \lambda \varphi(z) + i \sin \lambda] \right\}.$$

Analogous to the well known Alexander equivalence [2], we have

$$(1.5) \quad f \in C_p^\lambda(\varphi) \Leftrightarrow -\frac{1}{p}zf' \in S_p^\lambda(\varphi) \quad (p \in \mathbb{N}).$$

Remark 1.1. For

$$(1.6) \quad \varphi(z) = \frac{1+Az}{1+Bz} \quad (-1 \leq B < A \leq 1),$$

we set

$$S_p^\lambda(\varphi) =: S_p^\lambda[A, B] \quad \text{and} \quad C_p^\lambda(\varphi) =: C_p^\lambda[A, B].$$

For $\lambda = 0$, we write

$$S_p^0(\varphi) =: S_p^*(\varphi) \quad \text{and} \quad C_p^0(\varphi) =: C_p(\varphi),$$

$$S_p^0[A, B] =: S_p[A, B] \quad \text{and} \quad C_p^0[A, B] =: C_p[A, B].$$

For $0 \leq \alpha < 1$, the classes $S_p^\lambda[1-2\alpha, -1]$ and $C_p^\lambda[1-2\alpha, -1]$ reduces to the classes $S_p^\lambda(\alpha)$ and $C_p^\lambda(\alpha)$ of meromorphic p -valently λ -spirallike functions



Convolution Conditions for Spirallikeness and Convex Spirallikeness of Certain Meromorphic p -valent Functions

V. Ravichandran, S. Sivaprasad Kumar
and K.G. Subramanian

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

[Page 4 of 16](#)

of order α and meromorphic p -valently λ -convex spirallike functions of order α in \mathbb{E}^* respectively:

$$S_p^\lambda(\alpha) := \left\{ f \in \Sigma_p : \Re \left\{ e^{i\lambda} \frac{zf'(z)}{f(z)} \right\} < -p\alpha \cos \lambda \quad (0 \leq \alpha < 1; |\lambda| < \frac{\pi}{2}) \right\},$$

$$\begin{aligned} C_p^\lambda(\alpha) := \left\{ f \in \Sigma_p : \Re \left\{ e^{i\lambda} \left(1 + \frac{zf''(z)}{f'(z)} \right) \right\} \right. \\ \left. < -p\alpha \cos \lambda \quad (0 \leq \alpha < 1; |\lambda| < \frac{\pi}{2}) \right\}. \end{aligned}$$

The classes $S_p^0[1-2\alpha, -1]$ and $C_p^0[1-2\alpha, -1]$ reduces to the classes $S_p^*(\alpha)$ and $C_p(\alpha)$ of meromorphic p -valently starlike functions of order α and meromorphic p -valently convex functions of order α in \mathbb{E}^* respectively.


For two functions $f(z)$ given by (1.1) and

$$(1.7) \quad g(z) = \frac{1}{z^p} + \sum_{n=1-p}^{\infty} b_n z^n \quad (p \in \mathbb{N}),$$

the Hadamard product (or convolution) of f and g is defined by

$$(1.8) \quad (f * g)(z) := \frac{1}{z^p} + \sum_{n=1-p}^{\infty} a_n b_n z^n =: (g * f)(z).$$

Many important properties of certain subclasses of meromorphic p -valent functions were studied by several authors including Aouf and Srivastava [1],



Convolution Conditions for Spirallikeness and Convex Spirallikeness of Certain Meromorphic p -valent Functions

V. Ravichandran, S. Sivaprasad Kumar
and K.G. Subramanian

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

Page 5 of 16

Joshi and Srivastava [3], Liu and Srivastava [4], Liu and Owa [5], Liu and Srivastava [6], Owa *et al.* [7] and Srivastava *et al.* [9]. Motivated by the works of Silverman *et al.* [8], Liu and Owa [5] have obtained the following Theorem 1.1 with $\lambda = 0$ for the class $S_p^*(\alpha)$ and Liu and Srivastava [6] have obtained it for the classes $S_p^\lambda(\alpha)$ (with a slightly different definition of the class).

Theorem 1.1. [6, Theorem 1, p. 14] Let $f(z) \in \Sigma_p$. Then

$$f \in S_p^\lambda(\alpha) \quad (0 \leq \alpha < 1; |\lambda| < \frac{\pi}{2}; p \in \mathbb{N})$$


if and only if

$$f(z) * \left[\frac{1 - \Omega z}{z^p(1 - z)^2} \right] \neq 0 \quad (z \in \mathbb{E}^*),$$

where

$$\Omega := \frac{1 + x + 2p(1 - \alpha) \cos \lambda e^{-i\lambda}}{2p(1 - \alpha) \cos \lambda e^{-i\lambda}}, \quad |x| = 1.$$

In the present investigation, we extend the Theorem 1.1 for the above defined class $S_p^\lambda(\varphi)$. As a consequence, we obtain a convolution condition for the functions in the class $C_p^\lambda(\varphi)$. Also we apply our result to obtain a convolution condition for a class of meromorphic functions defined by a linear operator.



Convolution Conditions for Spirallikeness and Convex Spirallikeness of Certain Meromorphic p -valent Functions

V. Ravichandran, S. Sivaprasad Kumar
and K.G. Subramanian

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

[Page 6 of 16](#)

2. Convolution condition for the class $S_p^\lambda(\varphi)$

We begin with the following result for the general class $S_p^\lambda(\varphi)$:

Theorem 2.1. *Let φ be analytic in \mathbb{E} and be defined on $\partial\mathbb{E} := \{z \in \mathbb{C} : |z| = 1\}$. The function $f \in \Sigma_p$ is in the class $S_p^\lambda(\varphi)$ if and only if*

$$(2.1) \quad f(z) * \frac{1 - \Psi z}{z^p(1 - z)^2} \neq 0 \quad (z \in \mathbb{E}^*)$$

where

$$(2.2) \quad \Psi := \frac{1 + p \{1 - e^{-i\lambda}[\cos \lambda \varphi(x) + i \sin \lambda]\}}{p \{1 - e^{-i\lambda}[\cos \lambda \varphi(x) + i \sin \lambda]\}} \quad (|x| = 1; |\lambda| < \frac{\pi}{2}).$$

Proof. In view of (1.3), $f(z) \in S_p^\lambda(\varphi)$ if and only if


$$\frac{zf'(z)}{f(z)} \neq -pe^{-i\lambda}[\cos \lambda \varphi(x) + i \sin \lambda] \quad (z \in \mathbb{E}^*; |x| = 1; |\lambda| < \frac{\pi}{2})$$

or

$$(2.3) \quad zf'(z) + pe^{-i\lambda}[\cos \lambda \varphi(x) + i \sin \lambda]f(z) \neq 0 \quad (z \in \mathbb{E}^*; |x| = 1; |\lambda| < \frac{\pi}{2}).$$

For $f \in \Sigma_p$ given by (1.1), we have

$$(2.4) \quad f(z) = f(z) * \frac{1}{z^p(1 - z)} \quad (z \in \mathbb{E}^*)$$



Convolution Conditions for
Spirallikeness and Convex
Spirallikeness of Certain
Meromorphic p -valent
Functions

V. Ravichandran, S. Sivaprasad Kumar
and K.G. Subramanian

Title Page

Contents



Go Back

Close

Quit

Page 7 of 16

and

$$(2.5) \quad zf'(z) = f(z) * \left[\frac{1}{z^p(1-z)^2} - \frac{p+1}{z^p(1-z)} \right] \quad (z \in \mathbb{E}^*).$$

By making use of the convolutions (2.5) and (2.4) in (2.3), we have

$$f(z) * \left[\frac{1}{z^p(1-z)^2} - \frac{p+1}{z^p(1-z)} + \frac{pe^{-i\lambda}[\cos \lambda \varphi(x) + i \sin \lambda]}{z^p(1-z)} \right] \neq 0$$
$$(z \in \mathbb{E}^*; |x| = 1; |\lambda| < \frac{\pi}{2})$$

or

$$f(z) * \left[\frac{p \{ e^{-i\lambda}[\cos \lambda \varphi(x) + i \sin \lambda] - 1 \}}{z^p(1-z)^2} \right. \\ \left. + \frac{[1 + p \{ 1 - e^{-i\lambda}[\cos \lambda \varphi(x) + i \sin \lambda] \}] z}{z^p(1-z)^2} \right] \neq 0$$
$$(z \in \mathbb{E}^*; |x| = 1; |\lambda| < \frac{\pi}{2}),$$

which yields the desired convolution condition (2.1) of Theorem 2.1. \square

By taking $\lambda = 0$ in the Theorem 2.1, we obtain the following result for the class $S_p^*(\varphi)$.

Corollary 2.2. *Let φ be analytic in \mathbb{E} and be defined on $\partial\mathbb{E}$. The function $f \in \Sigma_p$ is in the class $f \in S_p^*(\varphi)$ if and only if*

$$(2.6) \quad f(z) * \frac{1 - \Upsilon z}{z^p(1-z)^2} \neq 0 \quad (z \in \mathbb{E}^*)$$



Convolution Conditions for
Spirallikeness and Convex
Spirallikeness of Certain
Meromorphic p -valent
Functions

V. Ravichandran, S. Sivaprasad Kumar
and K.G. Subramanian

Title Page

Contents



Go Back

Close

Quit

Page 8 of 16

where

$$(2.7) \quad \Upsilon := \frac{1 + p(1 - \varphi(x))}{p(1 - \varphi(x))} \quad (|x| = 1).$$

By taking $\varphi(z) = (1 + Az)/(1 + Bz)$, $-1 \leq B < A \leq 1$ in Theorem 2.1, we obtain the following result for the class $S_p^\lambda[A, B]$.

Corollary 2.3. *The function $f \in \Sigma_p$ is in the class $S_p^\lambda[A, B]$ if and only if*

$$(2.8) \quad f(z) * \frac{1 - \Upsilon z}{z^p(1 - z)^2} \neq 0 \quad (z \in \mathbb{E}^*)$$

where

$$(2.9) \quad \Upsilon := \frac{x - B + p(A - B) \cos \lambda e^{-i\lambda}}{p(A - B) \cos \lambda e^{-i\lambda}} \quad (|x| = 1).$$

Remark 2.1. By taking $A = 1 - 2\alpha$, $B = -1$ in the above Corollary 2.3, we obtain Theorem 1.1 of Liu and Srivastava [6].



Convolution Conditions for
Spirallikeness and Convex
Spirallikeness of Certain
Meromorphic p -valent
Functions

V. Ravichandran, S. Sivaprasad Kumar
and K.G. Subramanian

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

Page 9 of 16

3. Convolution condition for the class $C_p^\lambda(\varphi)$

By making use of Theorem 2.1, we obtain a convolution condition for functions in the class $C_p^\lambda(\varphi)$ in the following:

Theorem 3.1. *Let φ be analytic in \mathbb{E} and be defined on $\partial\mathbb{E}$. The function $f \in \Sigma_p$ is in the class $C_p^\lambda(\varphi)$ if and only if*

$$(3.1) \quad f(z) * \frac{p - [2 + p + (p - 1)\Psi]z + (p + 1)\Psi z^2}{z^p(1 - z)^2} \neq 0 \quad (z \in \mathbb{E}^*)$$

where Ψ is given by (2.2).

Proof. In view of the Alexander-type equivalence (1.5), we find from Theorem 2.1 that $f \in C_p^\lambda(\varphi)$ if and only if

$$zf'(z) * \frac{1 - \Psi z}{z^p(1 - z)^2} = f(z) * z \left(\frac{1 - \Psi z}{z^p(1 - z)^2} \right)' \neq 0 \quad (z \in \mathbb{E}^*)$$


which readily yields the desired assertion (3.1) of Theorem 3.1. \square

By taking $\lambda = 0$ in the Theorem 3.1, we obtain the following result of the class $C_p(\varphi)$.

Corollary 3.2. *Let φ be analytic in \mathbb{E} and be defined on $\partial\mathbb{E}$. The function $f \in \Sigma_p$ is in the class $f \in C_p(\varphi)$ if and only if*

$$(3.2) \quad f(z) * \frac{p - [2 + p + (p - 1)\Upsilon]z + (p + 1)\Upsilon z^2}{z^p(1 - z)^2} \neq 0 \quad (z \in \mathbb{E}^*)$$

where Υ is given by (2.7).



Convolution Conditions for
Spirallikeness and Convex
Spirallikeness of Certain
Meromorphic p -valent
Functions

V. Ravichandran, S. Sivaprasad Kumar
and K.G. Subramanian

Title Page

Contents



Go Back

Close

Quit

Page 10 of 16

4. Convolution conditions for a class of function defined by linear operator

We begin this section by defining a class $\mathcal{T}_{n+p-1}(\varphi)$. First of all for a function $f(z) \in \Sigma_p$, define $D^{n+p-1}f(z)$ by

$$\begin{aligned} D^{n+p-1}f(z) &= f(z) * \left[\frac{1}{z^p(1-z)^{n+p}} \right] \\ &= \frac{(z^{n+2p-1}f(z))^{(n+p-1)}}{(n+p-1)!z^p} \\ &= \frac{1}{z^p} + \sum_{m=1-p}^{\infty} \frac{(m+n+2p-1)!}{(n+p-1)!(m+p)!} a_m z^m. \end{aligned}$$

By making use of the operator $D^{n+p-1}f(z)$, we define the class $\mathcal{T}_{n+p-1}(\varphi)$ by


$$\mathcal{T}_{n+p-1}(\varphi) = \left\{ f(z) \in \Sigma_p : \frac{D^{n+p}f(z)}{D^{n+p-1}f(z)} \prec \varphi(z) \right\}.$$

When

$$\varphi(z) = \frac{1 + (1 - 2\gamma)z}{1 - z}$$

where

$$\gamma = \frac{n+p(2-\alpha)}{n+p} \quad (0 \leq \alpha < 1),$$



Convolution Conditions for
Spirallikeness and Convex
Spirallikeness of Certain
Meromorphic p -valent
Functions

V. Ravichandran, S. Sivaprasad Kumar
and K.G. Subramanian

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

Page 11 of 16

the class $\mathcal{T}_{n+p-1}(\varphi)$ reduces to the following class $\mathcal{T}_{n+p-1}(\alpha)$ studied by Liu and Owa [5]:

$$\mathcal{T}_{n+p-1}(\alpha) = \left\{ f(z) \in \Sigma_p : \Re \left(\frac{D^{n+p}f(z)}{D^{n+p-1}f(z)} - \frac{n+2p}{n+p} \right) < -\frac{p\alpha}{n+p} \right\}.$$

By making use of Corollary 2.2, we prove the following result for the class $\mathcal{T}_{n+p-1}(\varphi)$:

Theorem 4.1. *The function $f(z) \in \Sigma_p$ is in the class $\mathcal{T}_{n+p-1}(\varphi)$ if and only if*

$$(4.1) \quad f(z) * \frac{1 + [(n+p)(1-\Omega) - 1]z}{z^p(1-z)^{n+p+1}} \neq 0 \quad (z \in E^*; |x| = 1),$$

where Ω is given by

$$(4.2) \quad \Omega := \frac{1 + (n+p)(\varphi(x) - 1)}{(n+p)(\varphi(x) - 1)} \quad (|x| = 1).$$

Proof. By making use of the familiar identity

$$z(D^{n+p-1}f(z))' = (n+p)D^{n+p}f(z) - (n+2p)D^{n+p-1}f(z),$$

we have

$$\frac{z(D^{n+p-1}f(z))'}{D^{n+p-1}f(z)} = (n+p) \frac{D^{n+p}f(z)}{D^{n+p-1}f(z)} - (n+2p),$$

and therefore, by using the definition of the class $\mathcal{T}_{n+p-1}(\varphi)$, we see that $f(z) \in \mathcal{T}_{n+p-1}(\varphi)$ if and only if

$$D^{n+p-1}f(z) \in S_p^* \left(\frac{n+2p}{p} - \frac{n+p}{p} \varphi(z) \right).$$



Convolution Conditions for Spirallikeness and Convex Spirallikeness of Certain Meromorphic p -valent Functions

V. Ravichandran, S. Sivaprasad Kumar
and K.G. Subramanian

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

Page 12 of 16

Then, by applying Corollary 2.2 for the function $D^{n+p-1}f(z)$, we have

$$(4.3) \quad D^{n+p-1}f(z) * \frac{1 - \Omega z}{z^p(1 - z)^2} \neq 0,$$

where

$$\begin{aligned}\Omega &= \frac{1 + p \left[1 - \left\{ \frac{n+2p}{p} - \frac{n+p}{p} \varphi(x) \right\} \right]}{p \left[1 - \left\{ \frac{n+2p}{p} - \frac{n+p}{p} \varphi(x) \right\} \right]} \\ &= \frac{1 + (n + p)(\varphi(x) - 1)}{(n + p)(\varphi(x) - 1)}, \quad |x| = 1.\end{aligned}$$

Since

$$D^{n+p-1}f(z) = f(z) * \left[\frac{1}{z^p(1 - z)^{n+p}} \right],$$

the condition (4.3) becomes


$$(4.4) \quad f(z) * \left(g(z) * \frac{1 - \Omega z}{z^p(1 - z)^2} \right) \neq 0$$

where

$$g(z) = \frac{1}{z^p(1 - z)^{n+p}}.$$

By making use of the convolutions (2.5) and (2.4), it is fairly straight forward to show that

$$(4.5) \quad g(z) * \frac{1 - \Omega z}{z^p(1 - z)^2} = \frac{1 + [(n + p)(1 - \Omega) - 1]z}{z^p(1 - z)^{n+p+1}}.$$



Convolution Conditions for Spirallikeness and Convex Spirallikeness of Certain Meromorphic p -valent Functions

V. Ravichandran, S. Sivaprasad Kumar
and K.G. Subramanian

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

Page 13 of 16

By using (4.5) in (4.4), we see that the assertion in (4.1) follows and thus the proof of our Theorem 4.1 is completed. \square

By taking

$$\varphi(z) = \frac{1 + (1 - 2\gamma)z}{1 - z}$$

where

$$\gamma = \frac{n + p(2 - \alpha)}{n + p} \quad (0 \leq \alpha < 1)$$


in our Theorem 4.1, we obtain the following result of Liu and Owa [5]:

Corollary 4.2. *The function $f(z) \in \Sigma_p$ is in the class $\mathcal{T}_{n+p-1}(\alpha)$ if and only if*

$$f(z) * \frac{1 + [(n + p)(1 - \Omega) - 1]z}{z^p(1 - z)^{n+p+1}} \neq 0 \quad (z \in E^*; |x| = 1),$$

where Ω is given by

$$\Omega = \frac{1 + x + 2p(1 - \alpha)}{2p(1 - \alpha)} \quad (|x| = 1).$$



Convolution Conditions for Spirallikeness and Convex Spirallikeness of Certain Meromorphic p -valent Functions

V. Ravichandran, S. Sivaprasad Kumar
and K.G. Subramanian

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

Page 14 of 16

References

- [1] M.K. AOUF AND H.M. SRIVASTAVA, A new criterion for meromorphically p -valent convex functions of order α , *Math. Sci. Res. Hot. Line*, **1**(8) (1997), 7–12.
- [2] P.L. DUREN, Univalent Functions, In *Grundlehren der Mathematischen Wissenschaften, Bd., Volume 259*, Springer-Verlag, New York, (1983).
- [3] S.B. JOSHI AND H.M. SRIVASTAVA, A certain family of meromorphically multivalent functions, *Computers Math. Appl.* **38**(3/4) (1999), 201–211.
- [4] J.-L. LIU AND H.M. SRIVASTAVA, A linear operator and associated families of meromorphically multivalent functions, *J. Math. Anal. Appl.*, **259** (2001), 566–581.
- [5] J.-L. LIU AND S. OWA, On a class of meromorphic p -valent functions involving certain linear operators, *Internat. J. Math. Math. Sci.*, **32** (2002), 271–180.
- [6] J.-L. LIU AND H.M. SRIVASTAVA, Some convolution conditions for starlikeness and convexity of meromorphically multivalent functions, *Applied Math. Letters*, **16** (2003), 13–16.
- [7] S. OWA, H.E. DARWISH AND M.K. AOUF, Meromorphic multivalent functions with positive and fixed second coefficients, *Math. Japon.*, **46** (1997), 231–236.



Convolution Conditions for
Spirallikeness and Convex
Spirallikeness of Certain
Meromorphic p -valent
Functions

V. Ravichandran, S. Sivaprasad Kumar
and K.G. Subramanian

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

[Page 15 of 16](#)

- [8] H. SILVERMAN, E.M. SILVIA AND D. TELAGE, Convolution conditions for convexity, starlikeness and spiral-likeness, *Math. Zeitschr.*, **162** (1978), 125–130.
- [9] H.M. SRIVASTAVA, H.M. HOSSEN AND M.K. AOUF, A unified presentation of some classes of meromorphically multivalent functions, *Computers Math. Appl.* **38** (11/12) (1999), 63–70.



Convolution Conditions for
Spirallikeness and Convex
Spirallikeness of Certain
Meromorphic p -valent
Functions

V. Ravichandran, S. Sivaprasad Kumar
and K.G. Subramanian

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

Page 16 of 16