

Journal of Inequalities in Pure and Applied Mathematics

ON THE FEKETE-SZEGÖ PROBLEM FOR SOME SUBCLASSES OF ANALYTIC FUNCTIONS

T.N. SHANMUGAM AND S. SIVASUBRAMANIAN

Department of Mathematics
College of Engineering
Anna University, Chennai-600 025
Tamilnadu, India.
EMail: shan@annauniv.edu

Department of Mathematics
Easwari Engineering College
Ramapuram, Chennai-600 089
Tamilnadu, India.
EMail: sivasaisastha@rediffmail.com

©2000 Victoria University
ISSN (electronic): 1443-5756
165-05



volume 6, issue 3, article 71,
2005.

*Received 12 May, 2005;
accepted 24 May, 2005.*

Communicated by: S. Saitoh

Abstract

Contents




Home Page

Go Back

Close

Quit



Abstract

In this present investigation, the authors obtain Fekete-Szegő's inequality for certain normalized analytic functions $f(z)$ defined on the open unit disk for which $\frac{zf'(z)+\alpha z^2 f''(z)}{(1-\alpha)f(z)+\alpha z f'(z)}$ ($\alpha \geq 0$) lies in a region starlike with respect to 1 and is symmetric with respect to the real axis. Also certain applications of the main result for a class of functions defined by convolution are given. As a special case of this result, Fekete-Szegő's inequality for a class of functions defined through fractional derivatives is obtained. The Motivation of this paper is to give a generalization of the Fekete-Szegő inequalities obtained by Srivastava and Mishra .

2000 Mathematics Subject Classification: Primary 30C45

Key words: Analytic functions, Starlike functions, Subordination, Coefficient problem, Fekete-Szegő inequality.

Contents

1	Introduction	3
2	Fekete-Szegő Problem	6
3	Applications to Functions Defined by Fractional Derivatives	10
	References	

On the Fekete-Szegő Problem for Some Subclasses of Analytic Functions

T.N. Shanmugam and
S. Sivasubramanian

Title Page

Contents



Go Back

Close

Quit

Page 2 of 15

1. Introduction

Let \mathcal{A} denote the class of all *analytic* functions $f(z)$ of the form

$$(1.1) \quad f(z) = z + \sum_{k=2}^{\infty} a_k z^k \quad (z \in \Delta := \{z \in \mathbb{C} \mid |z| < 1\})$$

and \mathcal{S} be the subclass of \mathcal{A} consisting of univalent functions. Let $\phi(z)$ be an analytic function with positive real part on Δ with $\phi(0) = 1$, $\phi'(0) > 0$ which maps the unit disk Δ onto a region starlike with respect to 1 which is symmetric with respect to the real axis. Let $S^*(\phi)$ be the class of functions in $f \in \mathcal{S}$ for which


$$\frac{zf'(z)}{f(z)} \prec \phi(z), \quad (z \in \Delta)$$

and $C(\phi)$ be the class of functions in $f \in \mathcal{S}$ for which

$$1 + \frac{zf''(z)}{f'(z)} \prec \phi(z), \quad (z \in \Delta),$$

where \prec denotes the subordination between analytic functions. These classes were introduced and studied by Ma and Minda [10]. They have obtained the Fekete-Szegő inequality for the functions in the class $C(\phi)$. Since $f \in C(\phi)$ if and only if $zf'(z) \in S^*(\phi)$, we get the Fekete-Szegő inequality for functions in the class $S^*(\phi)$. For a brief history of the Fekete-Szegő problem for the class of starlike, convex and close-to-convex functions, see the recent paper by Srivastava *et al.* [7].

In the present paper, we obtain the Fekete-Szegő inequality for functions in a more general class $M_\alpha(\phi)$ of functions which we define below. Also we give



On the Fekete-Szegő Problem
for Some Subclasses of
Analytic Functions

T.N. Shanmugam and
S. Sivasubramanian

Title Page

Contents



Go Back

Close

Quit

Page 3 of 15

applications of our results to certain functions defined through convolution (or the Hadamard product) and in particular we consider a class $M_\alpha^\lambda(\phi)$ of functions defined by fractional derivatives. The motivation of this paper is to give a generalization of the Fekete-Szegő inequalities of Srivastava and Mishra [6].

Definition 1.1. Let $\phi(z)$ be a univalent starlike function with respect to 1 which maps the unit disk Δ onto a region in the right half plane which is symmetric with respect to the real axis, $\phi(0) = 1$ and $\phi'(0) > 0$. A function $f \in \mathcal{A}$ is in the class $M_\alpha(\phi)$ if

$$\frac{zf'(z) + \alpha z^2 f''(z)}{(1 - \alpha)f(z) + \alpha z f'(z)} \prec \phi(z) \quad (\alpha \geq 0).$$


For fixed $g \in \mathcal{A}$, we define the class $M_\alpha^g(\phi)$ to be the class of functions $f \in \mathcal{A}$ for which $(f * g) \in M_\alpha(\phi)$.

To prove our main result, we need the following:

Lemma 1.1. [10] If $p_1(z) = 1 + c_1z + c_2z^2 + \dots$ is an analytic function with positive real part in Δ , then

$$|c_2 - vc_1^2| \leq \begin{cases} -4v + 2 & \text{if } v \leq 0; \\ 2 & \text{if } 0 \leq v \leq 1; \\ 4v - 2 & \text{if } v \geq 1. \end{cases}$$

When $v < 0$ or $v > 1$, the equality holds if and only if $p_1(z)$ is $(1+z)/(1-z)$ or one of its rotations. If $0 < v < 1$, then the equality holds if and only if $p_1(z)$



On the Fekete-Szegő Problem
for Some Subclasses of
Analytic Functions

T.N. Shanmugam and
S. Sivasubramanian

Title Page

Contents



Go Back

Close

Quit

Page 4 of 15

is $(1 + z^2)/(1 - z^2)$ or one of its rotations. If $v = 0$, the equality holds if and only if

$$p_1(z) = \left(\frac{1}{2} + \frac{1}{2}\lambda\right) \frac{1+z}{1-z} + \left(\frac{1}{2} - \frac{1}{2}\lambda\right) \frac{1-z}{1+z} \quad (0 \leq \lambda \leq 1)$$


or one of its rotations. If $v = 1$, the equality holds if and only if p_1 is the reciprocal of one of the functions such that the equality holds in the case of $v = 0$.

Also the above upper bound is sharp, and it can be improved as follows when $0 < v < 1$:

$$|c_2 - vc_1^2| + v|c_1|^2 \leq 2 \quad \left(0 < v \leq \frac{1}{2}\right)$$

and

$$|c_2 - vc_1^2| + (1-v)|c_1|^2 \leq 2 \quad \left(\frac{1}{2} < v \leq 1\right).$$



**On the Fekete-Szegő Problem
for Some Subclasses of
Analytic Functions**

T.N. Shanmugam and
S. Sivasubramanian

Title Page

Contents



Go Back

Close

Quit

Page 5 of 15

2. Fekete-Szegő Problem

Our main result is the following:

Theorem 2.1. Let $\phi(z) = 1 + B_1z + B_2z^2 + B_3z^3 + \dots$. If $f(z)$ given by (1.1) belongs to $M_\alpha(\phi)$, then

$$|a_3 - \mu a_2^2| \leq \begin{cases} \frac{B_2}{2(1+2\alpha)} - \frac{\mu}{(1+\alpha)^2} B_1^2 + \frac{1}{2(1+2\alpha)(1+\alpha)^2} B_1^2 & \text{if } \mu \leq \sigma_1; \\ \frac{B_1}{2(1+2\alpha)} & \text{if } \sigma_1 \leq \mu \leq \sigma_2; \\ -\frac{B_2}{2(1+2\alpha)} + \frac{\mu}{(1+\alpha)^2} B_1^2 - \frac{1}{2(1+2\alpha)(1+\alpha)^2} B_1^2 & \text{if } \mu \geq \sigma_2, \end{cases}$$

where

$$\sigma_1 := \frac{(1+\alpha)^2(B_2 - B_1) + (1+\alpha^2)B_1^2}{2(1+2\alpha)B_1^2},$$

$$\sigma_2 := \frac{(1+\alpha)^2(B_2 + B_1) + (1+\alpha^2)B_1^2}{2(1+2\alpha)B_1^2}.$$


The result is sharp.

Proof. For $f(z) \in M_\alpha(\phi)$, let

$$(2.1) \quad p(z) := \frac{zf'(z) + \alpha z^2 f''(z)}{(1-\alpha)f(z) + \alpha z f'(z)} = 1 + b_1z + b_2z^2 + \dots$$

From (2.1), we obtain

$$(1+\alpha)a_2 = b_1 \quad \text{and} \quad (2+4\alpha)a_3 = b_2 + (1+\alpha^2)a_2^2.$$



On the Fekete-Szegő Problem
for Some Subclasses of
Analytic Functions

T.N. Shanmugam and
S. Sivasubramanian

Title Page

Contents



Go Back

Close

Quit

Page 6 of 15

Since $\phi(z)$ is univalent and $p \prec \phi$, the function

$$p_1(z) = \frac{1 + \phi^{-1}(p(z))}{1 - \phi^{-1}(p(z))} = 1 + c_1z + c_2z^2 + \dots$$

is analytic and has a positive real part in Δ . Also we have

$$(2.2) \quad p(z) = \phi \left(\frac{p_1(z) - 1}{p_1(z) + 1} \right)$$

and from this equation (2.2), we obtain

$$b_1 = \frac{1}{2}B_1c_1$$

and

$$b_2 = \frac{1}{2}B_1(c_2 - \frac{1}{2}c_1^2) + \frac{1}{4}B_2c_1^2.$$

Therefore we have


$$(2.3) \quad a_3 - \mu a_2^2 = \frac{B_1}{4(1 + 2\alpha)} \{c_2 - \nu c_1^2\},$$

where

$$\nu := \frac{1}{2} \left[1 - \frac{B_2}{B_1} + \frac{(2\mu - 1) + \alpha(4\mu - \alpha)}{(1 + \alpha)^2} B_1 \right].$$

Our result now follows by an application of Lemma 1.1. To show that the bounds are sharp, we define the functions $K_\alpha^{\phi_n}$ ($n = 2, 3, \dots$) by

$$\frac{z[K_\alpha^{\phi_n}]'(z) + \alpha z^2[K_\alpha^{\phi_n}]''(z)}{(1 - \alpha)[K_\alpha^{\phi_n}](z) + \alpha z[K_\alpha^{\phi_n}]'(z)} = \phi(z^{n-1}), \quad K_\alpha^{\phi_n}(0) = 0 = [K_\alpha^{\phi_n}]'(0) - 1$$



**On the Fekete-Szegő Problem
for Some Subclasses of
Analytic Functions**

T.N. Shanmugam and
S. Sivasubramanian

Title Page

Contents



Go Back

Close

Quit

Page 7 of 15

and the function F_α^λ and G_α^λ ($0 \leq \lambda \leq 1$) by

$$\frac{z[F_\alpha^\lambda]'(z) + \alpha z^2[F_\alpha^\lambda]''(z)}{(1-\alpha)[F_\alpha^\lambda](z) + \alpha z[F_\alpha^\lambda]'(z)} = \phi\left(\frac{z(z+\lambda)}{1+\lambda z}\right), \quad F^\lambda(0) = 0 = (F^\lambda)'(0) - 1$$

and

$$\frac{z[G_\alpha^\lambda]'(z) + \alpha z^2[G_\alpha^\lambda]''(z)}{(1-\alpha)[G_\alpha^\lambda](z) + \alpha z[G_\alpha^\lambda]'(z)} = \phi\left(-\frac{z(z+\lambda)}{1+\lambda z}\right), \quad G^\lambda(0) = 0 = (G^\lambda)'(0).$$

Clearly the functions $K_\alpha^{\phi n}, F_\alpha^\lambda, G_\alpha^\lambda \in M_\alpha(\phi)$. Also we write $K_\alpha^\phi := K_\alpha^{\phi 2}$.


If $\mu < \sigma_1$ or $\mu > \sigma_2$, then the equality holds if and only if f is K_α^ϕ or one of its rotations. When $\sigma_1 < \mu < \sigma_2$, the equality holds if and only if f is $K_\alpha^{\phi 3}$ or one of its rotations. If $\mu = \sigma_1$ then the equality holds if and only if f is F_α^λ or one of its rotations. If $\mu = \sigma_2$ then the equality holds if and only if f is G_α^λ or one of its rotations. \square

Remark 1. If $\sigma_1 \leq \mu \leq \sigma_2$, then, in view of Lemma 1.1, Theorem 2.1 can be improved. Let σ_3 be given by

$$\sigma_3 := \frac{(1+\alpha)^2 B_2 + (1+\alpha^2) B_1^2}{2(1+2\alpha) B_1^2}.$$

If $\sigma_1 \leq \mu \leq \sigma_3$, then

$$\begin{aligned} |a_3 - \mu a_2^2| + \frac{(1+\alpha)^2}{2(1+2\alpha) B_1^2} \left[B_1 - B_2 + \frac{(2\mu-1) + \alpha(4\mu-\alpha)}{(1+2\alpha)} B_1^2 \right] |a_2|^2 \\ \leq \frac{B_1}{2(1+2\alpha)}. \end{aligned}$$



On the Fekete-Szegő Problem
for Some Subclasses of
Analytic Functions

T.N. Shanmugam and
S. Sivasubramanian

Title Page

Contents



Go Back


Close

Quit

Page 8 of 15

If $\sigma_3 \leq \mu \leq \sigma_2$, then

$$|a_3 - \mu a_2^2| + \frac{(1 + \alpha)^2}{2(1 + 2\alpha)B_1^2} \left[B_1 + B_2 - \frac{(2\mu - 1) + \alpha(4\mu - \alpha)}{(1 + 2\alpha)^2} B_1^2 \right] |a_2|^2 \leq \frac{B_1}{2(1 + 2\alpha)}.$$



**On the Fekete-Szegő Problem
for Some Subclasses of
Analytic Functions**

T.N. Shanmugam and
S. Sivasubramanian

Title Page

Contents



Go Back

Close

Quit

Page 9 of 15

3. Applications to Functions Defined by Fractional Derivatives

In order to introduce the class $M_\alpha^\lambda(\phi)$, we need the following:

Definition 3.1 (see [3, 4]; see also [8, 9]). *Let $f(z)$ be analytic in a simply connected region of the z -plane containing the origin. The fractional derivative of f of order λ is defined by*

$$D_z^\lambda f(z) := \frac{1}{\Gamma(1-\lambda)} \frac{d}{dz} \int_0^z \frac{f(\zeta)}{(z-\zeta)^\lambda} d\zeta \quad (0 \leq \lambda < 1),$$


where the multiplicity of $(z-\zeta)^\lambda$ is removed by requiring that $\log(z-\zeta)$ is real for $z-\zeta > 0$.

Using the above Definition 3.1 and its known extensions involving fractional derivatives and fractional integrals, Owa and Srivastava [3] introduced the operator $\Omega^\lambda : \mathcal{A} \rightarrow \mathcal{A}$ defined by

$$(\Omega^\lambda f)(z) = \Gamma(2-\lambda) z^\lambda D_z^\lambda f(z), \quad (\lambda \neq 2, 3, 4, \dots).$$

The class $M_\alpha^\lambda(\phi)$ consists of functions $f \in \mathcal{A}$ for which $\Omega^\lambda f \in M_\alpha(\phi)$. Note that $M_0^0(\phi) \equiv S^*(\phi)$ and $M_\alpha^\lambda(\phi)$ is the special case of the class $M_\alpha^g(\phi)$ when

$$(3.1) \quad g(z) = z + \sum_{n=2}^{\infty} \frac{\Gamma(n+1)\Gamma(2-\lambda)}{\Gamma(n+1-\lambda)} z^n.$$



On the Fekete-Szegő Problem
for Some Subclasses of
Analytic Functions

T.N. Shanmugam and
S. Sivasubramanian

Title Page

Contents



Go Back

Close

Quit

Page 10 of 15

Let

$$g(z) = z + \sum_{n=2}^{\infty} g_n z^n \quad (g_n > 0).$$

Since

$$f(z) = z + \sum_{n=2}^{\infty} a_n z^n \in M_{\alpha}^g(\phi)$$


if and only if

$$(f * g) = z + \sum_{n=2}^{\infty} g_n a_n z^n \in M_{\alpha}(\phi),$$

we obtain the coefficient estimate for functions in the class $M_{\alpha}^g(\phi)$, from the corresponding estimate for functions in the class $M_{\alpha}(\phi)$. Applying Theorem 2.1 for the function $(f * g)(z) = z + g_2 a_2 z^2 + g_3 a_3 z^3 + \dots$, we get the following Theorem 3.1 after an obvious change of the parameter μ :

Theorem 3.1. *Let the function $\phi(z)$ be given by $\phi(z) = 1 + B_1 z + B_2 z^2 + B_3 z^3 + \dots$. If $f(z)$ given by (1.1) belongs to $M_{\alpha}^g(\phi)$, then*

$$|a_3 - \mu a_2^2| \leq \begin{cases} \frac{1}{g_3} \left[\frac{B_2}{2(1+2\alpha)} - \frac{\mu g_3}{(1+\alpha)^2 g_2^2} B_1^2 + \frac{1}{2(1+2\alpha)(1+\alpha)^2} B_1^2 \right] & \text{if } \mu \leq \sigma_1; \\ \frac{1}{g_3} \left[\frac{B_1}{2(1+2\alpha)} \right] & \text{if } \sigma_1 \leq \mu \leq \sigma_2; \\ \frac{1}{g_3} \left[-\frac{B_2}{2(1+2\alpha)} + \frac{\mu g_3}{(1+\alpha)^2 g_2^2} B_1^2 - \frac{1}{2(1+2\alpha)(1+\alpha)^2} B_1^2 \right] & \text{if } \mu \geq \sigma_2, \end{cases}$$



**On the Fekete-Szegő Problem
for Some Subclasses of
Analytic Functions**

T.N. Shanmugam and
S. Sivasubramanian

Title Page

Contents



Go Back

Close

Quit

Page 11 of 15

where

$$\sigma_1 := \frac{g_2^2 (1 + \alpha)^2 (B_2 - B_1) + (1 + \alpha^2) B_1^2}{g_3 \quad 2(1 + 2\alpha) B_1^2}$$

$$\sigma_2 := \frac{g_2^2 (1 + \alpha)^2 (B_2 + B_1) + (1 + \alpha^2) B_1^2}{g_3 \quad 2(1 + 2\alpha) B_1^2}.$$

The result is sharp.

Since

$$(\Omega^\lambda f)(z) = z + \sum_{n=2}^{\infty} \frac{\Gamma(n+1)\Gamma(2-\lambda)}{\Gamma(n+1-\lambda)} a_n z^n,$$

we have

$$(3.2) \quad g_2 := \frac{\Gamma(3)\Gamma(2-\lambda)}{\Gamma(3-\lambda)} = \frac{2}{2-\lambda}$$


and

$$(3.3) \quad g_3 := \frac{\Gamma(4)\Gamma(2-\lambda)}{\Gamma(4-\lambda)} = \frac{6}{(2-\lambda)(3-\lambda)}.$$

For g_2 and g_3 given by (3.2) and (3.3), Theorem 3.1 reduces to the following:

Theorem 3.2. Let the function $\phi(z)$ be given by $\phi(z) = 1 + B_1 z + B_2 z^2 + B_3 z^3 + \dots$. If $f(z)$ given by (1.1) belongs to $M_\alpha^\lambda(\phi)$, then

$$|a_3 - \mu a_2^2| \leq \begin{cases} \frac{(2-\lambda)(3-\lambda)}{6} \gamma & \text{if } \mu \leq \sigma_1; \\ \frac{(2-\lambda)(3-\lambda)}{6} \cdot \frac{B_1}{2(1+2\alpha)} & \text{if } \sigma_1 \leq \mu \leq \sigma_2; \\ \frac{(2-\lambda)(3-\lambda)}{6} \gamma & \text{if } \mu \geq \sigma_2, \end{cases}$$



On the Fekete-Szegő Problem
for Some Subclasses of
Analytic Functions

T.N. Shanmugam and
S. Sivasubramanian

Title Page

Contents



Go Back

Close

Quit

Page 12 of 15

where


$$\gamma := \frac{B_2}{2(1+2\alpha)} - \frac{3(2-\lambda)}{2(3-\lambda)} \frac{\mu}{(1+\alpha)^2} B_1^2 + \frac{1}{2(1+2\alpha)(1+\alpha)^2} B_1^2,$$

$$\sigma_1 := \frac{2(3-\lambda)}{3(2-\lambda)} \cdot \frac{(1+\alpha)^2(B_2 - B_1) + (1+\alpha^2)B_1^2}{2(1+2\alpha)B_1^2}$$

$$\sigma_2 := \frac{2(3-\lambda)}{3(2-\lambda)} \cdot \frac{(1+\alpha)^2(B_2 + B_1) + (1+\alpha^2)B_1^2}{2(1+2\alpha)B_1^2}.$$

The result is sharp.

Remark 2. When $\alpha = 0$, $B_1 = 8/\pi^2$ and $B_2 = 16/(3\pi^2)$, the above Theorem 3.1 reduces to a recent result of Srivastava and Mishra [6, Theorem 8, p. 64] for a class of functions for which $\Omega^\lambda f(z)$ is a parabolic starlike function [2, 5].



On the Fekete-Szegő Problem
for Some Subclasses of
Analytic Functions

T.N. Shanmugam and
S. Sivasubramanian

Title Page

Contents



Go Back


Close

Quit

Page 13 of 15

References

- [1] B.C. CARLSON AND D.B. SHAFFER, Starlike and prestarlike hypergeometric functions, *SIAM J. Math. Anal.*, **15** (1984), 737–745.
- [2] A.W. GOODMAN, Uniformly convex functions, *Ann. Polon. Math.*, **56** (1991), 87–92.
- [3] S. OWA AND H.M. SRIVASTAVA, Univalent and starlike generalized hypergeometric functions, *Canad. J. Math.*, **39** (1987), 1057–1077.
- [4] S. OWA, On the distortion theorems I, *Kyungpook Math. J.*, **18** (1978), 53–58.
- [5] F. RØNNING, Uniformly convex functions and a corresponding class of starlike functions, *Proc. Amer. Math. Soc.*, **118** (1993), 189–196.
- [6] H.M. SRIVASTAVA AND A.K. MISHRA, Applications of fractional calculus to parabolic starlike and uniformly convex functions, *Computer Math. Appl.*, **39** (2000), 57–69.
- [7] H.M. SRIVASTAVA, A.K. MISHRA AND M.K. DAS, The Fekete-Szegő problem for a subclass of close-to-convex functions, *Complex Variables, Theory Appl.*, **44** (2001), 145–163.
- [8] H.M. SRIVASTAVA AND S. OWA, An application of the fractional derivative, *Math. Japon.*, **29** (1984), 383–389.
- [9] H.M. SRIVASTAVA AND S. OWA, *Univalent functions, Fractional Calculus, and their Applications*, Halsted Press/John Wiley and Sons, Chichester/New York, (1989).



On the Fekete-Szegő Problem
for Some Subclasses of
Analytic Functions

T.N. Shanmugam and
S. Sivasubramanian

Title Page

Contents




Go Back

Close

Quit

Page 14 of 15

- [10] W. MA AND D. MINDA, A unified treatment of some special classes of univalent functions, in: *Proceedings of the Conference on Complex Analysis*, Z. Li, F. Ren, L. Yang, and S. Zhang(Eds.), Int. Press (1994), 157–169.



**On the Fekete-Szegő Problem
for Some Subclasses of
Analytic Functions**

T.N. Shanmugam and
S. Sivasubramanian

Title Page

Contents



Go Back

Close

Quit

Page 15 of 15