

CHAO-PING CHEN

College of Mathematics and Informatics  
Henan Polytechnic University  
Jiaozuo City, Henan 454010, China.

*E*Mail: [chenchaoping@hpu.edu.cn](mailto:chenchaoping@hpu.edu.cn)



---

volume 6, issue 4, article 100,  
2005.

*Received 02 July, 2005;*  
*accepted 08 September, 2005.*

*Communicated by: A. Laforgia*

---

Abstract

Contents




Home Page

Go Back

Close

Quit



## Abstract

Let  $a$  and  $b$  be given real numbers with  $0 \leq a < b < a + 1$ . Then the function  $\theta_{a,b}(x) = [\Gamma(x + b)/\Gamma(x + a)]^{1/(b-a)} - x$  is strictly convex and decreasing on  $(-a, \infty)$  with  $\theta_{a,b}(\infty) = \frac{a+b-1}{2}$  and  $\theta_{a,b}(-a) = a$ , where  $\Gamma$  denotes the Euler's gamma function.

*2000 Mathematics Subject Classification:* 33B15, 26A48.

*Key words:* Gamma function; monotonicity; convexity.

The author was supported in part by the Science Foundation of the Project for Fostering Innovation Talents at Universities of Henan Province, China

## Contents

1	Introduction .....	3
2	Proof of Theorem 1.1 .....	9
	References	

### Monotonicity and Convexity for the Gamma Function

Chao-Ping Chen

Title Page

Contents



Go Back

Close

Quit

Page 2 of 13

# 1. Introduction

Kazarinoff [10] proved that the function  $\theta(n)$ ,

$$\frac{1 \cdot 3 \cdot 5 \cdots (2n-1)}{2 \cdot 4 \cdot 6 \cdots (2n)} = \frac{1}{\sqrt{\pi(n + \theta(n))}},$$

satisfies

$$(1.1) \quad \frac{1}{4} < \theta(n) < \frac{1}{2}, \quad n \in \mathbb{N}.$$

More generally, set

$$\theta(x) = \left[ \frac{\Gamma(x+1)}{\Gamma(x+\frac{1}{2})} \right]^2 - x, \quad x > -\frac{1}{2},$$


where

$$\Gamma(x) = \int_0^\infty t^{x-1} e^{-t} dt \quad (x > 0)$$

is the Euler's gamma function. Watson [15] proved that the function  $\theta$  is strictly decreasing on  $(-1/2, \infty)$ . Applying this result, together with the observation that  $\theta(\infty) = 1/4$ ,  $\theta(-1/2) = 1/2$  and  $\theta(1) = 4\pi^{-1} - 1$ , we obviously imply sharper inequalities:

$$(1.2) \quad \frac{1}{4} < \theta(x) < \frac{1}{2} \quad \text{for } x > -\frac{1}{2},$$

$$(1.3) \quad \frac{1}{4} < \theta(x) \leq 4\pi^{-1} - 1 \quad \text{for } x \geq 1.$$



Monotonicity and Convexity for  
the Gamma Function

Chao-Ping Chen

Title Page

Contents



Go Back

Close

Quit

Page 3 of 13

In particular, take in (1.3)  $x = n$ , we get

$$(1.4) \quad \frac{1}{\sqrt{\pi(n + 4\pi^{-1} - 1)}} \leq \frac{1 \cdot 3 \cdot 5 \cdots (2n - 1)}{2 \cdot 4 \cdot 6 \cdots (2n)} < \frac{1}{\sqrt{\pi(n + 1/4)}},$$

and the constants  $4\pi^{-1} - 1$  and  $1/4$  are the best possible.

The inequality (1.4) is called Wallis' inequality. For more information on Wallis' inequality, please refer to the paper [6] and the references therein.

H. Alzer [2] proved that the function  $\theta$  is strictly decreasing on  $[0, \infty)$ . Applying this result, he showed that for all integers  $n \geq 1$ ,

$$(1.5) \quad \sqrt{\frac{n + A}{2\pi}} < \frac{\Omega_{n-1}}{\Omega_n} \leq \sqrt{\frac{n + B}{2\pi}}$$

with the best possible constants


$$A = \frac{1}{2} \quad \text{and} \quad B = \frac{\pi}{2} = 0.57079\dots,$$

where  $\Omega_n = \pi^{n/2}/\Gamma(1 + n/2)$  denotes the volume of the unit ball in  $\mathbf{R}^n$ . (1.5) is an improvement of the following result given by Borewardt [5, p. 253]

$$(1.6) \quad \sqrt{\frac{n}{2\pi}} \leq \frac{\Omega_{n-1}}{\Omega_n} \leq \sqrt{\frac{n + 1}{2\pi}}.$$

If we denote by

$$\theta_{a,b}(x) = \left[ \frac{\Gamma(x + b)}{\Gamma(x + a)} \right]^{\frac{1}{b-a}} - x,$$



## Monotonicity and Convexity for the Gamma Function

Chao-Ping Chen

Title Page

Contents



Go Back

Close

Quit

Page 4 of 13

then we conclude from the representations [1, p. 257]

$$(1.7) \quad x^{b-a} \frac{\Gamma(x+a)}{\Gamma(x+b)} = 1 + \frac{(a-b)(a+b-1)}{2x} + O(x^{-2}) \quad (x \rightarrow \infty),$$

that

$$(1.8) \quad \theta_{a,b}(x) = x \left\{ \frac{1}{x} \left[ \frac{\Gamma(x+b)}{\Gamma(x+a)} \right]^{1/(b-a)} - 1 \right\} \rightarrow \frac{a+b-1}{2} \quad \text{as } x \rightarrow \infty.$$

Hence it is of interest to investigate the possible monotonic character of the function  $x \mapsto \theta_{a,b}(x)$ .


**Theorem 1.1.** *Let  $a$  and  $b$  be given real numbers with  $0 \leq a < b < a + 1$ . Then the function  $\theta_{a,b}(x) = [\Gamma(x+b)/\Gamma(x+a)]^{1/(b-a)} - x$  is strictly convex and decreasing on  $(-a, \infty)$ .*

Since  $\theta_{a,b}(x) = \theta_{b,a}(x)$ , it is clear that  $x \mapsto \theta_{a,b}(x)$  is strictly convex and decreasing on  $(-b, \infty)$  for  $0 \leq b < a < b + 1$ .

From  $\theta_{a,b}(\infty) = \frac{a+b-1}{2}$ ,  $\theta_{a,b}(-a) = a$  and the monotonicity of  $x \mapsto \theta_{a,b}(x)$ , we obtain the following

**Corollary 1.2.** *Let  $a$  and  $b$  be given real numbers with  $0 \leq a < b < a + 1$ , then for  $x > -a$ ,*

$$(1.9) \quad \left( x + \frac{a+b-1}{2} \right)^{b-a} < \frac{\Gamma(x+b)}{\Gamma(x+a)} < (x+a)^{b-a}.$$



A proof of the theorem above has been shown in [8], here we provide another proof. The ratio of two gamma functions has been investigated intensively by many authors. For example, Gautschi [9] proved the following inequalities

$$(1.10) \quad x^{1-x} < \frac{\Gamma(x+1)}{\Gamma(x+s)} < (x+1)^{1-s}, \quad 0 < s < 1, x = 1, 2, \dots$$

Kershaw [11] has given some improvements of these inequalities such as

$$(1.11) \quad \left(x + \frac{s}{2}\right)^{1-s} < \frac{\Gamma(x+1)}{\Gamma(x+s)} < \left(x - \frac{1}{2} + \sqrt{x + \frac{1}{4}}\right)^{1-s}$$

for real  $x > 0$  and  $0 < s < 1$ .


Inequalities for the ratio  $\Gamma(x+1)/\Gamma(1+\lambda)$  ( $x > 0; \lambda \in (0, 1)$ ) have a remarkable application, they can be used to obtain estimates for ultraspherical polynomials. The ultraspherical polynomials are defined by

$$P_n^{(\lambda)}(x) = \sum_{k=0}^{[n/2]} (-1)^k \frac{\Gamma(n-k+\lambda)}{\Gamma(\lambda)\Gamma(k+1)\Gamma(n-2k+1)} (2x)^{n-2k},$$

where  $n \geq 0$  is an integer and  $\lambda > 0$  is a real number.

In 1931, S. Bernstein [4] proved the following inequality for ultraspherical polynomials: If  $0 < \lambda < 1$ ,  $n \geq 1$ , and  $0 < \theta < \pi$ , then

$$(1.12) \quad (\sin \theta)^\lambda |P_n^{(\lambda)}(\cos \theta)| < \frac{2^{1-\lambda}}{\Gamma(\lambda)} n^{\lambda-1},$$



Title Page

Contents



Go Back

Close

Quit

Page 6 of 13

where the constant  $2^{1-\lambda}/\Gamma(\lambda)$  cannot be replaced by a smaller one.

We note that inequality (1.12) with  $\lambda = 1/2$  leads to a well-known inequality for the Legendre polynomials  $P_n = P_n^{(1/2)}$ :

$$(1.13) \quad (\sin \theta)^{1/2} |P_n \cos \theta| < \left(\frac{2}{\pi}\right)^{1/2} n^{-1/2}.$$

Several authors presented remarkable refinements of (1.12). They proved that in (1.12) the factor  $n^{\lambda-1}$  can be replaced by smaller expressions. The left-hand inequality of (1.11) was also considered in 1984 by L. Lorch [13]. He obtained the following results for integer  $x > 0$ :


$$(1.14) \quad \frac{\Gamma(x+1)}{\Gamma(x+s)} > \left(x + \frac{s}{2}\right)^{1-s} \quad \text{for } 0 < s < 1 \quad \text{or } s > 2,$$

$$(1.15) \quad \frac{\Gamma(x+1)}{\Gamma(x+s)} > \left(x + \frac{s}{2}\right)^{1-s} \quad \text{for } 1 < s < 2.$$

Lorch [13] used (1.14) to prove a sharpened inequality for ultraspherical polynomials:

$$(1.16) \quad (\sin \theta)^\lambda |P_n^{(\lambda)} \cos \theta| < \frac{2^{1-\lambda}}{\Gamma(\lambda)} (n + \lambda)^{\lambda-1}.$$

In 1992, A. Laforgia [12] proved in (1.12) that the term  $n^{\lambda-1}$  can be replaced by  $\Gamma(n + \lambda)/\Gamma(n + 1)$ . Since  $\Gamma(n + \lambda)/\Gamma(n + 1) < n^{\lambda-1}$ , see [9], this provides another refinement of Bernstein's inequality. In 1994, Y. Chow et al. [7] showed




---

**Monotonicity and Convexity for  
the Gamma Function**

Chao-Ping Chen

---

Title Page

Contents



Go Back

Close

Quit

Page 7 of 13


that (1.12) holds with  $\Gamma(n + 2\lambda)(\Gamma(n + 1))^{-1}(n + \lambda)^{-\lambda}$  instead of  $n^{\lambda-1}$ . This sharpens Lorch's result for all  $\lambda \in (0, 1/2)$ , since the inequality

$$\frac{\Gamma(n + 2\lambda)}{\Gamma(n + 1)(n + \lambda)^\lambda} < (n + \lambda)^{\lambda-1}$$

is valid for all  $\lambda \in (0, 1/2)$ . In 1997, H. Alzer [3] showed the following inequality

$$(1.17) \quad (\sin \theta)^\lambda |P_n^{(\lambda)} \cos \theta| < \frac{2^{1-\lambda}}{\Gamma(\lambda)} \cdot \frac{\Gamma(n + \frac{3}{2}\lambda)}{\Gamma(n + 1 + \frac{1}{2}\lambda)}.$$

Inequality (1.17) refines the results given by S. Bernstein, L. Lorch and A. Laforgia.




---

**Monotonicity and Convexity for  
the Gamma Function**

Chao-Ping Chen

---

Title Page

Contents



Go Back

Close

Quit

Page 8 of 13



## 2. Proof of Theorem 1.1

Easy computation yields

$$\begin{aligned}\theta'_{a,b}(x) &= \frac{1}{b-a}[\psi(x+b) - \psi(x+a)](\theta_{a,b}(x) + x) - 1, \\ \frac{(b-a)\theta''_{a,b}(x)}{\theta_{a,b}(x) + x} &= \psi'(x+b) - \psi'(x+a) + \frac{1}{b-a}[\psi(x+b) - \psi(x+a)]^2.\end{aligned}$$

Using the representations [14, p. 16]


$$\begin{aligned}\psi(x) &= \frac{\Gamma'(x)}{\Gamma(x)} = -\gamma + \int_0^\infty \frac{e^{-t} - e^{-xt}}{1 - e^{-t}} dt, \\ \psi'(x) &= \int_0^\infty \frac{t}{1 - e^{-t}} e^{-xt} dt\end{aligned}$$

for  $x > 0$ ,  $\gamma = 0.57721\dots$  is the Euler-Mascheroni constant, it follows that

$$\frac{(b-a)\theta''_{a,b}(x)}{\theta_{a,b}(x) + x} = - \int_0^\infty t\delta(t)e^{-(x+a)t} dt + \frac{1}{b-a} \left( \int_0^\infty \delta(t)e^{-(x+a)t} dt \right)^2,$$

where

$$\delta(t) = \frac{1 - e^{-(b-a)t}}{1 - e^{-t}} \quad \text{and} \quad \delta(0) = b - a.$$



Monotonicity and Convexity for  
the Gamma Function

Chao-Ping Chen

Title Page

Contents



Go Back

Close

Quit

Page 9 of 13

By using the convolution theorem for Laplace transforms, we have

$$\begin{aligned}
 \frac{(b-a)\theta''_{a,b}(x)}{\theta_{a,b}(x)+x} &= -\int_0^\infty t\delta(t)e^{-(x+a)t} dt \\
 (2.1) \qquad \qquad \qquad &+ \frac{1}{b-a} \int_0^\infty \left[ \int_0^t \delta(s)\delta(t-s) ds \right] e^{-(x+a)t} dt \\
 &= \int_0^\infty e^{-(x+a)t} \omega(t) dt,
 \end{aligned}$$

where

$$(2.2) \qquad \qquad \omega(t) = \int_0^t \left[ \frac{1}{b-a} \delta(s)\delta(t-s) - \delta(t) \right] ds.$$

Now we are in a position to prove that


$$(2.3) \qquad \frac{1}{b-a} \delta(s)\delta(t-s) - \delta(t) > 0 \quad \text{for } t > s > 0.$$

Define for  $t > s > 0$ ,

$$\varphi(t) = \ln \frac{\delta(s)}{b-a} + \ln \delta(t-s) - \ln \delta(t).$$

Elementary calculations reveal that

$$\begin{aligned}
 \varphi'(t) &= \frac{\delta'(t-s)}{\delta(t-s)} - \frac{\delta'(t)}{\delta(t)}, \\
 \left( \frac{\delta'(t)}{\delta(t)} \right)' &= (\ln \delta(t))'' = \frac{e^t}{(e^t - 1)^2} - \frac{(b-a)^2 e^{(b-a)t}}{[e^{(b-a)t} - 1]^2}.
 \end{aligned}$$




---

**Monotonicity and Convexity for  
the Gamma Function**

Chao-Ping Chen

---

Title Page

Contents



Go Back

Close

Quit

Page 10 of 13

Defined for  $r \in (0, 1)$ ,

$$g(r) = \frac{r^2 e^{rt}}{(e^{rt} - 1)^2} = \frac{1}{t^2} \left( \frac{rt/2}{\sinh(rt/2)} \right)^2.$$

Since  $x \mapsto \frac{\sinh x}{x}$  is strictly increasing with  $x \in (0, \infty)$ , we get  $g$  is strictly decreasing with  $r \in (0, 1)$ . This implies that  $\left(\frac{\delta'(t)}{\delta(t)}\right)' < 0$  for  $t > 0$  and  $0 < b - a < 1$ , and then,  $\varphi'(t) > 0$  and  $\varphi(t) > \varphi(s) = 0$ . This means (2.3) holds, and thus,  $\theta''_{a,b}(x) > 0$  ( $x > -a$ ) follows from (2.1), (2.2) and (2.3).

From the representations (1.7) and

$$\psi(x) = \ln x - \frac{1}{2x} + O(x^{-2}) \quad (x \rightarrow \infty),$$

(see [1, p. 259]), we conclude that


$$(2.4) \quad \lim_{x \rightarrow \infty} x^{a-b} \frac{\Gamma(x+b)}{\Gamma(x+a)} = 1,$$

$$(2.5) \quad \lim_{x \rightarrow \infty} x \left[ \psi(x+b) - \psi(x+a) \right] = \frac{1}{b-a}.$$

From (2.4), (2.5) and the monotonicity of the function  $x \mapsto \theta'_{a,b}(x)$ , we imply

$$\begin{aligned} \theta'_{a,b}(x) &< \lim_{x \rightarrow \infty} \theta'_{a,b}(x) \\ &= \lim_{x \rightarrow \infty} \frac{1}{b-a} \left[ x^{a-b} \frac{\Gamma(x+b)}{\Gamma(x+a)} \right]^{\frac{1}{b-a}} x [\psi(x+b) - \psi(x+a)] - 1 \\ &= 0. \end{aligned}$$

The proof is complete.



## Monotonicity and Convexity for the Gamma Function

Chao-Ping Chen

Title Page

Contents



Go Back


Close

Quit

Page 11 of 13

## References

- [1] M. ABRAMOWITZ AND I.A. STEGUN (Eds), *Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables*, National Bureau of Standards, Applied Mathematics Series 55, 4th printing, with corrections, Washington, 1965.
- [2] H. ALZER, Inequalities for the volume of unit ball in  $\mathbb{R}^n$ , *J. Math. Anal. Appl.*, **252** (2000), 353–363.
- [3] H. ALZER, On Bernstein’s inequality for ultraspherical polynomials, *J. Math. Anal. Appl.*, **252** (2000), 353–363.
- [4] S. BERNSTEIN, Sur les polynômes orthogonaux relatifs à un segment fini, *J. Math.*, **10** (1931), 219–286.
- [5] K.H. BORGWARDT, *The Simplex Method*, Springer-Verlag, Berlin, 1987.
- [6] CHAO-PING CHEN AND FENG QI, The best bounds in Wallis’ inequality, *Proc. Amer. Math. Soc.*, **133**(2) (2005), 397–401.
- [7] Y. CHOW, L. GATTESCHI AND R. WONG, A Bernstein-type for the Jacobi polynomial, *Proc. Amer. Math. Soc.*, **121**(2) (1994), 703–709.
- [8] N. ELEZOVIĆ, C. GIORDANO AND J. PEČARIĆ, The best bounds in Gautschi’s inequality, *Math. Inequal. Appl.*, **3**(2) (2000), 239–252.
- [9] W. GAUTSCHI, Some elementary inequalities relating to the gamma and incomplete gamma function, *J. Math. Phys.*, **38** (1959), 77–81.



---

### Monotonicity and Convexity for the Gamma Function

Chao-Ping Chen

---

Title Page

Contents




Go Back

Close

Quit

Page 12 of 13

- [10] D.K. KAZARINOFF, On Wallis' formula, *Edinburgh. Math. Soc. Notes*, No. 40 (1956), 19–21.
- [11] D. KERSHAW, Some extensions of W. Gautschi's inequalities for gamma function, *Math. Comp.*, **41** (164) (1983), 607–611.
- [12] A. LAFORGIA, A simple proof of the Bernstein inequality for ultraspherical polynomials, *Boll. Un. Mat. Ital.*, **6A** (1992), 267–269.
- [13] L. LORCH, Inequalities for ultraspherical polynomials and the gamma function, *J. Approx. Theory*, **40** (1984), 115–120.
- [14] W. MAGNUS, F. OBERHETTINGER AND R.P. SONI, *Formulas and Theorems for the Special Functions of Mathematical Physics*, Springer, Berlin, 1966.
- [15] G.N. WATSON, A note on Gamma functions, *Proc. Edinburgh Math. Soc.*, (2) **11** 1958/1959 Edinburgh Math. Notes No. 42 (misprinted 41) (1959), 7–9.




---

**Monotonicity and Convexity for  
the Gamma Function**

Chao-Ping Chen

---

Title Page

Contents



Go Back

Close

Quit

Page 13 of 13