

SOME RESULTS ASSOCIATED WITH FRACTIONAL CALCULUS OPERATORS INVOLVING APPELL HYPERGEOMETRIC FUNCTION

R. K. RAINA

10/11 Ganpati Vihar, Opposite Sector 5
Udaipur 313002, Rajasthan, India
EMail: rkraina_7@hotmail.com

Received: 14 August, 2008

Accepted: 11 January, 2009

Communicated by: **H.M. Srivastava**

2000 AMS Sub. Class.: 26A33, 30C45.

Key words: Analytic functions, Hardy space, Fractional derivatives and fractional integrals, Appell hypergeometric function, Inclusion relation.

Abstract: A class of fractional derivative operators (with the Appell hypergeometric function in the kernel) is used here to define a new subclass of analytic functions and a coefficient bound inequality is established for this class of functions. Also, an inclusion theorem for a class of fractional integral operators involving the Hardy space of analytic functions is proved. The concluding remarks briefly mentions the relevances of the main results and possibilities of further work by using these new classes of fractional calculus operators.

Fractional Calculus Operators

R. K. Raina

vol. 10, iss. 1, art. 14, 2009

[Title Page](#)

[Contents](#)

[◀◀](#) [▶▶](#)

[◀](#) [▶](#)

Page 1 of 16

[Go Back](#)

[Full Screen](#)

[Close](#)

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756

Contents

1	Introduction, Definitions and Preliminaries	3
2	A Set of Coefficient Bounds	7
3	Inclusion Relations	9
4	Concluding Remarks	14



Fractional Calculus Operators

R. K. Raina

vol. 10, iss. 1, art. 14, 2009

[Title Page](#)

[Contents](#)



Page 2 of 16


[Go Back](#)

[Full Screen](#)

[Close](#)

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756

[Title Page](#)[Contents](#)

Page 3 of 16

[Go Back](#)[Full Screen](#)[Close](#)

1. Introduction, Definitions and Preliminaries

Let $\mathcal{A}(n)$ denote the class of functions $f(z)$ normalized by

$$(1.1) \quad f(z) = z + \sum_{k=n+1}^{\infty} a_k z^k \quad (n \in \mathbb{N}),$$

which are analytic in the open unit disk

$$\mathbb{U} = \{z : z \in \mathbb{C} \quad \text{and} \quad |z| < 1\}.$$

We denote by $\Delta_n^{(\alpha, \alpha', \beta, \beta', \gamma)}(\sigma)$ the subclass of functions in $\mathcal{A}(n)$ which also satisfy the inequality:


$$(1.2) \quad \operatorname{Re} \left\{ \chi_1(\alpha, \alpha', \beta, \beta', \gamma) z^{\alpha+\alpha'+\gamma-1} D_{0,z}^{(\alpha, \alpha', \beta, \beta', \gamma)} f(z) \right\} > \sigma \quad (z \in \mathbb{U}),$$

where $D_{0,z}^{(\alpha, \alpha', \beta, \beta', \gamma)}$ is the generalized fractional derivative operator (defined below), and (for convenience)

$$(1.3) \quad \chi_m(\alpha, \alpha', \beta, \beta', \gamma) = \frac{\Gamma(1+m+\beta')\Gamma(1+m-\alpha-\alpha'-\gamma)\Gamma(1+m-\alpha'-\beta-\gamma)}{\Gamma(1+m)\Gamma(1+m-\alpha'+\beta')\Gamma(1+m-\alpha-\alpha'-\beta-\gamma)} \quad (m \in \mathbb{N}),$$

provided that

$$(1.4) \quad \begin{aligned} 0 \leq \sigma < 1; 0 \leq \gamma < 1; \\ \gamma < \min(-\alpha - \alpha', -\alpha' - \beta, -\alpha - \alpha' - \beta) + m + 1; \\ \beta' > \max(0, \alpha') - m - 1. \end{aligned}$$

[Title Page](#)[Contents](#)

Page 4 of 16

[Go Back](#)[Full Screen](#)[Close](#)

Following [8], a function $f(z)$ is said to be in the class $\mathcal{V}_n(\theta_k)$ if $f(z) \in \mathcal{A}(n)$ satisfies the condition that

$$\arg(a_k) = \theta_k \quad (k \geq n + 1; n \in \mathbb{N})$$

and if there exists a real number ρ such that

$$(1.5) \quad \theta_k + (k - 1)\rho \equiv \pi \pmod{2\pi} \quad (k \geq n + 1; n \in \mathbb{N}),$$


then we say that $f(z)$ is in the class $\mathcal{V}_n(\theta_k; \rho)$. Suppose $\mathcal{V}_n = \cup \mathcal{V}_n(\theta_k; \rho)$ over all possible sequences θ_k with ρ satisfying (1.5), then we denote by $\nabla_n^{(\alpha, \alpha', \beta, \beta', \gamma)}(\sigma)$ the subclass of \mathcal{V}_n which consists of functions $f(z)$ belonging to the class $\Delta_n^{(\alpha, \alpha', \beta, \beta', \gamma)}(\sigma)$.

We present here the following family of fractional integral (and derivative) operators which involve the familiar Appell hypergeometric function F_3 (see also Kiryakova [4] and Saigo and Maeda [9]).

Definition 1.1. Let $\gamma > 0$ and $\alpha, \alpha', \beta, \beta' \in \mathbb{R}$. Then the fractional integral operator $I_{0,z}^{(\alpha, \alpha', \beta, \beta', \gamma)}$ of a function $f(z)$ is defined by

$$(1.6) \quad I_{0,z}^{(\alpha, \alpha', \beta, \beta', \gamma)} f(z) = \frac{z^{-\alpha}}{\Gamma(\gamma)} \int_0^z (z - \zeta)^{\gamma-1} \zeta^{-\alpha'} F_3 \left(\alpha, \alpha', \beta, \beta'; \gamma; 1 - \frac{\zeta}{z}, 1 - \frac{z}{\zeta} \right) f(\zeta) d\zeta \quad (\gamma > 0),$$

where the function $f(z)$ is analytic in a simply-connected region of the complex z -plane containing the origin, and it is understood that $(z - \zeta)^{\gamma-1}$ denotes the principal value for $0 \leq \arg(z - t) < 2\pi$. The function F_3 occurring in the kernel of (1.6) is the familiar Appell hypergeometric function of third type (also known as Horn's F_3



[Title Page](#)

[Contents](#)



Page 5 of 16

[Go Back](#)

[Full Screen](#)

[Close](#)

- function; see, for example, [10]) defined by

$$(1.7) \quad F_3(\alpha, \alpha', \beta, \beta'; \gamma; z, \xi) = \sum_{m=0}^{\infty} \sum_{n=0}^{\infty} \frac{(\alpha)_m (\alpha')_n (\beta)_m (\beta')_n}{(\gamma)_{m+n}} \frac{z^m \xi^n}{m! n!} \quad (|z| < 1, |\xi| < 1),$$

which is related to the Gaussian hypergeometric function ${}_2F_1(\alpha, \beta; \gamma; z)$ by the following relationship:

$$\begin{aligned} {}_2F_1(\alpha, \beta; \gamma; z) &= F_3(\alpha, \alpha', \beta, \beta'; \gamma; z, 0) \\ &= F_3(\alpha, 0, \beta, \beta'; \gamma; z, \xi) = F_3(\alpha, \alpha', \beta, 0; \gamma; z, \xi). \end{aligned}$$

Definition 1.2. The fractional derivative operator $D_{0,z}^{(\alpha, \alpha', \beta, \beta', \gamma)}$ of a function $f(z)$ is defined by

$$(1.8) \quad D_{0,z}^{(\alpha, \alpha', \beta, \beta', \gamma)} f(z) = \frac{d^n}{dz^n} I_{0,z}^{(\alpha, \alpha', \beta-n, \beta', n-\gamma)} f(z) \quad (n-1 \leq \gamma < n; n \in \mathbb{N}).$$

It may be observed that for


$$(1.9) \quad \alpha = \lambda + \mu, \quad \alpha' = \beta' = 0, \quad \beta = -\eta, \quad \gamma = \lambda,$$

we obtain the relationship

$$(1.10) \quad I_{0,z}^{(\lambda+\mu, 0, -\eta, 0, \lambda)} = I_{0,z}^{\lambda, \mu, \eta}$$

in terms of the Saigo type fractional integral operator $I_{0,z}^{\lambda, \mu, \eta}$ ([12]). On the other hand, if

$$(1.11) \quad \alpha = \mu - \lambda, \quad \alpha' = \beta' = 0, \quad \beta = -\eta, \quad \gamma = \lambda,$$



[Title Page](#)

[Contents](#)

[◀◀](#) [▶▶](#)

[◀](#) [▶](#)

Page 6 of 16

[Go Back](#)

[Full Screen](#)

[Close](#)

then we get

$$(1.12) \quad D_{0,z}^{(\mu-\lambda,0,-\eta,0,\lambda)} = J_{0,z}^{\lambda,\mu,\eta},$$

where $J_{0,z}^{(\lambda,\mu,\eta)}$ is the Saigo type fractional derivative operator ([6]; see also [7]). Further, when

$$(1.13) \quad \alpha = \beta' = 0, \quad \alpha' = 1 - \mu, \quad \gamma = \lambda \text{ (or } -\lambda),$$

then the operators $I_{0,z}^{(0,1-\mu,0,0,\lambda)}$ and $D_{0,z}^{(0,1-\mu,0,0,-\lambda)}$ correspond to the differential-integral operators Q_μ^λ due to Dziok [2].


Let \mathcal{H}^p ($0 \leq p < \infty$) be the class of analytic functions in \mathbb{U} such that

$$(1.14) \quad \|f\|_p = \lim_{r \rightarrow 1^-} \{M_p(r, f)\} < \infty,$$

where

$$(1.15) \quad \|f\|_p = \begin{cases} \left(\frac{1}{2\pi} \int_0^{2\pi} |f(re^{i\theta})|^p \right)^{\frac{1}{p}} & (0 < p < \infty), \\ \sup_{|z| \leq r} |f(z)|. \end{cases}$$

In this paper we first define a new function class in terms of the fractional derivative operators (with the Appell hypergeometric function in the kernel) and then establish a coefficient bound inequality for this function class. Also, we prove an inclusion theorem for a class of fractional integral operators involving the Hardy space of analytic functions. The relevance of the main results and possibilities of further work by using the new classes of fractional calculus operators are briefly pointed out in the concluding section of this paper.

[Title Page](#)[Contents](#)

Page 7 of 16

[Go Back](#)[Full Screen](#)[Close](#)

2. A Set of Coefficient Bounds

We begin by proving the following coefficient bounds inequality for a function $f(z)$ to be in the class $\Delta_n^{(\alpha, \alpha', \beta, \beta', \gamma)}(\sigma)$.

Theorem 2.1. Let $f(z)$ defined by (1.1) be in the class $\Delta_n^{(\alpha, \alpha', \beta, \beta', \gamma)}(\sigma)$, then

$$(2.1) \quad \sum_{k=n+1}^{\infty} \frac{|a_k|}{\chi_k(\alpha, \alpha', \beta, \beta', \gamma)} \leq \frac{1 - \sigma}{\chi_1(\alpha, \alpha', \beta, \beta', \gamma)},$$

where $\chi_m(\alpha, \alpha', \beta, \beta', \gamma)$ is defined by (1.3). The result is sharp.

Proof. Assume that

$$\operatorname{Re} \left\{ \chi_1(\alpha, \alpha', \beta, \beta', \gamma) z^{\alpha + \alpha' - \gamma - 1} D_{0,z}^{(\alpha, \alpha', \beta, \beta', \gamma)} f(z) \right\} > \sigma \quad (z \in \mathbb{U}).$$

Using (1.1) and the formula (see, e.g. [9, p. 394]):

$$(2.2) \quad \begin{aligned} & D_{0,z}^{(\alpha, \alpha', \beta, \beta', \gamma)} z^q \\ &= \frac{\Gamma(1+q)\Gamma(1+q-\alpha'+\beta')\Gamma(1+q-\alpha-\beta-\gamma)}{\Gamma(1+q+\beta')\Gamma(1+q-\alpha'-\beta-\gamma)\Gamma(1+q-\alpha-\alpha'-\gamma)} z^{q-\alpha-\alpha'-\gamma}, \\ & (0 \leq \gamma < 1; \alpha, \alpha', \beta, \beta' \in \mathbb{R}; q > \max(0, \alpha' - \beta', \alpha + \beta + \gamma) - 1) \end{aligned}$$

we obtain

$$(2.3) \quad \operatorname{Re} \left\{ 1 + \sum_{k=n+1}^{\infty} \frac{\chi_1(\alpha, \alpha', \beta, \beta', \gamma)}{\chi_k(\alpha, \alpha', \beta, \beta', \gamma)} a_k z^{k-1} \right\} > \sigma \quad (z \in \mathbb{U}),$$

and for $f(z) \in \mathcal{V}_n(\theta_k; \rho)$ ($z = re^{i\theta}$), the inequality thus obtainable from (2.3) on letting $r \rightarrow 1-$ therein, readily yields

$$(2.4) \quad \operatorname{Re} \left\{ 1 + \sum_{k=n+1}^{\infty} \frac{\chi_1(\alpha, \alpha', \beta, \beta', \gamma)}{\chi_k(\alpha, \alpha', \beta, \beta', \gamma)} |a_k| \exp(i(\theta_k + (k-1)\rho)) \right\} > \sigma.$$

If we apply (1.5), then (2.4) gives

$$(2.5) \quad 1 - \sum_{k=n+1}^{\infty} \frac{\chi_1(\alpha, \alpha', \beta, \beta', \gamma)}{\chi_k(\alpha, \alpha', \beta, \beta', \gamma)} |a_k| > \sigma,$$

which leads to the desired inequality (2.1). We also observe that the equality sign in (2.1) is attained for the function $f(z)$ defined by

$$(2.6) \quad f(z) = z + \frac{(1-\sigma)\chi_k(\alpha, \alpha', \beta, \beta', \gamma)}{\chi_1(\alpha, \alpha', \beta, \beta', \gamma)} z^k \exp(i\theta_k) \quad (k \geq n+1; n \in \mathbb{N}),$$

and this completes the proof of Theorem 2.1. □



[Title Page](#)

[Contents](#)




Page 8 of 16

[Go Back](#)

[Full Screen](#)

[Close](#)



[Title Page](#)

[Contents](#)



Page 9 of 16

[Go Back](#)

[Full Screen](#)

[Close](#)

3. Inclusion Relations

Under the hypotheses of Definition 1.1, let

$$(3.1) \quad \gamma > 0; \min(\gamma - \alpha - \alpha', \gamma - \alpha' - \beta, \beta', \gamma - \alpha - \alpha' - \beta, \beta' - \alpha') > -2; \\ \alpha, \alpha', \beta, \beta' \in \mathbb{R},$$

then the fractional integral operator

$$\Omega_z^{(\alpha, \alpha', \beta, \beta', \gamma)} : \mathcal{A} \rightarrow \mathcal{A} \quad (\mathcal{A}(1) = \mathcal{A})$$

is defined by

$$(3.2) \quad \Omega_z^{(\alpha, \alpha', \beta, \beta', \gamma)} f(z) = \chi_1(\alpha, \alpha', \beta, \beta', -\gamma) z^{\alpha + \alpha' + \gamma} I_{(0, z)}^{(\alpha, \alpha', \beta, \beta', \gamma)} f(z).$$

where $\chi_1(\alpha, \alpha', \beta, \beta', -\gamma)$ is given by (1.3).

By using the formula ([9, p. 394]; see also [4, p. 170, Lemma 9])


$$(3.3) \quad I_{0, z}^{(\alpha, \alpha', \beta, \beta', \gamma)} z^q \\ = \frac{\Gamma(1 + q)\Gamma(1 + q - \alpha' + \beta')\Gamma(1 + q - \alpha - \alpha' - \beta + \gamma)}{\Gamma(1 + q + \beta')\Gamma(1 + q - \alpha' - \beta + \gamma)\Gamma(1 + q - \alpha - \alpha' + \gamma)} z^{q - \alpha - \alpha' + \gamma}, \\ (\gamma > 0; \alpha, \alpha', \beta, \beta' \in \mathbb{R}; q > \max(0, \alpha' - \beta', \alpha + \beta - \gamma) - 1)$$

it follows from (1.1), (3.2) and (3.3) that

$$(3.4) \quad \Omega_z^{(\alpha, \alpha', \beta, \beta', \gamma)} f(z) = z + \chi_1(\alpha, \alpha', \beta, \beta', -\gamma) \sum_{k=2}^{\infty} \frac{a_k}{\chi_k(\alpha, \alpha', \beta, \beta', -\gamma)} z^k,$$

where (as before) $\chi_k(\alpha, \alpha', \beta, \beta', -\gamma)$ is given by (1.3).

Before stating and proving our main inclusion theorem, we recall here the following known results concerning the class $\mathcal{R}(\rho)$ in \mathcal{A} which satisfies the inequality that $\Re\{f'(z)\} > \rho$ ($0 \leq \rho < 1$), where $\mathcal{R}(1)$ is denoted by \mathcal{R} .



Title Page

Contents

◀▶

◀▶

Page 10 of 16

Go Back

Full Screen

Close

Lemma 3.1 ([3, p. 141]). Let $f(z) \in \mathcal{R}$, then

$$(3.5) \quad f(z) \in \mathcal{H}^p \quad : (0 < p < \infty).$$

Lemma 3.2 ([5, p. 533]). Let $f(z)$ defined by (1.1) be in the class $\mathcal{R}(\rho)$ ($0 \leq \rho < 1$), then

$$(3.6) \quad |a_k| \leq \frac{2}{k} \quad (k = 2, 3, 4, \dots).$$

Theorem 3.3. Let $f(z) \in \mathcal{R}$, then (under the constraints stated in (3.1))

$$(3.7) \quad \Omega_z^{(\alpha, \alpha', \beta, \beta', \gamma)} f(z) \in \mathcal{H}^p \quad (0 < p < \infty)$$

and


$$(3.8) \quad \Omega_z^{(\alpha, \alpha', \beta, \beta', \gamma)} f(z) \in \mathcal{H}^\infty \quad (\gamma > 1).$$

Proof. In view of (1.6) and (3.2), we obtain

$$(3.9) \quad \Omega_z^{(\alpha, \alpha', \beta, \beta', \gamma)} f(z) = \chi_1(\alpha, \alpha', \beta, \beta', -\gamma) \\ \times \int_0^1 (1-t)^{\gamma-1} t^{-\alpha'} F_3 \left(\alpha, \alpha', \beta, \beta'; \gamma; 1-t, 1-\frac{1}{t} \right) f(zt) dt.$$

This implies that

$$(3.10) \quad \operatorname{Re} \left\{ \frac{d}{dz} \Omega_z^{(\alpha, \alpha', \beta, \beta', \gamma)} f(z) \right\} = \chi_1(\alpha, \alpha', \beta, \beta', -\gamma) \\ \times \int_0^1 (1-t)^{\gamma-1} t^{1-\alpha'} F_3 \left(\alpha, \alpha', \beta, \beta'; \gamma; 1-t, 1-\frac{1}{t} \right) \Re \{ f'(zt) \} dt.$$



[Title Page](#)

[Contents](#)



Page 11 of 16

[Go Back](#)

[Full Screen](#)

[Close](#)

Since $f(z) \in \mathcal{R}$, therefore, we infer from (3.10) that

$$(3.11) \quad \Omega_z^{(\alpha, \alpha', \beta, \beta', \gamma)} f(z) \in \mathcal{R},$$

and applying Lemma 3.1, (3.11) gives the inclusion relation (3.7) under the conditions stated in (3.1).

To prove the result (3.8), we observe the following three-term recurrence relation:

$$(3.12) \quad \frac{d}{dz} \Omega_z^{(\alpha, \alpha', \beta, \beta', \gamma)} f(z) = z^{-1} \left\{ (\gamma - \alpha' - \beta + 1) \Omega_z^{(\alpha, \alpha', \beta, \beta', \gamma-1)} f(z) - (\gamma - \alpha' - \beta) \Omega_z^{(\alpha, \alpha', \beta, \beta', \gamma)} f(z) \right\},$$


which yields the inequality

$$(3.13) \quad \left| \frac{d}{dz} \Omega_z^{(\alpha, \alpha', \beta, \beta', \gamma)} f(z) \right|^p \leq r^{-p} \left\{ (\gamma - \alpha' - \beta + 1)^p \left| \Omega_z^{(\alpha, \alpha', \beta, \beta', \gamma-1)} f(z) \right|^p - (\gamma - \alpha' - \beta)^p \left| \Omega_z^{(\alpha, \alpha', \beta, \beta', \gamma)} f(z) \right|^p \right\} \quad (|z| = r),$$

provided that

$$(3.14) \quad \gamma > 1; \quad \min(1 + \gamma - \alpha - \alpha', 1 + \gamma - \alpha' - \beta, 1 + \beta', 1 + \gamma - \alpha - \alpha' - \beta, 1 + \beta' - \alpha') > -1; \quad \alpha, \alpha', \beta, \beta' \in \mathbb{R}$$

and $0 < p < \infty$.



[Title Page](#)

[Contents](#)



Page 12 of 16

[Go Back](#)

[Full Screen](#)

[Close](#)

Making use of (1.14) and (1.15), the above inequality (3.13) (with $p = 1$) yields

$$(3.15) \quad M_1 \left(r, \frac{d}{dz} \Omega_z^{(\alpha, \alpha', \beta, \beta', \gamma)} f(z) \right) \leq r^{-1} \left\{ (\gamma - \alpha' - \beta + 1) M_1 \left(r, \Omega_z^{(\alpha, \alpha', \beta, \beta', \gamma-1)} f(z) \right) - (\gamma - \alpha' - \beta) M_1 \left(r, \Omega_z^{(\alpha, \alpha', \beta, \beta', \gamma)} f(z) \right) \right\}$$

and

$$(3.16) \quad \left\| \frac{d}{dz} \Omega_z^{(\alpha, \alpha', \beta, \beta', \gamma)} f(z) \right\|_1 \leq (\gamma - \alpha' - \beta + 1) \left\| \Omega_z^{(\alpha, \alpha', \beta, \beta', \gamma-1)} f(z) \right\|_1 - (\gamma - \alpha' - \beta) \left\| \Omega_z^{(\alpha, \alpha', \beta, \beta', \gamma)} f(z) \right\|_1.$$


Applying (3.7), we infer (under the constraints stated in (3.14)) that

$$(3.17) \quad \Omega_z^{(\alpha, \alpha', \beta, \beta', \gamma-1)} f(z) \in \mathcal{H}^1 \quad \text{and} \quad \Omega_z^{(\alpha, \alpha', \beta, \beta', \gamma)} f(z) \in \mathcal{H}^1 \quad (\gamma > 1),$$

and consequently (3.16) implies that

$$\frac{d}{dz} \Omega_z^{(\alpha, \alpha', \beta, \beta', \gamma)} f(z) \in \mathcal{H}^1,$$

provided that the conditions stated in (3.14) are satisfied. By appealing to a known result [1, p. 42, Theorem 3.11], we infer from (3.17) that $\Omega_z^{(\alpha, \alpha', \beta, \beta', \gamma)} f(z)$ is continuous in $\mathbb{U}^* = \{z : z \in \mathbb{C} \text{ and } |z| \geq 1\}$. But \mathbb{U}^* being compact, we finally conclude that $\Omega_z^{(\alpha, \alpha', \beta, \beta', \gamma)} f(z)$ is a bounded analytic function in \mathbb{U} , and the proof of the second assertion (3.8) of Theorem 3.3 is complete.



Title Page

Contents



Page 13 of 16

Go Back

Full Screen

Close

The assertion (3.8) of Theorem 3.3 can also be proved by applying Lemma 3.2 (see also [3, p. 145]). Indeed, it follows from (3.4) and (3.6) that

$$\begin{aligned} \left| \Omega_z^{(\alpha, \alpha', \beta, \beta', \gamma)} f(z) \right| &\leq |z| + \chi_1(\alpha, \alpha', \beta, \beta', -\gamma) \sum_{k=2}^{\infty} \frac{|a_k|}{\chi_k(\alpha, \alpha', \beta, \beta', -\gamma)} |z^k| \\ &\leq 1 + 2 : \chi_1(\alpha, \alpha', \beta, \beta', -\gamma) \sum_{k=2}^{\infty} \frac{\Gamma(k)}{\Gamma(k+1)\chi_k(\alpha, \alpha', \beta, \beta', -\gamma)} \\ &= 1 + \frac{2(2 - \alpha' + \beta')(2 - \alpha - \alpha' - \beta + \gamma)}{(2 + \beta')(2 - \alpha - \alpha' + \gamma)(2 - \alpha' - \beta + \gamma)} \\ &\quad \times {}_4F_3 \left[\begin{matrix} 1, 2, 3 - \alpha' + \beta', 3 - \alpha - \alpha' - \beta + \gamma; \\ 3 + \beta', 3 - \alpha - \alpha' + \gamma, 3 - \alpha' - \beta + \gamma; \end{matrix} \middle| 1 \right] \end{aligned}$$

in terms of the generalized hypergeometric function.

Now, for fixed values of the parameters $\alpha, \alpha', \beta, \beta', \gamma$ satisfying the conditions stated in (3.1), we observe that by using the asymptotic formula [10, p. 109],

$$\frac{\Gamma(k)}{\Gamma(k+1)\chi_k(\alpha, \alpha', \beta, \beta', -\gamma)} = o(k^{-\gamma-1}) \quad (k \rightarrow \infty),$$

and since $\gamma > 1$, this proves our assertion (3.8). □

4. Concluding Remarks

In view of the relationships (1.10) and (1.12), the main results (Theorems 2.1 and 3.3) of this paper would correspond to the results due to Raina and Srivastava [8, p. 75, Theorem 1; p. 79, Theorem 7]. Furthermore, in view of the relationship (1.13), we can easily apply Theorems 2.1 and 3.3 to obtain the corresponding results associated with Dziok's differential-integral operators [2]. The family of fractional calculus operators (fractional integrals and fractional derivatives) defined by (1.6) and (1.8) can fruitfully be used in Geometric Function Theory. Several new analytic, multivalent (or meromorphic) function classes can be defined and the various properties of coefficient estimates, distortion bounds, radii of starlikeness, convexity and close to convexity for such contemplated classes investigated.



Fractional Calculus Operators

R. K. Raina

vol. 10, iss. 1, art. 14, 2009

Title Page

Contents



Page 14 of 16

Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756

References

- [1] P.L. DUREN, *Theory of H^p Spaces*, Vol. 38, A series of monographs and textbooks in pure and applied mathematics, Academic Press, New York, 1970.
- [2] J. DZIOK, Applications of the Jack lemma, *Acta Math. Hungar.*, **105** (2004), 93–102.
- [3] I. B. JUNG, Y. C. KIM AND H. M. SRIVASTAVA, The Hardy space of analytic functions associated with certain one-parameter families of integral operators, *J. Math. Anal. Appl.*, **176** (1993), 138–147.
- [4] V. KIRYAKOVA, On two Saigo's fractional integral operators in the class of univalent functions, *Frac. Cal. Appl. Math.*, **9** (2006), 159–176.
- [5] T.H. MACGREGOR, Functions whose derivative has a positive real part, *Trans. Amer. Math. Soc.*, **104** (1962), 532–537.
- [6] R.K. RAINA AND T.S. NAHAR, Characterization properties for starlikeness and convexity of some subclasses of analytic functions involving a class of fractional derivative operators, *Acta Math. Univ. Comenianae*, **69** (2000), 1–8.
- [7] R.K. RAINA AND H.M. SRIVASTAVA, A certain subclass of analytic functions associated with operators of fractional calculus, *Comput. Math. Appl.*, **32** (1996), 13–19.
- [8] R.K. RAINA AND H.M. SRIVASTAVA, Some subclasses of analytic functions associated with fractional calculus operators, *Comput. Math. Appl.*, **37** (1999), 73–84.
- [9] M. SAIGO AND N. MAEDA, More generalization of fractional calculus, in: *Transform Methods and Special Functions, Varna'96 (Proc. Second Internat.*



Fractional Calculus Operators

R. K. Raina

vol. 10, iss. 1, art. 14, 2009

Title Page

Contents



Page 15 of 16

Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756

Workshop), Science Culture Technology Publishing, Singapore (1998), 386-400.

- [10] H.M. SRIVASTAVA AND P.W. KARLSSON, *Multiple Gaussian Hypergeometric Series*, Halsted Press (Ellis Horwood Limited, Chichester), John Wiley and Sons, New York, 1985.
- [11] H.M. SRIVASTAVA AND S. OWA, *Current Topics in Analytic Function Theory*, World Scientific Publishing Company, Singapore, New Jersey, London and Hongkong, 1992.
- [12] H.M. SRIVASTAVA, M. SAIGO AND S. OWA, A class of distortion theorems involving certain operators of fractional calculus, *J. Math. Anal. Appl.*, **131** (1988), 412–420.



Fractional Calculus Operators

R. K. Raina

vol. 10, iss. 1, art. 14, 2009

Title Page

Contents



Page 16 of 16

Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756