

Journal of Inequalities in Pure and Applied Mathematics

A STOCHASTIC GRONWALL INEQUALITY AND ITS APPLICATIONS

KAZUO AMANO

Department of Mathematics
Faculty of Engineering
Gunma University
Kiryu, Tenjin 1-5-1
376-8515 Japan

*E*Mail: kamano@math.sci.gunma-u.ac.jp

©2000 Victoria University
ISSN (electronic): 1443-5756
242-04



volume 6, issue 1, article 17,
2005.

*Received 15 December, 2004;
accepted 1 February, 2005.*

Communicated by: S.S. Dragomir

Abstract

Contents



Home Page

Go Back

Close

Quit

Abstract


In this paper, we show a Gronwall type inequality for Itô integrals (Theorems 1.1 and 1.2) and give some applications. Our inequality gives a simple proof of the existence theorem for stochastic differential equation (Example 2.1) and also, the error estimate of Euler-Maruyama scheme follows immediately from our result (Example 2.2).

2000 Mathematics Subject Classification: 26D10, 26D20, 60H05, 60H35

Key words: Gronwall inequality, Itô integral

Contents

1	A Stochastic Gronwall type inequality	3
2	Applications	7
	References	



A Stochastic Gronwall Inequality and Its Applications

Kazuo Amano

Title Page

Contents



Go Back

Close

Quit

Page 2 of 11

1. A Stochastic Gronwall type inequality

Let $w(t), t \geq 0$ be a standard Brownian motion on a probability space (Ω, \mathcal{F}, P) and $\mathcal{F}_t, t \geq 0$ be the natural filtration of \mathcal{F} . For a positive number T , $M_w^2[0, T]$ denotes the set of all separable nonanticipative functions $f(t)$ with respect to \mathcal{F}_t defined on $[0, T]$ satisfying

$$E \left[\int_0^T f^2(t) dt \right] < \infty.$$

Theorem 1.1. Assume that $\xi(t)$ and $\eta(t)$ belong to $M_w^2[0, T]$. If there exist functions $a(t)$ and $b(t)$ belonging to $M_w^2[0, T]$ such that

$$(1.1) \quad |\xi(t)| \leq \left| \int_0^t a(s) ds + \int_0^t b(s) dw(s) \right|$$

and if there are nonnegative constants $\alpha_0, \alpha_1, \beta_0$ and β_1 such that

$$(1.2) \quad |a(t)| \leq \alpha_0|\eta(t)| + \alpha_1|\xi(t)|, \quad |b(t)| \leq \beta_0|\eta(t)| + \beta_1|\xi(t)|$$


for $0 \leq t \leq T$, then we have

$$(1.3) \quad E\xi^2(t) \leq 4(\alpha_0\sqrt{t} + \beta_0)^2 \exp\left(4t(\alpha_1\sqrt{t} + \beta_1)^2\right) \int_0^t E\eta^2(s) ds$$

for $0 \leq t \leq T$.

Proof. Since

$$E \left(\int_0^t b(s) dw(s) \right)^2 = E \int_0^t b^2(s) ds,$$



Title Page

Contents



Go Back

Close

Quit

Page 3 of 11

(1.1) implies, by Minkowsky and Schwarz inequalities,

$$(E\xi^2(t))^{\frac{1}{2}} \leq \left(t \int_0^t Ea^2(s) ds \right)^{\frac{1}{2}} + \left(\int_0^t Eb^2(s) ds \right)^{\frac{1}{2}}.$$

Direct computation gives, by (1.2),

$$\begin{aligned} \left(t \int_0^t Ea^2(s) ds \right)^{\frac{1}{2}} &\leq \sqrt{2t}\alpha_0 \left(\int_0^t E\eta^2(s) ds \right)^{\frac{1}{2}} + \sqrt{2t}\alpha_1 \left(\int_0^t E\xi^2(s) ds \right)^{\frac{1}{2}}, \\ \left(\int_0^t Eb^2(s) ds \right)^{\frac{1}{2}} &\leq \sqrt{2}\beta_0 \left(\int_0^t E\eta^2(s) ds \right)^{\frac{1}{2}} + \sqrt{2}\beta_1 \left(\int_0^t E\xi^2(s) ds \right)^{\frac{1}{2}}. \end{aligned}$$


Combining the above estimates, we obtain

$$(1.4) \quad E\xi^2(t) \leq 4(\alpha_0\sqrt{t} + \beta_0)^2 \int_0^t E\eta^2(s) ds + 4(\alpha_1\sqrt{t} + \beta_1)^2 \int_0^t E\xi^2(s) ds$$

for $0 \leq t \leq T$.

Let us fix a nonnegative number $t_0 \leq T$ arbitrarily. Then, for any $\delta > 0$, the last inequality (1.4) shows

$$\begin{aligned} \frac{d}{dt} \log \left(4(\alpha_0\sqrt{t_0} + \beta_0)^2 \int_0^{t_0} E\eta^2(s) ds + 4(\alpha_1\sqrt{t_0} + \beta_1)^2 \int_0^t E\xi^2(s) ds + \delta \right) \\ \leq 4(\alpha_1\sqrt{t_0} + \beta_1)^2 \end{aligned}$$



**A Stochastic Gronwall
Inequality and Its Applications**

Kazuo Amano

Title Page

Contents



Go Back

Close

Quit

Page 4 of 11

almost everywhere in $[0, t_0]$. Integrating this estimate from 0 to t_0 with respect to t , we get

$$\log \left(\frac{4(\alpha_0\sqrt{t_0} + \beta_0)^2 \int_0^{t_0} E\eta^2(s) ds + 4(\alpha_1\sqrt{t_0} + \beta_1)^2 \int_0^{t_0} E\xi^2(s) ds + \delta}{4(\alpha_0\sqrt{t_0} + \beta_0)^2 \int_0^{t_0} E\eta^2(s) ds + \delta} \right) \leq 4t_0(\alpha_1\sqrt{t_0} + \beta_1)^2.$$

Therefore, by (1.4), we have

$$E\xi^2(t_0) \leq \exp(4t_0(\alpha_1\sqrt{t_0} + \beta_1)^2) \left(4(\alpha_0\sqrt{t_0} + \beta_0)^2 \int_0^{t_0} E\eta^2(s) ds + \delta \right).$$

Now, letting $\delta \rightarrow 0$, we obtain (1.3). □


In case $\xi(t)$ is a step function, a weak assumption (1.6) will be enough to show the inequality (1.3), which would play an important role in the error analysis of the numerical solutions of stochastic differential equations.

Theorem 1.2. *Assume that $\xi(t)$ and $\eta(t)$ belong to $M_w^2[0, T]$ and $\xi(t)$ is a step function such that*

$$(1.5) \quad \xi(t) = \xi(t_n) \quad \text{when} \quad t_n \leq t < t_{n+1}$$

for $n = 0, 1, 2, \dots, N - 1$, where N is a positive integer and $\{t_n\}_{n=0}^N$ is a partition of the interval $[0, T]$ satisfying $0 = t_0 < t_1 < t_2 < \dots < t_{N-1} < t_N = T$. If there exist functions $a(t)$ and $b(t)$ belonging to $M_w^2[0, T]$ such that

$$(1.6) \quad |\xi(t_n)| \leq \left| \int_0^{t_n} a(s) ds + \int_0^{t_n} b(s) dw(s) \right|$$



**A Stochastic Gronwall
Inequality and Its Applications**

Kazuo Amano

Title Page

Contents



Go Back

Close

Quit

Page 5 of 11

is valid for each $n = 0, 1, 2, \dots, N$ and if there are nonnegative constants α_0 , α_1 , β_0 and β_1 satisfying (1.2) for $0 \leq t \leq T$, then we have (1.3) for $0 \leq t \leq T$.

Proof. As in the proof of Theorem 1.1, we have


$$E\xi^2(t_n) \leq 4(\alpha_0\sqrt{t_n} + \beta_0)^2 \int_0^{t_n} E\eta^2(s) ds + 4(\alpha_1\sqrt{t_n} + \beta_1)^2 \int_0^{t_n} E\xi^2(s) ds$$

for $n = 0, 1, 2, \dots, N$; this implies, by (1.5),

$$E\xi^2(t) \leq 4(\alpha_0\sqrt{t} + \beta_0)^2 \int_0^t E\eta^2(s) ds + 4(\alpha_1\sqrt{t} + \beta_1)^2 \int_0^t E\xi^2(s) ds$$

for $0 \leq t \leq T$.

The remaining part of the proof is exactly same as that of Theorem 1.1. \square



**A Stochastic Gronwall
Inequality and Its Applications**

Kazuo Amano

Title Page

Contents



Go Back

Close

Quit

Page 6 of 11

2. Applications

Throughout this section, we assume that $\xi(t) \in M_w^2[0, T]$ is a solution of the stochastic differential equation

$$d\xi(t) = a(t, \xi(t)) dt + b(t, \xi(t)) dw(t), \quad 0 \leq t \leq T$$

satisfying the initial condition $\xi(0) = \xi_0$, where $a(t, x)$ and $b(t, x)$ are real-valued functions defined in $[0, T]$ such that

$$\begin{aligned} |a(t, x)|, |b(t, x)| &\leq K(1 + |x|), \\ |a(t, x) - a(s, y)|, |b(t, x) - b(s, y)| &\leq L(|t - s| + |x - y|). \end{aligned}$$

Here K and L are nonnegative constants.


Example 2.1. *Theorem 1.1 gives a simple proof of the existence theorem for stochastic differential equations.*

We use Picard's method. Let us consider a sequence $\{\xi_n(t)\}$ defined by $\xi_0(t) = \xi_0$ and

$$\xi_{n+1}(t) = \xi_0 + \int_0^t a(s, \xi_n(s)) ds + \int_0^t b(s, \xi_n(s)) dw(s)$$

for $n = 0, 1, 2, \dots$. Then, we easily have

$$\begin{aligned} \xi_{n+1}(t) - \xi_n(t) &= \int_0^t (a(s, \xi_n(s)) - a(s, \xi_{n-1}(s))) ds \\ &\quad + \int_0^t (b(s, \xi_n(s)) - b(s, \xi_{n-1}(s))) dw(s) \end{aligned}$$



Title Page

Contents



Go Back

Close

Quit

Page 7 of 11

and the Lipschitz continuity of $a(t, x)$ and $b(t, x)$ implies

$$\begin{aligned} |a(s, \xi_n(s)) - a(s, \xi_{n-1}(s))| &\leq L|\xi_n(s) - \xi_{n-1}(s)|, \\ |b(s, \xi_n(s)) - b(s, \xi_{n-1}(s))| &\leq L|\xi_n(s) - \xi_{n-1}(s)|. \end{aligned}$$

Hence, Theorem 1.1 with $\alpha_0 = \beta_0 = L$ and $\alpha_1 = \beta_1 = 0$ shows

$$E|\xi_{n+1}(t) - \xi_n(t)|^2 \leq 4L^2(\sqrt{t} + 1)^2 \int_0^t E|\xi_n(s) - \xi_{n-1}(s)|^2 ds$$

for $n = 1, 2, 3, \dots$; the recursive use of this estimate gives

$$E|\xi_{n+1}(t) - \xi_n(t)|^2 \leq \frac{(4L^2(\sqrt{t} + 1)^2 t)^n}{n!} \sup_{0 \leq s \leq t} E|\xi_1(s) - \xi_0(s)|^2.$$


Consequently, as is well-known, the convergence of $\{\xi_n(t)\}$ follows.

By virtue of Theorem 1.1 with $\alpha_0 = \beta_0 = 0$ and $\alpha_1 = \beta_1 = L$, the uniqueness of the solution is clear.

Example 2.2. *The error estimate of the Euler-Maruyama scheme*

$$\xi_{n+1} = \xi_n + a(t_n, \xi_n)\Delta t + b(t_n, \xi_n)\Delta w_n, \quad n = 0, 1, 2, \dots, N - 1$$

follows immediately from Theorem 1.2, where N is a sufficiently large positive integer, $\Delta t = T/N$, $t_n = n\Delta t$ and $\Delta w_n = w(t_{n+1}) - w(t_n)$ for $n = 0, 1, 2, \dots, N - 1$.



A Stochastic Gronwall
Inequality and Its Applications

Kazuo Amano

Title Page

Contents



Go Back

Close

Quit

Page 8 of 11

Since

$$\xi_{n+1} = \xi_n + \int_{t_n}^{t_{n+1}} a(t_n, \xi_n) ds + \int_{t_n}^{t_{n+1}} b(t_n, \xi_n) dw(s),$$

$$\xi(t_{n+1}) = \xi(t_n) + \int_{t_n}^{t_{n+1}} a(s, \xi(s)) ds + \int_{t_n}^{t_{n+1}} b(s, \xi(s)) dw(s),$$

we have

$$\xi_{n+1} - \xi(t_{n+1}) = \xi_n - \xi(t_n) + \int_{t_n}^{t_{n+1}} (a(t_n, \xi_n) - a(s, \xi(s))) ds$$

$$+ \int_{t_n}^{t_{n+1}} (b(t_n, \xi_n) - b(s, \xi(s))) dw(s).$$

Now, for $n = 0, 1, 2, \dots, N - 1$, if we put

$$\varepsilon(s) = \xi_n - \xi(t_n),$$

$$f(s) = a(t_n, \xi_n) - a(s, \xi(s)),$$


$$g(s) = b(t_n, \xi_n) - b(s, \xi(s))$$

when $t_n \leq s < t_{n+1}$ and

$$\varepsilon(t_N) = \xi_N - \xi(t_N),$$

$$f(t_N) = a(t_N, \xi_N) - a(t_N, \xi(t_N)),$$

$$g(t_N) = b(t_N, \xi_N) - b(t_N, \xi(t_N)),$$



**A Stochastic Gronwall
Inequality and Its Applications**

Kazuo Amano

Title Page

Contents



Go Back

Close

Quit

Page 9 of 11

then we obtain

$$\varepsilon(t_n) = \int_0^{t_n} f(s) ds + \int_0^{t_n} g(s) dw(s)$$

for $n = 0, 1, 2, \dots, N$. The Lipschitz continuity of $a(t, x)$ and $b(t, x)$ shows

$$|f(s)| \leq L(\Delta t + |\xi(s) - \tilde{\xi}(s)| + |\varepsilon(s)|),$$

$$|g(s)| \leq L(\Delta t + |\xi(s) - \tilde{\xi}(s)| + |\varepsilon(s)|),$$

where $\tilde{\xi}(s) = \xi(t_n)$ when $t_n \leq s < t_{n+1}$ for $n = 0, 1, 2, \dots, N - 1$ and $\tilde{\xi}(t_N) = \xi(t_N)$. Hence, Theorem 1.2 with $\alpha_0 = \alpha_1 = \beta_0 = \beta_1 = L$ shows

$$E\varepsilon^2(t) \leq 4L^2(\sqrt{t} + 1)^2 \exp(4L^2(\sqrt{t} + 1)^2 t) \int_0^t E(\Delta t + |\xi(s) - \tilde{\xi}(s)|)^2 ds.$$

It follows from the fundamental property of Itô integrals that


$$E|\xi(s) - \tilde{\xi}(s)|^2 \leq C\Delta t,$$

where C is a nonnegative constant depending only on T , K and L . Combining the above estimates, we obtain

$$E\varepsilon^2(t) = O(\Delta t)$$

for any $0 \leq t \leq T$ when $\Delta t \rightarrow 0$; the error estimate of the Euler-Maruyama scheme is proved.

Our Gronwall type inequality works for other numerical solutions of stochastic differential equations.



A Stochastic Gronwall Inequality and Its Applications

Kazuo Amano

Title Page

Contents



Go Back


Close

Quit

Page 10 of 11

References

- [1] A. FRIEDMAN, *Stochastic Differential Equations and Applications, Volume I*, Academic Press, 1975.
- [2] K. ITÔ AND H. P. MCKEAN, *Diffusion Processes and Their Sample Paths*, Springer-Verlag, Berlin, 1965.
- [3] P.E. KLOEDEN AND E. PLATEN, *Numerical Solution of Stochastic Differential Equations*, Springer-Verlag, 1992.
- [4] D.W. STROOCK AND S.R.S. VARADHAN, *Multidimensional Diffusion Processes*, Springer, 1982.



A Stochastic Gronwall
Inequality and Its Applications

Kazuo Amano

Title Page

Contents



Go Back

Close

Quit

Page 11 of 11