

AN INEQUALITY ABOUT ${}_3\phi_2$ AND ITS APPLICATIONS

MINGJIN WANG AND HONGSHUN RUAN

Department of Information Science

Jiangsu Polytechnic University

Changzhou City 213164

Jiangsu Province, P.R. China.

E-Mail: wmj@jpu.edu.cn rhs@em.jpu.edu.cn

Received: 17 August, 2007

Accepted: 18 May, 2008


Communicated by: S.S. Dragomir

2000 AMS Sub. Class.: Primary 26D15; Secondary 33D15.

Key words: Basic hypergeometric function ${}_3\phi_2$; q -binomial theorem; q -Chu-Vandermonde formula; Grüss inequality.

Abstract: In this paper, we use the terminating case of the q -binomial formula, the q -Chu-Vandermonde formula and the Grüss inequality to drive an inequality about ${}_3\phi_2$. As applications of the inequality, we discuss the convergence of some q -series involving ${}_3\phi_2$.

Acknowledgements: The author would like to express deep appreciation to the referee for the helpful suggestions. In particular, the author thanks the referee for helping to improve the presentation of the paper.





Inequality About ${}_3\phi_2$
Mingjin Wang and
Hongshun Ruan

vol. 9, iss. 2, art. 48, 2008

Title Page

Contents



Page 1 of 14

Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756

Contents

1	Statement of Main Results	3
2	Notations and Known Results	4
3	Proof of the Theorem	6
4	Some Applications of the Inequality	10



Inequality About ${}_3\phi_2$
Mingjin Wang and
Hongshun Ruan

vol. 9, iss. 2, art. 48, 2008

Title Page

Contents



Page 2 of 14


Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756



Title Page

Contents



Page 3 of 14

Go Back

Full Screen

Close

1. Statement of Main Results

q -series, which are also called basic hypergeometric series, play a very important role in many fields, such as affine root systems, Lie algebras and groups, number theory, orthogonal polynomials and physics, etc. Inequalities techniques provide useful tools in the study of special functions (see [1, 6, 7, 8, 9, 10]). For example, Ito used inequalities techniques to give a sufficient condition for convergence of a special q -series called the Jackson integral [6]. In this paper, we derive the following new inequality about q -series involving ${}_3\phi_2$.


Theorem 1.1. *Let a_1, a_2, b_1, b_2 be some real numbers such that $b_i < 1$ for $i = 1, 2$. Then for all positive integers n , we have:*

$$(1.1) \quad \left| {}_3\phi_2 \left(\begin{matrix} b_1/a_1, b_2/a_2, q^{-n} \\ b_1, b_2 \end{matrix}; q, -a_1 a_2 q^n \right) - \frac{(a_1, a_2; q)_n}{(-1, b_1, b_2; q)_n} \right| \leq \lambda \mu (-1; q)_n,$$

where

$$\lambda = \max\{1, M^n\}, \quad M = \max \left\{ |a_1|, \frac{|a_1 - b_1|}{1 - b_1} \right\},$$
$$\mu = \max\{1, N^n\}, \quad N = \max \left\{ |a_2|, \frac{|a_2 - b_2|}{1 - b_2} \right\}.$$

Applications of this inequality are also given.



2. Notations and Known Results

We recall some definitions, notations and known results which will be used in the proof. Throughout this paper, it is supposed that $0 < q < 1$. The q -shifted factorials are defined as

$$(2.1) \quad (a; q)_0 = 1, \quad (a; q)_n = \prod_{k=0}^{n-1} (1 - aq^k), \quad (a; q)_\infty = \prod_{k=0}^{\infty} (1 - aq^k).$$

We also adopt the following compact notation for multiple q -shifted factorials:

$$(2.2) \quad (a_1, a_2, \dots, a_m; q)_n = (a_1; q)_n (a_2; q)_n \cdots (a_m; q)_n,$$

where n is an integer or ∞ .

The q -binomial theorem (see [2, 3, 4]) is given by

$$(2.3) \quad \sum_{k=0}^{\infty} \frac{(a; q)_k z^k}{(q; q)_k} = \frac{(az; q)_\infty}{(z; q)_\infty}, \quad |z| < 1.$$

When $a = q^{-n}$, where n denotes a nonnegative integer, we have

$$(2.4) \quad \sum_{k=0}^n \frac{(q^{-n}; q)_k z^k}{(q; q)_k} = (zq^{-n}; q)_n.$$

Heine introduced the ${}_{r+1}\phi_r$ basic hypergeometric series, which is defined by

$$(2.5) \quad {}_{r+1}\phi_r \left(\begin{matrix} a_1, a_2, \dots, a_{r+1} \\ b_1, b_2, \dots, b_r \end{matrix}; q, x \right) = \sum_{n=0}^{\infty} \frac{(a_1, a_2, \dots, a_{r+1}; q)_n x^n}{(q, b_1, b_2, \dots, b_r; q)_n}.$$

Title Page

Contents



Page 4 of 14


Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756



[Title Page](#)

[Contents](#)



Page 5 of 14

[Go Back](#)

[Full Screen](#)

[Close](#)

The q -Chu-Vandermonde sums (see [2, 3, 4]) are

$$(2.6) \quad {}_2\phi_1 \left(\begin{matrix} a, q^{-n} \\ c \end{matrix}; q, q \right) = \frac{a^n (c/a; q)_n}{(c; q)_n}$$

and, reversing the order of summation, we have

$$(2.7) \quad {}_2\phi_1 \left(\begin{matrix} a, q^{-n} \\ c \end{matrix}; q, cq^n/a \right) = \frac{(c/a; q)_n}{(c; q)_n}.$$

At the end of this section, we recall the Grüss inequality (see [5]):

$$(2.8) \quad \left| \frac{1}{b-a} \int_a^b f(x)g(x)dx - \left(\frac{1}{b-a} \int_a^b f(x)dx \right) \left(\frac{1}{b-a} \int_a^b g(x)dx \right) \right| \leq \frac{(M-m)(N-n)}{4},$$


provided that $f, g : [a, b] \rightarrow \mathbb{R}$ are integrable on $[a, b]$ and $m \leq f(x) \leq M$, $n \leq g(x) \leq N$ for all $x \in [a, b]$, where m, M, n, N are given constants.

By simple calculus, one can prove that the discrete version of the Grüss inequality can be stated as:

if $a \leq \lambda_i \leq A$ and $b \leq \mu_i \leq B$, $i = 1, 2, \dots, n$, then for all sequences $(p_i)_{0 \leq i \leq n}$ of nonnegative real numbers satisfying $\sum_{i=1}^n p_i = 1$, we have

$$(2.9) \quad \left| \sum_{i=1}^n \lambda_i \mu_i p_i - \left(\sum_{i=1}^n \lambda_i p_i \right) \cdot \left(\sum_{i=1}^n \mu_i p_i \right) \right| \leq \frac{(A-a)(B-b)}{4},$$

where a, A, b, B are some given real constants.



Title Page

Contents



Page 6 of 14

Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756

© 2007 Victoria University. All rights reserved.

3. Proof of the Theorem

In this section, we use the terminating case of the q -binomial formula, the q -Chu-Vandermonde formula and the Grüss inequality to prove (1.1). For this purpose, we need the following lemma.

Lemma 3.1. *Let a and b be two real numbers such that $b < 1$, and let $0 \leq t \leq 1$. Then,*

$$(3.1) \quad \left| \frac{a - bt}{1 - bt} \right| \leq \max \left\{ |a|, \frac{|a - b|}{1 - b} \right\}.$$

Proof. Let

$$f(t) = \frac{a - bt}{1 - bt}, \quad 0 \leq t \leq 1,$$

then

$$f'(t) = \frac{b(a - 1)}{(1 - bt)^2}, \quad 0 \leq t \leq 1.$$


So $f(t)$ is a monotonic function with respect to $0 \leq t \leq 1$. Since $f(0) = a$ and $f(1) = \frac{a-b}{1-b}$, (3.1) holds. \square

Now, we are in a position to prove the inequality (1.1).

Proof. Put

$$(3.2) \quad p_k = \frac{(q^{-n}; q)_k (-q^n)^k}{(q; q)_k (-1; q)_n}, \quad k = 0, 1, \dots, n.$$

It is obvious that $p_k \geq 0$.



[Title Page](#)

[Contents](#)

[◀◀](#) [▶▶](#)

[◀](#) [▶](#)

Page 7 of 14

[Go Back](#)

[Full Screen](#)

[Close](#)

On the other hand, using (2.4), we obtain

$$\sum_{k=0}^n p_k = \frac{1}{(-1; q)_n} \sum_{k=0}^n \frac{(q^{-n}; q)_k (-q^n)^k}{(q; q)_k} = 1.$$

Let

$$(3.3) \quad \lambda_k = \frac{(-a_1)^k (b_1/a_1; q)_k}{(b_1; q)_k},$$

and

$$(3.4) \quad \mu_k = \frac{(-a_2)^k (b_2/a_2; q)_k}{(b_2; q)_k},$$

where $k = 0, 1, \dots, n$.

According to the definitions of M , N , λ and μ , it is easy to see that


$$M^k \leq \lambda \quad \text{and} \quad N^k \leq \mu, \quad 0 \leq k \leq n.$$

Using the lemma, one can get for all $0 \leq k \leq n$,

$$(3.5) \quad |\lambda_k| = \left| \frac{a_1 - b_1}{1 - b_1} \cdot \frac{a_1 - b_1 q}{1 - b_1 q} \cdot \dots \cdot \frac{a_1 - b_1 q^{k-1}}{1 - b_1 q^{k-1}} \right| \leq M^k \leq \lambda$$

and

$$(3.6) \quad |\mu_k| = \left| \frac{a_2 - b_2}{1 - b_2} \cdot \frac{a_2 - b_2 q}{1 - b_2 q} \cdot \dots \cdot \frac{a_2 - b_2 q^{k-1}}{1 - b_2 q^{k-1}} \right| \leq N^k \leq \mu.$$



Substitution of (3.2), (3.3), (3.4), (3.5) and (3.6) into (2.9), gives

$$(3.7) \quad \left| \sum_{k=0}^n \frac{(q^{-n}; q)_k (-q^n)^k}{(q; q)_k (-1; q)_n} \cdot \frac{(-a_1)^k (b_1/a_1; q)_k}{(b_1; q)_k} \cdot \frac{(-a_2)^k (b_2/a_2; q)_k}{(b_2; q)_k} \right. \\
\left. - \sum_{k=0}^n \frac{(q^{-n}; q)_k (-q^n)^k}{(q; q)_k (-1; q)_n} \cdot \frac{(-a_1)^k (b_1/a_1; q)_k}{(b_1; q)_k} \right. \\
\left. \times \sum_{k=0}^n \frac{(q^{-n}; q)_k (-q^n)^k}{(q; q)_k (-1; q)_n} \cdot \frac{(-a_2)^k (b_2/a_2; q)_k}{(b_2; q)_k} \right| \leq \lambda \mu.$$

Using (2.5) and (2.7), we get

$$(3.8) \quad \sum_{k=0}^n \frac{(q^{-n}; q)_k (-q^n)^k}{(q; q)_k (-1; q)_n} \cdot \frac{(-a_1)^k (b_1/a_1; q)_k}{(b_1; q)_k} \cdot \frac{(-a_2)^k (b_2/a_2; q)_k}{(b_2; q)_k} \\
= \frac{1}{(-1; q)_n} \sum_{k=0}^n \frac{(q^{-n}, b_1/a_1, b_2/a_2; q)_k}{(q, b_1, b_2; q)_k} (-a_1 a_2 q^n)^k \\
= \frac{1}{(-1; q)_n} {}_3\phi_2 \left(\begin{matrix} b_1/a_1, b_2/a_2, q^{-n} \\ b_1, b_2 \end{matrix}; q, -a_1 a_2 q^n \right),$$

$$(3.9) \quad \sum_{k=0}^n \frac{(q^{-n}; q)_k (-q^n)^k}{(q; q)_k (-1; q)_n} \cdot \frac{(-a_1)^k (b_1/a_1; q)_k}{(b_1; q)_k} = \frac{(a_1; q)_n}{(-1, b_1; q)_n}$$

and

Title Page

Contents




Page 8 of 14

Go Back

Full Screen

Close



$$(3.10) \quad \sum_{k=0}^n \frac{(q^{-n}; q)_k (-q^n)^k}{(q; q)_k (-1; q)_n} \cdot \frac{(-a_2)^k (b_2/a_2; q)_k}{(b_2; q)_k} = \frac{(a_2; q)_n}{(-1, b_2; q)_n}.$$

Substituting (3.8), (3.9) and (3.10) into (3.7), we obtain (1.1). □

Taking $a_2 = 1$ in (1.1), we get the following corollary.

Corollary 3.2. *We have*

$$(3.11) \quad \left| {}_2\phi_1 \left(\begin{matrix} b_1/a_1, q^{-n} \\ b_1 \end{matrix}; q, -a_1 q^n \right) \right| \leq \lambda(-1; q)_n.$$

Title Page

Contents




Page 9 of 14

Go Back

Full Screen

Close



4. Some Applications of the Inequality

Convergence of q -series is an important problem which is at times very difficult. As applications of the inequality derived in this paper, we obtain some results about the convergence of the q -series involving ${}_3\phi_2$. In this section, we mainly discuss the convergence of the following q -series:

$$(4.1) \quad \sum_{n=0}^{\infty} c_n {}_3\phi_2 \left(\begin{matrix} b_1/a_1, b_2/a_2, q^{-n} \\ b_1, b_2 \end{matrix}; q, -a_1 a_2 q^n \right).$$

Theorem 4.1. Suppose $|a_i| \leq 1$ and $b_i < \frac{a_i+1}{2}$ for $i = 1, 2$. Let $\{c_n\}$ be a real sequence satisfying

$$\lim_{n \rightarrow \infty} \left| \frac{c_{n+1}}{c_n} \right| = p < 1.$$

Then the series (4.1) is absolutely convergent.

Proof. It is obvious that $b_i < 1$ for $i = 1, 2$. Combining the following inequality

$$(4.2) \quad \left| {}_3\phi_2 \left(\begin{matrix} b_1/a_1, b_2/a_2, q^{-n} \\ b_1, b_2 \end{matrix}; q, -a_1 a_2 q^n \right) \right| - \left| \frac{(a_1, a_2; q)_n}{(-1, b_1, b_2; q)_n} \right| \\ \leq \left| {}_3\phi_2 \left(\begin{matrix} b_1/a_1, b_2/a_2, q^{-n} \\ b_1, b_2 \end{matrix}; q, -a_1 a_2 q^n \right) - \frac{(a_1, a_2; q)_n}{(-1, b_1, b_2; q)_n} \right|$$

with (2.1), shows that

$$(4.3) \quad \left| {}_3\phi_2 \left(\begin{matrix} b_1/a_1, b_2/a_2, q^{-n} \\ b_1, b_2 \end{matrix}; q, -a_1 a_2 q^n \right) \right| \leq \left| \frac{(a_1, a_2; q)_n}{(-1, b_1, b_2; q)_n} \right| + \lambda \mu (-1; q)_n.$$

Since

$$|a_i| \leq 1, \quad b_i < \frac{a_i + 1}{2}, \quad i = 1, 2,$$

Title Page

Contents

◀◀ ▶▶

◀ ▶

Page 10 of 14


Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756



Title Page	
Contents	
◀◀	▶▶
◀	▶
Page 11 of 14	
Go Back	
Full Screen	
Close	

which is equivalent to

$$|a_i| \leq 1, \quad \frac{|a_i - b_i|}{1 - b_i} < 1, \quad i = 1, 2,$$

then

$$(4.4) \quad \lambda = \mu = 1.$$

Substituting (4.4) into (4.3), we obtain

$$(4.5) \quad \left| {}_3\phi_2 \left(\begin{matrix} b_1/a_1, b_2/a_2, q^{-n} \\ b_1, b_2 \end{matrix}; q, -a_1 a_2 q^n \right) \right| \leq \left| \frac{(a_1, a_2; q)_n}{(-1, b_1, b_2; q)_n} \right| + (-1; q)_n.$$

Multiplication of the two sides of (4.5) by $|c_n|$ gives

$$(4.6) \quad \left| c_n {}_3\phi_2 \left(\begin{matrix} b_1/a_1, b_2/a_2, q^{-n} \\ b_1, b_2 \end{matrix}; q, -a_1 a_2 q^n \right) \right| \leq \left| \frac{c_n (a_1, a_2; q)_n}{(-1, b_1, b_2; q)_n} \right| + |c_n (-1; q)_n|.$$

The ratio test shows that both


$$\sum_{n=0}^{\infty} \frac{c_n (a_1, a_2; q)_n}{(-1, b_1, b_2; q)_n} \quad \text{and} \quad \sum_{n=0}^{\infty} c_n (-1; q)_n$$

are absolutely convergent. From (4.6), we get that (4.1) is absolutely convergent. \square

Theorem 4.2. Suppose $|a_1| > 1$ or $a_1 < 2b_1 - 1$, $b_1 < 1$, $|a_2| \leq 1$ and $b_2 \leq \frac{a_2+1}{2}$. Let $\{c_n\}$ be a real sequence satisfying

$$\lim_{n \rightarrow \infty} \left| \frac{c_{n+1}}{c_n} \right| = p < \frac{1}{M},$$

where $M = \max \left\{ |a_1|, \frac{|a_1 - b_1|}{1 - b_1} \right\}$. Then the series (4.1) is absolutely convergent.



Title Page

Contents



Page 12 of 14

Go Back

Full Screen

Close

Proof. First we point out that $a_1 < 2b_1 - 1$ implies

$$\frac{|a_1 - b_1|}{1 - b_1} > 1.$$

So, under the conditions of the theorem, we know

$$\lambda = M^n \quad \text{and} \quad \mu = 1.$$

Multiplying both sides of (4.3) by $|c_n|$, one gets

$$(4.7) \quad \left| c_n {}_3\phi_2 \left(\begin{matrix} b_1/a_1, b_2/a_2, q^{-n} \\ b_1, b_2 \end{matrix}; q, -a_1 a_2 q^n \right) \right| \leq \left| \frac{c_n(a_1, a_2; q)_n}{(-1, b_1, b_2; q)_n} \right| + |c_n M^n (-1; q)_n|.$$

The ratio test shows that both

$$\sum_{n=0}^{\infty} \frac{c_n(a_1, a_2; q)_n}{(-1, b_1, b_2; q)_n} \quad \text{and} \quad \sum_{n=0}^{\infty} c_n M^n (-1; q)_n$$

are absolutely convergent. From (4.7), we get that (4.1) is absolutely convergent. \square

Similarly, we have

Theorem 4.3. Suppose $|a_i| > 1$ or $a_i < 2b_i - 1$, $b_i < 1$ with $i = 1, 2$. Let $\{c_n\}$ be a real sequence satisfying

$$\lim_{n \rightarrow \infty} \left| \frac{c_{n+1}}{c_n} \right| = p < \frac{1}{MN},$$

where $M = \max \left\{ |a_1|, \frac{|a_1 - b_1|}{1 - b_1} \right\}$ and $N = \max \left\{ |a_2|, \frac{|a_2 - b_2|}{1 - b_2} \right\}$. Then the series (4.1) is absolutely convergent.



References



- [1] G.D. ANDERSON, R.W. BARNARD, K.C. VAMANAMURTHY AND M. VUORINEN, Inequalities for zero-balanced hypergeometric functions, *Transactions of the American Mathematical Society*, **347**(5), 1995.
- [2] G.E. ANDREWS, *The Theory of Partitions*, Encyclopedia of Mathematics and Applications; Volume 2. Addison-Wesley Publishers, 1976.
- [3] W.N. BAILEY, Generalized hypergeometric series, *Cambridge Math. Tract* No.32, Cambridge University Press, London and New York.1960.
- [4] G. GASPER AND M. RAHMAN, *Basic Hypergeometric Series*, Cambridge Univ.Press, Cambridge, MA, 1990.
- [5] G. GRÜSS. Über das maximum des absoluten Betrages von $\frac{1}{b-a} \int_a^b f(x)g(x)dx - (\frac{1}{b-a} \int_a^b f(x)dx)(\frac{1}{b-a} \int_a^b g(x)dx)$, *Math.Z.*, **39** (1935), 215–226.
- [6] M. ITO, Convergence and asymptotic behavior of Jackson integrals associated with irreducible reduced root systems, *Journal of Approximation Theory*, **124** (2003) 154–180.
- [7] MINGJIN WANG, An inequality about q -series, *J. Ineq. Pure and Appl. Math.*, **7**(4) (2006), Art. 136. [ONLINE: <http://jipam.vu.edu.au/article.php?sid=756>].
- [8] MINGJIN WANG, An Inequality and its q -analogue, *J. Ineq. Pure and Appl. Math.*, **8**(2) (2007), Art. 50. [ONLINE: <http://jipam.vu.edu.au/article.php?sid=853>].

Inequality About ${}_3\phi_2$
Mingjin Wang and
Hongshun Ruan

vol. 9, iss. 2, art. 48, 2008

Title Page

Contents



Page 13 of 14


Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756



- [9] MINGJIN WANG, An inequality for ${}_{r+1}\phi_r$ and its applications, *Journal of Mathematical Inequalities*, **1**(2007) 339–345.
- [10] MINGJIN WANG, Two inequalities for ${}_r\phi_r$ and applications, *Journal of Inequalities and Applications*, 2008, Article ID 471527.

Inequality About ${}_3\phi_2$
Mingjin Wang and
Hongshun Ruan

vol. 9, iss. 2, art. 48, 2008

Title Page

Contents



Page 14 of 14

Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756