



On the Binomial Identities of Frisch and Klamkin

H. W. Gould

Department of Mathematics
West Virginia University
Morgantown, WV 26506
USA

henrygou@gmail.com

Jocelyn Quaintance

Department of Computer Science
University of Pennsylvania
Philadelphia, PA 19104
USA

jocelynq@seas.upenn.edu

Abstract

In this paper we investigate two somewhat similar identities for sums of ratios of binomial coefficients. We give several proofs, and note that the identities all follow from a hypergeometric identity of Gauss. Inverse identities are also given.

1 Introduction

Klamkin [3] stated the following identity in a letter to Gould in 1966:

$$\sum_{k=0}^n \frac{\binom{n}{k}}{\binom{n+a}{k+b}} = \frac{a+1+n}{(a+1)\binom{a}{b}} \quad (1)$$

Identity (4.6) in Gould's book [2] is a special case of this, corresponding to $a = 2x$ and $b = x$.

Frisch [1], in his dissertation in 1926, gave the curious formula

$$\sum_{k=0}^n (-1)^k \binom{n}{k} \frac{1}{\binom{b+k}{c}} = \frac{c}{n+c} \frac{1}{\binom{n+b}{b-c}}, \quad b \geq c > 0. \quad (2)$$

This was cited and proved by Netto [4, pp. 337–338] and is tabulated as Formula (4.2) in Gould’s book [2].

(It is interesting to note that Frisch’s research laid the foundations for modern econometrics theory and micro- and macro-economics, work for which he later received the Nobel Prize.)

Klamkin’s identity is actually a special case of the identity

$$\sum_{k=0}^n \frac{\binom{n}{k}}{\binom{x}{k+b}} = \frac{x+1}{(x-n+1)\binom{x-n}{b}}. \quad (3)$$

Here is how we obtain Eq. (3). Let $t_k = \frac{\binom{n}{k}}{\binom{x}{k+b}}$. Then

$$\frac{t_{k+1}}{t_k} = \frac{\binom{n}{k+1}\binom{x}{k+b}}{\binom{x}{k+1+b}\binom{n}{k}} = \frac{(n-k)(k+1+b)}{(k+1)(x-k-b)}.$$

Therefore,

$$\sum_{k=0}^n \frac{\binom{n}{k}}{\binom{x}{k+b}} = \frac{1}{\binom{x}{b}} {}_2F_1 \left[\begin{matrix} 1+b, & -n \\ -x+b & |1 \end{matrix} \right]. \quad (4)$$

Applying Gauss’s ${}_2F_1$ formula, namely,

$${}_2F_1 \left[\begin{matrix} a, & b \\ c & |1 \end{matrix} \right] = \frac{\Gamma(c-a-b)\Gamma(c)}{\Gamma(c-a)\Gamma(c-b)},$$

we have

$$\begin{aligned} \sum_{k=0}^n \frac{\binom{n}{k}}{\binom{x}{k+b}} &= \frac{1}{\binom{x}{b}} \cdot \frac{\Gamma(-x+n-1)\Gamma(-x+b)}{\Gamma(-x-1)\Gamma(-x+b+n)} \\ &= \frac{b!}{x(x-1)\dots(x-b+1)} \cdot \frac{(x+1)x(x-1)\dots(x-n+3)(x-n+2)}{(x-b)(x-b-1)\dots(x-n-b+2)(x-n-b+1)} \\ &= \frac{b!(x+1)\prod_{j=0}^{n-2}(x-j)}{\prod_{j=0}^{n+b-1}(x-j)} \\ &= \frac{b!(x+1)}{\prod_{j=n-1}^{n+b-1}(x-j)} \\ &= \frac{x+1}{x-n+1} \cdot \frac{b!}{(x-n)(x-n-1)\dots(x-n-b+1)} \\ &= \frac{x+1}{x-n+1} \cdot \frac{1}{\binom{x-n}{b}}. \end{aligned}$$

In this proof we assumed that $n \geq 2$. If $n = 0$, Eq. (1) is obviously true, while if $n = 1$, we obtain easily verified identity

$$\frac{1}{\binom{x}{b}} + \frac{1}{\binom{x}{b+1}} = \frac{x+1}{x\binom{x-1}{b}}.$$

2 Proof of Klamkin's and Frisch's formulas

We shall now prove Klamkin's formula directly from Formula (7.1) in Gould [2], namely

$$\sum_{k=0}^n (-1)^k \binom{n}{k} \frac{\binom{z}{k}}{\binom{y}{k}} = \frac{\binom{y-z}{n}}{\binom{y}{n}}, \quad (5)$$

and the easy binomial identity

$$\binom{x}{k+b} \binom{k+b}{k} = \binom{x}{b} \binom{x-b}{k} \quad (6)$$

Indeed, using these we have

$$\begin{aligned} \sum_{k=0}^n \frac{\binom{n}{k}}{\binom{x}{k+b}} &= \frac{1}{\binom{x}{b}} \sum_{k=0}^n \binom{n}{k} \frac{\binom{k+b}{k}}{\binom{x-b}{k}} \\ &= \frac{1}{\binom{x}{b}} \sum_{k=0}^n (-1)^k \binom{n}{k} \frac{\binom{-1-b}{k}}{\binom{x-b}{k}} \\ &= \frac{1}{\binom{x}{b}} \frac{\binom{x+1}{n}}{\binom{x-b}{n}} = \frac{x+1}{(x+1-n)\binom{x-n}{b}} \end{aligned}$$

by Formula (7.1), and our proof is complete. We remark that Formula (7.1) is equivalent to using the Gauss ${}_2F_1$ formula.

Simple binomial inversion yields the inverse Klamkin identity

$$\sum_{k=0}^n (-1)^{n-k} \binom{n}{k} \frac{x+1}{(x+1-k)\binom{x-k}{b}} = \frac{1}{\binom{x}{n+b}}. \quad (7)$$

We now proceed in a similar way to prove Frisch's identity. It is easy to verify the binomial identity

$$\frac{1}{\binom{k+b}{c}} = \frac{1}{\binom{b}{c}} \frac{\binom{c-b-1}{k}}{\binom{-b-1}{k}},$$

so that

$$\sum_{k=0}^n (-1)^k \binom{n}{k} \frac{1}{\binom{k+b}{c}} = \frac{1}{\binom{b}{c}} \sum_{k=0}^n (-1)^k \binom{n}{k} \frac{\binom{c-b-1}{k}}{\binom{-b-1}{k}}.$$

Upon applying Eq. (5), we find the sum equals

$$\frac{1}{\binom{b}{c}} \frac{\binom{-c}{n}}{\binom{-b-1}{n}} = \frac{c}{n+c} \frac{1}{\binom{n+b}{b-c}},$$

as desired to show.

This new proof of Frisch's identity should be compared to Gould's original proof [5, Section 7.2], a two-page calculation involving an application of Melzak's formula.

An inverse Frisch identity then follows by simple binomial inversion, and we have

$$\sum_{k=0}^n (-1)^k \binom{n}{k} \frac{c}{k+c} \frac{1}{\binom{k+b}{b-c}} = \frac{1}{\binom{n+b}{c}}. \quad (8)$$

References

- [1] Ragnar Frisch, Sur les semi-invariants et moments employés dans l'étude des distributions statistiques, *Skrifter utgitt av Det Norske Videnskaps-Akademi i Oslo, II, Historisk-Filosofisk Klasse* **3** (1926), 1–87. Quoted by Th. Skolem, p. 337, in Netto's *Lehrbuch*.
- [2] H. W. Gould, *Combinatorial Identities*, published by the author, Morgantown, WV, 1972.
- [3] Murray S. Klamkin, Letter to H. W. Gould, May 16, 1966.
- [4] Eugen Netto, *Lehrbuch der Kombinatorik*, 2nd edition, Chelsea Publications, 1927.
- [5] Jocelyn Quaintance and H. W. Gould, *Combinatorial Identities for Stirling Numbers*, World Scientific Press, 2016.

2010 *Mathematics Subject Classification*: Primary 05A19; Secondary 11B65, 33C10.

Keywords: binomial identity, hypergeometric series, Gauss' hypergeometric summation formula, binomial inversion.

Received July 4 2016; revised versions received August 31 2016; September 1 2016; September 7 2016. Published in *Journal of Integer Sequences*, September 7 2016.

Return to [Journal of Integer Sequences home page](#).