

DOUBLE n -ARY RELATIONAL STRUCTURES

JIŘÍ KARÁSEK, Brno

(Received January 22, 1996)

Abstract. In [7], V. Novák and M. Novotný studied ternary relational structures by means of pairs of binary structures; they obtained the so-called double binary structures. In this paper, the idea is generalized to relational structures of any finite arity.

Keywords: n -ary relation, n -ary structure, binding relation, double n -ary structure

MSC 1991: 04A05, 08A02

Let G be a set, let $n \geq 2$ be an integer. As usual, an n -ary relation on G is defined as a set $R \subseteq G^n$. The pair $\mathbf{G} = (G, R)$ is then called an n -ary relational structure (or briefly an n -ary structure). An n -ary structure $\mathbf{G} = (G, R)$ (and the relation R on G as well) is called

symmetric if $(x_1, x_2, \dots, x_n) \in R$ implies $(x_n, x_{n-1}, \dots, x_1) \in R$ for any $x_1, x_2, \dots, x_{n-1}, x_n \in G$;

asymmetric if $(x_1, x_2, \dots, x_n) \in R$ implies $(x_n, x_{n-1}, \dots, x_1) \notin R$ for any $x_1, x_2, \dots, x_{n-1}, x_n \in G$;

cyclic if $(x_1, x_2, \dots, x_n) \in R$ implies $(x_2, x_3, \dots, x_n, x_1) \in R$ for any $x_1, x_2, x_3, \dots, x_n \in G$;

transitive if $(x_1, x_2, \dots, x_n) \in R$, $(x_n, x_{n-1}, \dots, x_2, x_{n+1}) \in R$ imply $(x_1, x_2, \dots, x_{n-1}, x_{n+1}) \in R$ for any $x_1, x_2, \dots, x_{n-1}, x_n, x_{n+1} \in G$;

weakly transitive if $(x, y, y, \dots, y) \in R$, $(y, y, \dots, y, z) \in R$ imply $(x, y, y, \dots, y, z) \in R$ for any $x, y, z \in G$.

For any $\alpha = (x_1, x_2, \dots, x_n) \in G^n$, put $\alpha^{-1} = (x_n, x_{n-1}, \dots, x_1)$, $\alpha' = (x_{n-1}, x_{n-2}, \dots, x_1, x_n)$.

Let ϱ be an n -ary relation on G , let r be a binary relation on ϱ with the property: If $\alpha = (x_1, x_2, \dots, x_n) \in \varrho$, $\beta = (y_1, y_2, \dots, y_n) \in \varrho$, $(\alpha, \beta) \in r$, then $x_{j+1} = y_j$ for $j = 1, 2, \dots, n-1$. Then r is called a *binding relation* on ϱ .

Let ϱ be an n -ary relation on G , let r be a binding relation on ϱ . Then the triple $\mathbf{G} = (G, \varrho, r)$ is called a *double n -ary relational structure* (or briefly a *double n -ary structure*). An element $\alpha \in \varrho$ is called *isolated in G* if $(\alpha, \beta) \notin r$ and $(\beta, \alpha) \notin r$ for any $\beta \in \varrho$. The set of all isolated elements in G is denoted by ϱ_i .

A double n -ary structure $\mathbf{G} = (G, \varrho, r)$ (and its binary relation r) is called

inversely symmetric if $(\alpha, \beta) \in r$ implies $(\beta^{-1}, \alpha^{-1}) \in r$ for any $\alpha, \beta \in \varrho$;

inversely asymmetric if $(\alpha, \beta) \in r$ implies $(\beta^{-1}, \alpha^{-1}) \notin r$ for any $\alpha, \beta \in \varrho$;

transferable if $(\alpha, \beta) \in r$ implies the existence of elements $\alpha_1, \alpha_2, \dots, \alpha_{n-1} \in \varrho$ such that $(\beta, \alpha_1) \in r$, $(\alpha_j, \alpha_{j+1}) \in r$ for $j = 1, 2, \dots, n-2$, $(\alpha_{n-1}, \alpha) \in r$ for any $\alpha, \beta \in \varrho$;

reversely transitive if $(\alpha, \beta) \in r$, $(\beta^{-1}, \gamma') \in r$ imply $(\alpha, \gamma) \in r$ for any $\alpha, \beta, \gamma \in \varrho$.

Let $\mathbf{G} = (G, \varrho, r)$ be a double n -ary structure. Define an $(n+1)$ -ary relation R on G as follows:

$(x_1, x_2, \dots, x_n, x_{n+1}) \in R \iff (x_1, x_2, \dots, x_n) = \alpha \in \varrho, (x_2, x_3, \dots, x_n, x_{n+1}) = \beta \in \varrho, (\alpha, \beta) \in r$ for any $x_1, x_2, x_3, \dots, x_n, x_{n+1} \in G$. Denote $U(\mathbf{G}) = (G, R)$. Then $U(\mathbf{G})$ is an $(n+1)$ -ary structure.

If we denote by ${}_2\mathcal{R}_n$ the class of all double n -ary structures, and by \mathcal{R}_{n+1} the class of all $(n+1)$ -ary structures, then U is a map of ${}_2\mathcal{R}_n$ into \mathcal{R}_{n+1} .

Now, let $\mathbf{G} = (G, R)$ be an $(n+1)$ -ary structure. Define an n -ary relation ϱ on G as follows:

$(x_1, x_2, \dots, x_n) \in \varrho \iff$ there exists $t \in G$ such that $(x_1, x_2, \dots, x_n, t) \in R$ or $(t, x_1, x_2, \dots, x_n) \in R$ for any $x_1, x_2, \dots, x_n \in G$; further, define a binary relation r on ϱ as follows:

$(\alpha, \beta) \in r \iff \alpha = (x_1, x_2, \dots, x_n) \in \varrho, \beta = (x_2, x_3, \dots, x_{n+1}) \in \varrho, (x_1, x_2, \dots, x_n, x_{n+1}) \in R$ for any $x_1, x_2, \dots, x_n, x_{n+1} \in G$. Denote $L(\mathbf{G}) = (G, \varrho, r)$. Then $L(\mathbf{G})$ is a double n -ary structure and L is a map of \mathcal{R}_{n+1} into ${}_2\mathcal{R}_n$.

Moreover, denote by ${}_2\mathcal{R}'_n$ the class of all double n -ary structures without isolated elements.

1. Theorem. *Let \mathbf{G} be an $(n+1)$ -ary structure. Then $(U \cdot L)(\mathbf{G}) = \mathbf{G}$, i.e. $U \cdot L = \text{id}_{\mathcal{R}_{n+1}}$.*

Proof. Let $\mathbf{G} = (G, R)$, $L(\mathbf{G}) = (G, \varrho, r)$, $(U \cdot L)(\mathbf{G}) = (G, R')$. Let $(x_1, x_2, \dots, x_n, x_{n+1}) \in R$. By the definition of L , we have $(x_1, x_2, \dots, x_n) = \alpha \in \varrho$, $(x_2, x_3, \dots, x_n, x_{n+1}) = \beta \in \varrho$, $(\alpha, \beta) \in r$. By the definition of U , we have $(x_1, x_2, \dots, x_n, x_{n+1}) \in R'$. Thus $R \subseteq R'$. Let $(x_1, x_2, \dots, x_n, x_{n+1}) \in R'$. Then, by the definition of U , $(x_1, x_2, \dots, x_n) = \alpha \in \varrho$, $(x_2, x_3, \dots, x_n, x_{n+1}) = \beta \in \varrho$, $(\alpha, \beta) \in r$. By the definition of L , $(x_1, x_2, \dots, x_{n+1}) \in R$. Hence $R' \subseteq R$. Summarizing, we conclude $R = R'$. \square

2. Theorem. Let $\mathbf{G} = (G, \varrho, r)$ be a double n -ary structure and let $(L \cdot U)(\mathbf{G}) = (G, \varrho', r')$. Then $\varrho' = \varrho - \varrho_i$, $r' = r$, i.e. $L \cdot U|_{2\mathcal{R}'_n} = \text{id}_{2\mathcal{R}'_n}$.

Proof. Denote $U(\mathbf{G}) = (G, R)$. Let $(x_1, x_2, \dots, x_n) \in \varrho'$. Then, by the definition of L , there exists $t \in G$ such that $(x_1, x_2, \dots, x_n, t) \in R$ or $(t, x_1, x_2, \dots, x_n) \in R$. In the first case, by the definition of U , we have $(x_1, x_2, \dots, x_n) = \alpha \in \varrho$, $(x_2, x_3, \dots, x_n, t) = \beta \in \varrho$, $(\alpha, \beta) \in r$, thus the element $\alpha \in \varrho$ is not isolated, so that $\alpha \in \varrho - \varrho_i$. In the second case, $(t, x_1, x_2, \dots, x_{n-1}) = \alpha \in \varrho$, $(x_1, x_2, \dots, x_{n-1}, x_n) = \beta \in \varrho$, $(\alpha, \beta) \in r$, hence the element $\beta \in \varrho$ is not isolated and $\beta \in \varrho - \varrho_i$. We have $\varrho' \subseteq \varrho - \varrho_i$. Let, on the contrary, $\alpha = (x_1, x_2, \dots, x_n) \in \varrho - \varrho_i$. Then there exists $\beta \in \varrho$ such that $(\alpha, \beta) \in r$ or $(\beta, \alpha) \in r$. In the first case we have $\beta = (x_1, x_2, \dots, x_n, t)$ for some $t \in G$, therefore, by the definition of U , $(x_1, x_2, \dots, x_n, t) \in R$ and, by the definition of L , $\alpha \in \varrho'$. The second case is analogous. Hence $\varrho - \varrho_i \subseteq \varrho'$. Altogether, we have $\varrho' = \varrho - \varrho_i$.

Let $(\alpha, \beta) \in r'$. By the definition of L , $\alpha = (x_1, x_2, \dots, x_n)$, $\beta = (x_2, x_3, \dots, x_n, x_{n+1}) \in R$ for some $x_1, x_2, x_3, \dots, x_n, x_{n+1} \in G$, $(x_1, x_2, \dots, x_n, x_{n+1}) \in R$. This implies, by the definition of U , $\alpha \in \varrho$, $\beta \in \varrho$, $(\alpha, \beta) \in r$. Thus $r' \subseteq r$. Let $(\alpha, \beta) \in r$. Then $\alpha = (x_1, x_2, \dots, x_n) \in \varrho$, $\beta = (x_2, x_3, \dots, x_n, x_{n+1}) \in \varrho$ for some $x_1, x_2, x_3, \dots, x_n, x_{n+1} \in G$, hence, by the definition of U , we have $(x_1, x_2, \dots, x_n, x_{n+1}) \in R$. Consequently, by the definition of L , $\alpha \in \varrho'$, $\beta \in \varrho'$, $(\alpha, \beta) \in r'$, and $r \subseteq r'$. Summarizing, we obtain $r = r'$.

In the case that G contains no isolated elements, we have $\varrho_i = \emptyset$, thus $\varrho = \varrho'$, $r = r'$, so that $L \cdot U|_{2\mathcal{R}'_n} = \text{id}_{2\mathcal{R}'_n}$.

Denote by ${}_2\mathbf{R}_n$ the category whose class of objects is ${}_2\mathcal{R}_n$ and whose morphisms are maps preserving both relations, i.e., for $\mathbf{G} = (G, \varrho, r)$, $\mathbf{H} = (H, \sigma, s) \in {}_2\mathcal{R}_n$, a map $f : G \rightarrow H$ is a morphism if $(x_1, x_2, \dots, x_n) \in \varrho$ implies $(f(x_1), f(x_2), \dots, f(x_n)) \in \sigma$, and $((x_1, x_2, \dots, x_n), (x_2, x_3, \dots, x_{n+1})) \in r$ implies $((f(x_1), f(x_2), \dots, f(x_n)), (f(x_2), f(x_3), \dots, f(x_{n+1}))) \in s$ for any $x_1, x_2, x_3, \dots, x_n, x_{n+1} \in G$.

Further, denote by \mathbf{R}_{n+1} the category whose class of objects is \mathcal{R}_{n+1} and whose morphisms are maps preserving the relation, i.e., for $\mathbf{G} = (G, H)$, $\mathbf{H} = (H, S) \in \mathcal{R}_{n+1}$ a map $f : G \rightarrow H$ is a morphism if $(x_1, x_2, \dots, x_n, x_{n+1}) \in R$ implies $(f(x_1), f(x_2), \dots, f(x_n), f(x_{n+1})) \in S$ for any $x_1, x_2, \dots, x_n, x_{n+1} \in G$.

Moreover, for any morphism $f \in \text{Hom}_{{}_2\mathbf{R}_n}(\mathbf{G}, \mathbf{H})$, where $\mathbf{G} = (G, \varrho, r)$, $\mathbf{H} = (H, \sigma, s)$, denote $U(f) = f$. Similarly, for any morphism $f \in \text{Hom}_{\mathbf{R}_{n+1}}(\mathbf{G}, \mathbf{H})$, denote $L(f) = f$. \square

3. Theorem. U is a covariant functor from the category ${}_2\mathbf{R}_n$ to the category \mathbf{R}_{n+1} , L is a covariant functor from the category \mathbf{R}_{n+1} to the category ${}_2\mathbf{R}_n$.

Proof. Let $f \in \text{Hom}_{{}_2\mathbf{R}_n}(\mathbf{G}, \mathbf{H})$, where $\mathbf{G} = (G, \varrho, r)$, $\mathbf{H} = (H, \sigma, s)$, $U(\mathbf{H}) = (H, S)$. Let $(x_1, x_2, \dots, x_n, x_{n+1}) \in R$. Then $(x_1, x_2, \dots, x_n) \in \varrho$,

$(x_2, x_3, \dots, x_n, x_{n+1}) \in \varrho$, $((x_1, x_2, \dots, x_n), (x_2, x_3, \dots, x_n, x_{n+1})) \in r$, so that $(f(x_1), f(x_2), \dots, f(x_n)) \in \sigma$, $(f(x_2), f(x_3), \dots, f(x_n), f(x_{n+1})) \in \sigma$, $((f(x_1), f(x_2), \dots, f(x_n)), (f(x_2), f(x_3), \dots, f(x_n), f(x_{n+1}))) \in s$, thus $(f(x_1), f(x_2), \dots, f(x_n), f(x_{n+1})) \in S$ and $U(f) \in \text{Hom}_{\mathbf{R}_{n+1}}(U(\mathbf{G}), U(\mathbf{H}))$. It is easy to show that $U(\text{id}_{\mathbf{G}}) = \text{id}_{U(\mathbf{G})}$ for any $\mathbf{G} \in {}_2\mathcal{R}_n$ and $U(g \cdot f) = U(g) \cdot U(f)$ for any $f \in \text{Hom}_{\mathbf{R}_n}(\mathbf{G}, \mathbf{H})$, $g \in \text{Hom}_{\mathbf{R}_n}(\mathbf{H}, \mathbf{K})$, $\mathbf{G}, \mathbf{H}, \mathbf{K} \in {}_2\mathcal{R}_n$. Analogously for L . \square

4. Theorem. *Let \mathbf{G} be a double n -ary structure. Then the following assertions hold:*

- (i) \mathbf{G} is inversely symmetric if and only if $U(\mathbf{G})$ is symmetric.
- (ii) \mathbf{G} is inversely asymmetric if and only if $U(\mathbf{G})$ is asymmetric.

Proof. Let $\mathbf{G} = (G, \varrho, r)$, $U(\mathbf{G}) = (G, R)$.

(i) Let G be inversely symmetric and let $(x_1, x_2, \dots, x_n, x_{n+1}) \in R$. Then $(x_1, x_2, \dots, x_n) = \alpha \in \varrho$, $(x_2, x_3, \dots, x_n, x_{n+1}) = \beta \in \varrho$, $(\alpha, \beta) \in r$. This implies $(\beta^{-1}, \alpha^{-1}) \in r$, thus $\beta^{-1} = (x_{n+1}, x_n, \dots, x_3, x_2) \in \varrho$, $\alpha^{-1} = (x_n, \dots, x_2, x_1) \in \varrho$, so that $(x_{n+1}, x_n, \dots, x_2, x_1) \in R$ and $U(\mathbf{G})$ is symmetric. Let $U(\mathbf{G})$ be symmetric and let $(\alpha, \beta) \in r$. Then there exist elements $x_1, x_2, \dots, x_n, x_{n+1} \in G$ such that $\alpha = (x_1, x_2, \dots, x_n) \in \varrho$, $\beta = (x_2, x_3, \dots, x_n, x_{n+1}) \in \varrho$. This implies $(x_1, x_2, \dots, x_n, x_{n+1}) \in R$, so that $(x_{n+1}, x_n, \dots, x_2, x_1) \in R$, i.e. $(x_{n+1}, x_n, \dots, x_3, x_2) = \beta^{-1} \in \varrho$, $(x_n, \dots, x_2, x_1) = \alpha^{-1} \in \varrho$, hence $(\beta^{-1}, \alpha^{-1}) \in r$ and \mathbf{G} is inversely symmetric.

(ii) Let \mathbf{G} be inversely asymmetric and let $(x_1, x_2, \dots, x_n, x_{n+1}) \in R$. Then again $(x_1, x_2, \dots, x_n) = \alpha \in \varrho$, $(x_2, x_3, \dots, x_n, x_{n+1}) = \beta \in \varrho$, $(\alpha, \beta) \in r$. This implies $(\beta^{-1}, \alpha^{-1}) \notin r$. But $\beta^{-1} = (x_{n+1}, x_n, \dots, x_3, x_2)$, $\alpha^{-1} = (x_n, \dots, x_2, x_1)$, thus $(x_{n+1}, x_n, \dots, x_2, x_1) \notin R$ and $U(\mathbf{G})$ is asymmetric. Let $U(\mathbf{G})$ be asymmetric and let $(\alpha, \beta) \in r$. Then there exist elements $x_1, x_2, x_3, \dots, x_n, x_{n+1} \in G$ such that $(x_1, x_2, \dots, x_n) = \alpha \in \varrho$, $(x_2, x_3, \dots, x_n, x_{n+1}) = \beta \in \varrho$. This implies $(x_1, x_2, \dots, x_n, x_{n+1}) \in R$, so that $(x_{n+1}, x_n, \dots, x_2, x_1) \notin R$. Consequently $(x_{n+1}, x_n, \dots, x_3, x_2) = \beta^{-1} \notin \varrho$ or $(x_n, \dots, x_2, x_1) = \alpha^{-1} \notin \varrho$ or $\beta^{-1}, \alpha^{-1} \in \varrho$, but $(\beta^{-1}, \alpha^{-1}) \notin r$. In all three cases, however, we have $(\beta^{-1}, \alpha^{-1}) \notin r$, and \mathbf{G} is inversely asymmetric. \square

5. Theorem. *Let \mathbf{G} be an $(n+1)$ -ary structure. Then the following assertions hold:*

- (i) \mathbf{G} is symmetric if and only if $L(\mathbf{G})$ is inversely symmetric.
- (ii) \mathbf{G} is asymmetric if and only if $L(\mathbf{G})$ is inversely asymmetric.

Proof. (i) If $L(\mathbf{G})$ is inversely symmetric, then, by 4, $U(L(\mathbf{G}))$ is symmetric. But, by 1, $U(L(\mathbf{G})) = \mathbf{G}$. If $\mathbf{G} = U(L(\mathbf{G}))$ is symmetric, then, by 4, $L(\mathbf{G})$ is inversely symmetric.

(ii) If $L(\mathbf{G})$ is inversely asymmetric, then, by 4, $U(L(\mathbf{G}))$ is asymmetric. But $U(L(\mathbf{G})) = \mathbf{G}$. If $\mathbf{G} = U(L(\mathbf{G}))$ is asymmetric, then, by 4, $L(\mathbf{G})$ is inversely asymmetric. \square

6. Theorem. *Let \mathbf{G} be a double n -ary structure. Then \mathbf{G} is transferable if and only if $U(\mathbf{G})$ is cyclic.*

Proof. Let $\mathbf{G} = (G, \varrho, r)$, $U(\mathbf{G}) = (G, R)$. Let \mathbf{G} be transferable and let $(x_1, x_2, \dots, x_n, x_{n+1}) \in R$. Then $(x_1, x_2, \dots, x_n) = \alpha \in \varrho$, $(x_2, x_3, \dots, x_n, x_{n+1}) = \beta \in \varrho$, $(\alpha, \beta) \in r$. Thus, there exist $\alpha_1, \alpha_2, \dots, \alpha_{n-1} \in \varrho$ such that $(\beta, \alpha_1) \in r$, $(\alpha_j, \alpha_{j+1}) \in r$ for $j = 1, 2, \dots, n-2$ and $(\alpha_{n-1}, \alpha) \in r$. Denote $\alpha_0 = \beta$, $\alpha_n = \alpha$. Then we have (α_j, α_{j+1}) for $j = 0, 1, 2, \dots, n-1$. We shall show by induction that $\alpha_j = (x_{j+2}, x_{j+3}, \dots, x_n, x_{n+1}, x_1, x_2, \dots, x_j)$ for $j = 0, 1, 2, \dots, n$. For $j = 0$ it is true. Let $0 < j_0 \leq n$. Let the preceding hold for each j , $0 \leq j < j_0$. As $(\alpha_{j_0-1}, \alpha_{j_0}) \in r$ and r is binding, there exists $y \in G$ such that $\alpha_{j_0} = (x_{j_0+2}, x_{j_0+3}, \dots, x_1, \dots, x_{j_0-1}, y)$. We shall show by another induction that α_{j_0+k} has y on the $(n-k)$ -th position, for $k = 0, 1, 2, \dots, n-j_0$. For $k = 0$ it is true. Let $0 < k_0 \leq n-j_0$. As $(\alpha_{j_0+k_0-1}, \alpha_{j_0+k_0}) \in r$, $\alpha_{j_0+k_0-1}$ has y on the $(n-k_0+1)$ -th position, and r is binding, $\alpha_{j_0+k_0}$ has y on the $(n-k_0)$ -th position. Particularly, α_n has y on the j_0 -th position, hence $y = x_{j_0}$. Thus, we have $\beta = (x_2, x_3, \dots, x_n, x_{n+1}) \in \varrho$, $\alpha_1 = (x_3, x_4, \dots, x_n, x_{n+1}, x_1) \in \varrho$, $(\beta, \alpha_1) \in r$, so that $(x_2, x_3, \dots, x_n, x_{n+1}, x_1) \in R$ and $U(\mathbf{G})$ is cyclic.

Let, on the contrary, $U(\mathbf{G})$ be cyclic and let $(\alpha, \beta) \in r$. Then there exist elements $x_1, x_2, \dots, x_n, x_{n+1} \in G$ such that $\alpha = (x_1, x_2, \dots, x_n) \in \varrho$, $\beta = (x_2, x_3, \dots, x_n, x_{n+1}) \in \varrho$, thus $(x_1, x_2, \dots, x_n, x_{n+1}) \in R$. Hence $(x_2, x_3, \dots, x_n, x_{n+1}, x_1) \in R$, $(x_3, x_4, \dots, x_n, x_{n+1}, x_1, x_2) \in R, \dots, (x_{n+1}, x_1, x_2, \dots, x_n) \in R$. Denote $\alpha_1 = (x_3, x_4, \dots, x_n, x_{n+1}, x_1)$, $\alpha_2 = (x_4, x_5, \dots, x_{n+1}, x_1, x_2), \dots, \alpha_{n-1} = (x_{n+1}, x_1, x_2, \dots, x_{n-1})$. Then $\alpha_j \in \varrho$ for $j = 1, 2, \dots, n-1$, $(\beta, \alpha_1) \in r$, $(\alpha_j, \alpha_{j+1}) \in r$ for $j = 1, 2, \dots, n-2$, $(\alpha_{n-1}, \alpha) \in r$. Consequently, \mathbf{G} is transferable. \square

7. Theorem. *Let $L(\mathbf{G})$ be an $(n+1)$ -ary structure. Then \mathbf{G} is cyclic if and only if $L(\mathbf{G})$ is transferable.*

Proof. Let $L(\mathbf{G})$ be transferable. By 6, $U(L(\mathbf{G}))$ is cyclic. But, by 1, $\mathbf{G} = U(L(\mathbf{G}))$.

Let $\mathbf{G} = U(L(\mathbf{G}))$ be cyclic. Then, by 6, $L(\mathbf{G})$ is transferable. \square

8. Theorem. *Let $\mathbf{G} = (G, \varrho, r)$ be a double n -ary structure. If the binary relation r is transitive, then $U(\mathbf{G})$ is weakly transitive.*

Proof. Let $U(\mathbf{G}) = (G, R)$ and let $(x, y, y, \dots, y) \in R$, $(y, y, \dots, y, z) \in R$. Then $\alpha = (x, y, y, \dots, y) \in \varrho$, $\beta = (y, y, \dots, y) \in \varrho$, $\gamma = (y, y, \dots, y, z) \in \varrho$, $(\alpha, \beta) \in$

$r, (\beta, \gamma) \in r$. Hence $(\alpha, \gamma) \in r$, so that $(x, y, y, \dots, y, z) \in R$ and $U(\mathbf{G})$ is weakly transitive. \square

9. Remark. The converse of 8 does not hold, which can be easily shown by a counterexample.

10. Theorem. *Let \mathbf{G} be a double n -ary structure. Then \mathbf{G} is reversely transitive if and only if $U(\mathbf{G})$ is transitive.*

Proof. Let $\mathbf{G} = (G, \varrho, r)$, $U(\mathbf{G}) = (G, R)$. Let \mathbf{G} be reversely transitive, let $(x_1, x_2, \dots, x_n, x_{n+1}) \in R$, $(x_{n+1}, x_n, \dots, x_2, x_{n+2}) \in R$. Then, by the definition of U , $(x_1, x_2, \dots, x_n) = \alpha \in \varrho$, $(x_2, x_3, \dots, x_n, x_{n+1}) = \beta \in \varrho$, $(\alpha, \beta) \in r$, $(x_{n+1}, x_n, \dots, x_2) = \beta^{-1} \in \varrho$, $(x_n, x_{n-1}, \dots, x_2, x_{n+2}) = \gamma' \in \varrho$, $(\beta^{-1}, \gamma') \in r$. As \mathbf{G} is reversely transitive, we have $(\alpha, \gamma) \in r$. But $\gamma = (x_2, x_3, \dots, x_n, x_{n+2}) \in \varrho$, hence $(x_1, x_2, \dots, x_n, x_{n+2}) \in R$ and $U(\mathbf{G})$ is transitive.

Let $U(\mathbf{G})$ be transitive and let $\alpha, \beta, \gamma \in \varrho$, $(\alpha, \beta) \in r$, $(\beta^{-1}, \gamma') \in r$. There exist elements $x_1, x_2, \dots, x_n, x_{n+1}, x_{n+2} \in G$ such that $\alpha = (x_1, x_2, \dots, x_n)$, $\beta = (x_2, x_3, \dots, x_n, x_{n+1})$ (for r is binding), $\gamma = (x_2, x_3, \dots, x_n, x_{n+2})$ (for $\beta^{-1} = (x_{n+1}, x_n, \dots, x_3, x_2)$, $\gamma' = (x_n, x_{n-1}, \dots, x_3, x_2, x_{n+2})$ and r is binding). Hence $(x_1, x_2, \dots, x_n, x_{n+1}) \in R$, $(x_{n+1}, x_n, \dots, x_3, x_2, x_{n+2}) \in R$, so that $(x_1, x_2, \dots, x_n, x_{n+2}) \in R$, for $U(\mathbf{G})$ is transitive. Consequently, $(\alpha, \gamma) \in r$ and \mathbf{G} is reversely transitive. \square

11. Theorem. *Let \mathbf{G} be an $(n + 1)$ -ary structure. Then \mathbf{G} is transitive if and only if $L(\mathbf{G})$ is reversely transitive.*

Proof. By 1, $U(L(\mathbf{G})) = \mathbf{G}$. Hence $L(\mathbf{G})$ is reversely transitive if and only if $U(L(\mathbf{G})) = \mathbf{G}$ is transitive, by 10. \square

References

- [1] V. Novák: Cyclically ordered sets. Czechoslovak Math. J. 32 (1982), 460–473.
- [2] V. Novák, M. Novotný: On determination of a cyclic order. Czechoslovak Math. J. 33 (1983), 555–563.
- [3] V. Novák: Dimension theory for cyclically and cocyclically ordered sets. Czechoslovak Math. J. 33 (1983), 647–653.
- [4] V. Novák: On some minimal problem. Arch. Math. (Brno) 20 (1984), 95–99.
- [5] V. Novák: Cuts in cyclically ordered sets. Czechoslovak Math. J. 34 (1984), 322–333.
- [6] V. Novák, M. Novotný: Universal cyclically ordered sets. Czechoslovak Math. J. 35 (1985), 158–161.
- [7] V. Novák, M. Novotný: Binary and ternary relations. Math. Bohem. 117 (1992), 283–292.

Author's address: Jiří Karásek, Technical University, Technická 2, 616 69 Brno, Czech Republic.