

DOMINATING FUNCTIONS OF GRAPHS WITH TWO VALUES

BOHDAN ZELINKA, Liberec

(Received March 6, 1997)

Abstract. The Y -domination number of a graph for a given number set Y was introduced by D. W. Bange, A. E. Barkauskas, L. H. Host and P. J. Slater as a generalization of the domination number of a graph. It is defined using the concept of a Y -dominating function. In this paper the particular case where $Y = \{0, 1/k\}$ for a positive integer k is studied.

Keywords: Y -dominating function of a graph, Y -domination number of a graph

MSC 1991: 05C35

This paper will concern a certain generalization of the domination number of a graph. All graphs considered will be finite undirected graphs without loops and multiple edges.

A subset D of the vertex set $V(G)$ of a graph G is called dominating in G , if for each vertex $x \in V(G) - D$ there exists a vertex $y \in D$ adjacent to x . The minimum number of vertices of a dominating set in G is called the domination number of G and denoted by $\gamma(G)$.

This well-known concept can be defined in another way, using domination functions. We will speak about functions f which map $V(G)$ into some set of numbers. If $S \subseteq V(G)$, then we denote $f(S) = \sum_{x \in S} f(x)$. If $x \in V(G)$, then by $N[x]$ we denote the closed neighbourhood of x in G , i.e. the set consisting of x and of all vertices which are adjacent to x in G . Besides, we will also consider the open neighbourhood $N(x) = N[x] - \{x\}$. Now we can formulate the alternative definition of the domination number.

A function $f: V(G) \rightarrow \{0, 1\}$ is called a dominating function of G , if $f(N[x]) \geq 1$ for each $x \in V(G)$. The minimum sum $f(V(G)) = \sum_{x \in V(G)} f(x)$ taken over all dominating functions f of G is called the domination number of G and denoted by $\gamma(G)$.

It is evident that these two definitions are equivalent. Namely, if D is a dominating set in G , then the function f defined so that $f(x) = 1$ for $x \in D$ and $f(x) = 0$ for $x \in V(G) - D$ is a dominating function of G . Conversely, if f is a dominating function of G , then the set $D = \{x \in V(G); f(x) = 1\}$ is a dominating set in G .

The concept of a dominating function and obviously also the related concept of the domination number were generalized by some authors in such a way that the set of values $\{0, 1\}$ was replaced by another number set. In [1] the signed dominating function and the signed domination number were defined by replacing the set $\{0, 1\}$ by $\{-1, 1\}$ and in [2] the minus dominating function and the minus domination number were defined by using the set $\{-1, 0, 1\}$. The fractional dominating function and the fractional domination number were introduced in [3] by using the set of real numbers. The most general case is the Y -dominating function and the Y -domination number, where a quite arbitrary set Y of values of f is used [4].

Therefore, following [4], a function $f: V(G) \rightarrow Y$, where Y is a given set of numbers, is called a Y -dominating function of G , if $f(N[x]) \geq 1$ for each $x \in V(G)$. The minimum of $f(V(G))$ taken over all Y -dominating functions f of G is called the Y -domination number of G and is denoted by $\gamma_Y(G)$.

We will not treat the domination in such a general way. We restrict our considerations to natural generalizations of the set $\{0, 1\}$, namely to two-element number sets $\{0, t\}$, where t is a positive real number.

The following proposition is easy to prove.

Proposition 1. *Let $Y = \{0, t\}$, where t is a positive real number. Let G be a graph. The Y -domination number $\gamma_Y(G)$ of G is defined and at least one Y -dominating function of G exists if and only if $\delta(G) \geq 1/t - 1$, where $\delta(G)$ denotes the minimum degree of a vertex of G .*

Let f be a function which maps $V(G)$ into the set of real numbers and let $x \in V(G)$. The vertex set x will be called a zero vertex of f , if $f(x) = 0$.

The following theorem enables us to restrict our consideration to numbers t which are inverses of positive integers.

Theorem 1. *Let t be a positive real number, let G be a graph with $\delta(G) \geq 1/t - 1$. Let $k = \lceil 1/t \rceil$ and $Y_1 = \{0, t\}$, $Y_2 = \{0, 1/k\}$. Then $\gamma_{Y_1}(G) = kt\gamma_{Y_2}(G)$ and there exists a one-to-one correspondence between Y_1 -dominating functions of G and Y_2 -dominating functions of G such that the corresponding functions have the same set of zero vertices.*

Proof. Let $f: V(G) \rightarrow Y_1$, $g: V(G) \rightarrow Y_2$ and suppose that f, g have the same set of zero vertices. Then $f(x) = ktg(x)$ and also $f(N[x]) = ktg(N[x])$ for

each $x \in V(G)$. Suppose that g is a Y_2 -dominating function of G : then $g(N[x]) \geq 1$ for each $x \in V(G)$. Evidently $kt \geq 1$ and thus $f(N[x]) \geq g(N[x]) \geq 1$ for each $x \in V(G)$ and f is a Y_1 -dominating function of G . Now suppose that g is not a Y_2 -dominating function of G . There exists $x \in V(G)$ such that $g(N[x]) < 1$. If $k = 1$, then $g(N[x])$ must be a non-negative integer and therefore $g(N[x]) = 0$. This is possible only if $g(y) = 0$ for each $y \in N[x]$. But then also $f(y) = 0$ for each $y \in N[x]$ and $f(N[x]) = 0$; the function f is not a Y_1 -dominating function of G . If $k \geq 2$, then the number of vertices of $N[x]$ which are not zero vertices of g is at most $k - 1$. But these vertices are exactly those vertices which are not zero vertices of f . We have $f(N[x]) \leq (k - 1)t$. Evidently $1/t > k - 1$ and thus $f(N[x]) \leq (k - 1)t < 1$; the function f is not a Y_1 -dominating function of G . If g_0 is a minimal (i.e. with the minimum sum on $V(G)$) Y_2 -dominating function, then the corresponding function f_0 is a minimal Y_1 -dominating function. We have $\gamma_{Y_1}(G) = \sum_{x \in V(G)} f_0(x) = \sum_{x \in V(G)} ktg_0(x) = kt \sum_{x \in V(G)} g_0(x) = kt\gamma_{Y_2}(G)$. \square

For each positive integer k we denote $Y(k) = \{0, 1/k\}$ and $\gamma(k, G) = \gamma_{Y(k)}G$. From Proposition 1 we have the following corollary.

Corollary 1. *Let k be a positive integer, let G be a graph. The $Y(k)$ -domination number $\gamma(k, G)$ is defined and at least one $Y(k)$ -dominating function of G exists if and only if $\delta(G) \geq k - 1$.*

Note that $\gamma(1, G) = \gamma(G)$, the usual domination number of G .

If we speak about a function $f: V(G) \rightarrow Y(k)$, we will use the notation $V^0 = \{x \in V(G); f(x) = 0\}$, $V^+ = \{x \in V(G); f(x) = 1/k\}$.

Theorem 2. *Let G be a regular graph of degree $k - 1$ with n vertices. Then $\gamma(k, G) = n/k$.*

Proof. The neighbourhood $N[x]$ for each $x \in V(G)$ has exactly k vertices. If f is a $Y(k)$ -dominating function, then f must assign the value $1/k$ to all vertices of $N[x]$. As x was chosen arbitrarily, it assigns $1/k$ to all vertices of G , which implies the assertion. \square

By G^2 we denote the square of the graph G , i.e. the graph such that $V(G^2) = V(G)$ and two vertices are adjacent in G^2 if and only if their distance in G is at most 2. The symbol $\alpha_0(G)$ denotes the independence number of G , i.e. the maximum number of pairwise non-adjacent vertices in G .

Theorem 3. *Let G be a regular graph of degree k with n vertices. Then $\gamma(k, G) = (n - \alpha_0(G^2))/k$.*

Proof. For each vertex x of G the set $N[x]$ has $k + 1$ vertices. If f is a $Y(k)$ -dominating function of G , then $N[x]$ contains at most one zero vertex of f . The distance between two zero vertices of f cannot be 1; then the closed neighbourhood of either of them would contain them both. This distance cannot be 2; then there would exist a vertex adjacent to both of them and its closed neighbourhood would contain them both. Therefore the distance between two zero vertices of f in G is at least 3 and in G^2 at least 2; they form an independent set in G^2 . Therefore there are at most $\alpha_0(G^2)$ zero vertices of f and at least $n - \alpha_0(G^2)$ vertices x such that $f(x) = 1/k$. This implies the assertion. \square

Corollary 2. *Let C_n be the circuit of length n . Then $\gamma(3, C_n) = n/3$ and $\gamma(2, C_n) = n/3$ for $n \equiv 0(\text{mod } 3)$, $\gamma(2, C_n) = n/3 - 1/6$ for $n \equiv 1(\text{mod } 3)$, $\gamma(2, C_n) = n/3 + 1/3$ for $n \equiv 2(\text{mod } 3)$.*

A path is a similar case. If f is a $Y(2)$ -dominating function of a path P_n of length n , then again the distance between any two zero vertices of f is at least 3 and moreover neither the vertices of degree 1, nor the vertices adjacent to them may be zero vertices of f . This yields the result.

Proposition 2. *Let P_n be a path of length n . Then $\gamma(2, P_n) = n/3 + 1$ for $n \equiv 0(\text{mod } 3)$, $\gamma(2, P_n) = n/3 + 2/3$ for $n \equiv 1(\text{mod } 3)$, $\gamma(2, P_n) = n/3 + 5/6$ for $n \equiv 2(\text{mod } 3)$.*

Now we turn to complete graphs and complete bipartite graphs.

Theorem 4. *Let k, n be positive integers, $k \leq n$. Then $\gamma(k, K_n) = 1$.*

Proof. In the complete graph K_n we have $N[x] = V(K_n)$ for each vertex x . If f is a $Y(k)$ -dominating function, then $f(V(K_n)) = f(N[x]) \geq 1$. Moreover, there exists a function f which assigns the value $1/k$ to k vertices and the value 0 to the remaining $n - k$ vertices: then $f(V(K_n)) = 1$. \square

Theorem 5. *Let k, m, n be positive integers, $k - 1 \leq m \leq n$. If $k < m$, then $\gamma(k, K_{m,n}) = 2$. If $m = k - 1$, then $\gamma(k, K_{m,n}) = (m + n)/k = (k + n - 1)/k$. If $m = k$, then $\gamma(k, K_{m,n}) = 2 - 1/k$.*

Proof. Let $k < m$. Let A, B be the bipartition classes of K , $|A| = m$, $|B| = n$. For each vertex $x \in A$, its open neighbourhood satisfies $N(x) \subseteq B$. As $N[x] = \{x\} \cup N(x)$ and $f(N[x]) \geq 1$ for a $Y(k)$ -dominating function f , there are at least $k - 1$ vertices $y \in N(x) \subseteq B$ which are in V^+ . If moreover $f(x) = 0$, then there are at least k such vertices. Therefore either $f(x) = 1/k$ for all $x \in A$ and

$f(y) = 1/k$ for at least $k - 1$ vertices of B , or $f(y) = 1/k$ for at least k vertices of B . In the former case $f(V(K_{m,n})) \geq (m + k - 1)/k \geq 2$. In the latter case analogously either $f(x) = 1/k$ for all $x \in B$ and $f(y) = 1/k$ for at least $k - 1$ vertices of A , or $f(y) = 1/k$ for at least k vertices of A . In both these cases again $f(V(K_{m,n})) \geq 2$. A function f which assigns $1/k$ to exactly k vertices of A and to exactly k vertices of B has $f(V(K_{m,n})) = 2$.

Now suppose $m = k - 1$. Then $|A| = k - 1$. Let $x \in B$ and again let f be a $Y(k)$ -dominating function of $K_{m,n}$. The set $N[x]$ has exactly k vertices and thus $f(x) = 1/k$ for each $y \in N[x]$. This means that $f(y) = 1/k$ for each $y \in A$ and also $f(x) = 1/k$. As x is an arbitrary vertex of B , we have $f(x) = 1/k$ for all $x \in V(K_{m,n})$ and $f(V(K_{m,n})) = (k - 1 + n)/k$. Another $Y(k)$ -dominating function does not exist and thus $\gamma(k, K_{m,n}) = (k - 1 + n)/k$.

Finally, let $k = m$. If f is a $Y(k)$ -dominating function, then either $f(x) = 1/k$ for each $x \in A$ and for at least $k - 1$ vertices x of B , or $f(x) = 1/k$ for exactly $k - 1$ vertices of A and all vertices $x \in B$. In the former case $f(V(K_{m,n})) \geq (2k - 1)/k = 2 - 1/k$, in the latter case $f(V(K_{m,n})) \geq (k - 1 + n)/k \geq (2k - 1)/k = 2 - 1/k$. If f assigns the value $1/k$ to all vertices of A and to exactly $k - 1$ vertices of B , then $f(V(K_{m,n})) = 2 - 1/k$, therefore $\gamma(k, K_{m,n}) = 2 - 1/k$. \square

By the symbol $G \oplus H$ we denote the Zykov sum of graphs G and H , i.e. the graph obtained from vertex-disjoint graphs G and H by joining all vertices of G with all vertices of H by edges.

Theorem 6. *Let k, q be positive integers, let G, H be two graphs such that $\gamma(k, G), \gamma(k, H)$ are defined and $q \leq 1 + \min(\gamma(k, G), \gamma(k, H))$. Then $\gamma(kq, G \oplus H) \leq (\gamma(k, G) + \gamma(k, H))/q$.*

Proof. Let g and h be minimal $Y(k)$ -dominating functions of G and H , respectively. Let $f: V(G) \cup V(H) \rightarrow Y(kq)$ be defined so that $f(x) = g(x)/q$ for $x \in V(G)$ and $f(x) = h(x)/q$ for $x \in V(H)$. Consider $x \in V(G)$. The closed neighbourhood of x in $G \oplus H$ is the disjoint union of the closed neighbourhood of x in G and of $V(H)$. The sum of values of f over the closed neighbourhood of x in G is at least $1/q$, its sum over $V(H)$ is at least $\gamma(k, H)/q$. It follows from the assumption that $1/q + \gamma(k, H)/q \geq 1$. For $x \in V(H)$ this may be proved quite analogously. Therefore f is a $Y(kq)$ -dominating function of $G \oplus H$. This implies the assertions. \square

For the particular case $k = 1$ we have a corollary.

Corollary 3. *Let q be a positive integer, let G, H be two graphs such that $q \leq 1 + \min(\gamma(G), \gamma(H))$. Then $\gamma(q, G \oplus H) \leq (\gamma(G) + \gamma(H))/q$.*

A similar assertion holds for $G \oplus K_1$, i.e. the graph which is obtained from G by adding a new vertex and joining it with all vertices of G by edges.

Theorem 7. *Let k be a positive integer, let G be a graph for which $\gamma(k, G)$ is defined. Then*

$$\gamma(k+1, G \oplus K_1) = \gamma(k, G) \cdot \frac{k}{k+1} + \frac{1}{k+1}.$$

Proof. Let f be a minimal $Y(k)$ -dominating function of G . Let w be the added vertex. Let $g: V(G) \cup \{w\} \rightarrow Y(k+1)$ be defined so that $g(x) = kf(x)/(k+1)$ for $x \in V(G)$ and $g(w) = 1/(k+1)$. Then the sum of $g(x)$ over the closed neighbourhood of x in $G \oplus K_1$ is equal to the sum of g over the closed neighbourhood of x in G plus $g(w)$. The sum of g over the closed neighbourhood of x in G is at least $k/(k+1)$ and $g(w) = 1/(k+1)$, therefore the sum of g over the closed neighbourhood of x in $G \oplus K_1$ is at least 1. The closed neighbourhood of w in $G \oplus K_1$ is $V(G) \cup \{w\}$ and the sum of g over it is greater than or equal to this sum over the closed neighbourhood of any other vertex, therefore it is also at least 1 and

$$\sum_{x \in V(G) \cup \{w\}} g(x) = \frac{k}{k+1} \sum_{x \in V(G)} f(x) + g(w) = \frac{k}{k+1} \gamma(k, G) + \frac{1}{k+1}.$$

Hence $\gamma(k+1, G \oplus K_1) \leq \frac{k}{k+1} \gamma(k, G) + \frac{1}{k+1}$. On the other hand, let g_0 be a minimal $Y(k+1)$ -dominating function of $G \oplus K_1$ and let $f_0: V(G) \rightarrow Y(k)$ be defined so that $f_0(x) = (k+1)g_0(x)/k$ for each $x \in V(G)$. The sum of values of g_0 over the closed neighbourhood of any vertex $x \in V(G)$ in $G \oplus K_1$ is at least $1 - 1/(k+1)$ and thus such a sum of f_0 is at least 1. We have $\sum_{x \in V(G)} f_0(x) = \sum_{x \in V(G)} (k+1)g_0(x)/k = \frac{k+1}{k} \sum_{x \in V(G)} g_0(x) = \frac{k+1}{k} \gamma(k+1, G \oplus K_1) - g_0(w) = \frac{k+1}{k} \gamma(k+1, G \oplus K_1) - \frac{1}{k}$ and thus $\gamma(k, G) \leq \frac{k+1}{k} \gamma(k+1, G \oplus K_1) - \frac{1}{k}$, which yields $\gamma(k+1, G \oplus K_1) \geq \frac{k}{k+1} \gamma(k, G) + \frac{1}{k+1}$. Hence we have the equality $\gamma(k+1, G \oplus K_1) = \frac{k}{k+1} \gamma(k, G) + \frac{1}{k+1}$. \square

In the end we will consider the number $\gamma(k, G)$ for different numbers k and for the same graph G .

Theorem 8. *Let k, q be positive integers. Then there exists a graph G such that $\gamma(k+1, G) - \gamma(k, G) = q$.*

Proof. Denote $p = kq + q + 1$ and let G be the Zykov sum $K_k \oplus \bar{K}_p$, where \bar{K}_p denotes the complement of the complete graph K_p , i.e. the graph consisting of p isolated vertices. If f is a function such that $f(x) = 0$ for $x \in V(\bar{K}_p)$ and $f(x) = 1/k$ for $x \in V(K_k)$, then f is a $Y(k)$ -dominating function of G ; namely, we have $V(K_k) \subseteq N[x]$ for each $x \in V(G)$ and $f(V(K_k)) = 1$. We have $\gamma(k, G) = 1$. Each vertex of

\bar{K}_p has degree k in G and therefore for each $Y(k+1)$ -dominating function g we have $g(y) = 1/(k+1)$ for each $y \in V(G)$ and $\gamma(k+1, G) = (p+k)/(k+1) = q+1$. \square

The next theorem is not expressed for k in general, but only for $\gamma(1, G)$ and $\gamma(2, G)$.

Theorem 9. *Let q be a positive integer. Then there exists a graph G such that $\gamma(1, G) - \gamma(2, G) = q$.*

Proof. Let H be a graph obtained from the circuit of length 4 by adding a new vertex u and joining it to a vertex v of the circuit by an edge. Take $2q$ pairwise vertex-disjoint copies H_1, \dots, H_{2q} of H . Take a vertex w and join it by edges with the vertex corresponding to u in each of the graphs H_1, \dots, H_{2q} . Finally, take a new vertex x and join it with w by an edge. The resulting graph will be G . For $q = 4$ this graph is shown Fig. 1. The number $\gamma(1, G)$ is the usual domination number $\gamma(G)$ of G ,




Fig. 1

i.e. the minimum number of vertices of a dominating set D in G . Evidently such a dominating set must contain at least one of the vertices w, x and at least two vertices from each H for $i = 1, \dots, 2q$: hence $\gamma(G) \geq 4q + 1$. If D consists of w , of the vertices corresponding to v in H and of one other vertex of the circuit in H for $i = 1, \dots, 2q$, then D is dominating in G and $|D| = 4q + 1$, which implies $\gamma(G) = 4q + 1$. Now let V^+ be the set consisting of all vertices of D and, moreover, of x and of one more vertex of the circuit in each H for $i = 1, \dots, 2q$. We have $|V^+| = 6q + 2$. If $f(x) = \frac{1}{2}$ for $x \in V^+$ and $f(x) = 0$ for $x \in V(G) - V^+$, then f is a $V(2)$ -dominating function of G and is evidently minimal. We have $\gamma(2, G) = f(V(G)) = \frac{1}{2}|V^+| = 3q + 1$. Hence $\gamma(1, G) - \gamma(2, G) = q$. \square

Problem. Can Theorem 10 be generalized to a theorem analogous to Theorem 9?

A final remark. The $Y(k)$ -domination number of a graph can be defined in another way, without using the concept of a $Y(k)$ -dominating function:

A subset D of $V(G)$ is called k -tuply dominating in G , if for each $x \in V(G) - D$ there exist k vertices y_1, \dots, y_k of D adjacent to x and for each $y \in D$ there exist $k - 1$ vertices z_1, \dots, z_{k-1} adjacent to y . The minimum number of vertices of a k -tuply dominating set in G is called the $Y(k)$ -domination number of G .

A k -tuply dominating set was defined and used also in [5], but in a weaker form: the requirement of existence of z_1, \dots, z_{k-1} for $y \in D$ was not used there.

References

- [1] Dunbar J.; Hedetniemi S. T.; Henning M. A.; Slater P. J.: Signed domination in graphs. In: Graph Theory, Combinatorics and Applications (ed. Y. Alavi, A. Schwenk). Wiley, New York, 1995, pp. 311–321.
- [2] Dunbar J.; Hedetniemi S. T.; Henning M. A.; McRae A.: Minus domination in graphs. Computers Math. Appl. To appear.
- [3] Grinstead D. L.; Slater P. J.: Fractional domination and fractional packing in graphs. Congr. Numer. 71 (1990), 153–172.
- [4] Bange D. W.; Barkauskas A. E.; Host L. H.; Slater P. J.: Generalized domination and efficient domination in graphs. Discrete Math. 159 (1996), 1–11.
- [5] Zelinka B.: On k -ply domatic numbers of graphs. Math. Slovaca 34 (1984), 313–318.

Author's address: Bohdan Zelinka, Katedra diskrétní matematiky a statistiky Technické university, Hálkova 6, 461 17 Liberec 1, Czech Republic, e-mail: bohdan.zelinka@vslib.cz.