

Note on the cortex of some exponential Lie groups

Bécher Dali

ABSTRACT. In this paper, we built a family of $4d$ -dimensional two-step nilpotent Lie algebras $(\mathfrak{g}_d)_{d \geq 2}$ so that the cortex of the dual of each \mathfrak{g}_d is a projective algebraic set. We also give a complete description of the cortex of the exponential connected and simply connected Lie group $G = \mathbb{R}^n \rtimes \mathbb{R}$.

CONTENTS

1. Introduction	1247
2. Background material and notations	1248
3. The two-step nilpotent Lie algebras	1250
3.1. Main example	1252
4. The Lie group $\mathbb{R}^n \rtimes \mathbb{R}$	1256
References	1260

1. Introduction

The cortex of general locally compact group G was defined in [9] as

$$\text{cor}(G) = \{\pi \in \widehat{G}, \pi \text{ is not Hausdorff-separated}$$

from the identity representation $\mathbf{1}_G\}$,

where \widehat{G} is the dual of G (set of equivalence classes of unitary irreducible representations of G). Note that \widehat{G} is equipped with the topology of Fell which can be described in terms of weak containment (see [6]) and, in general, is not separated. However, if G is abelian, then \widehat{G} is separated and hence $\text{cor}(G) = \{\mathbf{1}_G\}$.

When G is a connected and simply connected nilpotent Lie group with Lie algebra \mathfrak{g} , the Kirillov theory says that $\mathfrak{g}^*/\text{Ad}^*(G)$ and \widehat{G} are homeomorphic,

Received April 29, 2015.

2010 *Mathematics Subject Classification.* 22E25, 22E15, 22D10.

Key words and phrases. Nilpotent and solvable Lie groups, orbit method, unitary representations of locally compact Lie groups.

The author would like to extend his sincere appreciation to the Deanship of Scientific Research at King Saud University for its funding through Research group RG-1435-069.

where $\text{Ad}^*(G)$ denotes the coadjoint representation of G on the dual \mathfrak{g}^* of \mathfrak{g} . Hence, for this class of Lie groups, $\text{cor}(G)$ can be identified with a certain $\text{Ad}^*(G)$ -invariant subset of \mathfrak{g}^* . From [2], one introduces the cortex of \mathfrak{g}^* as

$$\text{Cor}(\mathfrak{g}^*) = \left\{ \ell = \lim_{m \rightarrow \infty} \text{Ad}_{s_m}^*(\ell_m), \text{ where } \{s_m\} \subset G \right. \\ \left. \text{and } \{\ell_m\} \subset \mathfrak{g}^* \text{ such that } \lim_{m \rightarrow \infty} \ell_m = 0 \right\}$$

and we have $\pi_\ell \in \text{cor}(G)$ if and only if $\ell \in \text{Cor}(\mathfrak{g}^*)$. Note that in the case of general Lie groups, the two definitions are not so easily related. Motivated by this situation, the authors in [3] define the cortex $C_V(G)$ of a representation of a locally compact group G on a finite-dimensional vector space V as the set of all $v \in V$ for which $G.v$ and $\{0\}$ cannot be Hausdorff-separated in the orbit-space V/G . They give a precise description of $C_V(G)$ in the case $G = \mathbb{R}$. Moreover, they consider the subset $IC_V(G)$ of V consisting of the common zeroes of all G -invariant polynomials P on V with $P(0) = 0$. Note that when G is a nilpotent Lie group, one has $IC_V(G) \subset C_V(G)$ and they show that $IC_V(G) = C_V(G)$ when G is a nilpotent Lie group of the form $G = \mathbb{R}^n \rtimes \mathbb{R}$ and $V = \mathfrak{g}^*$ the dual of the Lie algebra \mathfrak{g} . This fails for a general nilpotent Lie group, even in the case of two-step nilpotent Lie group (see [2]). In [7], the authors show that the cortex of a connected and simply connected nilpotent Lie group is a semi-algebraic set. In [5] one gives an explicit description of the cortex of certain class of exponential Lie algebras (using the results of parametrization in [1]).

Fixing the class of two-step nilpotent Lie algebras, we see that each coadjoint orbit is a flat (affine) symplectic manifold, however the cortex of that class of Lie algebras may not be flat and in this paper, we give a generalization of the example given in [2] p. 210. Our example consists of a family of $4d$ -dimensional two-step nilpotent Lie algebras $(\mathfrak{g}_d)_{d \geq 2}$ such that the cortex of each \mathfrak{g}_d^* is the zero set of a homogeneous polynomial of degree d in the complement \mathfrak{z}_d^\perp of the center \mathfrak{z}_d of \mathfrak{g}_d . Finally we give some remarks on the cortex of $\mathbb{R}^n \rtimes \mathbb{R}$.

The paper is organized as follows: The next section is a review of the mathematics and basic tools used throughout the rest of the text. In the third section, we focus on the class of two-step nilpotent Lie algebras \mathfrak{g} , and we give a refinement of Theorem 4.5 ([1] p. 548) by which we give a description of the algebra of G -invariant polynomials on \mathfrak{g}^* (G is the corresponding Lie group of \mathfrak{g}). Next we give an interesting example of a family of two-step nilpotent Lie algebras $(\mathfrak{g}_d)_{d \geq 2}$ for which the cortex of the dual \mathfrak{g}_d^* of each \mathfrak{g}_d is the zero set of homogeneous polynomials of degree d . In the final section, we consider the exponential nonnilpotent Lie group $G = \mathbb{R}^n \rtimes \mathbb{R}$ and we give a complete and explicit description of the cortex of the dual of its Lie algebra.

2. Background material and notations

If G is a locally compact group, Vershik and Karpushev [9] introduce the notion of cortex of G as the set of all unitary irreducible representations of

G that cannot be Hausdorff separated from the trivial representation. If G a Lie group with Lie algebra \mathfrak{g} , it's known that G acts on \mathfrak{g} by the adjoint action denoted by Ad and on \mathfrak{g}^* by the coadjoint action denoted by Ad^* . Following [3], we recall the following:

Definition 2.1. Let π be a continuous representation of a locally compact Lie group G on a finite-dimensional (real) space V we define

$$C_V(\pi) = \{v = \lim_{m \rightarrow \infty} \pi(s_m)v_m, \lim_{m \rightarrow \infty} v_m = 0, \{s_m\}_m \subset G\},$$

and the cortex of invariants of π as

$$IC_V(\pi) = \{v \in V : p(v) = p(0) \text{ for all } G\text{-invariant polynomials on } V\}.$$

In particular when G is a locally compact Lie group and π is the contra-gradient representation of G on the dual \mathfrak{g}^* of the Lie algebra \mathfrak{g} of G , one has:

Definition 2.2. We define the cortex of \mathfrak{g}^* as

$$\text{Cor}(\mathfrak{g}^*) = \left\{ \lim_{m \rightarrow \infty} \text{Ad}_{s_m}^*(\ell_m) \mid (s_m)_m \subset G, (\ell_m)_m \subset \mathfrak{g}^* \text{ with } \lim_{m \rightarrow \infty} \ell_m = 0 \right\},$$

and the cortex of invariants

$$\text{ICor}(\mathfrak{g}^*) = \{\ell \in \mathfrak{g}^* : p(\ell) = p(0), \text{ for all } G\text{-invariant polynomial } p \text{ on } \mathfrak{g}^*\}.$$

When G is a nilpotent connected and simply connected Lie group, Kirillov's theory establishes a bijection between $\mathfrak{g}^*/\text{Ad}^*(G)$ (the orbit space of the coadjoint representation of G on \mathfrak{g}^*) and \widehat{G} (the unitary dual of G). More precisely, associated to $\ell \in \mathfrak{g}^*$ is an irreducible representation π_ℓ of G , and π_f and π_ℓ ($f \in \mathfrak{g}^*$) are equivalent if and only if $f \in \text{Ad}^*(G)\ell$. The Kirillov correspondence is a homeomorphism provided that $\mathfrak{g}^*/\text{Ad}^*(G)$ is endowed with the quotient topology [4]. In that case, the unitary dual \widehat{G} of G can be parameterized via the orbit-method. More precisely, let $\ell \in \mathfrak{g}^*$ and \mathfrak{p}_ℓ be a Pukanszky polarization at ℓ , we define the representation $\pi_{\ell, \mathfrak{p}_\ell}$ by

$$\pi_{\ell, \mathfrak{p}_\ell} := \text{ind}_{\mathcal{P}_\ell}^G \chi_\ell,$$

where $\mathcal{P}_\ell = \exp \mathfrak{p}_\ell$ and χ_ℓ is the unitary character associated with \mathcal{P}_ℓ given by

$$\chi_\ell(\exp X) = e^{-i\langle \ell, X \rangle}, \quad X \in \mathfrak{p}_\ell.$$

Then:

Theorem 2.1 (A. A. Kirillov). *Let G be a simply connected nilpotent real Lie group with Lie algebra \mathfrak{g} . If $\ell \in \mathfrak{g}^*$, there exists a polarization $\mathfrak{p}(\ell)$ of \mathfrak{g} for ℓ such that the monomial representation $\pi_{\ell, \mathfrak{p}(\ell)} := \text{ind}_{\exp \mathfrak{p}(\ell)}^G \chi_\ell$ is irreducible and of trace class. If ℓ' is an element of \mathfrak{g}^* which belongs to the coadjoint orbit of ℓ and $\mathfrak{p}_{\ell'}$ is a polarization of \mathfrak{g} for ℓ' , then the monomial representations $\pi_{\ell, \mathfrak{p}_\ell}$ and $\pi_{\ell', \mathfrak{p}_{\ell'}}$ are unitarily equivalent. Conversely, if \mathfrak{h} and \mathfrak{h}' are polarizations of \mathfrak{g} for $\ell \in \mathfrak{g}^*$ and $\ell' \in \mathfrak{g}^*$ respectively such that the monomial representations $\pi_{\ell, \mathfrak{h}}$ and $\pi_{\ell', \mathfrak{h}'}$ of G are unitarily equivalent, then*

ℓ and ℓ' belong to the same coadjoint orbit of G in \mathfrak{g}^* . Finally, for each irreducible unitary representation π of G , there exists a unique coadjoint orbit \mathcal{O} of G in \mathfrak{g}^* such that for any linear form ℓ and each polarization \mathfrak{h} of \mathfrak{g} for ℓ , the representations π and $\text{ind}_{\exp \mathfrak{h}}^G \chi_\ell$ are unitarily equivalent. Any irreducible unitary representation of G is strongly trace class. Moreover the mapping

$$\begin{aligned} K : \mathfrak{g}^*/\text{Ad}^*(G) &\longrightarrow \widehat{G} \\ \mathcal{O}_\ell &\longmapsto [\pi_{\ell, \mathfrak{p}(\ell)}] \end{aligned}$$

is a homeomorphism (the Kirillov correspondence).

The above Kirillov’s result was generalized immediately to the class known as exponential solvable Lie groups, the Kirillov correspondence is still a bijection. For more details, see [8]. With this in mind, we see that if G is an exponential Lie group, then $\pi := \pi_{\ell, \mathfrak{p}_\ell} \in \text{cor}(G)$ (cortex of G) if and only if $\ell \in \text{Cor}(\mathfrak{g}^*)$. However if G is exponential nonnilpotent, $\text{ICor}(\mathfrak{g}^*)$ may not be defined.

Throughout, G will always denote a connected and simply connected Lie group with (real) Lie algebra \mathfrak{g} . We denote by \mathfrak{z} the center of \mathfrak{g} (if it exists) and \mathfrak{g}^* denotes the dual of \mathfrak{g} . If $\ell \in \mathfrak{g}^*$, \mathcal{O}_ℓ denotes the coadjoint orbit of ℓ .

3. The two-step nilpotent Lie algebras

Let G be a connected and simply connected two-step nilpotent Lie group with Lie algebra \mathfrak{g} , then if $\mathcal{O}_\ell = \text{Ad}^*(G)\ell$, one has

$$\mathcal{O}_\ell = \{\ell\} + T_\ell \mathcal{O}_\ell,$$

and

$$T_\ell \mathcal{O}_\ell = \mathfrak{g}(\ell)^\perp,$$

where $T_\ell \mathcal{O}_\ell$ is the tangent space of \mathcal{O}_ℓ at ℓ , by which we see that the coadjoint orbits in two-step nilpotent Lie algebras are flat (and symplectic) manifolds. In [2], the authors show the following:

Proposition 3.1. *Let \mathfrak{g} be a nilpotent Lie algebra of class 2 (i.e., $[\mathfrak{g}, [\mathfrak{g}, \mathfrak{g}]] = 0$), and let $G = \exp \mathfrak{g}$ be the associated Lie group. Denote by ad^* the coadjoint representation of \mathfrak{g} on \mathfrak{g}^* . Let $f \in \mathfrak{g}^*$. Then the corresponding representation π_f of G belongs to $\text{cor}(G)$ if and only if f belongs to the closure of the subset $\{\text{ad}_X^*(\ell), X \in \mathfrak{g}, \ell \in \mathfrak{g}^*\}$ of \mathfrak{g}^* .*

From this we can conclude the following:

Corollary 3.2. *Let \mathfrak{g} is a two-step nilpotent Lie algebra. If $T_\ell \mathcal{O}_\ell$ denotes the tangent space to the coadjoint orbit \mathcal{O}_ℓ at ℓ , then the $\text{Cor}(\mathfrak{g}^*)$ is the closure in \mathfrak{g}^* of the set*

$$\bigcup_{\ell \in \mathfrak{g}^*} T_\ell \mathcal{O}_\ell = \bigcup_{\mathcal{O}_\ell \in \mathfrak{g}^*/\text{Ad}^*(G)} T\mathcal{O}_\ell,$$

where $T\mathcal{O}_\ell$ is the fiber tangent of \mathcal{O}_ℓ and $\mathfrak{g}^*/\text{Ad}^*(G)$ is the space of coadjoint orbits in \mathfrak{g}^* .

Proof. Indeed, for any $\ell \in \mathfrak{g}^*$, one has

$$\{ad_X^*(\ell); X \in \mathfrak{g}\} = T_\ell \mathcal{O}_\ell,$$

and hence with Proposition (3.1), the conclusion yields. □

Here we give a refinement of Theorem 4.5 ([1] p. 548).

Proposition 3.3. *Let G be a two-step nilpotent Lie group with Lie algebra \mathfrak{g} , choose a real Jordan–Hölder basis $\{X_j\}$. Let \mathcal{P} be the corresponding fine stratification of \mathfrak{g}^* , and let Ω be a layer belonging to \mathcal{P} . Then there is an explicit construction of an open set U in \mathfrak{g}^* and real-valued functions $p_1, p_2, \dots, p_d, q_1, q_2, \dots, q_d$ on U , such that U contains Ω , and such that for each coadjoint orbit \mathcal{O}_ℓ in Ω , $p_1|_{\mathcal{O}_\ell}, p_2|_{\mathcal{O}_\ell}, \dots, p_d|_{\mathcal{O}_\ell}, q_1|_{\mathcal{O}_\ell}, q_2|_{\mathcal{O}_\ell}, \dots, q_d|_{\mathcal{O}_\ell}$ are real-valued, global canonical coordinates for \mathcal{O}_ℓ . Moreover, for each $1 \leq j \leq n$, $0 \leq u \leq d$, there are rational functions $\alpha_{j,u}$ and $\beta_{j,u}$ such that for each $1 \leq j \leq n$ and $\ell \in \Omega$ one has*

$$\ell_j := \ell(X_j) = \sum_{u: j_u \leq j} \alpha_{j,u}(\ell) p_u + \sum_{r=1}^d \beta_{j,u}(\ell) q_u.$$

Proof. Recall that the construction of p_r, q_r depends on the flag

$$(\mathfrak{g}_j = \text{span}\{X_1, \dots, X_j\})_{1 \leq j \leq n}.$$

More precisely if $j_t = \min\{j_r, 1 \leq r \leq d\}$, then:

$$p_1^{(1)} = \ell_{i_t}, \quad q_1^{(1)} = \frac{\ell_{j_t}}{\ell[X_{j_t}, X_{i_t}]}.$$

Now suppose we have built $p_1^{(m)}, \dots, p_k^{(m)}, \dots, q_1^{(m)}, \dots, q_k^{(m)}$, then for \mathfrak{g}_{m+1} one has either $m + 1 \notin \mathbf{e}$ and in this case $p_r^{(m+1)} = p_r^{(m)}, q_r^{(m+1)} = q_r^{(m)}$ or $m + 1 = j_{k+1} \in \mathbf{e}$ and in this case

$$q_r^{(m+1)}(\ell) = q_r^{(m)}(\exp -qX_{m+1}\ell) = q_r^{(m)}(\ell) - q\{x_{m+1}, q_r^{(m)}\},$$

and

$$p_r^{(m+1)}(\ell) = p_r^{(m)}(\exp -qX_{m+1}\ell) = p_r^{(m)}(\ell) - q\{x_{m+1}, p_r^{(m)}\},$$

with $q = \frac{y}{\ell[X_{m+1}, y]}$, where y is a G_m -invariant and non- G_{m+1} -invariant polynomial function such that $\{x_{m+1}, y\}$ is nonvanishing on Ω (here $G_j = \exp \mathfrak{g}_j$). □

Corollary 3.4. *Let $\mathbf{e} = \{e_1 < \dots < e_{2d}\}$ be the set of jump indices corresponding to the minimal layer in \mathfrak{g}^* . Let F be the cross-section mapping associated with the minimal layer Ω then $F(\ell) = (F_1(\ell), \dots, F_n(\ell))$ and let $\mathbf{e} = \{e_1 < \dots < e_{2d}\}$ be the corresponding jump indices then*

$$F_k(\ell) = \begin{cases} \ell_k, & \text{if } k = 1, \dots, e_1 - 1; \\ 0, & \text{if } k \in \mathbf{e}; \\ \ell_k + \sum_{j: e_j \leq k-1} a_j(\ell) \ell_{e_j}, & \text{if } k \notin \mathbf{e}, k \geq e_1, \end{cases}$$

where each of $a_1(\ell), \dots, a_{k-1}(\ell)$ is (nontrivial) a rational regular function on the minimal layer depending only upon $\ell^0 = \ell|_{\mathfrak{z}}$ (\mathfrak{z} is the center of \mathfrak{g}).

Proof. For each layer in \mathfrak{g}^* , the mapping $(\ell_{e_1}, \dots, \ell_{e_{2d}}) \mapsto (p_i(\ell), q_i(\ell))_{1 \leq i \leq d}$ is a rational diffeomorphism whose inverse is also rational on any layer, then we consider the minimal layer and by Proposition 3.3, we can write

$$p_i(\ell) = \sum_j u_j(\ell_1, \dots, \ell_p)\ell_{e_j}, \quad q_i(\ell) = \sum_j v_j(\ell_1, \dots, \ell_p)\ell_{e_j}, \quad i = 1, \dots, d,$$

where u_j and v_j are rational regular functions on the minimal layer. Then after substituting each of the functions $(p_i, q_i)_i$ by the above expressions in the coordinate functions $(\ell_k)_{k \notin \mathbf{e}}$ we obtain the invariant functions of \mathfrak{g}^* and this ends the proof. \square

Corollary 3.5. *If \mathfrak{g} is a two-step nilpotent Lie algebra and \mathfrak{g}^* denotes its dual, then*

$$\text{ICor}(\mathfrak{g}^*) = \{\ell \in \mathfrak{g}^* : \ell(Z) = 0 \quad \forall Z \in \mathfrak{z}\}.$$

Proof. The nontrivial coordinates of the cross-section mapping

$$(F_k(\ell))_{k \geq p, k \notin \mathbf{e}}$$

associated with the minimal layer can be written as

$$F_k(\ell) = \frac{B(\ell^0)\ell_k + A_k(\ell^0)}{B(\ell^0)}, \quad k \notin \mathbf{e}, k > p := e_1,$$

where each of $B(\ell^0)$ and $B(\ell^0)\ell_k + A_k(\ell^0), (k \geq p, k \notin \mathbf{e})$ is a nontrivial G -invariant polynomial on \mathfrak{g}^* , with $\ell^0 = \ell|_{\mathfrak{z}}$. Note that these polynomials are homogeneous and for each $k \geq p, k \notin \mathbf{e}$, one has

$$\deg(B(\ell^0)\ell_k + A_k(\ell^0)) = \deg(B(\ell^0)) + 1.$$

Finally the ring $\text{Pol}(\mathfrak{g}^*)^G$ of G -invariant polynomials is spanned by the polynomials

$$\ell_1, \dots, \ell_p, B(\ell^0), (B(\ell^0)\ell_k + A_k(\ell^0))_{k \geq p, k \notin \mathbf{e}},$$

and this ends the proof. \square

3.1. Main example. In [2], one introduces an interesting example of 8-dimensional two-step nilpotent Lie algebra \mathfrak{g} so that the corresponding cortex in \mathfrak{g}^* is a projective algebraic set given by a quadric and such that $\text{Cor}(\mathfrak{g}^*) \subsetneq \text{ICor}(\mathfrak{g}^*)$. Here we give a generalization of that example. Let $d \in \mathbb{N}$ with $d \geq 2$ and let \mathfrak{g}_d be the Lie algebra with basis

$$(Z_1, \dots, Z_d, Y_1, Y_2, \dots, Y_{2d-1}, Y_{2d}, X_1, \dots, X_d),$$

and nontrivial brackets

$$\begin{aligned} [X_i, Y_{2i-1}] &= Z_1, & i &= 1, \dots, d, \\ [X_k, Y_{2k}] &= Z_{k+1}, & k &= 1, \dots, d-1, \\ [X_d, Y_{2d}] &= Z_2 + \dots + Z_d. \end{aligned}$$

Let's denote the center of \mathfrak{g}_d by $\mathfrak{z}_d = \text{span}\{Z_1, \dots, Z_d\}$ and G_d the corresponding connected and simply connected Lie group.

Proposition 3.6. *For each Lie algebra \mathfrak{g}_d ($d \geq 2$), one has:*

(i) *The minimal layer in \mathfrak{g}_d^* is given by*

$$\Omega_d = \{\ell \in \mathfrak{g}_d^* : \ell(Z_1) \neq 0\}.$$

(ii) *The coadjoint orbits in Ω_d are 2d-dimensional and if*

$$\begin{aligned} \ell &= \sum_{k=1}^d (\lambda_k Z_k^* + \beta_k X_k^*) + \sum_{k=1}^{2d} \gamma_k Y_k^* \in \Omega_d, \\ \xi &= ((z_i)_{1 \leq i \leq d}, (y_j)_{1 \leq j \leq 2d}, (x_k)_{1 \leq k \leq d}) \in Gl, \end{aligned}$$

then

$$\xi = \begin{cases} z_k = \lambda_k, & \text{if } k = 1, \dots, d; \\ y_{2k-1} = \gamma_{2k-1} + s_j \lambda_1, & \text{if } k = 1, \dots, d-1; \\ y_{2k} = \gamma_{2k} + s_j \lambda_{k+1}, & \text{if } k = 1, \dots, d-1; \\ y_{2d-1} = \gamma_{2d-1} + s_d \lambda_1; \\ y_{2d} = \gamma_{2d} + s_d (\lambda_2 + \dots + \lambda_d); \\ x_k = \beta_k + t_k & \text{if } k = 1, \dots, d. \end{cases}$$

(iii) *The algebra of G-invariant polynomials is*

$$\text{Pol}(\mathfrak{g}_d^*)^{G_d} = \mathbb{R}[z_1, \dots, z_d, z_1 y_2 - z_2 y_1, \dots, z_1 y_{2d-2} - z_{d-1} y_{2d-3}, \\ z_1 y_{2d} - (z_2 + \dots + z_d) y_{2d-1}].$$

Proof. Let $\mathcal{B}_d = (U_1, \dots, U_{4d})$ be the Jordan–Hölder basis defined by

$$U_i = \begin{cases} Z_i, & \text{if } 1 \leq i \leq d, \\ Y_{i-d}, & \text{if } d+1 \leq i \leq 3d; \\ X_{i-3d}, & \text{if } 3d+1 \leq i \leq 4d. \end{cases}$$

Using the methods of [1], we can see that the minimal layer in \mathfrak{g}_d^* is

$$\Omega_d = \{\ell \in \mathfrak{g}_d^* : \ell(U_1) = \ell(Z_1) \neq 0\},$$

which corresponds to the set of jump indices $\mathbf{e}_d = \mathbf{i}_d \cup \mathbf{j}_d$ with

$$\begin{aligned} \mathbf{i}_d &= \{d+1 < d+3 < \dots < 3d-1\}, \\ \mathbf{j}_d &= \{3d+1, 3d+2, \dots, 4d\}. \end{aligned}$$

Then by using the methods of [1] (the parametrization of coadjoint orbits) we can deduce the results of (ii) and (iii). □

Remark 3.1.

- (i) If Ω_d is the minimal layer given as above, then the canonical coordinates on Ω_d (see [1]) are given by

$$p_i(\ell) = x_i, q_i(\ell) = \frac{y_{2i-1}}{z_1}, \quad i = 1, \dots, d.$$

- (ii) The cross-section Σ_d is given by

$$\Sigma_d = \left(\sum_{k \notin \mathbf{e}} \mathbb{R}U_k^* \right) \cap \Omega_d = \left(\sum_{k=1}^d \mathbb{R}Z_k^* + \mathbb{R}Y_{2k}^* \right) \cap \Omega_d.$$

- (iii) The cross-section mapping $F_d : \Omega_d \rightarrow \Sigma_d$ is as follows

$$F_d(z_i, y_j, x_k) = \sum_{i=1}^d z_i Z_i^* + \sum_{i=1}^{d-1} \left(y_{2i} - \frac{z_{i+1}}{z_i} y_{2i-1} \right) Y_{2i}^* + \left(y_{2d} - \frac{z_2 + \dots + z_d}{z_1} \right) Y_{2d}^*,$$

where $(Z_1^*, \dots, Z_d^*, Y_1^*, \dots, Y_{2d}^*, X_1^*, \dots, X_d^*)$ is the dual basis of \mathcal{B} .

Proposition 3.7. *Let's denote $\ell = \sum_{i=1}^d (z_i Z_i^* + x_i X_i^*) + \sum_{j=1}^{2d} y_j Y_j^* \in \mathfrak{g}^*$ by $\ell = (z_i, y_j, x_k)$, where $(Z_1^*, \dots, Z_d^*, Y_1^*, \dots, Y_{2d}^*, X_1^*, \dots, X_d^*)$ is the dual basis in \mathfrak{g}_d^* . Then the cortex of \mathfrak{g}_d^* is the projective algebraic set given by*

$$\text{Cor}(\mathfrak{g}_d^*) = \left\{ \ell = (z_i, y_j, x_k) : z_1 = \dots = z_d = y_{2d-1} \left(\sum_{i=1}^{d-1} y_{2i} \prod_{j=1, j \neq i}^{d-1} y_{2j-1} \right) - y_{2d} \prod_{j=1}^{d-1} y_{2j-1} = 0 \right\}.$$

Proof. Note that since Ω_d is dense in \mathfrak{g}_d^* (Zariski open subset in \mathfrak{g}_d^*) then

$$\text{Cor}(\mathfrak{g}_d^*) = \left\{ \lim_m \text{Ad}_{\exp X_m}^* \ell_m, (\ell_m) \in \mathfrak{g}_d^*, (\ell_m)_m \in \Omega_d, \text{ and } \lim_m \ell_m = 0 \right\}.$$

On the other hand if $\mathcal{O}_\ell = G\ell$, then the tangent space $T_\ell \mathcal{O}_\ell$ at ℓ is

$$T_\ell \mathcal{O}_\ell = \{ ad_X^*(\ell), \ell \in \Omega_d, X \in \text{Vect}\{Y_{2k-1}, X_k, 1 \leq k \leq d\} \}.$$

Now if $\ell = (\lambda_i, \gamma_j, \beta_k) \in \Omega_d$ and $\xi \in T_\ell \mathcal{O}_\ell$, with

$$\xi = \sum_{i=1}^d (z_i Z_i^* + x_i X_i^*) + \sum_{j=1}^{2d} y_i Y_i^*,$$

then

$$\begin{cases} z_i = 0, & \text{if } i = 1, \dots, d; \\ y_{2j-1} = s_j \lambda_1, & \text{if } j = 1, \dots, d-1; \\ y_{2j} = s_j \lambda_{j+1}, & \text{if } j = 1, \dots, d-1; \\ y_{2d-1} = s_d \lambda_1; \\ y_{2d} = s_d (\lambda_2 + \dots + \lambda_d); \\ x_k = t_k, & \text{if } k = 1, \dots, d. \end{cases}$$

From which we can see that $\xi = (z_i, y_j, x_k) \in T_\ell \mathcal{O}$ if and only if

$$\begin{cases} z_i = 0, & \text{if } i = 1, \dots, d; \\ y_{2j} = y_{2j-1} \frac{\lambda_{j+1}}{\lambda_1}, & \text{if } j = 1, \dots, d-1; \\ y_{2d} = y_{2d-1} \frac{\lambda_2 + \dots + \lambda_d}{z_1}. \end{cases}$$

with y_{2j-1}, x_j are free variables in \mathbb{R} ($j = 1, \dots, d$). Then we see that a.e. $\xi \in T_\ell \mathcal{O}$ satisfies

$$\frac{y_{2d}}{y_{2d-1}} = \sum_{j=1}^{d-1} \frac{y_{2j}}{y_{2j-1}},$$

and hence

$$\text{Cor}(\mathfrak{g}_d^*) = \left\{ \ell = (z_i, y_j, x_k) \in \mathfrak{g}_d^* : z_i = 0, \right. \\ \left. y_{2d-1} \left(\sum_{i=1}^{d-1} y_{2i} \prod_{j=1, j \neq i}^{d-1} y_{2j-1} \right) - y_{2d} \prod_{j=1}^{d-1} y_{2j-1} = 0 \right\}. \quad \square$$

Corollary 3.8. *For each integer $d \geq 2$ if \mathfrak{z}_d denotes the center of the Lie algebra \mathfrak{g}_d , then*

$$\text{Cor}(\mathfrak{g}_d^*) \subsetneq \text{ICor}(\mathfrak{g}_d^*) = \mathfrak{z}_d^\perp.$$

Proof. The ring of G -invariant polynomials on \mathfrak{g}^* is given by

$$\text{Pol}(\mathfrak{g}_d^*)^{G_d} = \mathbb{R} \left[z_1, \dots, z_d, (z_1 y_{2i} - z_{i+1} y_{2i-1})_{1 \leq i \leq d-1}, z_1 y_{2d} - (z_2 + \dots + z_d) y_{2d-1} \right],$$

where G_d is the connected and simply connected (nilpotent) Lie group corresponding to \mathfrak{g}_d . Thus

$$\{\ell \in \mathfrak{g}_d^* : P(\ell) = P(0), \forall P \in \text{Pol}(\mathfrak{g}_d^*)^{G_d}\} = \mathfrak{z}_d^\perp,$$

where

$$\mathfrak{z}_d^\perp = \{\ell \in \mathfrak{g}_d^* : \ell(Z) = 0, \forall Z \in \mathfrak{z}_d\}.$$

hence with Proposition 3.7, we conclude that

$$\text{Cor}(\mathfrak{g}_d^*) \subsetneq \text{ICor}(\mathfrak{g}_d^*) = \mathfrak{z}_d^\perp. \quad \square$$

4. The Lie group $\mathbb{R}^n \rtimes \mathbb{R}$

In [3], in there is a study of the cortex of the nilpotent Lie group

$$G = \mathbb{R}^n \rtimes \mathbb{R},$$

the authors show that

$$\begin{aligned} \text{Cor}(\mathfrak{g}^*) &= \text{ICor}(\mathfrak{g}^*) \\ &= \{\ell \in \mathfrak{g}^* : P(\ell) = P(0), P \text{ is } G \text{ invariant polynomial on } \mathfrak{g}^*\}. \end{aligned}$$

The definition of $\text{ICor}(\mathfrak{g}^*)$ may not exist if G is not nilpotent but we can define G -invariant (or semi-invariant) functions. Let's consider the following example:

Example 4.1. Let (X_1, X_2, A) be a basis in \mathfrak{g} with

$$[A, X_1,] = X_1, [A, X_2] = -2X_2.$$

Let's identify \mathfrak{g}^* with \mathbb{R}^3 under the dual basis (X_1^*, X_2^*, A^*) , and denote $x = (x_1, x_2, a) \in \mathfrak{g}^*$, then the minimal layer is

$$\Omega = \{\ell = (\ell_1, \ell_2, a) \in \mathfrak{g}^* : \ell_1 \neq 0\}.$$

If $\ell = (\ell_1, \ell_2, a) \in \mathfrak{g}^*$, then the coadjoint orbit of ℓ is given by

$$\mathcal{O}_\ell = \{x \in \mathfrak{g}^* : x = (\ell_1 e^t, \ell_2 e^{-2t}, a + s), t, s \in \mathbb{R}\},$$

that is,

$$\mathcal{O}_\ell = \{x = (x_1, x_2, x_3) \in \mathfrak{g}^* : \text{sign}(x_1) = \text{sign}(\ell_1), x_1^2 x_2 = \ell_1^2 \ell_2, x_3 \in \mathbb{R}\}.$$

We can check that the cortex of \mathfrak{g}^* is given by

$$\text{Cor}(\mathfrak{g}^*) = \{\ell = (\ell_1, \ell_2, \ell_3) \in \mathfrak{g}^* : \ell_1 \ell_2 = 0\}.$$

On other hand, the cross-section mapping is as follows

$$F : \Omega \rightarrow \Omega, \quad \ell \mapsto \left(\text{sign}(\ell_1) = \frac{\ell_1}{|\ell_1|}, \ell_1^2 \ell_2, 0 \right),$$

from which we see the existence of G -invariant polynomial $p(x) = x_1^2 x_2$ and we see that

$$\text{Cor}(\mathfrak{g}^*) = \{\ell \in \mathfrak{g}^* : p(\ell) = 0\}.$$

In this example if we let $[A, X_1] = X_1, [A, X_2] = -\sqrt{2}X_2$ then there are no G -invariant polynomials on \mathfrak{g}^* , however the function $x_1^{\sqrt{2}} x_2$ is G -invariant and the cortex is still the same. This example can be generalized. To this end, if $\mathfrak{g} = \mathbb{R}^n \oplus \mathbb{R}A$, we denote $sp(adA) = \{\lambda_1, \dots, \lambda_n\}$ the set of eigenvalues of adA , and for $\lambda \in sp(adA)$, we set.

$$E_\lambda = \bigcup_{m \in \mathbb{N}} \ker(adA - \lambda)^m,$$

and

$$E^+ = \bigcup_{\lambda \in sp(adA), \Re(\lambda) > 0} E_\lambda,$$

$$E^- = \bigcup_{\lambda \in sp(adA), \Re(\lambda) < 0} E_\lambda.$$

Then we have the following:

Proposition 4.2. *Let $G = \mathbb{R}^n \rtimes \mathbb{R}$ be the Lie group whose Lie algebra $\mathfrak{g} = \mathbb{R}^n \oplus \mathbb{R}A$. Suppose that adA is diagonalizable, and let $sp(adA) = \{\lambda_1, \dots, \lambda_n\}$ denote the set of eigenvalues of adA .*

(a) *If $\{\Re(\lambda_j)\}_{1 \leq j \leq n} \subset (0, \infty)$ or $\{\Re(\lambda_j)\}_{1 \leq j \leq n} \subset (-\infty, 0)$ then*

$$\text{Cor}(\mathfrak{g}^*) = \mathfrak{g}^*.$$

(b) *If $\prod_{j=1}^n \Re(\lambda_j) < 0$. Then the cortex of \mathfrak{g}^* is the union of two vector spaces. More precisely*

$$\text{Cor}(\mathfrak{g}^*) = (V^+ + \mathbb{R}A^*) \cup (V^- + \mathbb{R}A^*),$$

where

$$V^+ = (E^+)^*, \quad V^- = (E^-)^*.$$

Proof. If $sp(adA) = \{\lambda_1, \dots, \lambda_n\}$ denotes the set of eigenvalues of adA (restricted to \mathbb{R}^n). Then identifying \mathfrak{g} with \mathbb{R}^{n+1} (respectively \mathbb{C}^{n+1} if some of the eigenvalues of adA are nonreal), the coadjoint orbit of any $\ell \in \mathfrak{g}^*$ is parameterized as follows

$$\mathcal{O}_\ell = \{(\ell_1 e^{\lambda_1 t}, \dots, \ell_n e^{\lambda_n t}, \ell_{n+1} + s), \quad t, s \in \mathbb{R}\}.$$

Since $\Re(\lambda_j) \neq 0, j = 1, \dots, n$, then for any $(\alpha_1, \dots, \alpha_n) \in \mathbb{C}^n$ the linear system

$$\begin{pmatrix} e^{\lambda_1 t} & 0 & \dots \\ 0 & \ddots & 0 \\ \dots & 0 & e^{\lambda_n t} \end{pmatrix} \begin{pmatrix} \ell_1 \\ \vdots \\ \ell_n \end{pmatrix} = \begin{pmatrix} \alpha_1 \\ \vdots \\ \alpha_n \end{pmatrix},$$

has a unique solution $(\ell_1, \dots, \ell_n)^\top$ with

$$\|(\ell_1, \dots, \ell_n)^\top\| = \|(e^{-\lambda_1 t} \alpha_1, \dots, e^{-\lambda_n t} \alpha_n)^\top\|.$$

(a) If $\{\Re(\lambda_j)\}_{1 \leq j \leq n} \subset (0, \infty)$ or $\{\Re(\lambda_j)\}_{1 \leq j \leq n} \subset (-\infty, 0)$ then for any $(\alpha_1, \dots, \alpha_n, \beta) \in \mathfrak{g}^*$ it exists $\{x^{(m)} = x(t_m, \ell^{(m)})\}_m \in \mathfrak{g}^*$ with $\{\ell^{(m)}\}_m \subset \Omega$ and $\lim_{m: \Re(\lambda_1) t_m \rightarrow \infty} \ell^{(m)} = 0$ such that

$$\lim_{m: \Re(\lambda_1) t_m \rightarrow \infty} x^{(m)} = (\alpha_1, \dots, \alpha_n, \beta),$$

and hence the cortex is all of \mathfrak{g}^* .

(b) In that case, let's rearrange the basis (X_1, \dots, X_n) in \mathbb{C}^n such that the matrix of adA in this basis is $\text{diag}(\lambda_1, \dots, \lambda_{k_0}, \lambda_{k_0+1}, \dots, \lambda_n)$ with

$$\Re(\lambda_1) > 0, \dots, \Re(\lambda_{k_0}) > 0, \Re(\lambda_{k_0+1}) < 0, \dots, \Re(\lambda_n) < 0,$$

then for any $x = (x_1, \dots, x_n, x_{n+1}) \in \mathcal{O}_\ell$ one has

$$\begin{cases} |\ell_1|^{\lambda_k} |x_k|^{\lambda_1} = |\ell_k|^{\lambda_1} |x_1|^{\lambda_k}, & k \leq k_0, \\ |x_k|^{\lambda_1} |x_1|^{-\lambda_k} = |\ell_k|^{\lambda_1} |\ell_1|^{-\lambda_k}, & k \geq k_0 + 1. \end{cases}$$

Hence one has

$$\begin{aligned} \text{Cor}(\mathfrak{g}^*) &= \{\ell \in \mathfrak{g}^* : \ell = (\ell_1, \dots, \ell_{k_0}, 0, \dots, 0, \ell_{n+1})\} \\ &\cup \{\ell \in \mathfrak{g}^* : \ell = (0, \dots, 0, \ell_{k_0+1}, \dots, \ell_n, \ell_{n+1})\}. \quad \square \end{aligned}$$

Remark 4.1. Let $\mathfrak{g} = \mathbb{R}^n \oplus \mathbb{R}A$ be a real Lie algebra. Suppose that there exists a basis (X_1, \dots, X_n) in \mathbb{R}^n such that

$$[A, X_j] = m_j X_j, \quad j = 1, \dots, n,$$

with $\{m_1, \dots, m_n\} \subset \mathbb{R}^\times$.

- (a) If $\{\frac{m_k}{m_1}\}_{1 \leq k \leq n} \in \mathbb{N}$, then any generic coadjoint orbit of ℓ ($\ell_1 \neq 0$) is given by

$$\mathcal{O}_\ell = \left\{ x = (x_1, \dots, x_n, x_{n+1}) \in \mathfrak{g}^* : x_1 \ell_1 > 0, x_k = \frac{\ell_k}{\ell_1^{\frac{m_k}{m_1}}} x_1^{\frac{m_k}{m_1}}, \right. \\ \left. k = 2, \dots, n, x_{n+1} \in \mathbb{R} \right\},$$

and hence, it is an open semi-algebraic subset in \mathfrak{g}^* .

- (b) Now suppose that $\{m_1, \dots, m_n\} \subset \mathbb{Z}^\times$ with $\prod_{j=1}^n m_j < 0$. We can assume the existence of a basis (X_1, \dots, X_n) in \mathbb{R}^n so that with respect to this basis the matrix of adA is

$$adA = \text{diag}(m_1, \dots, m_{k_0}, m_{k_0+1}, \dots, m_n)$$

with $m_1 > 0, \dots, m_{k_0} > 0, m_{k_0+1} < 0, \dots, m_n < 0$. Then for any $x = (x_1, \dots, x_n, x_{n+1}) \in \mathcal{O}_\ell$ (with $\ell_1 \neq 0$) one has

$$\begin{cases} x_1 \ell_1 > 0, \\ \ell_1^{m_j} x_j^{m_1} = \ell_j^{m_1} x_1^{m_j}, & j = 2, \dots, k_0, \\ x_1^{-m_j} x_j^{m_1} = \ell_1^{-m_j} \ell_j^{m_1}, & j = k_0 + 1, \dots, n. \end{cases}$$

On other hand, the polynomials

$$p_{i,j}(\ell) = \ell_i^{-m_j} \ell_j^{m_i}, \quad i = 1, \dots, k_0, \quad j = k_0 + 1, \dots, n$$

are G -invariant on \mathfrak{g}^* and the cortex is the union of two vector spaces given by:

$$\text{Cor}(\mathfrak{g}^*) = \{\ell \in \mathfrak{g}^* : p_{i,j}(\ell) = 0 \quad \forall 1 \leq i \leq k_0, k_0 + 1 \leq j \leq n\}.$$

Corollary 4.3. Let $\mathfrak{g} = \mathbb{R}^n \oplus \mathbb{R}A$ be a real Lie algebra. Let's denote $sp(adA) = \{\lambda_1, \dots, \lambda_n\} \subset \mathbb{C}$ the set of eigenvalues of adA . If

$$\{\Re(\lambda_j)\}_{1 \leq j \leq n} \subset (0, \infty) \quad \text{or} \quad \{\Re(\lambda_j)\}_{1 \leq j \leq n} \subset (-\infty, 0),$$

then

$$\text{Cor}(\mathfrak{g}^*) = \mathfrak{g}^*.$$

Proof. First let's suppose that the real endomorphism adA has a single eigenvalue $\lambda \in \mathbb{C} \setminus i\mathbb{R}$. According to λ is real or complex, we can suppose the existence of a basis in \mathbb{R}^n (resp. in \mathbb{C}^n) such that the matrix of adA is written in a Jordan block form:

$$adA = J_\lambda = \begin{pmatrix} \lambda & 1 & 0 & \dots \\ 0 & \ddots & \ddots & \ddots \\ \vdots & \ddots & \ddots & 1 \\ 0 & \dots & 0 & \lambda \end{pmatrix}$$

The coadjoint orbit of $\ell = (\ell_1, \dots, \ell_n, \ell_{n+1})$ is given by

$$\mathcal{O}_\ell = \left\{ \left(\left(x_k = e^{\lambda t} \sum_{j \geq 1, i+j=k} \frac{t^i}{i!} \ell_j \right)_{1 \leq k \leq n}, x_{n+1} = \ell_{n+1} + s \right), t, s \in \mathbb{R} \right\}.$$

Now let's remark that for any $\alpha \in \mathbb{R}^n$ (resp. in \mathbb{C}^n), since $\Re(\lambda) \neq 0$, the linear system

$$e^{\lambda t} \begin{pmatrix} 1 & 0 & \dots & 0 \\ t & 1 & 0 & \ddots \\ \vdots & \ddots & \ddots & 0 \\ \frac{t^{n-1}}{(n-1)!} & \dots & t & 1 \end{pmatrix} \begin{pmatrix} \ell_1 \\ \ell_2 \\ \vdots \\ \ell_n \end{pmatrix} = \begin{pmatrix} \alpha_1 \\ \alpha_2 \\ \vdots \\ \alpha_n \end{pmatrix}$$

has a unique solution and if we let

$$M(t) = \begin{pmatrix} 1 & 0 & \dots & 0 \\ t & 1 & 0 & \ddots \\ \vdots & \ddots & \ddots & 0 \\ \frac{t^{n-1}}{(n-1)!} & \dots & t & 1 \end{pmatrix},$$

then $M(t)$ is a unipotent matrix whose inverse $M^{-1}(t) = (p_{i,j}(t))_{1 \leq i,j \leq n}$ is also unipotent and all its entries $p_{i,j}(t)$ are polynomial functions in t , then

$$[\ell_1, \dots, \ell_n]^\perp = e^{-\lambda t} M^{-1}(t) [\alpha_1, \dots, \alpha_n]^\perp$$

and

$$\|[\ell_1, \dots, \ell_n]^\perp\| = e^{-\Re(\lambda)t} F(t),$$

where $F(t)^2$ is polynomial function in t and then

$$\lim_{\Re(\lambda)t \rightarrow \infty} e^{-\Re(\lambda)t} F(t) = 0,$$

For instance for any $\alpha = [\alpha_1, \dots, \alpha_n] \in \mathbb{R}^n$ if λ is real (resp. $\alpha \in \mathbb{C}^n$ if $\lambda \in \mathbb{C} \setminus i\mathbb{R}$) and $\{t_m\}_m \subset \mathbb{R}$ such that $\lim_{m \rightarrow \infty} t_m \Re(\lambda) = \infty$ it exists $\{\ell_1^{(m)}, \dots, \ell_n^{(m)}\}_m$ such that

$$\lim_{m \rightarrow \infty} e^{\lambda t_m} M(t_m) [\ell_1^{(m)}, \dots, \ell_n^{(m)}]^\perp = [\alpha_1, \dots, \alpha_n]^\perp$$

and

$$\lim_{m \rightarrow \infty} \ell_1^{(m)} = \dots = \lim_{m \rightarrow \infty} \ell_n^{(m)} = 0.$$

This shows that the cortex of \mathfrak{g}^* coincides with \mathfrak{g}^* . Finally if adA has more than one single eigenvalue, we can write $adA = \text{diag}(J_{\lambda_1}, \dots, J_{\lambda_k})$ where each J_λ is a Jordan block matrix. \square

Remark 4.2. Let $\mathfrak{g} = \mathbb{R}X_1 \oplus \mathbb{R}X_2 \oplus \mathbb{R}A$ with

$$[A, X_1] = X_1, [A, X_2] = X_2.$$

In this example the cortex of \mathfrak{g}^* is \mathfrak{g}^* . The cross-section mapping of the minimal layer is given by

$$F(\ell_1, \ell_2, \ell_3) = \left(\frac{\ell_1}{|\ell_1|}, \frac{\ell_2}{|\ell_1|}, 0 \right), \quad \ell_1 \neq 0.$$

On other hand, we remark that the rational function $r(\ell_1, \ell_2, \ell_3) = \frac{\ell_2}{\ell_1}$ is G -invariant on $\Omega = \{\ell = (\ell_1, \ell_2, \ell_3) \in \mathfrak{g}^* : \ell_1 \neq 0\}$ and

$$\text{Cor}(\mathfrak{g}^*) \supseteq \overline{\{\ell \in \Omega : r(\ell) = 0\}}.$$

Corollary 4.4. Let $\mathfrak{g} = \mathbb{R}^n \oplus \mathbb{R}A$, be a real Lie algebra. Let's denote $sp(adA) = \{\lambda_1, \dots, \lambda_n\} \subset \mathbb{C}$ the set of eigenvalues of adA . If $\prod_{j=1}^n \Re(\lambda_j) < 0$, then the cortex of \mathfrak{g}^* is the union of two vector spaces. More precisely, with the notations of Proposition 4.2, one has

$$\text{Cor}(\mathfrak{g}^*) = (V^+ + \mathbb{R}A^*) \cup (V^- + \mathbb{R}A^*).$$

Remark 4.3. Let $\mathfrak{g} = \mathbb{R}^n \oplus \mathbb{R}A$ be a real Lie algebra, and assume that all the eigenvalues of adA are purely imaginary. Let's denote $\mathfrak{h} = \mathbb{R}^n \oplus \mathbb{R}N$ where N is the nilpotent part in the Jordan decomposition of A , then by [3], one has

$$\text{Cor}(\mathfrak{g}^*) = \text{Cor}(\mathfrak{h}^*).$$

References

- [1] ARNAL, DIDIER; BEN AMMAR, MABROUK; CURREY, BRADLEY N.; DALI, BÉCHIR. Canonical coordinates for coadjoint orbits of completely solvable groups. *J. Lie Theory* **15** (2005), no. 2, 521–560. [MR2147442](#) (2006d:22009), [Zbl 1074.22003](#).
- [2] BEKKA, MOHAMMED E. B.; KANIUTH, EBERHARD. Irreducible representations of locally compact groups that cannot be Hausdorff separated from the identity representation. *J. Reine Angew. Math.* **385** (1988), 203–220. [MR0931221](#) (89j:22005), [Zbl 0634.22002](#), doi: [10.1515/crll.1988.385.203](#).
- [3] BOIDOL, J.; LUDWIG, J.; MÜLLER, D. On infinitely small orbits. *Studia Math.* **88** (1988), no. 1, 1–11. [MR0932001](#) (89f:22009), [Zbl 0662.22006](#).
- [4] BROWN, IAN D. Dual topology of a nilpotent Lie group. *Ann. Sci. École Norm. Sup.* (4) **6** (1973), 407–411. [MR0352326](#) (50 #4813), [Zbl 0284.57026](#).
- [5] DALI, BÉCHIR. On the cortex of a class of exponential Lie algebras. *J. Lie Theory* **22** (2012), no. 3, 845–867. [MR3012158](#), [Zbl 1247.22013](#).
- [6] FELL, J. M. G. Weak containment and induced representations of groups. *Canad. J. Math.* **14** (1962), 237–268. [MR0150241](#) (27 #242), [Zbl 0138.07301](#), doi: [10.4153/CJM-1962-016-6](#).

- [7] KÉDIM, IMED; HATEM, MEGDICHE. Sur le cortex d'un groupe de Lie nilpotent. *J. Math. Kyoto Univ.* **49** (2009), no. 1, 161–172. [MR2531133](#) (2010g:22015), [Zbl 1170.22004](#).
- [8] LEPTIN, HORST; LUDWIG, JEAN. Unitary representation theory of exponential Lie groups. de Gruyter Expositions in Mathematics, 18. *Walter de Gruyter & Co., Berlin*, 1994. x+200 pp. ISBN: 3-11-013938-3. [MR1307383](#) (96e:22001), [Zbl 0833.22012](#).
- [9] VERSHIK, A. M.; KARPUSHEV, S. I.. Cohomology of groups in unitary representations, neighborhood of the identity and conditionally positive definite functions. *Math. Sb.* **47** (1984), 513–526. [Zbl 0513.43009](#).

(Béchir Dali) DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCES OF BIZERTE, 7021 ZARZOUNA, BIZERTE, TUNISIA

Current address: King Saud University, college of science, Department of Mathematics, Riyadh, P.O Box 2455, Riyadh 11451, K.S.A.

bechir.dali@fss.rnu.tn

This paper is available via <http://nyjm.albany.edu/j/2015/21-55.html>.