

ON PARA-A-EISTEIN MANIFOLDS

B. B. Sinha and Ramesh Sharma

1. Introduction. An n -dimensional smooth manifold M_n with a tensor field f of type (1.1.), a vector field T , a l -form A and a Riemannian metric g is said to be an almost paracontact Riemannian manifold if [2]

$$(1.1) \quad (a) \quad f^2 = I - A \otimes T$$

$$(1.1) \quad (b) \quad g(fX, fY) = g(X, Y) - A(X)A(Y).$$

It can be shown that

$$(1.2) \quad A(T) = 1, \quad fT = 0, \quad Af = 0, \quad \text{rank}(f) = n - 1$$

whence it follows that

$$g(T, T) = 1, \quad 'f(X, Y) = 'f(Y, X)$$

' f being defined by ' $f(X, Y) = g(fX, Y)$ and X, Y standing for arbitrary vector fields on M_n .

If D be the Riemannian connexion induced on M_n by g such that [3]

$$(1.3) \quad (D_X A)Y + (D_Y A)X = 2'f(X, Y)$$

then the almost paracontact Riemannian manifold M_n is termed a paracontact Riemannian manifold. A paracontact Riemannian manifold M_n whose l -form A is closed, that is

$$(1.4) \quad (D_X A)Y - (D_Y A)X = 0$$

$$(1.5) \quad (D_X f)Y = 2A(X)A(Y)T - g(X, Y)T - A(Y)X$$

is called a normal paracontact Riemannian manifold. It is easy to show that the torsion tensor $N - (dA) \otimes T = 0$, where N is the Nijenhuis tensor of f .

2. Para-A-Einstein manifold. We define a para-A-Einstein manifold as a normal para contact Riemannian manifold whose Ricci tensor is given by [1]

$$(2.1) \quad \text{Ric}(X, Y) = a g(X, Y) + cA(X)A(Y)$$

where a and c are scalar functions. Obviously, we have

$$(2.2) \quad \text{Ric}(fX, Y) = \text{Ric}(X fY),$$

$$(2.3) \quad \text{Ric}(T, T) = a + c.$$

THEOREM 2.1. *The Riccian curvature of a para-A-Einstein manifold in the direction of T is equal to $-(n-1)$.*

Proof. From (1.3) and (1.4), we get

$$(2.4) \quad D_X T = fX.$$

From (1.5) and (2.4) we find the curvature tensor K satisfying

$$(2.5) \quad K(X, Y, T) = A(X)Y - A(Y)X.$$

Contracting (2.5), we get

$$(2.6) \quad \text{Ric}(Y, T) = -(n-1)A(Y).$$

Substituting T for Y in it we have the theorem.

THEOREM 2.2. *The functions a and c of the defining equation (2.1) are constants, provided $\text{tr. } f = 0$.*

Proof. Equation (2.3) and theorem (2.1) imply $a + c = 1 - n$. So we need only to show that a is constant. From (2.1), on contraction, we get $r = na + c$ which, on differentiation, yields

$$(2.7) \quad X_r = nX_a + X_c = (n-1)X_a,$$

where r is the scalar curvature. Again from (2.1) we have $R(X) = aX + cA(X)T$ which, on differentiation along Y , yields

$$(D_Y R)X = (Y a)X + (Y c)A(X)T + c(D_Y A)(X)T + cA(X)D_Y T.$$

The above equation assumes the form

$$(D_Y R)X = Y a + (Y c)A(X)T + c' f(X, Y)T + cA(X)fY$$

due to (2.4). Contracting in with respect to Y , we get $(\text{div } R)X = X_a + (T_c)A(X)$.

Using the identity $(\text{div } R)X = X_r/2$ and (2.7), we get

$$(2.8) \quad (n-3)X_a = 2(T_c)A(X).$$

Putting $X = T$ in it, we get

$$(n - 3)T_a = 2T_c = -2T_a$$

giving $T_a = 0$ and hence $T_c = 0$. Consequently (2.8) yields $X_a = 0$.

We now give a condition for a normal paracontact Riemannian manifold to be a para-A-Einstein manifold. With the help of (1.5) we can show for a normal paracontact Riemannian manifold that

$$(2.9) \quad \begin{aligned} K(X, Y, fZ) &= f(K(X, Y, Z)) + 2\{A(Y)'f(X, Z)T - A(X)'f(Y, Z)T \\ &\quad + A(Y)A(Z)fX - A(X)A(Z)fY\} - 'f(X, Z)Y + 'f(Y, Z)X \\ &\quad - g(Y, Z)fX + g(X, Z)fY. \end{aligned}$$

Contracting it with respect to X we find

$$(2.10) \quad \text{Ric}(Y, fZ) = (C'_1 \overline{K})(Y, Z) + (n-2)'f(Y, Z) + (C'_1 f)\{2A(Y)Z(Z) - g(Y, Z)\}$$

where C'_1 denotes contraction at the first slot and $\overline{K} \stackrel{\text{def}}{=} fK$.

$$(2.11) \quad (C'_1 \overline{K})(Y, Z) = (C'_1 \overline{K})(Z, Y),$$

From (2.10) and (2.11) it is obvious that

$$(2.12) \quad \text{Ric}(X, fY) = \text{Ric}(fX, Y).$$

THEOREM 2.3. *In order that a normal paracontact Riemannian manifold M_n may be a para-A-Einstein manifold it is necessary and sufficient that the symmetric tensors $C'_1 \overline{K}$ and $'f$ should be linearly dependent.*

Proof. The theorem follows in consequence of equations (2.10), (2.1), (1.1)a, (2.6) and Theorem 2.2.

THEOREM 2.4. *In a para-A-Einstein manifold, the symmetric $(0, 2)$ -tensor $C'_1 K$ is parallel along the vector field T .*

Proof. We have

$$(2.13) \quad (C'_1 K)(Y, Z) = (a - n + 2)'f(Y, Z)$$

due to (2.10) and (2.1). Differentiating it along T we have $(D_T C'_1 \overline{K})(Y, Z) = 0$ due to (1.5).

THEOREM 2.5. *For a para-A-Einstein manifold, the Lie-derivatives of the Ricci tensor and $C'_1 \overline{K}$ are given by*

$$(2.14) \quad L_T \text{Ric} = (2a/(a - n + 2))C'_1 K,$$

$$(2.15) \quad L_T(C'_1 \overline{K}) = 2(a - n + 2)(g - A \otimes A).$$

Proof. It is easy to show for a normal paracontact Riemannian manifold that $L_T A = 0$, $L_T f = 0$, $L_T g = 2'f$, $L_T'f = 2(g - A \otimes A)$

From these relations and Lie-derivation of the equations (2.1) and (2.13) along T , the theorem follows.

3. Examples. *Example (3.1)* From [3] it is known that a neighborhood of each point of a manifold of constant curvature is a normal paracontact Einstein manifold which is therefore a trivial example of a para-A-Einstein manifold with $c = 0$.

Example (3.2). Next, we give an example of non-trivial para-A-Einstein manifold. Consider a $2(m+1)$ -dimensional almost product and almost decomposable manifold $M_{2(m+1)}$ with structure tensor F such that the complementary distributions (having no common direction) may be of the same real dimension $m+1$. Suppose that $M_{2(m+1)}$ is of almost constant curvature [5]. Then its curvature tensor K is given by

$$(3.1) \quad {}^1\tilde{K}(\tilde{X}, \tilde{Y}, \tilde{Z}, \tilde{W}) = k[G(\tilde{X}, \tilde{W})G(\tilde{Y}, \tilde{Z}) - G(\tilde{X}, \tilde{Z})G(\tilde{Y}, \tilde{W}) \\ + {}^1F(\tilde{X}, \tilde{W}){}^1F(\tilde{Y}, \tilde{Z}) - {}^1F(\tilde{X}, \tilde{Z}) - {}^1F(\tilde{X}, \tilde{Z}){}^1F(\tilde{Y}, \tilde{W})]$$

where k is a constant, G is the metric tensor of $M_{2(m+1)}$ and $\tilde{X}, \tilde{Y}, \tilde{Z}, \tilde{W}$ are arbitrary vector fields on it. Let M_{2m+1} be a hypersurface in $M_{2(m+1)}$ and M_{2m+1} be a normal paracontact Riemannian manifold with structure tensors f, T, A, g . Then it can be shown [4] that

$$(3.2(a)) \quad HX = -X + A(X)T,$$

$$(3.2(b)) \quad C'H = -2n,$$

where H is the second fundamental tensor of type (1.1) of the hypersurface. Since the dimension of the hypersurface is odd we can adapt an orthonormal frame $e_1, \dots, e_m, fe_1, \dots, fe_m, T$ on M_{2m+1} , with respect to which C'_1f vanishes. Consequently $\text{div}T$ vanishes in view of (2.4) [3]. If B be the differential of the inclusion map $b: M_{2m+1} \rightarrow M_{2(m+1)}$, substituting BX, BY, BZ, BW for $\tilde{X}, \tilde{Y}, \tilde{Z}, \tilde{W}$ in (3.1), we have

$$(3.3) \quad {}^1\tilde{K}(BX, BY, BZ, BW) = [g(X, W)g(Y, Z) - g(X, Z)g(Y, W) \\ + {}^1f(X, W){}^1f(Y, Z) - {}^1f(X, Z){}^1f(Y, W)],$$

where we have used the transformation $FBX = BfX + A(X)N$, N being the unit normal vector field to the hypersurface.

Using Gauss characteristic equation in (3.3) and contractin, we get

$$\text{Ric}(Y, Z) + h(HZ) - (C'_1H)h(Y, Z) = k[A(Y)A(Z) + (2m-1)g(Y, Z)].$$

Using (3.2)(a), (3.2)(b) frequently in the above equation, we find

$$\text{Ric}(Y, Z) - (2m-1)\{g(Y, Z) - A(Y)A(Z)\} = k\{(2m-1)g(Y, Z) + A(Y)A(Z)\}$$

which implies $\text{Ric}(Y, Z) = (k+1)(2m-1)g(Y, Z) + (k+1-2m)A(Y)A(Z)$, showing that the normal paracontact Riemannian hypersurface of almost product and

almost decomposable manifold of almost constant curvature and whose complementary distributions have equal dimensions is a para-A-Einstein manifold.

It is notable that the scalar curvature of the enveloping manifold $M_{2(m+1)}$ is equal to $4n(n+1)k$.

REFERENCES

- [1] S. Sasaki, *Almost contact manifolds* (A lecture note), Mathematical institute, Tohoku University, 1965.
- [2] I. Sato, *On a structure similar to almost contact structures*, Tensor, N.S. **30** (1976), 219–224.
- [3] I. Sato, *On a structure similar to almost contact structures II*, Tensor, N.S. **31** (1977) 199–205.
- [4] B. B. Sinha, R. Sharma, *Infinitesimal variation of hypersurfaces of an almost product and almost decomposable manifold*, Indian J. Pure Appl. Math. **10** (8) (1979), 1009–1019.
- [5] K. Yano, *Differential Geometry on Complex and Almost Complex Spaces*, Pergamon Press, Oxford, 1965.

Department of Mathematics
Banaras Hindu University
Varanasi 221005, India.