

**A NOTE ON THE INDEPENDENCE NUMBER OF
AN IDENTICALLY SELF-DUAL PERFECT MATROID DESIGN**

Dănut Marcu

Abstract. Let E be a finite set and $M(E, r)$ an identically self-dual perfect matroid design on E , with hyperplane cardinality $c(M)$, and r as a rank function. If M is not the $r(E)$ -uniform matroid, we show that its independence number equals $c(M) - 1$.

Introduction. Our matroid-theoretic terminology is based essentially on the books of Rado [1] and Welsh [2]. Throughout, E will denote an n -set, $|S|$ denotes the cardinality of the set S , and $M := M(E, r)$ a *matroid* on E with r as a *rank function*. A subset $s \subseteq E$ is called *independent* if $r(s) = |s|$, a *basis* of M being a maximal independent subset of E . A subset $S \subseteq E$ is called *dependent* if $r(S) < |S|$, a *circuit* of M being a minimal dependent subset of E . It is well-known that a subset is independent iff does not contain a circuit. If $\{B_i\}$ is the set of bases of M , then $\{E - B_i\}$ is the set of bases of the *dual-matroid* M^* on E . If r^* denotes the rank function of M^* , then the following holds:

$$(1) \quad r(E) + r^*(E) = |E|.$$

A *cocircuit* of M is a circuit of M^* , and a hyperplane of M is a complement of a cocircuit in E . A matroid M on E is *identically self-dual* if $M = M^*$, i.e., every circuit is a cocircuit and vice versa.

The *span (closure)* \overline{S} of a subset $S \subseteq E$ is defined by

$$\overline{S} = \{e \in E : r(S \cup \{e\}) = r(S)\},$$

and S is called a *closed set* or *flat*, if $S = \overline{S}$. An *m-flat* is defined as a closed subset of E having rank m .

A *perfect matroid design* is a matroid in which each m -flat has a common cardinal, $1 \leq m \leq r(E)$; in particular, all of its hyperplanes have the same cardinality, which we denote by $c(M)$. If $k(M)$ is the minimum cardinality of a circuit

of the matroid M , it follows that $k(M) - 1$ is the maximum integer t such that every t -subset of E is independent; $k(M) - 1$ is called the *independence number* of M and will be denoted by $i(M)$.

The main result. Let M be a matroid on E and M^* its dual matroid. The family of independent sets of M^* is

$$(2) \quad \{S \subseteq E : r(E - S) = r(E)\}$$

The *k-uniform* matroid on E is defined by taking the family of bases to be $\{S \subseteq E : |S| = k\}$.

Let t be an integer greater than 1. A $t - (v, k, \lambda)$ *design* (V, β) is a v -set V and a system β of k -subsets of V ($k < v$), called blocks, such that every t -subset of V is contained in exactly λ blocks (repeated blocks are admissible in the system β). For example, $1 - (4, 2, 1)$, $3 - (8, 4, 1)$ and $5 - (12, 6, 1)$ are identically self dual perfect matroid designs. Moreover, no one of these is an $r(E)$ -uniform matroid. Thus, the following theorem seems to be of interest.

THEOREM. *Let M be an identically self dual perfect matroid design. If M is not the $r(E)$ -uniform matroid, then $i(M) = c(M) - 1$.*

Proof. From [2, Theorem 6, p. 212] we have

$$(3) \quad c(M) \leq n/2.$$

Since M is identically self-dual, then, by (1), we obtain

$$(4) \quad r(E) = n/2.$$

On the other hand, for each hyperplane H of M , the following holds:

$$(5) \quad r(E) - 1 = r(H) \leq |H| = c(M).$$

Hence, from (3) – (5), it follows that

$$(6) \quad n/2 - 1 \leq c(M) \leq n/2.$$

Suppose that $c(M) = n/2 - 1$. Thus, by (4) and (5), it follows that every hyperplane is an independent set, i.e., by (2), for each hyperplane H we have

$$(7) \quad r(E - H) = r(E).$$

Now, let $S \subseteq E$ be such that $|S| = r(E)$. If S is not an independent set, then S contains a circuit, i.e., since M is identically self dual, there exists a hyperplane H such that

$$(8) \quad E - H \subseteq S.$$

From (7) and (8), since $r(S) \leq r(E)$, we obtain $r(E) \leq r(S) \leq r(E)$, i.e., $r(S) = |S|$. Thus, S is an independent set. Moreover, because $|S| = r(E)$, it follows that S is a

basis of M . Consequently, every subset of cardinality $r(E)$ is a basis of M , i.e., M is the $r(E)$ -uniform matroid, contrary to the hypothesis. Hence, in (6), we must have

$$(9) \quad c(M) = n/2.$$

Let C be an arbitrary circuit of M (C is at the same time a cocircuit of M). By (9), we have $|C| = |E| - c(M) = c(M)$, i.e., all the circuits of M have cardinality $c(M)$, and therefore, every $[c(M) - 1]$ -subset of M is independent. Thus, $i(M) = c(M) - 1$.

COROLLARY 1. *Let S be a subset of E . If \bar{S} is a hyperline of M (M being like as in the Theorem), and S is not independent, then S is closed.*

Proof. Suppose that $S \neq \bar{S}$. Since $S \subseteq \bar{S}$, it follows that $|S| < |\bar{S}| - c(M)$, i.e., $|S| \leq c(M) - 1 = i(M)$. Thus (because every set contained in an independent set is independent) S is independent, contradicting the hypothesis, and the corollary is proved.

COROLLARY 2. *If S a subset of E such that $0 \leq r(S) \leq c(M) - 2$, then S is independent in M (M taken as in Theorem).*

Proof. Suppose that S is not independent, i.e., that there exists a circuit C such that $C \subseteq S$. Thus, $c(M) - 1 = r(C) \leq c(M) - 2$, which is absurd. Hence the corollary is proved.

REFERENCES

- [1] R. Randow, *Introduction to the Theory of Matroids*, Lecture Notes in Economics and Mathematical Systems, Springer-Verlag, 1975.
- [2] D. J. A. Welsh, *Matroid Theory*, Academic Prees, New York, 1976.

Fakulty of Mathematics
University of Bucharest
Academiei 14, 70109-Bucharest
Romania

(Received 29 01 1985)