

ON DISTANCES IN SOME BIPARTITE GRAPHS

Ivan Gutman

Abstract. Let $d(v | G)$ be the sum of the distances between a vertex v of a graph G and all other vertices of G . Let $W(G)$ be the sum of the distances between all pairs of vertices of G . A class $\mathbf{C}(k)$ of bipartite graphs is found, such that $d(v | G) \equiv 1 \pmod{k}$ holds for an arbitrary vertex of an arbitrary member of $\mathbf{C}(k)$. Further, for two members G and H of $\mathbf{C}(k)$, having equal cyclomatic number, $W(G) \equiv W(H) \pmod{2k^2}$.

Introduction

In the present paper we establish certain properties of the vertex distances of some bipartite graphs. If G is a (connected) graph and u and v are its two vertices, then the length of the shortest path which connects u and v is denoted by $d(u, v)$ and is called the distance between u and v . The sum of the distances between the vertex v and all other vertices of G is denoted by $d(G | v)$. The sum of the distances between all pairs of vertices of G is denoted by $W(G)$ or simply by W . Hence,

$$W = W(G) = \sum_{\{u,v\}} d(u, v)$$

where $\{u, v\}$ runs over all two-element subsets of the vertex set of G .

We mention in passing that the quantity W plays some role in chemistry [1]. In the chemical literature $W(G)$ is called the Wiener number of the graph G .

Let G be a connected bipartite graph and X and Y its two pertinent vertex sets. Then one immediately sees that $d(u, v)$ is even if both u and v belong to either X or to Y . Otherwise, $d(u, v)$ is odd. This implies the following consequence.

LEMMA 1. $d(v | G) \equiv 1 \pmod{2}$ iff either $v \in X$ and $|Y|$ is odd or $v \in Y$ and $|X|$ is odd. Further, $W(G) \equiv 1 \pmod{2}$ iff both $|X|$ and $|Y|$ are odd.

In the present paper we prove a number of additional congruence statements for the numbers $d(v | G)$ and $W(G)$, which hold for the elements of the sets $\mathbf{C}(h, k)$ and $\mathbf{C}(k)$.

Definition. Let k be a positive integer. If $h > 1$, then every element of $\mathbf{C}(h, k)$ is a graph obtained by joining the endpoints of a path with $2k$ vertices to a pair of adjacent vertices of some graph from $\mathbf{C}(h - 1, k)$. The set $\mathbf{C}(1, k)$ consists of one element only— the circuit with $2k + 2$ vertices.

It is both consistent and convenient to define $\mathbf{C}(0, k)$ as the one-element set, containing graph on two vertices.

The union of the sets $\mathbf{C}(h, k)$, $h = 0, 1, 2, \dots$ is denoted by $\mathbf{C}(k)$

For example, $\mathbf{C}(4, 1)$ consists of eight elements, namely the eight graphs depicted in Fig. 1.




Fig. 1

The basic properties of the above defined classes of graphs are collected in the following lemma.

LEMMA 2. *If G is a graph from $\mathbf{C}(h, K)$, then*

- (a) *G is a connected bipartite graph with $|X| = |Y|$;*
- (b) *the cyclomatic number of G is h ;*
- (c) *the girth of G is $2k + 2$ and every edge of G belongs to a $(2k + 2)$ -membered circuit;*
- (d) *G has $|G| = 2kh + 2$ vertices.*

The main results

THEOREM 1. *If graph from $\mathbf{C}(k)$ and v is its arbitrary vertex, then*

$$d(v | G) \equiv 1 \pmod{k}. \quad (1)$$

If, further, k is even, then

$$d(v | G) \equiv 1 \pmod{2k}. \quad (2)$$

THEOREM 2. If G and H are graphs from $\mathbf{C}(h, k)$, then

$$W(G) \equiv W(H) \pmod{2k^2}. \quad (3)$$

Proof of Theorem 1. We demonstrate the validity of Theorem 1 for $G \in \mathbf{C}(h, k)$ by induction on h . For $h = 0$, (1) and (2) hold in a trivial manner since then $W = 1$. For the unique graph from $\mathbf{C}(1, k)$, namely the circuit with $2k + 2$ vertices, it is easy to show that $d(v | G) = (k + 1)^2$. Whence (1) and (2) are satisfied.

Suppose now that G^* is an element of $\mathbf{C}(h - 1, k)$ and that G can be obtained by joining the endpoints u_1 and u_{2k} of a path with $2k$ vertices with the vertices p and q of G^* (see Fig. 2).




Fig. 2

The newly introduced vertices of G are labeled by u_1, u_2, \dots, u_{2k} .

From the construction of the graph G it is evident that

$$d(v | G) = d(v | G^*) + \sum_{i=1}^{2k} d(v, u_i).$$

Assuming that $d(v, p) < d(v, q)$, we have

$$\sum_{i=1}^k d(v, u_i) = kd(v, p) + k(k + 1)/2,$$

$$\sum_{i=k+1}^{2k} d(v, u_i) = k + kd(v, p) + k(k + 1)/2,$$

and

$$d(v | G) = d(v | G^*) + 2kd(v, p) + k(k + 2).$$

Consequently,

$$d(v | G) \equiv d(v | G^*) \pmod{k}$$

and, if k is even,

$$d(v | G) \equiv d(v | G^*) \pmod{2k}.$$

Therefore if (1) and (2) hold for the vertex v of G^* , then they also hold for the vertex v of G .

In order to complete the proof of Theorem 1, we have to show that (1) and (2) hold also for the vertices $u_i, i = 1, 2, \dots, 2k$ of G . Let u stand for one of these vertices. Then

$$d(u | G) = \sum_{i=1}^{2k} d(u, u_i) + d(u, p) + d(u, q) + \sum_v d(u, v) \quad (4)$$

with the second summation of the r.h.s. of (4) running over the vertices of G^* different than p and q . The vertices $u_1, u_2, \dots, u_{2k}, q, p$ form a circuit of size $2k+2$ in G and therefore

$$\sum_{i=1}^{2k} d(u, u_i) + d(u, p) + d(u, q) = k(k+1)^2.$$

Further,

$$\sum_v d(u, v) = d(x | G^*) - 1 + (|G^*| - 2)d(u, x)$$

where $x = p$ if $d(u, p) < d(u, q)$ and $x = q$ otherwise. According to Lemma 2, $|G^*| = 2k(h-1) + 2$.

Taking all this into account, eq. (4) becomes

$$d(u | G) = d(x | G^*) + k(k+2) + 2k(h-1)d(u, x),$$

Hence,

$$d(u | G) \equiv d(x | G^*) \pmod{k}$$

and, if k is even,

$$d(u | G) \equiv d(x | G^*) \pmod{2k}.$$

This means that if (1) and (2) hold for all vertices of $G^* \in \mathbf{C}(h-1, k)$, then they also hold for all vertices of $G \in \mathbf{C}(h, k)$. Consequently, they hold for all vertices of all graphs from $\mathbf{C}(k)$. \square

Proof of Theorem 2. The sets $\mathbf{C}(0, k)$, $\mathbf{C}(1, k)$ and $\mathbf{C}(2, k)$ contain one element each and therefore for $h = 0, 1$ and 2 Theorem 2 holds in a trivial manner. Direct calculation confirms the validity of Theorem 2 also in the case of $\mathbf{C}(3, k)$ (which contains $k+2$ elements).

In order to complete a proof by induction, assume that (3) is obeyed for all graphs from $\mathbf{C}(h-1, k)$ and in particular for G^* and H^* . Let $G \in \mathbf{C}(h, k)$ be obtained from G^* in the previously described way (see Fig. 2). Let $H \in \mathbf{C}(h, k)$ be obtained from H^* in a fully analogous manner.

Now, from Fig. 2 we see that

$$W(G) = W(G^*) + k(4k^2 - 1)/3 + k(k + 1) | G^* | + k[d(p | G^*) + d(q | G^*)]. \quad (5)$$

Namely, $k(4k^2 - 1)/3$ is the W number of the path with $2k$ vertices, where as the sum of the distances between the vertices of G^* and u_1, u_2, \dots, u_k is

$$kd(p | G^*) | G^* | \sum_{i=1}^k i$$

and, similiary, the sum of the distances between the vertices of G^* and u_{k+1}, \dots, u_{2k} is

$$kd(q | G^*) + | G^* | \sum_{i=0}^k i.$$

An analogous equality will hold for the graph H , viz.,

$$W(H) = W(H^*) + k(4k^2 - 1)/3 + k(k + 1) | H^* | + k | d(p | H^*) + d(q | H^*) |. \quad (6)$$

Bearing in mind that by Lemma 2, $| G^* | = | H^* |$, the identities (5) and (6) yield

$$W(G) - W(H) = W(G^*) - W(H^*) + k[d(p | G^*) - d(p | H^*) + d(q | G^*) - d(q | H^*)].$$

We now have to distinguish between two cases.

Case a: k is even. Then because of Theorem 1, $d(p | G^*) - d(p | H^*) + d(q | G^*) - d(q | H^*)$ is divisible by $2k$. Therefore the induction hypothesis that $W(G^*) - W(H^*)$ is divisible by $2k^2$ leads to the conclusion that then also $W(G) - W(H)$ is divisible by $2k^2$.

Theorem 2 follows for the case of even k .

Case b: k is odd. Then by Theorem 1, $d(p | G^*) - d(p | H^*) + d(q | G^*) - d(q | H^*)$ is divisible only by k and the above reasoning leads to the conclusion that $W(G) - W(H)$ is divisible by k^2 . On the other hand, according to Lemma 1, $W(G) - W(H)$ must be divisible by two. Since k^2 is assumed to be odd, $W(G) - W(H)$ must be divisible by $2k^2$.

This proves Theorem 2 also for odd values of k . \square

Discussion

1° In the set $\mathbf{C}(3, k)$ there exist graphs G and H such that $W(G) - W(H) = 2k^2$. Therefore $2k^2$ is the greatest possible argument in a relation of the type (3) and Theorem 2, is, in a certain sense, the best possible congruence statement for the W numbers of the members of $\mathbf{C}(h, k)$.

2° Eq. (5) and the fact that $| G | = 2k(h - 1) + 2$ imply

$$W(G) - W(G^*) \equiv k(k^2 + 11)/3 \pmod{k^2}.$$

This means that by increasing the cyclomatic number by one, W increases by $k(k^2 + 11)/3$ modulo k^2 . Since for the (unique) element of $\mathbf{C}(0, k)$ it is $W = 1$, one concludes that for the members of $\mathbf{C}(h, k)$.

$$W \equiv 1 + hk(k^2 + 11)/3 \pmod{k^2}.$$

Analogously, if k is even, then

$$W \equiv 1 + hk(k^2 + 11)/3 \pmod{2k^2}.$$

3° There is a natural way to generalize the definition of the presently considered bipartite graphs to non-bipartite ones. Instead of the set $\mathbf{C}(h, k)$ we may define that the set $\mathbf{C}^*(h, k)$ whose elements are constructed by joining the endpoints of a path with $2k - 1$ vertices to a pair adjacent vertices of a graph from $\mathbf{C}^*(h - 1, k)$. Further, $\mathbf{C}^*(1, k)$ would consist of the circuit of the size $2k + 1$.

Unfortunately, neither Theorem 1 nor Theorem 2 could be extended to $\mathbf{C}^*(h, k)$, nor any other similar congruence statement could be established.

REFERENCES

- [1] I. Gutman and O. E. Polansky, *Mathematical Concepts in Organic Chemistry*, Springer-Verlag, Berlin, 1986, pp. 124–127.

Prirodno-matematički fakultet
34000 Kragujevac, pp. 60
Jugoslavija

(Received 16 01 1987)