

## ABSOLUTE AND ORDINARY KÖTHE-TOEPLITZ DUALS OF SOME SETS OF SEQUENCES AND MATRIX TRANSFORMATIONS

Eberhard Malkowsky

**Abstract.** We determine the ordinary Köthe-Toeplitz dual of the set  $\Delta l_\infty(p)$  and the absolute Köthe-Toeplitz duals of the sets  $\Delta l_\infty(p)$ ,  $\Delta c_0(p)$  and  $\Delta c(p)$  defined by Ahmad and Mursaleen. Further we investigate matrix transformations in these spaces and give a characterization of the class  $(\Delta l_\infty(p), l_\infty)$ .

### 1. Introduction

By  $\omega$  we denote the set of all complex sequences  $x = (x_k)_{k=1}^\infty$ . Throughout the paper  $p = (p_k)_{k=1}^\infty$  shall always be an arbitrary sequence of positive reals. The following sets were introduced and investigated by various authors:

$$\begin{aligned}l_\infty(p) &:= \{x \in \omega : \sup_k |x_k|^{p_k} < \infty\}, \\c(p) &:= \{x \in \omega : |x_k - l|^{p_k} \rightarrow 0 \text{ for some complex } l\}, \\c_0(p) &:= \{x \in \omega : |x_k|^{p_k} \rightarrow 0\}, \\l(p) &:= \left\{x \in \omega : \sum_{k=1}^\infty |x_k|^{p_k} < \infty\right\} \quad (\text{cf. [2], [3], [5], and [7]}).\end{aligned}$$

Given any sequence  $x \in \omega$  we shall write  $\Delta x := (x_k - x_{k+1})$ . In a recent paper (cf. [1]), Ahmad and Mursaleen defined the following sets:

$$\begin{aligned}\Delta l_\infty(p) &:= \{x \in \omega : \Delta x \in l_\infty(p)\}, \\ \Delta c(p) &:= \{x \in \omega : \Delta x \in c(p)\}, \\ \Delta c_0(p) &:= \{x \in \omega : \Delta x \in c_0(p)\}.\end{aligned}$$

In the determination of the absolute Köthe-Toeplitz duals of  $\Delta l_\infty(p)$  and  $\Delta c_0(p)$ , they applied some arguments which do not seem to hold:

(i)  $x \in \Delta l_\infty(p)$  does not imply in general the existence of a finite number  $N > \sup_k k^{-1}|x_k|$ , as the following counterexample will show: If we put  $p_k := k^{-1}$  and  $x_k := k^2$  ( $k = 1, 2, \dots$ ) then  $|\Delta x_k|^{p_k} \rightarrow 1$  ( $k \rightarrow \infty$ ), hence  $x \in \Delta l_\infty(p)$ , and  $\sup_k k^{-1}|x_k| = \infty$ .

(ii) If  $a$  is a sequence such that

$$\sum_{k=1}^{\infty} k|a_k|N^{1/p_k} = \infty \quad \text{for some } N > 1, \quad (1.1)$$

then the sequence  $x$  defined by  $x_k := kN^{1/p_k} \operatorname{sgn} a_k$  is not in  $\Delta l_\infty(p)$ , in general. In order to see this, we put  $p_k := k$  and  $a_k := (-1)^k$  ( $k = 1, 2, \dots$ ). Then  $a$  satisfies (1.1) for all  $N > 1$  and  $|\Delta x_k|^{p_k} \rightarrow \infty$ , hence  $x \notin \Delta l_\infty(p)$ .

In this paper, we shall determine the absolute Köthe-Toeplitz duals of the sets  $\Delta l_\infty(p)$  and  $\Delta c_0(p)$ , and give new proofs for the characterizations of the matrix transformations considered in [1]. Further we shall state some new results.

## 2. Köthe-Toeplitz duals

For arbitrary set  $X$  of sequences, we define the ordinary and absolute Köthe-Toeplitz duals by

$$X^\dagger := \left\{ a \in \omega : \sum_{k=1}^{\infty} a_k x_k \text{ converges for all } x \in X \right\} \quad \text{and}$$

$$X^{|\dagger|} := \left\{ a \in \omega : \sum_{k=1}^{\infty} |a_k x_k| < \infty \text{ for all } x \in X \right\}$$

respectively; we shall write  $X^{\dagger\dagger} := (X^\dagger)^\dagger$  and  $X^{|\dagger\dagger|} := (X^{|\dagger|})^{|\dagger|}$ .

**THEOREM 2.1.** *For every strictly positive sequence  $p = (p_k)$ , we have*

$$(a) \quad (\Delta l_\infty(p))^{|\dagger|} = D_\infty^{(1)}(p) := \bigcap_{N=2}^{\infty} \left\{ a \in \omega : \sum_{k=1}^{\infty} |a_k| \sum_{j=1}^{k-1} N^{1/p_j} < \infty \right\},$$

$$(b) \quad (\Delta l_\infty(p))^{|\dagger\dagger|} = D_\infty^{(2)}(p) := \bigcup_{N=2}^{\infty} \left\{ a \in \omega : \sup_{k \geq 2} |a_k| \left[ \sum_{j=1}^{k-1} N^{1/p_j} \right]^{-1} < \infty \right\},$$

$$(c) \quad (\Delta c_0(p))^{|\dagger|} = D_0^{(1)}(p) := \bigcup_{N=2}^{\infty} \left\{ a \in \omega : \sum_{k=1}^{\infty} |a_k| \sum_{j=1}^{k-1} N^{-1/p_j} < \infty \right\},$$

$$(d) \quad (\Delta c_0(p))^{|\dagger\dagger|} = D_0^{(2)}(p) := \bigcap_{N=2}^{\infty} \left\{ a \in \omega : \sup_{k \geq 2} |a_k| \left[ \sum_{j=1}^{k-1} N^{-1/p_j} \right]^{-1} < \infty \right\}.$$

(We adopt the usual convention that  $\sum_{j=1}^m y_j = 0$  ( $m < 1$ ) for arbitrary  $y_i$ .)

*Proof:* (a) Let  $a \in D_\infty^{(1)}(p)$  and  $x \in \Delta l_\infty(p)$ . We choose  $N > \max\{1, \sup |\Delta x_k|^{p_k}\}$ . Then

$$\begin{aligned} \sum_{k=1}^{\infty} |a_k x_k| &\leq \sum_{k=1}^{\infty} |a_k| \left| \sum_{j=1}^{k-1} \Delta x_j \right| + |x_1| \sum_{k=1}^{\infty} |a_k| \\ &\leq \sum_{k=1}^{\infty} |a_k| \sum_{j=1}^{k-1} N^{1/p_j} + |x_1| \sum_{k=1}^{\infty} |a_k| < \infty. \end{aligned} \quad (2.1)$$

(Note: Since  $\sum_{j=1}^{k-1} N^{1/p_j} \geq 1$  for arbitrary  $N > 1$  ( $k = 2, 3, \dots$ ),  $a \in D_\infty^{(1)}(p)$  implies  $\sum_{k=1}^{\infty} |a_k| < \infty$ .)

Conversely let  $a \notin D_\infty^{(1)}(p)$ . Then we have  $\sum_{k=1}^{\infty} |a_k| \sum_{j=1}^{k-1} N^{1/p_j} = \infty$  for some integer  $N > 1$ .

We define the sequence  $x$  by  $x_k := \sum_{j=1}^{k-1} N^{1/p_j}$  ( $k = 1, 2, \dots$ ). Then it is easy to see that  $x \in \Delta l_\infty(p)$  and  $\sum_{k=1}^{\infty} |a_k x_k| = \infty$ , hence  $a \notin (\Delta l_\infty(p))^{|t|}$ .

(b) Let  $a \in D_\infty^{(2)}(p)$  and  $x \in (\Delta l_\infty(p))^{|t|} = D_\infty^{(1)}(p)$ , by part (a). Then for some  $N > 1$ , we have

$$\begin{aligned} \sum_{k=2}^{\infty} |a_k x_k| &= \sum_{k=2}^{\infty} |a_k| \left[ \sum_{j=1}^{k-1} N^{1/p_j} \right]^{-1} |x_k| \sum_{j=1}^{k-1} N^{1/p_j} \\ &\leq \sup_{k \geq 2} \left[ |a_k| \left[ \sum_{j=1}^{k-1} N^{1/p_j} \right]^{-1} \right] \sum_{k=2}^{\infty} |x_k| \sum_{j=1}^{k-1} N^{1/p_j} < \infty. \end{aligned}$$

Conversely let  $a \notin D_\infty^{(2)}(p)$ . Then for all integers  $N > 1$ , we have

$$\sup_{k \geq 2} |a_k| \left[ \sum_{j=1}^{k-1} N^{1/p_j} \right]^{-1} = \infty.$$

Hence there is a strictly increasing sequence  $(k(m))$  of integers  $k(m) \geq 2$  such that

$$|a_{k(m)}| \left[ \sum_{j=1}^{k(m)-1} m^{1/p_j} \right]^{-1} > m^2 \quad (m = 2, 3, \dots).$$

We define the sequence  $x$  by

$$x_k := \begin{cases} |a_{k(m)}|^{-1} & (k = k(m)) \\ 0 & (k \neq k(m)) \quad (m = 2, 3, \dots). \end{cases}$$

Then for all integers  $N \geq 2$ , we have

$$\begin{aligned} & \sum_{k=1}^{\infty} |x_k| \sum_{j=1}^{k-1} N^{1/p_j} \leq \sum_{m=2}^{\infty} |a_{k(m)}|^{-1} \sum_{j=1}^{k(m)-1} N^{1/p_j} \leq \\ & \leq \sum_{m=2}^{N-1} |a_{k(m)}|^{-1} \sum_{j=1}^{k(m)-1} N^{1/p_j} + \sum_{m=N}^{\infty} |a_{k(m)}|^{-1} \sum_{j=1}^{k(m)-1} m^{1/p_j} \leq \\ & \leq \sum_{m=2}^{N-1} |a_{k(m)}|^{-1} \sum_{j=1}^{k(m)-1} N^{1/p_j} + \sum_{m=N}^{\infty} m^{-2} < \infty, \end{aligned}$$

hence  $x \in (\Delta l_{\infty}(p))^{\dagger\dagger}$ , and

$$\sum_{k=1}^{\infty} |a_k x_k| = \sum_{N=2}^{\infty} 1 = \infty,$$

hence  $a \notin (\Delta l_{\infty}(p))^{\dagger\dagger\dagger}$ .

(c) Let  $a \in D_0^{(1)}(p)$ . Since  $|a_k| \leq |a_k| N^{1/p_1} \sum_{j=1}^{k-1} N^{-1/p_j}$  ( $k = 2, 3, \dots$ ), we have  $\sum_{k=1}^{\infty} |a_k| < \infty$ . Let  $x \in \Delta c_0(p)$ . Then there is an integer  $k_0$  such that  $\sup_{k > k_0} |\Delta x_k|^{p_k} \leq N^{-1}$ , where  $N$  is the number in  $D_0^{(1)}(p)$ . We put  $M := \max_{1 \leq k \leq k_0} |\Delta x_k|^{p_k}$ ,  $m := \min_{1 \leq k \leq k_0} p_k$ ,  $L := (M+1)N$  and define the sequence  $y$  by  $y_k := x_k L^{-1/m}$  ( $k = 1, 2, \dots$ ). Then it is easy to see that  $\sup_k |\Delta y|^{p_k} \leq N^{-1}$ , and as in (2.1) with  $N$  replaced by  $N^{-1}$ , we have

$$\sum_{k=1}^{\infty} |a_k x_k| = L^{1/m} \sum_{k=1}^{\infty} |a_k y_k| < \infty.$$

Conversely, let  $a \notin D_0^{(1)}(p)$ . Then we can determine a strictly increasing sequence  $(k(s))$  of integers such that  $k(1) := 1$  and

$$M_s := \sum_{k=k(s)}^{k(s+1)-1} |a_k| \sum_{j=1}^{k-1} (s+1)^{-1/p_j} > 1 \quad (s = 1, 2, \dots).$$

We define the sequence  $x$  by

$$\begin{aligned} x_k &:= \sum_{l=1}^{s-1} \sum_{j=k(l)}^{k(l+1)-1} (l+1)^{-1/p_j} + \sum_{j=k(s)}^{k-1} (s+1)^{-1/p_j} \\ & \quad (k(s) \leq k \leq k(s+1) - 1; s = 1, 2, \dots). \end{aligned}$$

Then it is easy to see that  $|\Delta x_k|^{p_k} = 1/(s+1)$  ( $k(s) \leq k \leq k(s+1) - 1; s = 1, 2, \dots$ ) hence  $x \in \Delta c_0(p)$ , and  $\sum_{k=1}^{\infty} |a_k x_k| \geq \sum_{s=1}^{\infty} 1 = \infty$ , i.e.  $a \notin (\Delta c_0(p))^{\dagger\dagger}$ .

(d) For  $N = 2, 3, \dots$ , we put

$$E_N := \left\{ a \in \omega : \sum_{k=1}^{\infty} |a_k| \sum_{j=1}^{k-1} N^{-1/p_j} < \infty \right\}$$

$$F_N := \left\{ a \in \omega : \sup_{k \geq 2} |a_k| \left[ \sum_{j=1}^{k-1} N^{-1/p_j} \right]^{-1} < \infty \right\}.$$

By a well known result (cf. [3, Lemma 4 (iv)]), we have to show  $F_N = E_N^{|\dagger|}$  ( $N = 2, 3, \dots$ ). The proof of this is standard and therefore omitted.

Now we shall give some new results:

**THEOREM 2.2.** *For every strictly positive sequence  $p = (p_k)$ , we have*

(a)  $(\Delta c(p))^{|\dagger|} = D^{(1)}(p) := D_0^{(1)} \cap \{a \in \omega : \sum_{k=1}^{\infty} |a_k| k < \infty\}$  and

(b)  $(\Delta l_{\infty}(p))^{\dagger} = D_{\infty}(p)$

$$:= \bigcap_{N=2}^{\infty} \left\{ a \in \omega : \sum_{k=1}^{\infty} a_k \sum_{j=1}^{k-1} N^{1/p_j} \text{ converges and } \sum_{k=1}^{\infty} N^{1/p_k} |R_k| < \infty \right\},$$

where  $R_k := \sum_{\nu=k+1}^{\infty} a_{\nu}$  ( $k = 1, 2, \dots$ ).

*Proof:* (a) Let  $a \in D^{(1)}(p)$  and  $x \in \Delta c(p)$ . Then there is a complex number  $l$  such that  $|\Delta x_k - l|^{p_k} \rightarrow 0$  ( $k \rightarrow \infty$ ). We define  $y$  by  $y_k := x_k + lk$  ( $k = 1, 2, \dots$ ). Then  $y \in \Delta c_0(p)$  and

$$\sum_{k=1}^{\infty} |a_k x_k| \leq \sum_{k=1}^{\infty} |a_k| \left| \sum_{j=1}^{k-1} \Delta y_j \right| + |l| \sum_{k=1}^{\infty} |a_k| k < \infty$$

by Theorem 2.1.(c) and since  $a \in D^{(1)}(p)$ . Now let  $a \in (\Delta c(p))^{|\dagger|} \subset (\Delta c_0(p))^{|\dagger|} = D_0^{(1)}(p)$  by Theorem 2.1.(c). Since the sequence  $x$  defined by  $x_k := k$  ( $k = 1, 2, \dots$ ) is in  $\Delta c(p)$  we have  $\sum_{k=1}^{\infty} |a_k| k < \infty$ .

(b) Let  $a \in D_{\infty}(p)$  and  $x \in \Delta l_{\infty}(p)$ . Then there is an integer  $N > \max\{1, \sup_k |\Delta x_k|^{p_k}\}$ . We have

$$\sum_{k=1}^n a_k x_k = - \sum_{j=1}^{n-1} \Delta x_j R_j + R_n \sum_{j=1}^{n-1} \Delta x_j + x_1 \sum_{k=1}^n a_k \quad (n = 1, 2, \dots). \quad (2.2)$$

Obviously the last term on the right in (2.2) is convergent. Since  $\sum_{j=1}^{\infty} |\Delta x_j| \times |R_j| \leq \sum_{j=1}^{\infty} N^{1/p_j} |R_j| < \infty$ , the first term on the right in (2.2) is absolutely convergent. Finally by Corollary 2 in [4], the convergence of  $\sum_{k=1}^{\infty} a_k \sum_{j=1}^{k-1} N^{1/p_j}$  implies  $\lim_{n \rightarrow \infty} R_n \sum_{j=1}^{n-1} N^{1/p_j} = 0$ . Conversely let  $a \in (\Delta l_{\infty}(p))^{\dagger}$ . Since  $e := (1, 1, \dots) \in \Delta l_{\infty}(p)$  and  $x = \left[ \sum_{j=1}^{k-1} N^{1/p_j} \right] \in \Delta l_{\infty}(p)$ , we conclude the convergence

of  $\sum_{k=1}^{\infty} a_k$  and  $\sum_{k=1}^{\infty} a_k \sum_{j=1}^{k-1} N^{1/p_j}$  respectively. Applying Corollary 2 in [4] again we have

$$\lim_{n \rightarrow \infty} R_n \sum_{j=1}^{k-1} N^{1/p_j} = 0.$$

From (2.2), we obtain the convergence of  $\sum_{k=1}^{\infty} \Delta x_k R_k$  for all  $x \in \Delta l_{\infty}(p)$ . Since  $x \in \Delta l_{\infty}(p)$  if and only if  $y := \Delta x \in l_{\infty}(p)$ , this implies  $R \in l_{\infty}^{\dagger}(p)$ , hence  $\sum_{k=1}^{\infty} N^{1/p_k} |R_k| < \infty$  for all  $N > 1$  by a well known theorem (cf. [2, Th. 2]).

### 3. Some matrix transformations

For any complex matrix  $A = (a_{nk})$ , we shall write  $A_n := (a_{nk})_k$  for the sequence in the  $n$ -th row of  $A$ . Given  $A$  we define the matrix  $B$  by

$$b_{nk} := a_{nk} - a_{n+1,k} \quad (n, k = 1, 2, \dots).$$

Let  $X, Y$  be two subsets of  $\omega$ . By  $(X, Y)$  we denote the class of all matrices  $A$  such that the series  $A_n x := \sum_{k=1}^{\infty} a_{nk} x_k$  converge for all  $x \in X$  ( $n = 1, 2, \dots$ ) and the sequence  $Ax := (A_n x)$  is in  $Y$  for all  $x \in X$ .

The following is obvious and therefore stated without proof:

LEMMA 3.1. *Let  $X, Y$  be linear sequence spaces. We put  $\Delta Y := \{y \in \omega : \Delta y \in Y\}$ . Then  $A \in (X, \Delta Y)$  if and only if  $B \in (X, Y)$  and  $A_1 \in X^{\dagger}$ .*

Lemma 3.1 and well known results together yield for instance the characterization of the following classes for strictly positive sequences  $q \in l_{\infty} : (l(p), \Delta l_{\infty}(q)), (l(p), \Delta c_0(q)), (l(p), \Delta c(q))$ , (cf. [5, Th. 5 (i), (ii) and (iii)] if  $0 < p_k \leq 1$  ( $k = 1, 2, \dots$ ), [5, Th. 8 and Th. 9] if  $1 < p_k \leq H < \infty$  ( $k = 1, 2, \dots$ )). Now we give a characterization for the class  $(\Delta l_{\infty}(p), l_{\infty})$ :

THEOREM 3.1. *For every strictly positive sequence  $p$ , we have  $A \in (\Delta l_{\infty}(p), l_{\infty})$  if and only if the following three conditions hold:*

- (i)  $M_1(N) := \sup_n \left| \sum_{k=1}^{\infty} a_{nk} \sum_{j=1}^{k-1} N^{1/p_j} \right| < \infty$  for all  $N > 1$ ,
- (ii)  $M_2(N) := \sup_n \left[ \sum_{\nu=1}^{\infty} N^{1/p_{\nu}} \left| \sum_{k=\nu+1}^{\infty} a_{nk} \right| \right] < \infty$  for all  $N > 1$ ,
- (iii)  $M_3 := \sup_n \left| \sum_{k=1}^{\infty} a_{nk} \right| < \infty$ .

*Proof:* Let conditions (i), (ii) and (iii) be satisfied. Then  $A_n \in (\Delta l_{\infty}(p))^{\dagger}$  ( $n = 1, 2, \dots$ ) by Theorem 2.2.(b). Hence the series  $A_n x$  converge for all  $x \in$

$\Delta l_\infty(p)$  ( $n = 1, 2, \dots$ ). Further as in the proof of Theorem 2.2.(b), we have for  $x \in \Delta l_\infty(p)$  such that  $\sup_k |\Delta x_k|^{p_k} < N$ :

$$\left| \sum_{k=1}^{\infty} a_{nk} x_k \right| \leq \sum_{\nu=1}^{\infty} N^{1/p_\nu} \left| \sum_{k=\nu+1}^{\infty} a_{nk} \right| + |x_1| \left| \sum_{k=1}^{\infty} a_{nk} \right| \leq M_2(N) + |x_1| M_3$$

$$(n = 1, 2, \dots),$$

hence  $Ax \in l_\infty$ .

Conversely let  $A \in (\Delta l_\infty(p), l_\infty)$ . The necessity of conditions (i) and (iii) follows from the fact that  $(x_k) := \left[ \sum_{j=1}^{k-1} N^{1/p_j} \right]$  and  $e$  are in  $\Delta l_\infty(p)$ . In order to show the necessity of condition (ii), we assume that  $M_2(N) = \infty$  for some  $N > 1$ .

Then for the matrix  $C$  defined by

$$c_{n\nu} := \sum_{k=\nu+1}^{\infty} a_{nk} \quad (n, \nu = 1, 2, \dots),$$

we have  $C \notin (l_\infty(p), l_\infty)$ . (cf. [2, Th. 3]) Hence there is a sequence  $x \in l_\infty(p)$  such that  $\sup_\nu |x_\nu|^{p_\nu} = 1$  and  $\sum_{\nu=1}^{\infty} c_{n\nu} x_\nu \neq O(1)$ . We define the sequence  $y$  by  $y_\nu := -\sum_{j=1}^{\nu-1} x_j + x_1$  ( $\nu = 1, 2, \dots$ ). Then  $y \in \Delta l_\infty(p)$  and  $\sum_{\nu=1}^{\infty} a_{n\nu} y_\nu = \sum_{\nu=1}^{\infty} c_{n\nu} x_\nu + x_1 \sum_{\nu=1}^{\infty} a_{n\nu} \neq 0(1)$ , a contradiction to the assumption  $A \in (\Delta l_\infty(p), l_\infty)$ . Therefore we must have  $M_2(N) < \infty$  for all  $N > 1$ .

#### REFERENCES

- [1] Z. U. Ahmad and Mursaleen, *Köthe-Toeplitz duals of some new sequence spaces and their matrix maps*, Publ. Inst. Math. (Beograd) **42** (56) (1987), 57–61.
- [2] C. G. Lascarides and I. J. Maddox, *Matrix transformations between some classes of sequences*, Proc. Cambridge Phil. Soc. **68** (1970), 99–104.
- [3] C. G. Lascarides, *A study of certain sequence spaces of Maddox and a generalization of a theorem of Iyer*, Pacific J. Math. **38** (1971), 487–500.
- [4] H. Kizmaz, *On certain sequence spaces*, Canadian Math. Bull. **24** (1981), 169–176.
- [5] I. J. Maddox, *Continuous and Köthe-Toeplitz duals of certain sequence spaces*, Proc. Cambridge Phil. Soc. **65** (1967), 471–475.
- [6] I. J. Maddox and M. A. L. Willey, *Continuous operators on paranormed spaces and matrix transformations*, Pacific J. Math. **53** (1974), 217–288.
- [7] S. Simons, *The sequence spaces  $l(p_\nu)$  and  $m(p_\nu)$* , Proc. London Math. Soc. **15** (1965), 422–436.

Mathematisches Institut  
Justus-Liebig-Universität  
Iheringstraße 6  
D-6300 Gießen  
West Germany

(Received 26 05 1988)  
(Revised 29 05 1989)